

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications from Nebraska Center for
Research on Children, Youth, Families, and
Schools

Children, Youth, Families & Schools, Nebraska
Center for Research on

2008

Food insecurity and maternal depression in rural, low-income families: A longitudinal investigation

Catherine Huddleston-Casas

University of Nebraska - Lincoln, chuddleston-casas2@unl.edu

Richard Charnigo

University of Kentucky

Leigh Ann Simmons

University of Kentucky, leighann@uky.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/cyfsfacpub>

 Part of the [Pre-Elementary, Early Childhood, Kindergarten Teacher Education Commons](#)

Huddleston-Casas, Catherine; Charnigo, Richard; and Simmons, Leigh Ann, "Food insecurity and maternal depression in rural, low-income families: A longitudinal investigation" (2008). *Faculty Publications from Nebraska Center for Research on Children, Youth, Families, and Schools*. 11.
<https://digitalcommons.unl.edu/cyfsfacpub/11>

This Article is brought to you for free and open access by the Children, Youth, Families & Schools, Nebraska Center for Research on at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications from Nebraska Center for Research on Children, Youth, Families, and Schools by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Food insecurity and maternal depression in rural, low-income families: a longitudinal investigation

Catherine Huddleston-Casas¹, Richard Charnigo² and Leigh Ann Simmons^{3,*}

¹Department of Child, Youth and Family Studies, University of Nebraska–Lincoln, Omaha Campus, 102 Arts and Sciences Hall, Omaha, NE 68182-0214, USA: ²College of Public Health 203-B, Department of Biostatistics, University of Kentucky, Lexington, KY 40536-0003, USA: ³Departments of Family Studies and Health Services Management, University of Kentucky, 315 Funkhouser Building, Lexington, KY 40506-0054, USA

Submitted 3 March 2008: Accepted 11 August 2008: First published online 15 September 2008

Abstract

Objective: The purpose of the present study was to examine the relationship between household food insecurity and maternal depression in a rural sample to determine whether food insecurity predicted mothers' depression over time or vice versa.

Design: The study employed a prospective design using three waves of data from 'Rural Families Speak', a multi-state study of low-income rural families in the USA. Food insecurity was measured using the Core Food Security Module and depression was measured using the Center for Epidemiologic Studies–Depression Scale. A structural equation model was fit to the data using the AMOS software package.

Setting: Sixteen states in the USA (California, Indiana, Kentucky, Louisiana, Massachusetts, Maryland, Michigan, Minnesota, Nebraska, New Hampshire, New York, Ohio, Oregon, South Dakota, West Virginia, Wyoming) between 2000 and 2002.

Subjects: Subjects included 413 women with at least one child under the age of 13 years living in the home.

Results: Findings based on the 184 subjects with complete data indicated that the causal relationship between household food insecurity and depression is bidirectional ($P=0.034$ for causation from depression to food insecurity, $P=0.003$ for causation from food insecurity to depression, $\chi^2/df=1.835$, root-mean-square error of approximation = 0.068, comparative fit index = 0.989). Findings based on all 413 subjects after imputation of missing values also indicated bidirectionality.

Conclusions: The recursive relationship between food insecurity and depression has implications for US nutrition, mental health and poverty policies. The study highlights the need to integrate programmes addressing food insecurity and poor mental health for the population of rural, low-income women.

Keywords
Food insecurity
Depression
Rural women
United States

Poverty is a significant social problem in the USA that is associated with a number of public health concerns, including poor mental and physical health status and disparities in health care⁽¹⁾. The most recent statistics show the national poverty rate increased to 9.8% between 2000 and 2006⁽²⁾. In rural areas this rate is even higher, with recent data showing that the rural poverty rate exceeds 14%⁽³⁾. These data reflect an ongoing trend of poverty and related problems faced by rural Americans over the last 50 years⁽⁴⁾.

One of the problems associated with rural poverty is food insecurity, which is defined as being uncertain of having or unable to acquire enough food to meet the needs of all family members due to insufficient money or other resources⁽⁵⁾. The rate of food insecurity nationwide increased from 9.8% to 11.0% between 2001 and 2006⁽⁶⁾,

and the rural food insecurity rate reached 12% in 2006⁽⁶⁾. The prevalence of food insecurity varies among household types. Households headed by single mothers and households with incomes below the federal poverty line have higher rates of food insecurity than those in the general population⁽⁶⁾. Minority households also have a higher rate of food insecurity. Recent data revealed Black and Latino households to have twice the rate of food insecurity as White households. Similarly, households with children were almost twice as likely to be food-insecure as households without children (15.6% *v.* 8.5%)⁽⁶⁾. In rural areas food insecurity is particularly concerning, because rural residents have more limited access to affordable food than their urban counterparts. Compared with urban consumers, rural residents face higher prices for food⁽⁷⁾ and are less likely to receive food stamps despite eligibility⁽⁸⁾.

*Corresponding author: Email leighann@uky.edu

A growing body of research has demonstrated a link between food insecurity and depression, although most of these studies were cross-sectional in nature. A study of California women showed that poor mental health was linked to food insecurity, but the authors noted that the direction of effect could not be determined⁽⁹⁾. In a qualitative study of poor women in Philadelphia, Chilton and Booth⁽¹⁰⁾ found that food-insecure women experienced 'hunger of the mind', which they defined as a sense of hopelessness and depression. In a study of women with young children, Casey and colleagues⁽¹¹⁾ found that nearly 53% of women who reported food insecurity scored positive on a depression screen. Similarly, a cross-sectional study of nearly 3000 mothers of 3-year-olds showed that the percentage of respondents with major depressive disorder increased with greater levels of food insecurity. Among food-secure mothers, 15.7% had major depressive disorder, compared with 20.2% of marginally food-secure mothers and 28.5% of food-insecure mothers. This research is underscored by findings from the Women's Employment Study, which revealed that urban welfare recipients in Michigan who experienced food insufficiency also were more likely to experience depression⁽¹²⁻¹⁴⁾. As other researchers have noted, the authors of the Michigan studies acknowledged that the direction of effect could not be ascertained.

The relationship between food insecurity and depression is concerning in light of the high rates of depression among rural women. Although assessments of the prevalence of depressive symptomatology and major depressive episodes (MDE) among rural populations are limited, Hauenstein and colleagues⁽¹⁵⁾ argue that population mental health deteriorates as the level of rurality increases. The prevalence of clinically depressive symptomatology among rural women has been shown to range from 24% to 69%⁽¹⁶⁻²²⁾, and low-income women appear to be particularly vulnerable^(16-18,21,22).

Given the high prevalence rates of both food insecurity and depression among rural women, and given previous studies suggesting that food insecurity and depression are associated, the purpose of the present study was to examine the longitudinal relationship between food insecurity and depression in a sample of low-income, rural women. Since previous studies have failed to document a direction of effect between food insecurity and depression, our goal was to examine whether: (i) food insecurity predicts depression; (ii) depression predicts food insecurity; or (iii) the relationship between food insecurity and depression is bidirectional. Additionally, to our knowledge this is the first longitudinal study of the relationship between food insecurity and depression among rural, low-income women in the USA. Findings have the potential to affect public policy and practice aimed at reducing the rates of both food insecurity and depression in this vulnerable population.

Methods

Sample

We utilized data from NC-223, 'Rural Families Speak' (RFS), a multi-state (California, Indiana, Kentucky, Louisiana, Massachusetts, Maryland, Michigan, Minnesota, Nebraska, New Hampshire, New York, Ohio, Oregon, South Dakota, West Virginia, Wyoming) study of rural, low-income women and their families. The study was approved by a research ethics board at all participating institutions, and all respondents provided informed consent. Between 2000 and 2002, three waves of data were collected in twenty-three rural counties on an annual basis at each participating research site. Investigators defined rural using Butler and Beale's⁽²³⁾ coding system for metropolitan and non-metropolitan areas. Of the participating counties, seventeen were coded as 6 (population of 2500 to 19 999, adjacent to a metropolitan area), 7 (population of 2500 to 19 999, not adjacent to a metropolitan area) or 8 (completely rural or less than 2500 urban population, adjacent to a metropolitan area). The remaining six counties were in states without counties having Beale codes of 6-8. Four counties were in rural areas with no urban centres of populations exceeding 10 000, and two counties had no population centre greater than 10 000.

A convenience sample of 413 respondents was recruited in wave 1 from programmes and places serving low-income families, including Welfare-to-Work programmes, Head Start programmes, the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC), public health clinics, and low-income housing. This approach to sample selection allowed the research team to access this hard-to-reach population through contacts known to and trusted by the participants. Trust is an extremely important issue for families in rural communities where anonymity and privacy are difficult to acquire and maintain. Thus, convenience sampling was preferable over randomized sampling. To be eligible for participation, women had to be at least 18 years of age, have at least one child aged 13 years or younger, and be eligible for food stamps in their state. All data were collected via an interviewer-administered questionnaire typically lasting 1.5-2.0 h. Of the original 413 respondents interviewed at wave 1, 325 were re-interviewed at wave 2 and 270 were re-interviewed at wave 3 for an overall attrition rate of 34.6%.

Measures

Food insecurity was measured at all three waves by the Core Food Security Module (CFSM), an eighteen-item scale with a 12-month time reference administered as part of the Current Population Survey conducted by the US Census Bureau for the Bureau of Labor Statistics. The scale has good validity and reliability, with a reliability coefficient of $\alpha = 0.81$ for households with children⁽²⁴⁾.

The CFSM is similarly reliable for populations of racial and ethnic minorities⁽²⁵⁾. Reliability for this rural sample matched reported reliability ($\alpha = 0.81$).

Depression was measured at all three waves using the Center for Epidemiologic Studies–Depression Scale (CES-D), which was designed to measure depressive symptomatology in the general population⁽²⁶⁾. The measure includes twenty items, each rated on a 4-point scale ranging from ‘rarely’ to ‘most or all of the time’. Respondents are asked to report on how they have been feeling over the last week. Statements include: (i) ‘I felt depressed’; (ii) ‘I felt everything I did was an effort’; and (iii) ‘I felt lonely’. There are four subscales, which include depressed affect, negative affect, interpersonal relationships and somatic symptoms. The four questions for negative affect are framed positively and then reverse scored. Reliability for the CES-D is $\alpha = 0.85$ in the general population and $\alpha = 0.90$ in psychiatric populations. Reliability in this sample matched the reliability reported for the general population ($\alpha = 0.85$).

Five sociodemographic variables also were measured at wave 1, including age, ethnicity, household income, marital status and education. Age and household income were measured continuously. Marital status was measured in five categories (single, married, living with partner, divorced, separated). Ethnicity was measured in four categories (White, Black, Latina, other). Education was measured in six categories (8th grade or less, some high school, high school degree or general education

equivalent, technical training, some college/associate’s degree, college degree or higher).

Data analysis

Since the food insecurity scores at each wave had a positively skewed distribution, we applied a logarithmic transformation to these scores before carrying out the following analyses. Bivariate associations among the depression scores and food insecurity scores were assessed using Pearson correlations. Each Pearson correlation was based on all subjects for whom measurements on both variables were available; the number of such subjects varied according to which two variables were considered. We then formulated a structural equation model encompassing all three waves of depression scores and food insecurity scores. A schematic for the model is presented in Fig. 1. The coefficients α_1 and α_2 link a depression score to a subsequent depression score, α_3 and α_4 link a food insecurity score to a subsequent food insecurity score, α_5 and α_6 link a depression score to a subsequent food insecurity score, and α_7 and α_8 link a food insecurity score to a subsequent depression score.

We fit two versions of the structural equation model. The primary version was based on the 184 subjects for whom depression scores were recorded at all three waves (i.e. were not missing at any wave), food insecurity scores were recorded at all three waves, and income was recorded at the first wave. We required the availability of income for a test of whether α_1 to α_8 were the same across the two median-determined income strata, as previous work⁽²⁷⁾




Fig. 1 Schematic for the structural equation model. Depression at wave 2 was modelled as a function of depression at wave 1, food insecurity at wave 1, and an error variable E_1 . Depression at wave 3 was modelled as a function of depression at wave 2, food insecurity at wave 2, and an error variable E_2 . Food insecurity at wave 2 was modelled as a function of depression at wave 1, food insecurity at wave 1, and an error variable E_3 . Food insecurity at wave 3 was modelled as a function of depression at wave 2, food insecurity at wave 2, and an error variable E_4 . Depression and food insecurity at wave 1 were assumed to be correlated, as were the two error variables affecting depression and the two error variables affecting food insecurity. The coefficients α_1 to α_8 were estimated using the structural equation modelling capabilities of AMOS version 7

had documented a significant association between income and depression scores. Besides testing whether the coefficients were the same across income strata, we tested two other null hypotheses. One of these was that $\alpha_5 = \alpha_6 = 0$, meaning that depression would not impact food insecurity; the other was that $\alpha_7 = \alpha_8 = 0$, meaning that food insecurity would not impact depression.

The secondary version of the structural equation model was based on all 413 subjects, with missing values estimated by regression imputation prior to fitting the model. Since standard errors of coefficient estimates are inversely related to the square root of the sample size, the secondary version of the structural equation model should offer more precise, although slightly biased, coefficient estimates. All analyses were carried out using the Statistical Package for the Social Sciences version 15 and AMOS version 7 statistical software packages (SPSS Inc., Chicago, IL, USA).

Three complementary fit statistics were utilized to evaluate the structural equation models. First is the χ^2/df lack-of-fit statistic. The χ^2 value pertains to a null hypothesis that the population covariance matrix of the observed variables equals the covariance matrix implied by the model. Because this null hypothesis is generally presumed to be false, most researchers do not formally accept or reject it. Rather, they divide the χ^2 value by its degrees of freedom. A large quotient (>5) indicates serious lack of fit and a small quotient (<3) suggests little lack of fit⁽²⁸⁾. Second is the root-mean-square error of approximation (RMSEA) lack-of-fit statistic, which is a non-negative quantity expressing the discrepancy between the population covariance matrix and that implied by the model. A large RMSEA (>0.10) reflects serious lack of fit, while a small RMSEA (<0.05) implies little lack of fit⁽²⁸⁾. Third is the comparative fit index (CFI) goodness-of-fit statistic, which is a quantity between 0 and 1 describing how much the model improves upon a straw-man 'independence model'. A large CFI (>0.95) indicates good fit⁽²⁹⁾.

Results

Demographic data have been summarized elsewhere⁽²⁷⁾. Briefly, the majority of respondents were married or living

with a partner, non-Hispanic White, and had a high school education, general education equivalent, or less. The mean age of participants was 30.1 years and median household income was \$14 826. There was a significant association between income and depression scores, but depression scores were not significantly associated with marital status, race, education or age.

Pearson correlations for the depression scores and food insecurity scores are presented in Table 1. All of the correlations were positive and significantly different from 0 (all $P < 0.01$). Particularly strong correlations were found between wave 1 depression scores and wave 2 depression scores (0.534), between wave 2 depression scores and wave 3 depression scores (0.620), and between any two sets of food insecurity scores (0.524, 0.540, 0.547).

The results for the primary version of the structural equation model, supposing that α_1 to α_8 are the same across income strata, are summarized in the 'Complete case analysis' columns of Table 2. The estimates of α_1 , α_2 , α_3 , α_4 and α_7 were positive and significantly different from 0 (all $P < 0.001$). The estimates of α_5 , α_6 and α_8 were not significantly different from 0 (all $P > 0.10$). However, coefficient estimates based on a sample size as small as 184 are fairly imprecise, so non-significant estimates do not imply that the coefficients equal 0. More importantly, a test of the null hypothesis $\alpha_5 = \alpha_6 = 0$ yielded $\chi^2 = 6.754$ on 2 df ($P = 0.034$). Thus, while we cannot make the specific claim that $\alpha_5 \neq 0$ and cannot make the specific claim that $\alpha_6 \neq 0$, we can infer that at least one of these two coefficients is non-zero. Hence, the model cannot be simplified by omitting the links from depression to food insecurity, which is to say that there is a causal relationship from depression to food insecurity. In addition, a test of the null hypothesis $\alpha_7 = \alpha_8 = 0$ yielded $\chi^2 = 11.786$ on 2 df ($P = 0.003$). Therefore, the model cannot be simplified by omitting the links from food insecurity to depression, which is to say that there is a causal relationship from food insecurity to depression. In sum, the model asserts a bidirectional relationship between food insecurity and depression, and fit statistics support this: $\chi^2/df = 1.835$, RMSEA = 0.068, CFI = 0.989. Following the guidelines in the data analysis section, we

Table 1 Pearson correlations for depression and food insecurity: rural, low-income women with at least one child <13 years of age living in the home, USA, 2000–2002

	Depression Wave 1	Depression Wave 2	Depression Wave 3	Food insecurity Wave 1	Food insecurity Wave 2	Food insecurity Wave 3
Depression Wave 1	1	0.534	0.383	0.305	0.270	0.227
Depression Wave 2	0.534	1	0.620	0.335	0.292	0.344
Depression Wave 3	0.383	0.620	1	0.246	0.229	0.321
Food insecurity Wave 1	0.305	0.335	0.246	1	0.524	0.540
Food insecurity Wave 2	0.270	0.292	0.229	0.524	1	0.547
Food insecurity Wave 3	0.227	0.344	0.321	0.540	0.547	1

Pearson correlations were calculated, using SPSS version 15, to quantify the bivariate associations for all pairs of variables that could be chosen from among depression at wave 1, depression at wave 2, depression at wave 3, food insecurity at wave 1, food insecurity at wave 2, and food insecurity at wave 3. Each Pearson correlation was based on all subjects for whom measurements on both variables were available. All Pearson correlations were significantly different from 0 at the 0.01 level.

Table 2 Structural equation modelling results for the relationship between depression and food insecurity: rural, low-income women with at least one child <13 years of age living in the home, USA, 2000–2002

Coefficient	Complete case analysis				Analysis with missing values imputed			
	Estimate	SE	Standardized estimate	<i>P</i>	Estimate	SE	Standardized estimate	<i>P</i>
α_1	0.426	0.061	0.442	<0.001	0.468	0.036	0.528	<0.001
α_2	1.015	0.146	0.858	<0.001	0.870	0.064	0.853	<0.001
α_3	0.486	0.060	0.505	<0.001	0.500	0.036	0.565	<0.001
α_4	1.034	0.144	1.053	<0.001	0.911	0.073	0.962	<0.001
α_5	0.006	0.004	0.077	0.132	0.006	0.003	0.082	0.023
α_6	0.008	0.005	0.103	0.102	0.011	0.003	0.139	<0.001
α_7	2.436	0.708	0.212	<0.001	2.103	0.437	0.193	<0.001
α_8	-0.826	0.885	-0.058	0.350	-0.379	0.477	-0.030	0.427

For the structural equation model depicted in Fig. 1, the coefficients α_1 to α_8 were estimated using AMOS version 7. The results under 'Complete case analysis' are based on the 184 subjects for whom there were no missing values on depression or food insecurity and for whom income at wave 1 was not missing. The results under 'Analysis with missing values imputed' are based on all 413 subjects, with any missing values imputed using AMOS version 7 prior to fitting the structural equation model. Each *P* value pertains to a null hypothesis that the corresponding coefficient is 0.

determined that the structural equation model adequately represents the empirical patterns among depression scores and food insecurity scores, and the assertion of a bidirectional relationship is credible.

Finally, to justify the supposition that α_1 to α_8 are the same across income strata, we refit the model allowing α_1 to α_8 (along with variance/covariance parameters) to be estimated separately for subjects with wave 1 income below the median (\$14 826) and for subjects with wave 1 income above the median. We then tested the null hypothesis that α_1 to α_8 (along with variance/covariance parameters) are the same across income strata. We obtained $\chi^2 = 18.074$ on 17 df ($P = 0.384$), which justifies the supposition.

The results for the secondary version of the structural equation model are summarized in the 'Analysis with missing values imputed' columns of Table 2. The estimates of α_1 , α_2 , α_3 , α_4 , α_5 , α_6 and α_7 were positive and significantly different from 0 (P for $\alpha_5 = 0.023$, all other $P < 0.001$). The estimate of α_8 was not significantly different from 0 ($P = 0.427$). A test of the null hypothesis $\alpha_5 = \alpha_6 = 0$ yielded $\chi^2 = 23.558$ on 2 df ($P < 0.001$), while a test of the null hypothesis $\alpha_7 = \alpha_8 = 0$ yielded $\chi^2 = 22.636$ on 2 df ($P < 0.001$). Hence, the secondary version of the model also supports the conclusion of a bidirectional relationship between food insecurity and depression, and model fit statistics confirm this: $\chi^2/df = 2.454$, RMSEA = 0.059, CFI = 0.994. These statistics suggest that the secondary version of the model adequately captures the empirical patterns among depression scores and food insecurity scores, adding further credibility to the conclusion of a bidirectional relationship.

Discussion

The present study found evidence of a simultaneous causal relationship between food insecurity and depression in a sample of rural, low-income women. To our knowledge, this study is the first of its kind to examine longitudinally the relationship between food insecurity

and depression in the USA. This finding is important, not only because it underscores previous research showing food insecurity and depression are closely related, but also because it demonstrates the relationship is recursive. Thus, interventions aimed at reducing the prevalence of food insecurity in low-income rural households and depression in women living in these households can be approached on multiple levels.

Interventions aimed at increasing household food security in rural areas must target barriers systemically, including addressing high prices and increasing access to food stamps. Rural supermarket prices are on average 4% higher than in suburban areas⁽⁷⁾. Further, there are fewer supermarkets and more small grocery stores, the latter of which have prices averaging 10% higher than chain supermarkets nationwide. Low-income rural residents are particularly vulnerable to paying higher prices for food. Thus, tax incentives should be provided to major supermarket chains to establish a presence in rural areas with low supermarket penetration.

Beyond paying higher prices for food, one-third of rural residents eligible to receive food stamps do not participate in the programme⁽⁸⁾, which further limits access. Food stamp programmes should include campaigns to educate rural residents about eligibility requirements and increase access to this important resource. Further, research suggests that the effectiveness of food stamp redemption could be increased. A US Department of Agriculture analysis of food stamp redemptions in low-income, rural areas revealed that supermarkets accounted for only 52.8% of redemptions, compared with 76.7% of redemptions in supermarkets nationwide⁽⁷⁾. These data suggest that rural families receiving food stamps are more likely than their urban counterparts to spend these benefits in smaller grocery stores where food is more expensive, thus decreasing the buying power of the stamps. Support for strategies to maximize the value of food stamps is needed. Travel vouchers to offset the expense of travelling to a supermarket in a neighbouring community in conjunction with

budgeting and meal planning would assist families in making the most of food stamp participation.

Given the recursive relationship between food insecurity and depression, other possible interventions might occur at the level of the food stamp nutrition education programmes (FSNEP). In light of data demonstrating that deficiencies in vitamin B₆⁽³⁰⁾ and folic acid^(31,32) contribute to depressive illness, FSNEP could educate participants on how to economically increase their intakes of these nutrients. Further, FSNEP could develop and test brief depression screening and referral programmes. Such programmes might improve identification of depression in a population whose members are less likely to be seen for mental health problems than their urban and suburban counterparts⁽³³⁾. Screening for maternal depression in other non-traditional settings has been shown to be beneficial⁽³⁴⁾.

As with food insecurity, research shows that rural women living in low-income households are vulnerable to depression, specifically undiagnosed depression. Hauenstein and Boyd⁽¹⁶⁾ found that 41.4% of 181 rural women interviewed by telephone reported clinically depressive symptomatology on the CES-D. Within their sample, young, unemployed and poorly educated women were most likely to report depressive symptoms. Sears *et al.*⁽¹⁷⁾ assessed depression in 136 rural, low-income, predominantly female primary care patients using the Primary Care Evaluation of Mental Disorders (PRIME-MD) and found that 46% screened positive for depression; fifty-three of those sixty-three patients (40% of the total sample) were further evaluated and classified as depressed. A qualitative investigation of the daily hassles faced by thirty-three rural, poor single mothers found that nearly 69% of the sample reported clinically depressive symptomatology (CES-D scores of 16 or greater)⁽¹⁸⁾. Hauenstein and Peddada⁽²¹⁾ sampled 982 impoverished rural women attending a community health centre for a routine visit and found approximately half screened positive for depressive symptoms. Follow-up assessments revealed that 14.3% of the positive screeners had current MDE.

Despite these high prevalence rates, rural residents often do not receive treatment. Research shows that rural residents with a history of depression are less likely to seek professional help^(35,36). Further, stigma^(37,38) and lack of anonymity⁽³⁹⁾ prevent many rural residents from seeking psychological care. These social barriers are compounded by structural barriers, including the shortage of qualified mental health professionals in rural areas and the lack of service outreach⁽³³⁾. Additionally, Hauenstein⁽³⁷⁾ found that even when care is available, rural women may not access it, because traditional mental health care is not congruent with how women discuss and understand their depression. Thus, mental health interventions in rural areas might include education to de-stigmatize depression and its treatments (both

pharmacological and talk therapy) as well as programmes to increase access to mental health care, such as mobile mental health services and tele-health programmes. Further, given the simultaneous relationship between food insecurity and depression, mental health care providers in rural areas might include as part of their psychosocial evaluation an assessment of food security and dietary status. Such screenings may identify poor nutritional intake as an important contributor to poor mental health status and result in improved access to important food supports, such as food stamps.

Limitations

Findings from the present study must be considered within the context of its limitations. Although these data highlight the complexity of the relationship between food insecurity and depression over time in an under-researched population, this sample is not nationally representative, thus limiting generalizability to the broader rural population in the USA. Further, because the sample consists of rural residents only, the recursive relationship between food insecurity and depression found here may not extend to urban/suburban populations. Additionally, although results were similar for the primary model (missing data excluded) and the secondary model (missing data imputed), there is always the possibility of bias with incomplete data. Similarly, bias is also possible with self-reported data despite the use of standardized instruments to measure food insecurity and depression.

Conclusions

The present study highlights the importance of food security to good mental health and the importance of good mental health to food security. This is especially relevant for professionals serving populations that are vulnerable to both food insecurity and depression, such as rural low-income women. Resources directed at the health and well-being of such populations may be made more effective by simultaneously addressing nutrition and mental health. Programmes directed at the food security and nutrition needs of low-income families could address the potential influence of depression on food insecurity by assessing the need for mental health services. Individuals requiring mental health care could be referred to appropriate providers or programmes. Reciprocally, programmes and services designed to enhance the mental health of rural low-income families could include food security and nutritional health evaluations as part of their psychological intake assessment. Mental health providers treating individuals confronting depression as well as food insecurity and nutrition issues could refer their clients to the appropriate programmes and services. Public health nutritionists would benefit from education about the association between food insecurity and depression

within rural populations. Additionally, these professionals could work with mental health professionals to simultaneously and comprehensively address the consequences of poor nutrition and poor mental health in rural settings. Future research focused on complementary nutrition and mental health interventions is warranted.

Acknowledgements

This research was supported in part by USDA/CSREES/NRICGP Grant 2001-35401-10215, 2002-35401-11591 and 2004-35401-14938, and by NIH/NIDA Grant 5 K12 DA014040-07 (to L.A.S.). The opinions expressed herein are solely those of the authors. There are no additional conflicts of interest. C.H.-C. contributed to the writing of the manuscript. R.C. conducted the analyses and contributed to the writing. L.A.S. conceptualized the study, contributed to the writing, and assisted with primary data collection for the 'Rural Families Speak' project.

References

1. Probst JC, Ladtika SB, Moore CG, Harun N, Powell MP & Baxley EG (2006) Rural-urban differences in depression prevalence: implications for family medicine. *Health Serv Res* **38**, 653-660.
2. DeNavas-Walt C, Proctor BD & Smith J, US Census Bureau (2007) *Income, Poverty, and Health Insurance Coverage in the United States: 2006*. Washington, DC: US Government Printing Office.
3. Economic Research Service (2005) *Rural America at a Glance: 2005. Economic Information Bulletin* no. 4. Washington, DC: US Department of Agriculture.
4. US Census (2001) *Survey of Income and Program Participation Users' Guide (Supplement to the Technical Documentation)*, 3rd ed. Washington, DC: US Department of Commerce, Economic and Statistics Administration.
5. Economic Research Service (2003) Food security in the United States: measuring food security. <http://www.ers.usda.gov/Briefing/FoodSecurity/measurement> (accessed November 2003).
6. Nord M, Andrews M & Carlson S (2007) *Household Food Security in the United States, 2006. Economic Research Report* no. 49. Washington, DC: US Department of Agriculture.
7. US Department of Agriculture (1999) The rural poor's access to supermarkets and large grocery stores. *Fam Econ Nutr Rev* **12**, 90-99.
8. Wolkwitz K (2007) *Technical Appendices to Trends in Food Stamp Participation Rates: 1999 to 2005: Final Report*. Report no. 6044-409. Princeton, NJ: Mathematica Policy Research, Inc.
9. Kaiser L, Baumrind N & Dumbauld S (2007) Who is food-insecure in California? Findings from the California Women's Health Survey, 2004. *Public Health Nutr* **10**, 574-581.
10. Chilton M & Booth S (2007) Hunger of the body and hunger of the mind: African American women's perceptions of food insecurity, health and violence. *J Nutr Educ Behav* **39**, 116-125.
11. Casey P, Goolsby S, Berkowitz C *et al.*; Children's Sentinel Nutritional Assessment Program Study Group (2004) Maternal depression, changing public assistance, food security, and child health status. *Pediatrics* **113**, 298-304.
12. Heflin CM, Siefert K & Williams DR (2005) Food insufficiency and women's mental health: findings from a 3-year panel of welfare recipients. *Soc Sci Med* **61**, 1971-1982.
13. Siefert K, Heflin CM, Corcoran ME & Williams DR (2001) Food insufficiency and the physical and mental health of low-income women. *Women Health* **32**, 159-177.
14. Siefert K, Heflin CM, Corcoran ME & Williams DR (2004) Food insufficiency and physical and mental health in a longitudinal survey of welfare recipients. *J Health Soc Behav* **45**, 171-186.
15. Hauenstein EJ, Petterson S, Merwin E, Rovnyak V, Heise B & Wagner D (2006) Rurality, gender, and mental health treatment. *Fam Community Health* **29**, 169-185.
16. Hauenstein EJ & Boyd MR (1994) Depressive symptoms in young women of Piedmont: prevalence in rural women. *Women Health* **21**, 105-123.
17. Sears SF, Danda CE & Evans GD (1999) PRIME-MD and rural primary care: detecting depression in a low-income rural population. *Prof Psychol Res Pr* **30**, 357-360.
18. Wijinberg M & Reding K (1999) Reclaiming a stress focus: the hassles of rural, poor single mothers. *Fam Soc* **80**, 506-515.
19. Brown AC, Brody GH & Stoneman Z (2000) Rural black women and depression: a contextual analysis. *J Marriage Fam* **62**, 187-198.
20. Carruth AK & Logan CA (2002) Depressive symptoms in farm women: effects of health status and farming lifestyle characteristics, behaviors, and beliefs. *J Community Health* **27**, 213-228.
21. Hauenstein EJ & Peddada SD (2007) Prevalence of major depressive episodes in rural women using primary care. *J Health Care Poor Underserved* **18**, 185-202.
22. Simmons LA, Huddleston-Casas CA & Berry A (2007) Low-income rural women and depression: factors associated with self-reporting. *Am J Health Behav* **31**, 657-666.
23. Butler MA & Beale C (1994) *Rural-Urban Continuum Codes for Metro and Nonmetro Counties, 1993*. Report no. 9425. Washington, DC: US Government Printing Office.
24. Hamilton WL, Cook JT & Thompson WW (1997) *Household Food Insecurity in the United States in 1995: Summary Report of the Food Security Measurement Project*. Alexandria, VA: Food and Consumer Service, US Department of Agriculture.
25. Frongillo EA (1999) Validation of measures of food insecurity and hunger. *J Nutr* **129**, 506S-509S.
26. Radloff LS (1977) The CES-D scale: a self-report depression scale for research in the general population. *Appl Psychol Meas* **1**, 385-401.
27. Simmons LA, Braun B, Charnigo R, Havens JR & Wright DW (2008) Depression and poverty among rural women: a relationship of social causation or social selection? *J Rural Health* **24**, 292-298.
28. Bollen KA & Long JS (editors) (1993) *Testing Structural Equation Models*. Newbury Park, CA: Sage Publications.
29. Mueller RO (1996) *Basic Principles of Structural Equation Modeling*. New York, NY: Springer-Verlag.
30. Hvas AM, Juul S, Bech P & Nexø E (2004) Vitamin B₆ level is associated with symptoms of depression. *Psychother Psychosom* **73**, 340-343.
31. Alpert JE & Fava M (1997) Nutrition and depression: the role of folate. *Nutr Rev* **55**, 145-149.
32. Alpert JE, Mischoulon D, Nierenberg AA & Fava M (2000) Nutrition and depression: focus on folate. *Nutrition* **16**, 544-581.
33. Spoth R (1997) Challenges in defining and developing the field of rural mental disorder preventive intervention research. *Am J Community Psychol* **25**, 425-448.

34. Olson AL, Dietrich AJ, Prazar G & Hurley J (2006) Brief maternal depression screening at well-child visits. *Pediatrics* **118**, 207–216.
35. Arons B (2002) Mental health services in rural America. *Vital Speeches Day* **66**, 369–371.
36. Rost K, Smith R & Taylor J (1993) Rural–urban differences in stigma and the use of care for depressive disorders. *J Rural Health* **9**, 57–62.
37. Hauenstein EJ (2003) No comfort in the rural south: women living depressed. *Arch Psychiatr Nurs* **17**, 3–11.
38. Hoyt D, Conger R, Valde J & Weihs K (1997) Psychological distress and help seeking in rural America. *Am J Community Psychol* **25**, 449–470.
39. Kelleher KJ & Robbins JM (1997) Social and economic consequences of rural alcohol use. *NIDA Res Monogr* **168**, 196–219.