

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

ANDRILL Research and Publications

Antarctic Drilling Program

2008

Palaeontological Characterisation and Analysis of the AND-2A Core, ANDRILL Southern McMurdo Sound Project, Antarctica

M. Taviani

Istituto di Scienze Marine (ISMAR)-CNR, marco.taviani@bo.ismar.cnr.it

M. Hannah

Victoria University of Wellington

D. M. Harwood

University of Nebraska at Lincoln, dharwood1@unl.edu

S. E. Ishman

Southern Illinois University

K. Johnson

Ohio State University

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/andrillrespub>



Part of the [Environmental Indicators and Impact Assessment Commons](#)

Taviani, M.; Hannah, M.; Harwood, D. M.; Ishman, S. E.; Johnson, K.; Olney, M.; Riesselman, C.; Tuzzi, E.; Beu, A. G.; Blair, S.; Cantarelli, V.; Ceregato, A.; Corrado, S.; Mohr, B.; Nielson, S. H.H.; Persico, D.; Petrushak, S.; Raine, J. I.; Warny, S.; and ANDRILL-SMS Science Team, "Palaeontological Characterisation and Analysis of the AND-2A Core, ANDRILL Southern McMurdo Sound Project, Antarctica" (2008). *ANDRILL Research and Publications*. 41.

<https://digitalcommons.unl.edu/andrillrespub/41>

This Article is brought to you for free and open access by the Antarctic Drilling Program at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in ANDRILL Research and Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

M. Taviani, M. Hannah, D. M. Harwood, S. E. Ishman, K. Johnson, M. Olney, C. Riesselman, E. Tuzzi, A. G. Beu, S. Blair, V. Cantarelli, A. Ceregato, S. Corrado, B. Mohr, S. H.H. Nielson, D. Persico, S. Petrushak, J. I. Raine, S. Warny, and ANDRILL-SMS Science Team

Palaeontological Characterisation and Analysis of the AND-2A Core, ANDRILL Southern McMurdo Sound Project, Antarctica

M. TAVIANI^{1*}, M. HANNAH², D.M. HARWOOD³, S.E. ISHMAN⁴, K. JOHNSON⁵, M. OLNEY⁶,
C. RIESSELMAN⁷, E. TUZZI³, R. ASKIN⁸, A.G. BEU⁹, S. BLAIR¹⁰, V. CANTARELLI¹¹,
A. CEREGATO¹, S. CORRADO¹¹, B. MOHR¹², S.H.H. NIELSEN¹⁰, D. PERSICO¹³,
S. PETRUSHAK¹⁰, J.I. RAINE⁹, S. WARNY¹⁴ & THE ANDRILL-SMS SCIENCE TEAM¹⁵

¹Istituto di Scienze Marine (ISMAR)-CNR, v. Gobetti 101, I-40129 Bologna – Italy

²Antarctic Research Centre, Victoria University of Wellington, P.O. Box 600, Wellington – New Zealand

³Dept. of Geosciences, ANDRILL SMO, 214 Bessey Hall, University of Nebraska, Lincoln, NE 68588-0340 – USA

⁴Dept. of Geology, Southern Illinois University, 1259 Lincoln Drive, Carbondale, IL 62901-6899 – USA

⁵School of Earth Sciences, The Ohio State University, 125 South Oval Mall, Columbus OH 43210 – USA

⁶University of South Florida, Dept. of Geology, 4202 E. Fowler Ave., SCA 528 Tampa, FL 33620-8100 – USA

⁷School of Earth Sciences, Stanford University, Braun Hall, Bldg. 320, Stanford, CA 94305-2115 – USA

⁸1930 Bunkhouse Dr., Jackson, WY 83001 – USA

⁹Paleontology & Environmental Change Section, GNS Science, 1 Fairway Drive, Lower Hutt 5040 – New Zealand

¹⁰Antarctic Marine Geological Research Facility, Florida State University, Tallahassee, FL 32306-4100 – USA

¹¹Dip. di Scienze Geologiche, Univ. degli Studi Roma Tre, Largo S. Leonardo Murialdo, 1, I-0014 Rome – Italy

¹²Naturhistorisches Forschungsinstitut, invalidenstrasse 43 D-10115, Berlin – Germany

¹³Dip. di Scienze della Terra, Università degli Studi di Parma, V.le G.P. Usberti 157A, I-40100 Parma – Italy

¹⁴Dept. of Geology and Geophysics, Louisiana State University, Baton Rouge, LA 70803 – USA

¹⁵<http://andriill.org/projects/sms/team.htm>

*Corresponding author (marco.taviani@bo.ismar.cnr.it)

Abstract – The palaeontological yield of the 1138.54 metre-long AND-2A sedimentary rock core provides unique documentation of Neogene environments in the Ross Sea region of Antarctica. Especially important is the biological legacy of the climatically crucial ‘mild’ middle Miocene phase. Diatom-bearing units provide key information for stratigraphic intervals never previously recovered from locations proximal to the Antarctic continent and constrain the age model for the AND-2A core. Benthic calcareous (and agglutinated) foraminifera were present at many levels; remarkable is the occurrence of planktonic taxa only seldom found in the Neogene nearshore record of the Ross Sea region. The sporadic occurrence of calcareous dinoflagellate remains (thoracosphaerids) is consistent with warmer-than-present seawater during the Miocene. Marine palynomorphs are almost ubiquitous, although their abundance and diversity are variable. Pollen and spores from the middle Miocene section suggest a mossy tundra vegetation and represent the first stratigraphically-constrained record of terrestrial vegetation in Victoria Land during this time. Fragments of lignin-rich organic matter (huminite-vitrinite and inertinite groups) are particularly predominant during the Miocene climatic optimum, and continue into the Pliocene. Macrofossils are reasonably common throughout the core. Polychaete worm tubes were almost ubiquitous. Especially remarkable is the bivalve record (mainly pectinids), with 4-5 different taxa pointing out a mild climatic situation in the Miocene nearshore.

INTRODUCTION

Drilling operations for the Southern McMurdo Sound (SMS) Project were conducted in the austral spring 2007 to recover the AND-2A core at 77°44.27'76"S and 165°17'18.24"E. This core comprises 1138.54 metres of various types of sedimentary rocks with some volcanics encompassing lower Miocene to Quaternary successions (Fielding et al., Fig. 1, this volume; Acton et al., this volume, b) (Fig. 1).

The Core Characterisation Phase of the SMS Project was conducted by an on-ice Palaeontology Team that included seven specialists of taxonomic groups of recognised biostratigraphic and environmental utility in the Antarctic Cenozoic, and an off-ice team who produced results reported herein after the

drilling phase was completed. The palaeontologic record of the AND-2A core is highly remarkable in terms of quality and quantity of fossils, with over 1200 samples collected on-ice, including those for off-ice specialists. The on-ice treatment of such a large set of samples was possible through the collaborative work of additional SMS on-ice personnel (J. Carnes, R. Frisch-Gleason, J. Hamre, L. Jovane, K. Mankhoff, K. Pound, J. Reed, E. Strada, P.-N. Webb, and R. Williams). It is worth mentioning the important role provided by using thin-sections (Fig. 2) to identify fossils in the AND-2A core, that were otherwise undetectable through visual inspection or washing; these on-ice thin sections were provided by S. Petrushak. Sample aliquots from intervals of presumptive critical palaeoclimatic significance were