

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Uniformed Services University of the Health
Sciences

U.S. Department of Defense

2011

Morphine-induced modulation of LTD at GABAergic synapses in the ventral tegmental area

Matthieu Dacher

Uniformed Services University of the Health Sciences

Fereshteh S. Nugent


Uniformed Services University of the Health Sciences, fnugent@usuhs.mil

Follow this and additional works at: <https://digitalcommons.unl.edu/usuhs>

 Part of the [Medicine and Health Sciences Commons](#)

Dacher, Matthieu and Nugent, Fereshteh S., "Morphine-induced modulation of LTD at GABAergic synapses in the ventral tegmental area" (2011). *Uniformed Services University of the Health Sciences*. 39.
<https://digitalcommons.unl.edu/usuhs/39>

This Article is brought to you for free and open access by the U.S. Department of Defense at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Uniformed Services University of the Health Sciences by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Morphine-induced modulation of LTD at GABAergic synapses in the ventral tegmental area

Matthieu Dacher, Fereshteh S. Nugent*

Uniformed Services University of the Health Sciences, Department of Pharmacology, Bethesda, MD 20814, USA

ARTICLE INFO

Article history:

Received 29 September 2010

Received in revised form

2 November 2010

Accepted 11 November 2010

Keywords:

LTP

LTD

Morphine

VTA

Addiction

Synaptic plasticity

ABSTRACT

Adaptive behaviors often require the learning of appropriate responses to rewarding stimuli, yet aberrant learning processes can lead to serious diseases such as addiction. Dopamine (DA) neurons of the ventral tegmental area (VTA) play an essential role in the treatment of rewarding stimuli, and they exhibit plasticity in response to such stimuli, but also to drugs of abuse. Previously we discovered a form of presynaptic nitric oxide (NO)-mediated long-term potentiation (LTP_{GABA}) at GABAergic synapses onto VTA DA neurons that is prevented with morphine *in vivo* 24 h after exposure. Here we investigated whether the same GABAergic synapses are capable of exhibiting long-term depression (LTD in addition to LTP_{GABA}) and its possible modulation by morphine *in vivo*. We found that indeed the efficacy of VTA GABAergic synapses can be down-regulated through induction of a novel form of LTD (i.e., LTD_{GABA}) in response to synaptic stimulation. Paired pulse ratio (PPR) and coefficient of variance (CV) analyses of evoked IPSCs confirmed that this plasticity may be postsynaptic. Consistently, LTD_{GABA} did not involve presynaptic cannabinoid CB₁ receptors (CB₁Rs). Moreover, NMDAR activation was not necessary for LTD_{GABA}. However, blockade of D₂ dopamine receptors (D₂R) significantly attenuated LTD_{GABA} proposing a novel synaptic mechanism for the regulation of excitability of DA neurons by endogenous DA and D₂R activation. Interestingly, 24 h after a single *in vivo* exposure to morphine, LTD_{GABA} was absent in slices from morphine-treated rats but unaffected in slices from saline-treated rats, confirming a bidirectional impact of morphine on GABAergic synaptic plasticity in the VTA. The control of bidirectional GABAergic plasticity by morphine in the VTA may represent the neural correlates necessary for the addictive properties of opiates.

This article is part of a Special Issue entitled 'Synaptic Plasticity and Addiction'.

Published by Elsevier Ltd.

1. Introduction

Drugs of abuse to include opiates, hijack synaptic plasticity mechanisms in reward-related brain areas. This highlights their value and reinforces their abuse (i.e., establishment of compulsive drug-seeking and drug-taking behaviors) (Kauer and Malenka, 2007). More recently, a critical link was found between losses of synaptic plasticity in the nucleus accumbens (NAc) of cocaine-addicted

animals to their inability to control their cocaine-intake (Kasanez et al., 2010). Such convincing evidence in line with previous findings supports the idea that synaptic plasticity is required not only for reinforcement of reward-based learning and motivated behaviors, but also for the addiction process. The well-studied forms of synaptic plasticity implicated in learning and addiction are LTP (strengthening of synapses) and LTD (weakening of synapses).

The VTA consisting of DA and GABA neurons is an area crucial for reward and drug reward. DA release originating from VTA DA neurons in a VTA target (specifically in the NAc) is believed to initiate drug-induced neuroplasticity necessary for the formation of addictive behaviors (Kauer and Malenka, 2007). Both glutamatergic and GABAergic synapses regulating VTA DA cell activity are found to be drug-sensitive and subject to plasticity (Nugent and Kauer, 2008). Glutamatergic synapses onto VTA DA neurons can be potentiated in response to a single *in vivo* exposure to different classes of addictive drugs including opiates which is believed to be essential for initiation of behavioral sensitization (Saal et al., 2003;

Abbreviations: LTP, Long-term potentiation; LTD, long-term depression; VTA, ventral tegmental area; NAc, nucleus accumbens; DA, dopamine; NO, nitric oxide; NMDAR, NMDA receptor; CB₁R, cannabinoid CB₁ receptor; eCB, endocannabinoids; D₂R, dopamine D₂ receptor; PPR, paired pulse ratio; CV, coefficient of variation; LFS, low frequency stimulation.

* Corresponding author. Edward Hebert School of Medicine, Department of Pharmacology, Uniformed Services University, 4301 Jones Bridge Road, Bethesda, MD, 20814, USA. Tel.: +1 301 295 3243; fax: +1 301 295 3220.

E-mail address: fnugent@usuhs.mil (F.S. Nugent).

Ungless et al., 2001). On the other hand, GABA enhancers have been found to be effective in impeding the learning of drug-seeking behaviors (Volkow et al., 2004) but many unacceptable side effects cannot be avoided. Interestingly, local facilitation of GABAergic inhibition in the VTA prevents addictive behaviors in animal models of addiction (Backes and Hemby, 2008; Lee et al., 2007; Xi and Stein, 2000) supporting the idea that selective GABAergic intervention in this area will likely offer new and promising treatments for drug addiction. In addition to drugs elevating GABA levels or GABA agonists directly targeting GABA receptors, an increase or a decrease in GABAergic inhibition can be achieved through synaptic plasticity. Our current knowledge about the control of GABAergic synaptic strength in the VTA is sparse but presents an important area of interest. Our previous work provided the first evidence of modulation of LTP at GABAergic synapses by morphine *in vivo*. We discovered a novel presynaptic form of LTP (LTP_{GABA}) at GABAergic synapses onto VTA DA neurons mediated by the NO/cGMP/cGMP-dependent protein kinase (PKG) signaling cascade. We found that 24 h after a single exposure to opiates, LTP_{GABA} is blocked presumably by opiate interaction at the level of guanylyl cyclase (Nugent et al., 2009, 2007). Opiate-induced blockade of LTP_{GABA} could emerge as early as 2 h after morphine exposure and could last up to 5 days. Intriguingly, LTP_{GABA} is also found to be a major target for other addictive drugs (nicotine, cocaine and ethanol) and even stress suggesting that GABAergic plasticity in the VTA may in fact be a common synaptic target for all addictive drugs (Guan and Ye, 2010; Niehaus et al., 2010). It is worthwhile to be mentioned that the common method for identification of DA neurons in the VTA slices is the measurement of a hyperpolarization-activated current (I_h) including the present study. Interestingly, it has been shown that non-DA neurons can also express I_h currents, therefore I_h alone may not reliably represent DA cells (Margolis et al., 2006). Nevertheless, a recent study also confirmed that the larger I_h is a consistent characteristic of DA neurons (Zhang et al., 2010). Therefore in our experiments, a small subset of the neurons recorded from and reported here are also non-DA neurons. Here we report that GABAergic synapses onto I_h-positive neurons, presumably DA neurons, exhibit a form of post-synaptic LTD (LTD_{GABA}) that can also be modulated by morphine *in vivo*. The bidirectional regulation of GABAergic plasticity in the VTA by morphine provides additional cellular correlates by which opiates and maybe other drugs of abuse could regulate DA cell excitability and DA release.

2. Materials and methods

2.1. Animals

All experiments were carried out in accordance with the National Institutes of Health guidelines for the care and use of laboratory animals and were approved by the Uniformed Services University Institutional Animal Care and Use Committee. All efforts were made to minimize animal suffering, and to reduce the number of animals used. Animals were 2–3 weeks old Sprague–Dawley rats provided food and water ad libitum, on a 12 h light/dark cycle.

2.2. Preparation of brain slices

Rats were anesthetized using isoflurane and quickly decapitated. The brain was rapidly removed into ice-cold artificial cerebrospinal fluid (ACSF) containing (in mM): 126 NaCl, 21.4 NaHCO₃, 2.5 KCl, 1.2 NaH₂PO₄, 2.4 CaCl₂, 1.2 MgSO₄, 11.1 glucose, 0.4 ascorbic acid, saturated with 95% O₂/5% CO₂ (pH 7.4). Horizontal midbrain slices containing the VTA (250 μm thick) were cut and incubated for at least 1 h at 34 °C, and then transferred to a recording chamber where the slice was submerged in warmed ACSF.

2.3. Electrophysiology

Whole cell recordings were performed on midbrain slices using a patch amplifier (Multiclamp 700B) under infrared-differential interference contrast

microscopy. Data acquisition and analysis were carried out using DigiData 1440A and pClamp 10 (molecular devices, Union City, CA). Signals were filtered at 3 kHz and digitized at 10 kHz. The recording ACSF was the same as the cutting solution except that it was ascorbic acid-free. Slices were perfused at 2–4 ml/min and recorded at 28–29 °C. To study GABA_A-mediated synaptic transmission, 6,7-dinitroquinoxaline-2,3-dione (DNQX; 10 μM), and strychnine (1 μM) were added to block AMPA-, and glycine-receptor-mediated synaptic currents, respectively. Patch pipettes were filled with (in mM): 125 KCl, 2.8 NaCl, 2 MgCl₂, 2 ATP–Na⁺, 0.3 GTP–Na⁺, 0.6 EGTA, and 10 HEPES (pH adjusted 7.28 with KOH, 275–280 mOsm).

Cells were voltage-clamped at –70 to –80 mV except during LTD protocol, and the cell input resistance and series resistance were monitored throughout the experiment; experiments were discarded if these values changed by more than 10% during the experiment. I_h current measurement was used to identify DA neurons by stepping cells from –50 mV to –100 mV. In our experimental conditions, the average of I_h current was 213 ± 27 pA, n = 62. However, in each set of our experiments there is a small subset of non-DA neurons that express I_h currents. Paired GABA_A IPSCs were stimulated using a bipolar stainless steel stimulating electrode placed 200–500 μm rostral to the recording site in the VTA at 0.1 Hz (100 μs). The stimulation intensity was adjusted so that the amplitude of IPSCs ranged between –200 and –600 pA. To induce LTD_{GABA}, afferents were stimulated at 1 Hz (low frequency stimulation, LFS) for 6 min while cells were held at –40 mV (voltage-clamp mode).

2.4. Drug application

Drugs were superfused at indicated concentrations for at least 15 min before LTD protocol. Interleaved control experiments were performed with experiments in which drugs were bath applied. Salts and all other drugs were obtained from Sigma–Research Biochemicals International or Tocris Bioscience.

2.5. Morphine treatment

For treatment with morphine, rats were injected once 24 h prior to sacrifice for slice preparation. Morphine sulfate was dissolved in 0.9% saline, and saline was used to treat control animals. Rats were brought up from the animal facility in the morning, weighed, and injected intraperitoneally with either 10 mg/kg morphine or a comparable volume of saline. They were then returned to their home cages after 2 h. Rats were brought back up to the lab for sacrifice and brain slice preparation 24 h post-morphine.

2.6. Statistics

Values are presented as means ± s.e.m. Statistical significance was determined using a two-tailed pair-sample or two-sample Student's *t*-test with significance level of *p* < 0.05. Levels of LTD are reported as averaged IPSC amplitudes for 5 min just before LTD induction compared with averaged IPSC amplitudes during the 5 min period from 20 to 25 min after LFS. PPR and CV were measured as previously described (Nugent et al., 2007). Briefly, PPR (50 ms interstimulus interval) and CV were measured over 5 min epochs of 30 IPSCs each. The peak values of the evoked paired IPSCs were measured relative to the same baseline. A stable baseline value was considered in each sweep of paired pulses starting at 20–50 ms right before the emergence of the inward IPSC current using pClamp 10 software. The average value for the amplitude of all 30 IPSC₂ was divided by the average value for the amplitude of the corresponding 30 IPSC₁ and reported as the mean PPR for that epoch. 1/CV₂ was measured by dividing the mean amplitude of IPSCs recorded over 5 min epochs by the variance of these IPSCs (only IPSC₁s were used to determine this value).

3. Results and discussion

3.1. LTD at GABAergic synapses onto VTA DA neurons (LTD_{GABA})

In continuation of our major finding of morphine-blockade of LTP_{GABA} (Nugent et al., 2007), we explored the possibility that GABAergic synapses originating from GABA interneurons in the VTA are capable of bidirectional plasticity (expression of LTD in addition to LTP). Using a pairing LTD protocol, low frequency synaptic stimulation with modest depolarization, we were able to reliably induce an enduring and stable LTD_{GABA} in DA cells (Fig. 1a–b, paired *t*-test: *p* < 0.000001). The same pairing stimulation paradigm has been shown to induce LTD of excitatory synapses onto DA neurons in the VTA (Jones et al., 2000; Thomas et al., 2000). The induction of such a novel inhibitory LTD in the VTA could provide a natural mechanism to increase DA neuron excitability by decreasing GABA inhibition and also promoting an excitatory glutamatergic LTP in the VTA (i.e., metaplasticity).




Fig. 1. GABAergic synapses onto DA neurons exhibit LTD in response to synaptic stimulation. During LTD induction protocol, LFS was paired with a modest depolarization for 6 min; for simplicity in this and other figures we only use LFS to show the LTD induction protocol. (a) Sample experiment of LTD_{GABA} and IPSCs recorded from a DA neuron in a control slice. LFS was delivered at the arrow. Inset: averaged IPSCs before (black) and 25 min after LFS (red). In this and all figures, ten consecutive IPSCs from each experiment were averaged for figure. Calibration for insets: 20 ms, 50 pA. The dotted line in this and other figures is an estimate of the mean response before LFS. (b) Average of 24 experiments from DA neurons expressing LTD_{GABA} in response to LTD stimulation protocol. LTD_{GABA} was reliably triggered in 22 out of 24 cells, but data from all cells are included in this and next figures (control LTD, filled symbols). (c) and (d) No changes in PPR or 1/CV² were detected after induction of LTD_{GABA} (data generated from all control cells). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article).

Remarkably, repeated exposure to cocaine *in vivo* is found to induce a form of presynaptic LTD of GABAergic synapses in the VTA. Cocaine-induced inhibitory LTD is induced using bath application of cocaine paired with synaptic stimulation (Pan et al., 2008b). The cocaine-induced LTD protocol is complex because cocaine inhibits the uptake of DA, norepinephrine and serotonin and each of these neurotransmitters could play a role in induction of such LTD that might only be specific to cocaine. To date, LTD_{GABA} is the first demonstration of an inhibitory LTD in the VTA in response to synaptic stimulation. Simultaneous induction of excitatory LTP and LTD_{GABA} in VTA DA neurons might work in concert to code for rewarding properties of natural rewarding stimuli as well as salience, therefore reinforcing the learning of reward-motivated behaviors. On the other hand, LTD_{GABA} if triggered by drugs of abuse along with excitatory LTP could code for drug-induced reward.

3.2. LTD_{GABA} is expressed postsynaptically

To identify the locus of LTD_{GABA} expression, PPR and CV analyses were performed in control experiments. PPR and CV are measures that reliably change with changes in probability of release and can be used to predict the nature of synaptic plasticity. The apparent advantage of CV is that it is independent of quantal size; therefore if CV changes similarly to the mean values of IPSCs, the locus of expression of plasticity is presynaptic, whereas if CV is unaffected, synaptic plasticity is expressed postsynaptically (Faber and Korn, 1991). In our experiments, we did not detect any significant changes associated with LTD_{GABA} in PPR or CV (Fig. 1c–d, paired *t*-test: $p = 0.121$ for PPR, and $p = 0.579$ for CV) which strongly points to a postsynaptic nature of LTD_{GABA}. This is in contrast to the nature of LTP_{GABA} which is induced postsynaptically but expressed presynaptically (i.e., through an increase in presynaptic GABA

release). Our results suggest that postsynaptic LTD_{GABA} may result from a decrease in number and/conductance of GABA_A receptors similar to a postsynaptic form of LTD of GABA inhibition in rat deep cerebellar nuclei (Morishita and Sastry, 1993, 1996). However, in contrast to deep cerebellar nuclear inhibitory LTD, LTD_{GABA} is NMDAR independent (see below).

3.3. LTD_{GABA} does not require NMDAR and CB₁R activation

NMDAR activation is required for induction of some forms of LTD at inhibitory GABAergic synapses in the brain (Komatsu and Iwakiri, 1993; Lu et al., 2000; McLean et al., 1996; Wang and Stelzer, 1996) and also in LTP_{GABA} of the VTA. In our LTD experiments, only AMPARs were blocked, therefore there was a possibility of NMDAR activation during LTD induction as cells were significantly depolarized during the induction protocol which could open NMDARs. Interestingly, blockade of NMDARs with d-APV (50 μM) did not block LTD_{GABA} suggesting that LTD_{GABA} in the VTA is NMDAR independent (Fig. 2a; paired *t*-test: $p = 0.007$ for control cells and $p = 0.012$ for APV cells). Similarly, a widespread form of presynaptic LTD at GABA_A synapses, i.e., endocannabinoid-mediated inhibitory LTD (eCB-LTD), is also NMDAR independent (Heifets and Castillo, 2009), although LTD_{GABA} is postsynaptic and does not require CB₁ receptors (see below).

Because of the importance and widespread expression of eCB-LTD at many synapses including eCB-LTD in the VTA induced by cocaine (Pan et al., 2008b), we explored whether eCBs also play a critical role in LTD_{GABA}. Unlike eCB-LTD, the CB₁R antagonist, AM251 (10 μM), did not have any effect on the induction of LTD_{GABA}, suggesting that LTD_{GABA} does not involve presynaptic CB₁Rs (Fig. 2b; paired *t*-test: $p = 0.007$ for control cells and $p = 0.004$ for AM251 cells). This is consistent with our results derived from PPR




Fig. 2. LTD_{GABA} is independent of NMDARs and CB1Rs. (a) APV (50 μM) did not affect LTD_{GABA} (control LTD, filled symbols: 76 ± 1.4% of pre-LFS values, *n* = 8; APV cells, open symbols: 82 ± 4% of pre-LFS values, *n* = 8). Insets: averaged IPSCs in APV before (black) and 25 min after LFS (red). Calibration for insets: 20 ms, 50 pA. (b) Bath-applied AM251 (10 μM) has no effect on LTD_{GABA} (control LTD, filled symbols: 72 ± 2% of pre-LFS values, *n* = 8; AM251 cells, open symbols: 72 ± 1% of pre-LFS values, *n* = 7). Insets: averaged IPSCs in AM251 before (black) and 25 min after LFS (red). Calibration for insets: 20 ms, 50 pA. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article).

and CV analyses which points to the postsynaptic nature of LTD_{GABA}. Given that CB₁Rs are exclusively expressed on GABAergic terminals in the VTA (Szabo et al., 2002), it is not surprising that these receptors are not required for expression of LTD_{GABA}.

3.4. D₂Rs are involved in induction of LTD_{GABA}

Somatodendritic release of DA in the VTA and its action on D₂ autoreceptors provides an endogenous feedback inhibitory mechanism for controlling the excitability of DA neurons. Interestingly, activation of D₂Rs (proposed to be located onto GABA terminals which is not anatomically verified) in the VTA could also facilitate the induction of cocaine-induced eCB-LTD through inhibition of the cAMP-PKA signaling cascade (Pan et al., 2008a). Given that drugs of abuse could increase somatodendritic release of DA (Bradberry and Roth, 1989; Campbell et al., 1996; Klitenick et al., 1992) and the critical role of D₂Rs in induction of cocaine-induced eCB-LTD and of

glutamatergic LTD in the VTA (Jones et al., 2000; Thomas et al., 2000), we next examined whether D₂R activation is also required for LTD_{GABA}. In the presence of D₂R antagonist, sulpiride (10 μM), we attempted to induce LTD_{GABA}. Sulpiride significantly attenuated LTD_{GABA} suggesting a critical role of D₂Rs in induction of this plasticity (Fig. 3a–b; paired *t*-test: *p* = 0.002 for control cells and *p* = 0.333 for sulpiride cells). It still remains to be determined whether post synaptic disruption of G protein signaling and also Ca²⁺ signaling block the induction of LTD_{GABA}. Based on the postsynaptic nature of LTD_{GABA}, we favor the idea that activation of D₂ autoreceptors triggers and/or interacts with postsynaptic signaling cascades essential for induction of synaptic plasticity. In fact, compelling evidence suggest a D₂R-triggered activation of calcineurin (i.e., protein phosphatase 2B) mediated by the Inositol 1, 4, 5-Trisphosphate (IP₃) signaling in the striatum (Hernandez-Lopez et al., 2000) that may also underlie LTD_{GABA} in the VTA. Based on this observation in the striatum, our hypothesis predicts that D₂Rs may activate phospholipase Cβ (PLCβ1) through Gβγ signaling, which in turn could trigger IP₃ signaling pathway and therefore, could increase intracellular levels of Ca²⁺. Activation of a Ca²⁺ dependent phosphatase such as calcineurin may eventually lead to dephosphorylation of GABA_A receptors or their associated regulatory subunit as shown previously in a form of LTD of GABAergic transmission in neurons of the deep cerebellar nuclei (Morishita and Sastry, 1996). Further work on the characterization of LTD_{GABA} and other forms of synaptic plasticity in the VTA should




Fig. 3. LTD_{GABA} requires D₂R activation. (a) Sample experiment of LTD_{GABA} and IPSCs recorded from a DA neuron in a bath-applied sulpiride slice. Inset: averaged IPSCs recorded from a DA neuron in a bath-applied sulpiride slice before (black) and 25 min after LFS (red). Calibration for insets: 20 ms, 50 pA. (b) Sulpiride blocked the induction of LTD_{GABA} (control LTD, filled symbols: 63 ± 1% of pre-LFS values, *n* = 8; sulpiride cells, open symbols: 89 ± 4% of pre-LFS values, *n* = 7). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article).

help elucidate the neural substrates of reward-motivated behaviors.

3.5. LFS-induced LTD_{GABA} is modulated by morphine *in vivo*

We have previously shown that 24 h after morphine *in vivo*, LTP_{GABA} is blocked providing a mechanism by which opiates could modulate DA cell excitability and DA release (Nugent et al., 2007). Modulation of LTD_{GABA} by opiates seemed to be very likely,




Fig. 4. *In vivo* exposure to morphine prevents LTD_{GABA} in slices prepared 24 h later. Sample experiments and IPSCs (insets) in slices from a saline-treated rat (a) and a morphine-treated rat (b). (c) Averaged LTD experiments in slices prepared 24 h after either from saline- or morphine-treated rats (saline, filled symbols: $73 \pm 2\%$ of pre-LFS values, $n = 10$; morphine, open symbols: $113 \pm 5\%$ of pre-LFS values, $n = 9$).

therefore we next examined whether 24 h after *in vivo* exposure to morphine, LTD_{GABA} is disrupted. We found that while slices from saline-treated rats exhibited LTD_{GABA} in response to LFS, LTD_{GABA} was absent in slices from morphine-treated rats (Fig. 4a–c, paired *t*-test: $p = 0.042$ for saline and $p = 0.507$ for morphine). This remarkable finding suggests that opiates can regulate a bidirectional inhibitory plasticity in the VTA. It is still not clear and yet to be determined whether morphine blocks or occludes LTD_{GABA}. It is tempting to favor the latter hypothesis where morphine induces LTD_{GABA} and therefore occludes LFS-induced LTD_{GABA}. Analyses of basal PPR and CV of GABA_A IPSCs in slices from saline and morphine animals showed no difference in these measures (data not shown, mean basal PPR values from saline-treated rats: 0.907 ± 0.16 , morphine-treated rats: 0.95 ± 0.28 , *t*-test: $p = 0.528$) which is consistent with our previous finding (Nugent et al., 2007). Based on this data, *in vivo* morphine does not seem to induce any form of presynaptic plasticity (LTP or LTD). However, because LTD_{GABA} is postsynaptic, it is likely that *in vivo* morphine could trigger or block LTD_{GABA}. In a scenario where morphine induces LTD_{GABA}, we assume that an increase in the somatodendritic release of DA by morphine through μ opioid receptors (Chefer et al., 2009) and DA's action on D₂ autoreceptors could trigger the morphine-induced plasticity. As a result, morphine-induced LTD_{GABA} in parallel with morphine-induced blockade of LTP_{GABA} and morphine-induced glutamatergic LTP could work together to over-excite DA neurons, thereby increasing DA release. On the other hand, if morphine prevents LTD_{GABA}, this may occur through a presynaptic inhibition of the somatodendritic release of DA by morphine through κ opioid receptors (Ford et al., 2007). In this situation, morphine-blockade of LTD_{GABA} would serve as a physiological brake to dampen the over-excitability of DA neurons produced by morphine-induced glutamatergic LTP and morphine-blockade of LTP_{GABA}. In sum, our finding of morphine-modulation of LTD_{GABA} is significant and provides an additional novel target for drugs of abuse to mediate their addictive properties.

4. Concluding remarks

Here we report a novel form of postsynaptic LTD (LTD_{GABA}) at GABAergic synapses onto VTA DA neurons that is independent of NMDARs and CB₁Rs but requires D₂R activation. Remarkably, we found that 24 h after a single *in vivo* exposure to morphine, LTD_{GABA} is absent from slices of morphine-treated rats. The induction of LTD at inhibitory GABAergic synapses provides an additional mechanism to promote the induction of synaptic plasticity at adjacent excitatory synapses. Moreover, the existence of LTD_{GABA} in addition to other forms of inhibitory plasticity in the VTA provides a physiological mechanism to reinforce the learning of reward-related behaviors. More importantly, drug-induced modulation of this plasticity could provide a means to promote or counteract maladaptive drug-induced changes at GABAergic synapses. The mechanisms by which morphine and other drugs of abuse block/trigger LTD_{GABA} are our next important questions. Our experiments further characterizing LTD_{GABA} will provide us avenues to pursue. Our results here provide new information on the learning mechanisms of the VTA inhibitory plasticity that is associated with reward and drug-induced reward and present possible novel therapeutic approaches to drug craving in addicts.

Acknowledgements

The opinions and assertions contained herein are the private opinions of the authors and are not to be construed as official or reflecting the views of the Uniformed Services University of the Health Sciences or the Department of Defense or the Government

of the United States. This work was supported by a DOD-intramural grant from the Uniformed Services University (USUHS), and the PhRMA Foundation Research Starter Grant to F.S.N. The funding agencies did not contribute to writing this article or deciding to submit it. Thanks to Drs. Brian Cox and Thomas Cote for their helpful and constructive discussions for the present article.

References

- Backes, E.N., Hemby, S.E., 2008. Contribution of ventral tegmental GABA receptors to cocaine self-administration in rats. *Neurochem. Res.* 33, 459–467.
- Bradberry, C.W., Roth, R.H., 1989. Cocaine increases extracellular dopamine in rat nucleus accumbens and ventral tegmental area as shown by *in vivo* microdialysis. *Neurosci. Lett.* 103, 97–102.
- Campbell, A.D., Kohl, R.R., McBride, W.J., 1996. Serotonin-3 receptor and ethanol-stimulated somatodendritic dopamine release. *Alcohol* 13, 569–574.
- Chefer, V.I., Denoroy, L., Zapata, A., Shippenberg, T.S., 2009. Mu opioid receptor modulation of somatodendritic dopamine overflow: GABAergic and glutamatergic mechanisms. *Eur. J. Neurosci.* 30, 272–278.
- Faber, D.S., Korn, H., 1991. Applicability of the coefficient of variation method for analyzing synaptic plasticity. *Biophys. J.* 60, 1288–1294.
- Ford, C.P., Beckstead, M.J., Williams, J.T., 2007. Kappa opioid inhibition of somatodendritic dopamine inhibitory postsynaptic currents. *J. Neurophysiol.* 97, 883–891.
- Guan, Y.Z., Ye, J.H., 2010. Ethanol blocks long-term potentiation of GABAergic synapses in the ventral tegmental area involving mu-opioid receptors. *Neuropsychopharmacology*.
- Heifets, B.D., Castillo, P.E., 2009. Endocannabinoid signaling and long-term synaptic plasticity. *Annu. Rev. Physiol.* 71, 283–306.
- Hernandez-Lopez, S., Tkatch, T., Perez-Garci, E., Galarraga, E., Bargas, J., Hamm, H., Surmeier, D.J., 2000. D2 dopamine receptors in striatal medium spiny neurons reduce L-type Ca^{2+} currents and excitability via a novel PLC[β]-IP3-calcineurin-signaling cascade. *J. Neurosci.* 20, 8987–8995.
- Jones, S., Kornblum, J.L., Kauer, J.A., 2000. Amphetamine blocks long-term synaptic depression in the ventral tegmental area. *J. Neurosci.* 20, 5575–5580.
- Kasanetz, F., Deroche-Gamonet, V., Berson, N., Balado, E., Lafourcade, M., Manzoni, O., Piazza, P.V., 2010. Transition to addiction is associated with a persistent impairment in synaptic plasticity. *Science* 328, 1709–1712.
- Kauer, J.A., Malenka, R.C., 2007. Synaptic plasticity and addiction. *Nat. Rev. Neurosci.* 8, 844–858.
- Klitenick, M.A., DeWitte, P., Kalivas, P.W., 1992. Regulation of somatodendritic dopamine release in the ventral tegmental area by opioids and GABA: an *in vivo* microdialysis study. *J. Neurosci.* 12, 2623–2632.
- Komatsu, Y., Iwakiri, M., 1993. Long-term modification of inhibitory synaptic transmission in developing visual cortex. *Neuroreport* 4, 907–910.
- Lee, D.Y., Guttilla, M., Fung, K.D., McFeron, S., Yan, J., Ranaldi, R., 2007. Rostral-caudal differences in the effects of intra-VTA muscimol on cocaine self-administration. *Pharmacol. Biochem. Behav.* 86, 542–549.
- Lu, Y.M., Mansuy, I.M., Kandel, E.R., Roder, J., 2000. Calcineurin-mediated LTD of GABAergic inhibition underlies the increased excitability of CA1 neurons associated with LTP. *Neuron* 26, 197–205.
- Margolis, E.B., Lock, H., Hjelmstad, G.O., Fields, H.L., 2006. The ventral tegmental area revisited: is there an electrophysiological marker for dopaminergic neurons? *J. Physiol.* 577, 907–924.
- McLean, H.A., Caillard, O., Ben-Ari, Y., Gaiarsa, J.L., 1996. Bidirectional plasticity expressed by GABAergic synapses in the neonatal rat hippocampus. *J. Physiol.* 496 (Pt 2), 471–477.
- Morishita, W., Sastry, B.R., 1993. Long-term depression of IPSPs in rat deep cerebellar nuclei. *Neuroreport* 4, 719–722.
- Morishita, W., Sastry, B.R., 1996. Postsynaptic mechanisms underlying long-term depression of GABAergic transmission in neurons of the deep cerebellar nuclei. *J. Neurophysiol.* 76, 59–68.
- Niehaus, J.L., Murali, M., Kauer, J.A., 2010. Drugs of abuse and stress impair LTP at inhibitory synapses in the ventral tegmental area. *Eur. J. Neurosci.* 32, 108–117.
- Nugent, F.S., Kauer, J.A., 2008. LTP of GABAergic synapses in the ventral tegmental area and beyond. *J. Physiol.* 586, 1487–1493.
- Nugent, F.S., Penick, E.C., Kauer, J.A., 2007. Opioids block long-term potentiation of inhibitory synapses. *Nature* 446, 1086–1090.
- Nugent, F.S., Niehaus, J.L., Kauer, J.A., 2009. PKG and PKA signaling in LTP at GABAergic synapses. *Neuropsychopharmacology* 34, 1829–1842.
- Pan, B., Hillard, C.J., Liu, Q.S., 2008a. D2 dopamine receptor activation facilitates endocannabinoid-mediated long-term synaptic depression of GABAergic synaptic transmission in midbrain dopamine neurons via cAMP-protein kinase A signaling. *J. Neurosci.* 28, 14018–14030.
- Pan, B., Hillard, C.J., Liu, Q.S., 2008b. Endocannabinoid signaling mediates cocaine-induced inhibitory synaptic plasticity in midbrain dopamine neurons. *J. Neurosci.* 28, 1385–1397.
- Saal, D., Dong, Y., Bonci, A., Malenka, R.C., 2003. Drugs of abuse and stress trigger a common synaptic adaptation in dopamine neurons. *Neuron* 37, 577–582.
- Szabo, B., Siemes, S., Wallmichrath, I., 2002. Inhibition of GABAergic neurotransmission in the ventral tegmental area by cannabinoids. *Eur. J. Neurosci.* 15, 2057–2061.
- Thomas, M.J., Malenka, R.C., Bonci, A., 2000. Modulation of long-term depression by dopamine in the mesolimbic system. *J. Neurosci.* 20, 5581–5586.
- Ungless, M.A., Whistler, J.L., Malenka, R.C., Bonci, A., 2001. Single cocaine exposure *in vivo* induces long-term potentiation in dopamine neurons. *Nature* 411, 583–587.
- Volkow, N.D., Fowler, J.S., Wang, G.J., 2004. The addicted human brain viewed in the light of imaging studies: brain circuits and treatment strategies. *Neuropharmacology* 47 (Suppl. 1), 3–13.
- Wang, J.H., Stelzer, A., 1996. Shared calcium signaling pathways in the induction of long-term potentiation and synaptic disinhibition in CA1 pyramidal cell dendrites. *J. Neurophysiol.* 75, 1687–1702.
- Xi, Z.X., Stein, E.A., 2000. Increased mesolimbic GABA concentration blocks heroin self-administration in the rat. *J. Pharmacol. Exp. Ther.* 294, 613–619.
- Zhang, T.A., Placzek, A.N., Dani, J.A., 2010. *In vitro* identification and electrophysiological characterization of dopamine neurons in the ventral tegmental area. *Neuropharmacology*.