

University of Nebraska - Lincoln

## DigitalCommons@University of Nebraska - Lincoln

---

Publications, Agencies and Staff of the U.S.  
Department of Commerce

U.S. Department of Commerce

---

2012

### Starvation causes disturbance in amino acid and fatty acid metabolism in *Diporeia*

Suman Maity

Purdue University, [smaity@purdue.edu](mailto:smaity@purdue.edu)

Amber Jannasch

Purdue University, [hopfas@purdue.edu](mailto:hopfas@purdue.edu)

Jiri Adamec

University of Nebraska-Lincoln, [jadamec2@unl.edu](mailto:jadamec2@unl.edu)

Thomas Nalepa

Great Lakes Environmental Research Laboratory, [thomas.nalepa@noaa.gov](mailto:thomas.nalepa@noaa.gov)

Tomas O. Höök

Purdue University, [thook@purdue.edu](mailto:thook@purdue.edu)

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/usdeptcommercepub>

 Part of the [Environmental Sciences Commons](#)

---

Maity, Suman; Jannasch, Amber; Adamec, Jiri; Nalepa, Thomas; Höök, Tomas O.; and Sepúlveda, Maria S., "Starvation causes disturbance in amino acid and fatty acid metabolism in *Diporeia*" (2012). *Publications, Agencies and Staff of the U.S. Department of Commerce*. 347.  
<https://digitalcommons.unl.edu/usdeptcommercepub/347>

This Article is brought to you for free and open access by the U.S. Department of Commerce at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications, Agencies and Staff of the U.S. Department of Commerce by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Suman Maity, Amber Jannasch, Jiri Adamec, Thomas Nalepa, Tomas O. Höök, and Maria S. Sepúlveda



## Starvation causes disturbance in amino acid and fatty acid metabolism in *Diporeia*

Suman Maity<sup>a</sup>, Amber Jannasch<sup>b</sup>, Jiri Adamec<sup>c</sup>, Thomas Nalepa<sup>d</sup>, Tomas O. Höök<sup>a</sup>, Maria S. Sepúlveda<sup>a,\*</sup>

<sup>a</sup> Department of Forestry and Natural Resources, Purdue University, 195 Marsteller St., West Lafayette, IN, 47907, United States

<sup>b</sup> Bindley Bioscience Center at Discovery Park, Purdue University, 1203 W. State St. West Lafayette, IN 47907, United States

<sup>c</sup> Department of Biochemistry, Beadle Center N151, University of Nebraska, 1901 Vine Street, Lincoln, NE, 68588, United States

<sup>d</sup> National Oceanic and Atmospheric Administration (NOAA), Great Lakes Environmental Research Laboratory, 4840 S. State Road, Ann Arbor, MI, 48108, United States

### ARTICLE INFO

#### Article history:

Received 6 October 2011

Received in revised form 22 December 2011

Accepted 29 December 2011

Available online 13 January 2012

#### Keywords:

Metabolomics

Two dimensional gas chromatography

Mass spectrometry

Amphipod

Great Lakes

### ABSTRACT

The benthic amphipod *Diporeia* spp. was once the predominant macroinvertebrate in deep, offshore regions of the Laurentian Great Lakes. However, since the early 1990s, *Diporeia* populations have steadily declined across the area. It has been hypothesized that this decline is due to starvation from increasing competition for food with invasive dreissenid mussels. In order to gain a better understanding of the changes in *Diporeia* physiology during starvation, we applied two-dimensional gas chromatography coupled with time of flight mass spectrometry (GCXGC/TOF-MS) for investigating the responses in *Diporeia* metabolome during starvation. We starved *Diporeia* for 60 days and collected five organisms every 12 days for metabolome analyses. Upon arrival to the laboratory, organisms were flash frozen and served as control (day 0). We observed an increase in lipid oxidation and protein catabolism with subsequent declines of essential amino acids (proline, glutamine, and phenylalanine), down-regulation of glycerophospholipid and sphingolipid metabolism, and decreased polyunsaturated fatty acid abundance in nutritionally stressed *Diporeia*. Abundance of 1-Iodo-2-methylundecane, a metabolite closely related to insect pheromones, also declined with starvation. This research has further substantiated the applicability of GCXGC/TOF-MS as a research tool in the field of environmental metabolomics. The next step is to apply this new knowledge for evaluating nutritional status of feral *Diporeia* to elucidate the underlying cause(s) responsible for their decline in the Great Lakes.

© 2012 Elsevier Inc. All rights reserved.

### 1. Introduction

*Diporeia* spp., a conglomerate of several poorly defined species, and glacial relict and freshwater amphipod, used to be the predominant benthic invertebrate in deep, offshore regions of the Laurentian Great Lakes (Cook and Johnson, 1974). The high lipid content in *Diporeia* makes it an ideal food resource for a number of fish species like lake whitefish (*Coregonus clupeaformis*), deepwater sculpin (*Myoxocephalus thompsoni*), and alewife (*Alosa pseudoharengus*). However, *Diporeia* populations have experienced precipitous declines throughout the Laurentian Great Lakes since the 1990s (Nalepa, 1998; Dermott, 2001; Lozano et al., 2001). The downfall in *Diporeia* numbers has been linked to the introduction and establishment of dreissenid mussels into the Great Lakes (Nalepa et al., 2005). Filter feeding activity by dreissenids may have removed substantial amounts of settling organic material (mainly diatoms) from the water column before reaching the lake bottom. Since *Diporeia* are

benthic feeders that inhabit the upper few centimeters of sediment (Marzolf, 1965; Johnson, 1987), it seems plausible that dreissenids are outcompeting *Diporeia* for the limited supply of available food. Lack of sufficient food could negatively impact *Diporeia*'s survival as was the case with *Monoporeia affinis*, a related amphipod species from the Baltic Sea (Wiklund et al., 2008).

Little information is available on the physiological response of amphipods during starvation. One study found no significant changes in total lipid concentration in *Diporeia* starved for a month (Gauvin et al., 1989), whereas another study reported that *M. affinis* switched to lipid-based energy metabolism during starvation (Lehtonen, 1994). Since one of the plausible causes of *Diporeia* declines is starvation driven by food competition with dreissenid mussels, research aimed at studying impacts of prolonged starvation in *Diporeia* would be useful in providing insights on how persistent food deprivation can affect biological functions in *Diporeia*. Using a metabolomics approach, we previously reported on impacts in non-polar metabolites (histidine, glycerolipids, and sphingolipids) during starvation in this amphipod (Maity et al., 2012). The present study complements the previous one by reporting on changes in polar metabolites in starved *Diporeia*.

Metabolomics is the qualitative and quantitative study of metabolites present at any given time within a cell, organ, or organism. It is a "discovery" driven holistic evaluation of changes in the expression

\* Corresponding author at: Department of Forestry and Natural Resources and School of Civil Engineering, Purdue University, 195 Marsteller St., West Lafayette, IN, 47907, United States. Tel.: +1 765 496 3428; fax: +1 765 496 2422.

E-mail addresses: [smaity@purdue.edu](mailto:smaity@purdue.edu) (S. Maity), [hopfas@purdue.edu](mailto:hopfas@purdue.edu) (A. Jannasch), [jadamec2@unl.edu](mailto:jadamec2@unl.edu) (J. Adamec), [thomas.nalepa@noaa.gov](mailto:thomas.nalepa@noaa.gov) (T. Nalepa), [thook@purdue.edu](mailto:thook@purdue.edu) (T.O. Höök), [msssepulv@purdue.edu](mailto:msssepulv@purdue.edu) (M.S. Sepúlveda).

patterns of hundreds of metabolites after exposure to environmental stimuli (Nicholson et al., 2002). Compared to traditional approaches, metabolomics deals with a lower number (hundreds vs. thousands) of molecules (Wishart, 2005) which can result in improved interpretation of biological data (Bino et al., 2004; Dunn and Ellis, 2005). The application of metabolomics to understand environmental problems is a relatively new field and expanding rapidly (Viant, 2008). Environmental metabolomics has been successfully applied to study the impacts of pollutants and pathogens in fish (Samuelsson and Larsson, 2008) and terrestrial and aquatic invertebrates (Lin et al., 2006; Bundy et al., 2009; Schock et al., 2010), including *Diporeia* (Ralston Hooper et al., 2011). Multiple technological platforms have been developed over the past few years for metabolomics research, including Nuclear Magnetic Resonance (NMR), Mass Spectrometry (MS)-based Gas Chromatography (GC), and Liquid Chromatography (LC) coupled with MS (Sepúlveda et al., 2011). Two-dimensional gas chromatography based mass spectrometry (GCXGC/TOF-MS) has been used in conjunction with NMR or LC-MS to provide a more inclusive picture of metabolome changes. Although GCXGC/TOF-MS has been implemented to collect metabolic profiles from plants (Pongsuwan et al., 2008), microbes (Garcia et al., 2008), yeast (Mohler et al., 2006), and mice (Shellie et al., 2005), this metabolomics platform has been used considerably less compared to the more classic approaches (e.g., NMR).

In this paper, we describe the application of GCXGC/TOF-MS-based metabolomics to investigate the physiological responses of *Diporeia* during starvation. We hypothesize that prolonged starvation will result in disruption of lipid metabolism and progressive protein catabolism for energy production. In addition, we attempt to develop a “starvation metabolic profile” which may have a practical application for health assessment of declining *Diporeia* populations in the Laurentian Great Lakes.

## 2. Materials and methods

### 2.1. Study animals

In summer 2008, we collected live *Diporeia* from southern Lake Michigan (site C-5, 42° 49.00' N, 86° 50.00' W) at a depth of 157 m during a field cruise onboard National Oceanic and Atmospheric Administration's (NOAA) R/V “The Laurentian”. Using ponar grabs, the top layer of the sediment was collected and sieved through a handheld mesh (0.5 mm) for the collection of live *Diporeia*. Animals were kept inside Nalgene® 1 L bottles filled with pre-chilled (4 °C) lake water. Bottles were placed inside coolers on wet ice at 4 °C and brought to the laboratory where animals were allowed to acclimate for 48 h prior to the initiation of the experiments.

### 2.2. Study design

Before the start of the starvation study, a group of live animals (> 4 mm) were flash frozen in liquid nitrogen and served as controls (“Day 0”). An ideal starvation experimental design would have also included a “positive control”, i.e., a group that was fed. The most commonly used diet for maintaining *Diporeia* is “Tetramin” (fish flakes). However, no studies have tested the quality of this diet nor have developed a more species-specific diet (in our case, it would consist mostly of diatoms). Since this study was conducted, we have been successful at feeding them lyophilized diatoms (Maity et al. in review), but additional studies are needed in order to determine more specific nutrient requirements for this amphipod. The remaining animals were randomly assigned to one of six replicates (n = 10/replicate). Each replicate consisted of a 1 L Pyrex® glass beaker with 50 g (~2 cm deep) autoclaved Lake Michigan sediment (collected from C-5) and 700 ml of reconstituted modified hard water (RMHW) (Ralston-Hooper et al., 2008). To simulate lake bottom

conditions, the experiment was set-up inside a walk-in cooler with an ambient temperature of 4 °C and complete absence of light. No food was provided during the entire duration (60 d) of the experiment. At regular intervals of 12 d, a subset of live specimens (4–6 total organisms per condition, one from each replicate and time period) was collected, flash frozen in liquid nitrogen and preserved at –80 °C for later metabolomic analyses.

### 2.3. Sample preparation

Specimens were processed for metabolomic analyses following a modified protocol from Ralston-Hooper et al. (2008) as described below. Each biological replicate was composed of a single *Diporeia* (0.0167 g – 0.0055 g). All reagents used were of analytical grade quality and were purchased from either Sigma Aldrich (St. Louis, MO, USA) or Regis Technologies (Morton Grove, IL, USA). Samples were prepared for metabolomics analyses as they thawed with the complete process, from retrieval of sample from sampling vial to homogenization, taking ~2 min, and carried while sample was kept on ice. Samples were homogenized for 20 s in pre-chilled (4 °C) methanol (300 µL) and MQ grade water (150 µL) using a 7 mm × 95 mm saw tooth stainless steel tissue homogenizer probe (Omni International, Marietta, GA, USA). Next, the homogenate was placed in a sonicator bath for 3 min to ensure uniform mixing. The solution was vortexed for 2 min after adding pre-chilled chloroform (450 µL) and then it was placed on wet ice for an additional 10 min. The chilled solution was then centrifuged at 3000 × g for 20 min to extract polar (methanol: water) and non-polar (chloroform) metabolites respectively. Each phase was pipetted into a new vial and was allowed to dry out at 45 °C using a Savant SPD 131DDA SpeedVac concentrator (Thermo Electron Corporation Milford, MA, USA). The chloroform phase was analyzed using LC/MS-TOF and the results of these analyses have been published elsewhere (Maity et al., 2012). The dried methanol: water extracted sample pellet (polar metabolites) was further derivatized using the following steps. First, methoxyamine hydrochloride (20 mg) was dissolved in 1 mL anhydrous pyridine and 30 µL of this solution was added to each pellet and mixed for 30 min at 60 °C. Next, 45 µL of N-methyl-N-(trimethylsilyl) trifluoroacetamide (MSTFA) was added and mixed for 1 h at 60 °C. The derivatized sample was transferred to an auto-sampler vial for GCXGC/TOF-MS analysis.

### 2.4. Instrumental conditions

Following derivatization, polar samples were analyzed on a Pegasus III GCXGC/TOF-MS (Leco Corporation, St. Joseph, MI, USA). Samples were injected (2 µL) onto a two dimensional GC column with an inlet split-mode ratio of 20 and pre-set flow rate (1.5 mL/min) of carrier gas (helium). The temperature in the inlet port was 280 °C. The dimensions of the two columns were as follows: first column, Restek Rtx-225 matrix (0.25 mm × 30 m, 0.25 µm), and second column, Agilent HP-5 matrix (0.32 mm × 2 m, 0.25 µm). The linear thermal gradient of the first column was between 50 °C and 240 °C (7 °C/min) followed by a 240 °C hold for 5 min. The temperature of the second column was set as the same gradient with an additional offset of +50 °C. The TOF-MS analysis utilized electron impact ionization with an ion source temperature of 200 °C under the following conditions: detector voltage 1700 V, electron energy –70 V, spectrum acquisition rate of 100 spectra/s, and a detectable mass range of 30–800 Da. An acquisition delay of 150 s was applied at the start of the run.

### 2.5. Data processing and statistical analysis

Initial mass spectra were collected and processed in ChromaTOF software (version 3.32) from Leco Corporation followed by analyses

using MSort©, an in-house software developed at the Bindley Biosciences Center at Purdue University (Oh et al., 2008). Since the same peak can generate multiple entries across different chromatograms, peaks were merged into a single entry and then aligned using retention time (RT) and similarity between fragmentation spectrums of merged peaks across different samples. The window size for the first and second RT dimensions were 1% and 5%, respectively, and a minimum correlational value of 0.95 was set as threshold criteria. Metabolites were identified by comparing each spectrum against the National Institute of Standards and Technology (NIST) database of GC–MS spectra (<http://www.nist.gov/srd/nist1.cfm>). Each individual peak was compared against the library and assigned a similarity value (SV, a value of 1000 denotes a complete match). For any given peak, we set up a lower threshold of  $\geq 750$  as potential match (Cristoni et al., 2009). Constant mean normalized data from GCXGC/TOF-MS was subjected to multivariate analyses using R® statistical software (version 2.9.2, R Foundation for Statistical Computing, Vienna, Austria). Missing values are a common challenge in “omics” datasets (Chich et al., 2007; Karpievitch et al., 2010). For our dataset, missing values were replaced by the group average if present in at least half of the total samples per group, otherwise missing values were replaced by 1 (Colinge et al., 2005). Principal Component Analysis (PCA) was performed including the complete metabolome data (not just the statistical significant ones) to highlight any underlying structure in the data. Metabolite abundance was compared across treatments using ANOVA followed by Tukey’s multiple comparison.

### 3. Results and discussion

Data from D36 are not shown since organisms were lost during the sampling processing. Overall, mortality was less than 75%. The results from the PCA revealed a distinct separation of metabolite expression patterns across starvation groups (Fig. 1). Regardless of the length of starvation (D12–D60), pre-starved animals (D0) were distantly grouped compared to the starved ones. The first three principal components helped explain more than half of the total variation (Fig. 2).

A total of eight metabolites were significantly changed during starvation (Table 1). These included L-proline, glutamine, L-threonine, L-phenylalanine, N-acetyl glucosamine, hexadecanoic acid, trans-9-octadecenoic acid, and 1-Iodo-2-methylundecane.




Fig. 1. Temporal changes in the metabolic profile of *Diporeia* under starvation. D = days in starvation.




Fig. 2. Bar graph showing proportion of variation explained by each principal component (PC). The first three PC explained over 50% of the total variation in the data.

Fig. 3 presents the results of the multiple comparison tests for these metabolites. Each horizontal bar represents the difference between two starvation groups with 95% confidence interval. If the interval excludes 0, then the difference is considered significant for that pairwise comparison. For example, abundance of glutamine and hexadecanoic acid (Fig. 3b and f) differed significantly in all starvation periods compared to Day 0 (in this case both increased, see Table 1).

Based on the significant metabolites detected using GCXGC/TOF-MS (present study) and LC-MS (Maity et al., 2012) a “starvation metabolic pathway map” was constructed using canonical maps available from the Kyoto Encyclopedia of Genes and Genomes (<http://www.genome.jp/kegg/>) (Fig. 4).

We have provided a detailed picture of changes in the metabolome of *Diporeia* during starvation. As would be expected, several metabolic pathways were impaired, including metabolism of amino acids, amino sugars, fatty acids, and complex lipids (Fig. 4). From our previous study on metabolome changes in *Diporeia* using LC-MS (targeting non-polar metabolites, Maity et al., 2012), we further observed alterations in glycerophospholipid metabolism. A down-regulation of glycerophospholipid and sphingolipid metabolism contributed to the disruption of the glycine-serine-threonine pathway

**Table 1**List of significantly altered ( $p < 0.05$ ) polar metabolites detected via GC×GC/TOF-MS in starved *Diporeia*.


Metabolite ID <sup>a</sup>	Peak ID <sup>b</sup>	SV <sup>c</sup>	Mean expression level <sup>d</sup> ( $\pm$ SE)				
			Day 0	Day 12	Day 24	Day 48	Day 60
<b>Amino acids</b>							
L-proline	315	815	16.00 $\pm$ 0.05	15.47 $\pm$ 0.19	15.05 $\pm$ 0.19	14.65 $\pm$ 0.17	15.51 $\pm$ 0.25
Glutamine	1063	819	4.52 $\pm$ 3.52	14.36 $\pm$ 0.14	14.00 $\pm$ 0.24	13.95 $\pm$ 0.09	14.06 $\pm$ 0.01
L-threonine	149	781	14.71 $\pm$ 0.15	14.62 $\pm$ 0.31	14.41 $\pm$ 0.14	BLD <sup>e</sup>	14.74 $\pm$ 0.11
L-phenylalanine	300	827	15.19 $\pm$ 0.05	14.52 $\pm$ 0.52	6.63 $\pm$ 3.16	12.95 $\pm$ 0.09	14.65 $\pm$ 0.07
<b>Amino sugars</b>							
N-acetylglucosamine	399	763	13.32 $\pm$ 0.29	14.23 $\pm$ 0.06	6.08 $\pm$ 2.84	4.21 $\pm$ 2.62	13.67 $\pm$ 0.10
<b>Fatty acids</b>							
Hexadecanoic acid	267	810	2.77 $\pm$ 1.77	8.62 $\pm$ 0.14	8.59 $\pm$ 0.13	9.12 $\pm$ 0.19	8.97 $\pm$ 0.24
Trans-9-octadecenoic acid	307	859	10.61 $\pm$ 0.06	10.72 $\pm$ 0.14	10.21 $\pm$ 0.14	10.04 $\pm$ 0.10	9.97 $\pm$ 0.09
<b>Pheromone</b>							
1-Iodo-2-methylundecane	123	827	4.45 $\pm$ 3.45	15.58 $\pm$ 0.04	3.98 $\pm$ 2.72	15.30 $\pm$ 0.02	6.99 $\pm$ 3.35

<sup>a</sup> Identification based on mass spectra match against National Institute of Standards and Technology (NIST) library.<sup>b</sup> Based on the alignment results of metabolic peaks across different samples.<sup>c</sup> Similarity value (SV) > 750 was considered a positive match.<sup>d</sup> Mean expression (log-transformed).<sup>e</sup> Below detection limit starvation groups.

in starved *Diporeia*. Furthermore, the metabolism of several other amino acids, including methionine, lysine, histidine and glutamate, were also negatively impacted. In addition, starvation induced lipid oxidation and a concomitant decrease in FA biosynthesis. All together, these results suggest that starving *Diporeia* switch to protein catabolism after lipid reserves have been exhausted.

Concentrations of several amino acids changed during starvation in a time-specific manner (Table 1 and Fig. 3). For instance, the amino-acid L-proline declined during starvation (0–48 d, Fig. 3a).

The role of proline as a fuel source has been reported for a number of invertebrates (Scaraffia and Wells, 2003; Giulivi et al., 2008; Laparie et al., 2011). In a study conducted with the larval dragonfly *Aeshna cyanea*, proline along with glutamine, was found to be one of the main contributors of the free amino acid reserves used to meet energy demands during starvation (Herzog and Liappis, 1987). In *Aeshna* larvae, proline levels dropped immediately after a short starvation period (10 d) followed by an eventual rise after 30 d of starvation (Herzog and Liappis, 1987). These authors proposed that



**Fig. 3.** Results of post hoc Tukey tests for group-wise comparisons of significant metabolites. Each horizontal bar represents the difference between two starvation groups with 95% confidence interval. If the interval excludes 0, then the difference is considered significant for that pair-wise comparison.







Fig. 3 (continued).

the larvae switched to protein-based metabolism after exhausting their carbohydrate reserves (Herzog and Liappis, 1987). We also observed a decrease in proline levels starting soon after the onset of starvation with a rise 60 d later. Proline is important for many other biological functions besides energy production. The presence of proline within the peptide chain regulates structural susceptibility to protease activity which in turn controls a multitude of biological functions (Yaron et al., 1993). Proline rich motifs are highly favorable as binding sites for modular domains (e.g., SH3, WW, EVH, WASP) which are important for intracellular communication. These proline binding domains form part of multi-domain molecules responsible for controlling cell growth, transcription rates, postsynaptic signaling, and cytoskeleton modifications. In relation to the latter, proline plays a crucial role in profilin function and actin polymerization which controls cytoskeletal movement. Proline rich motifs also control the degree of enzyme isomerization as well as offer phosphorylation and dephosphorylation reaction sites for proteins (Kay et al., 2000; Zarrinpar et al., 2003). Thus, disruptions in proline metabolism during starvation are likely to affect these processes in *Diporeia*. It is also important to note that alanine is the main precursor for proline synthesis in insects (Gade and Auerswald, 2002; Lorenz and Gade, 2009; Arrese and Soulages, 2010) and any changes in proline concentration are usually correlated with disruptions in alanine metabolism (Herzog and Liappis, 1987). Alanine is also an important osmolyte in crustaceans, essential for maintaining cellular ion balance (Abe et al., 1999). Thus, altered alanine metabolism could also result in an ionic imbalance in *Diporeia*.

The abundance of glutamine increased in starving *Diporeia* (Table 1, Fig. 3b). This amino acid is important as a stress response indicator (Tanguy et al., 2005; Hong et al., 2009; Leroy et al., 2010) and plays a crucial role in managing oxidative stress (Michaelis, 1998; Liu

et al., 2000; Matés et al., 2002). This is because glutamine is a major constituent of the glutathione biosynthetic pathway and glutathione is key a reducing agent against reactive oxygen species (ROS) protecting the cell from oxidative damage and apoptotic cell death. The oxidation of the prolyl residues produces the glutamyl residues in the cell (Farr and Kogoma, 1991). Thus as a result of increased protein catabolism, more glutamine is accumulated which might disrupt glutathione synthesis and increase the risk of cell death. Glutamine is also crucial for the synthesis of a number of amino acids like proline, alanine and glycine (Fujimori and Abe, 2002). Both proline and glutamine are linked with arginine metabolism which in turn is related to alanine-aspartate-glutamate metabolism. These metabolic pathways are often looped together and regulate each other in an intricate fashion. In crustaceans, elevated glutamine levels are associated with enhanced nitrogen detoxification from protein metabolism (Chen and Chen, 2000).

L-phenylalanine is an important component of tyrosine metabolism and a neurotransmitter precursor (Meyers, 2000; Van Ruitenbeek et al., 2009). Additionally, in marine mussels, phenylalanine is one of the constituents of a protein complex which acts like metallothionein by binding to heavy metal ions (Roesjyadi, 1981). Essential amino acids in crustaceans (valine, (iso)leucine, lysine, histidine, phenylalanine, threonine, arginine, methionine and tryptophan) are usually found in low abundance (Augusto et al., 2007). Of all these, phenylalanine is the most readily utilized amino acid in crustaceans. In the amphipod, *Echinogammarus stammeri* (freshwater scud), phenylalanine is an important controlling factor in the leucine transport pathway (Berra et al., 2006). The exact physiological role of phenylalanine in *Diporeia* is yet to be investigated but it is likely that phenylalanine can perform similar functions in neurotransmitter biosynthesis, regulate leucine uptake kinetics and/or



**Fig. 4.** Metabolic pathways impacted during starvation constructed using canonical biochemical pathways from the Kyoto Encyclopedia of Genes and Genomes (KEGG, (<http://www.genome.jp/kegg/>)). Included are both polar and non-polar metabolites detected by GCXGC/TOF-MS and LC-MS (from Maity et al., 2012). DG = diacylglycerols; DPA = docosapentaenoic acid; EPA = eicosapentaenoic acid; LsoPC = lysophosphatidylcholine; PC = phosphatidylcholine; PG = phosphatidylglycerol; PI = phosphatidylinositol; TG = triacylglycerols; Thr = threonine.

impact tyrosine and phenylalanine pathways. Both proline and phenylalanine increased in starved *Diporeia* after 60 d of starvation. This “approximation” to baseline values might be an indication that “acclimation” or “adaptation” has taken place. Therefore, it would be of interest to follow metabolomic changes during starvation on these amphipods for longer time periods in order to capture potential “adaptive” responses to starvation. A potential “adaptive” response of *Diporeia* to a stressor such as starvation could be of great significance when considering the uncertain future of this organism in the Great Lakes. However, amino acid concentrations in crustaceans are quite variable, being influenced by several factors besides starvation, including stress and temperature (Graney and Giesy, 1986). Therefore, we believe our results should be further validated before any major conclusion is made.

Another biologically relevant metabolite identified was N-acetylglucosamine, an amino sugar that plays a significant role in both inter- and intra-cellular interactions. It is a component of the glycocalyx, a cell surface component that participates in cell signaling. The charge distribution on these molecules as well as their flexibility to link up with one another has resulted in an excellent control of signal transduction. The ability of carbohydrate-based molecules (amino sugars) to maintain intercellular bonds and act as signal transduction mediators has been demonstrated in various taxa (Sharon, 2007; Netea et al., 2008; Sahly et al., 2008; Zhao et al., 2008; Dennis et al., 2009). N-acetylglucosamine is also found within the nuclear pore complex highlighting its importance for intracellular functions (Hanover, 2001). In most animals studied, N-acetylglucosamine is used to modify serine or threonine residues within the nuclear or cellular protein domains (Hart et al., 2007). Thus, N-acetylglucosamine can modify transcription factors and control protein trafficking. It can also act as regulator of cellular growth and proliferation in early

stages of embryonic development (Hart et al., 2007). In invertebrates, amino sugars are also thought to play a key role in differentiating between self and non-self-recognition protecting host cells from attacks by their own immune systems (Rosenstiel et al., 2009; Vazquez et al., 2009) and are important for the synthesis of chitin, an important component of invertebrate exoskeleton (Cauchie, 2002). Therefore, long-term starvation has the potential to cause perturbed amino sugar metabolism in *Diporeia* which can lead to alterations in immune system functioning, reduced cellular integrity and intercellular communication and compromise the physical protection of their exoskeleton by disrupting cuticle production.

An interesting outcome of our studies has been the characterization of amino acid declines during starvation. Amino acids are essential for several functions including energy production, osmoregulation and muscle growth (Carefoot et al., 1992; Rosas et al., 2001; Huong et al., 2010; Karanova and Andreev, 2010). Our studies are in accordance with similar findings which found decreased levels of amino acids in crustaceans in the wild possibly due to starvation-like conditions (Rosa and Nunes, 2003, 2005; Mente et al., 2010). Thus, it can be concluded that in general, starved crustaceans respond with increased degradation of proteins and decrease in protein synthesis (Helland et al., 2000; Mente et al., 2002).

Concentrations of fatty acids such as hexadecanoic and trans-9-octadecanoic acid also changed during starvation, but in opposite trends (Table 1, Fig. 3f and g). Compounds derived from hexadecanoic acid have been reported to perform a range of functions in insects. For example, both hexadecanoic acid and 3-oxo-(Z)-9-hexadecanal have been found in chemical secretions from termites (Prestwich and Collins, 1982; Chen et al., 1999), while triacontanyl hexadecanoate is important for cuticle formation in corn earworm (*Helicoverpa zea*) pupae (Buckner et al., 1996). Concentrations of hexadecanoic acid



could be used as an indicator of progressive use of glycerolipids for energy production in starving *Diporeia*. Trans-9-octadecenoic acid is a breakdown product of linoleic acid metabolism and concentrations decrease during stress (Gardner, 1979, 1995), including exposure to contaminants (Jones et al., 2008). These results are in agreement with our findings and would suggest an increase in lipid metabolism during starvation.

An interesting finding was the detection of 1-Iodo-2-methylundecane (1I2MU). As far as we know, this is the first time that 1I2MU has been reported from crustaceans. This metabolite is believed to function as a sex pheromone and its production is estrogen-dependent (Achiraman et al., 2010a). A similar volatile compound, 1-iodoundecane, has been reported from cows during estrus (Achiraman et al., 2010b). Levels of this metabolite fluctuated widely during starvation with elevated levels in d12 and d48 while dropping in all other groups (Table 1).

In sum, our results have shown that nutritionally stressed *Diporeia* respond with an increase in lipid oxidation and protein catabolism (denoted by an increase in glutamine); declines of essential amino acids (proline, threonine, and phenylalanine); down-regulation of glycerophospholipid and sphingolipid metabolism; and a decrease in PUFA abundance (Fig. 4). Abundance of 1-Iodo-2-methylundecane, a metabolite closely related to insect pheromones, also decreased during starvation (24 d). Declines of these metabolites could impair reproductive function, growth, and ultimately long-term survival. These metabolite changes also result in a decrease of *Diporeia*'s nutritional quality potentially impacting the health of fish that consume them. This research has further substantiated the applicability of GCXGC/TOF-MS as a research tool in the field of environmental metabolomics. The next step is to use this new knowledge for evaluating nutritional status of feral *Diporeia* and determine the underlying cause(s) responsible for their decline in the Great Lakes.

## Acknowledgements

This research was supported by the Great Lakes Fisheries Trust (Grant # 2008.886). We would also like to thank the crews of "The Laurentian" for their help during sample collection; Kimberly Ralston-Hooper for providing valuable input for the experimental set-up during its initial phase; Michael Gribskov for providing assistance with the statistical analyses; and Nicholas McCann and Payel Acharya for providing thoughtful suggestions for improving the manuscript.

## References

- Abe, H., Okuma, E., Amano, H., Noda, H., Watanabe, K., 1999. Role of free- and -alanine in the Japanese mitten crab *Eriocheir japonicus* to intracellular osmoregulation during downstream spawning migration. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 123, 55–59.
- Achiraman, S., Archunan, G., Ponmanickam, P., Rameshkumar, K., Kannan, S., John, G., 2010a. 1-Iodo-2 methylundecane [1I2MU]: An estrogen-dependent urinary sex pheromone of female mice. *Theriogenology* 74, 345–353.
- Achiraman, S., Ponmanickam, P., Ganesh, D., Archunan, G., 2010b. Detection of estrus by male mice: synergistic role of olfactory-vomeroneasal system. *Neurosci. Lett.* 477, 144–148.
- Arrese, E.L., Soulages, J.L., 2010. Insect fat body: energy, metabolism, and regulation. *Annu. Rev. Entomol.* 55, 207–225.
- Augusto, A., Greene, L.J., Laure, H.J., McNamara, J.C., 2007. The ontogeny of isosmotic intracellular regulation in the diadromous, freshwater palaemonid shrimps, *Macrobrachium amazonicum* and *M. olfersi* (Decapoda). *J. Crustacean Biol.* 27, 626–634.
- Berra, E., Forcella, M., Giacchini, R., Parenti, P., 2006. Leucine transport across plasmamembranes from the scud *Echinogammarus stammeri* (Amphipoda: Gammaridae). *Int. J. Limnol.* 42, 79–85.
- Bino, R.J., Hall, R.D., Fiehn, O., Kopka, J., Saito, K., Draper, J., Nikolau, B.J., Mendes, P., Roessner-Tunali, U., Beale, M.H., Trethewey, R.N., Lange, B.M., Wurtele, E.S., Sumner, L.W., 2004. Potential of metabolomics as a functional genomics tool. *Trends Plant Sci.* 9, 418–425.
- Buckner, J.S., Mardaus, M.C., Nelson, D.R., 1996. Cuticular lipid composition of *Heliothis virescens* and *Helicoverpa zea* pupae. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 114, 207–216.
- Bundy, J.G., Davey, M.P., Viant, M.R., 2009. Environmental metabolomics: a critical review and future perspectives. *Metabolomics* 5, 3–21.
- Carefoot, T., Taylor, B., Kalwa, S., Somogyi, G., Hughes, B., 1992. Effect of dietary d-amino acids on growth, survival, ammonia excretion and specific dynamic action in the supralittoral isopod, *Ligia pallasii*. *Comp. Biochem. Physiol. A Physiol.* 103, 559–563.
- Cauchie, H.M., 2002. Chitin production by arthropods in the hydrosphere. *Hydrobiologia* 470, 63–95.
- Chen, J.M., Chen, J.C., 2000. Study on the free amino acid levels in the hemolymph, gill, hepatopancreas and muscle of *Penaeus monodon* exposed to elevated ambient ammonia. *Aquat. Toxicol.* 50, 27–37.
- Chen, J., Henderson, G., Laine, R.A., 1999. Lignoceric acid and hexacosanoic acid: major components of soldier frontal gland secretions of the formosan subterranean termite (*Coptotermes formosanus*). *J. Chem. Ecol.* 25, 817–824.
- Chich, J.F., David, O., Villers, F., Schaeffer, B., Lutomska, D., Huet, S., 2007. Statistics for proteomics: experimental design and 2-DE differential analysis. *J. Chromatogr. B* 849, 261–272.
- Colinge, J., Chiappe, D., Lagache, S., Moniatte, M., Bougueleret, L., 2005. Differential proteomics via probabilistic peptide identification scores. *Anal. Chem.* 77, 596–606.
- Cook, D., Johnson, M., 1974. Benthic macroinvertebrates of the St-Lawrence Great Lakes. *J. Fish. Res. Board Can.* 3, 763–782.
- Cristoni, S., Molin, L., Lai, A., Bernardi, L.R., Pucciarelli, S., Agostini, M., Bedin, C., Nitti, D., Seraglia, R., Repetto, O., Dibari, V.F., Orlandi, R., Sinues, P.M.L., Traldi, P., 2009. MALDI-MS-NIST library approach for colorectal cancer diagnosis. *Rapid Commun. Mass Spectrom.* 23, 2839–2845.
- Dennis, J.W., Lau, K.S., Demetriou, M., Nabi, I.R., 2009. Adaptive regulation at the cell surface by N-glycosylation. *Traffic* 10, 1569–1578.
- Dermott, R., 2001. Sudden disappearance of the amphipod *Diporeia* from Eastern Lake Ontario, 1993–1995. *J. Great Lakes Res.* 27, 423–433.
- Dunn, W.B., Ellis, D.I., 2005. Metabolomics: current analytical platforms and methodologies. *Trends Anal. Chem.* 24, 285–294.
- Farr, S.B., Kogoma, T., 1991. Oxidative stress responses in *Escherichia coli* and *Salmonella typhimurium*. *Microbiol. Mol. Biol. Rev.* 55, 561–585.
- Fujimori, T., Abe, H., 2002. Physiological roles of free D- and L-alanine in the crayfish *Procambarus clarkii* with special reference to osmotic and anoxic stress responses. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 131, 893–900.
- Gade, G., Auerswald, L., 2002. Beetles' choice—proline for energy output: control by AKHs. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 132, 117–129.
- Garcia, D.E., Baidoo, E.E., Benke, P.L., Pingitore, F., Tang, Y.J., Villa, S., Keasling, J.D., 2008. Separation and mass spectrometry in microbial metabolomics. *Curr. Opin. Microbiol.* 11, 233–239.
- Gardner, H., 1979. Lipid hydroperoxide reactivity with proteins and amino acids: a review. *J. Agric. Food Chem.* 27, 220–229.
- Gardner, H., 1995. Biological roles and biochemistry of the lipoxygenase pathway. *Hortscience* 30, 197–204.
- Gauvin, J.M., Gardner, W.S., Quigley, M.A., 1989. Effects of food removal on nutrient release rates and lipid content of Lake Michigan *Pontoporeia hoyi*. *Can. J. Fish. Aquat. Sci.* 46, 1125–1130.
- Giulivi, C., Ross-Inta, C., Horton, A., Luckhart, S., 2008. Metabolic pathways in *Anopheles stephensi* mitochondria. *Biochem. J.* 415, 309–316.
- Graney, R.L., Giesy Jr., J.P., 1986. Seasonal changes in the free amino acid pool of the freshwater amphipod *Gammarus pseudolimnaeus* Bousfield (Crustacea: Amphipoda). *Comp. Biochem. Physiol. A Physiol.* 85, 535–543.
- Hanover, J.A., 2001. Glycan-dependent signaling: O-linked N-acetylglucosamine. *FASEB J.* 15, 1865–1876.
- Hart, G.W., Housley, M.P., Slawson, C., 2007. Cycling of O-linked -N-acetylglucosamine on nucleocytoplasmic proteins. *Nature* 446, 1017–1022.
- Helland, S., Triantaphyllidis, G., Fyhn, H., Evjen, M., Lavens, P., Sorgeloos, P., 2000. Modulation of the free amino acid pool and protein content in populations of the brine shrimp *Artemia* spp. *Mar. Biol.* 137, 1005–1016.
- Herzog, H.U., Liappis, N., 1987. The plasma composition of larval *Aeshna cyanea* Müller-II. The effect of the nutritional state on the free amino acids. *Comp. Biochem. Physiol. A Physiol.* 87, 47–52.
- Hong, M., Chen, L., Qin, J.G., Sun, X., Li, E., Gu, S., Yu, N., 2009. Acute tolerance and metabolic responses of Chinese mitten crab (*Eriocheir sinensis*) juveniles to ambient nitrite. *Comp. Biochem. Physiol. B Toxicol. Pharmacol.* 149, 419–426.
- Huong, D.T.T., Wang, T., Bayley, M., Phuong, N.T., 2010. Osmoregulation, growth and molting cycles of the giant freshwater prawn (*Macrobrachium rosenbergii*) at different salinities. *Aquacult. Res.* 41, e135–e143.
- Johnson, R.K., 1987. The life history, production and food habits of *Pontoporeia affinis* Lindström (Crustacea: Amphipoda) in mesotrophic Lake Erken. *Hydrobiologia* 144, 277–283.
- Jones, O., Spurgeon, D., Svendsen, C., Griffin, J., 2008. A metabolomics based approach to assessing the toxicity of the polyaromatic hydrocarbon pyrene to the earthworm *Lumbricus rubellus*. *Chemosphere* 71, 601–609.
- Karanova, M., Andreev, A., 2010. Free amino acids and reducing sugars in the freshwater shrimp *Gammarus lacustris* (Crustacea, Amphipoda) at the initial stage of preparation to winter season. *J. Evol. Biochem. Physiol.* 46, 335–340.
- Karpievitch, Y.V., Polpitiya, A.D., Anderson, G.A., Smith, R.D., Dabney, A.R., 2010. Liquid chromatography mass spectrometry-based proteomics: biological and technological aspects. *Ann. Appl. Stat.* 4, 1797–1823.
- Kay, B.K., Williamson, M.P., Sudol, M., 2000. The importance of being proline: the interaction of proline-rich motifs in signaling proteins with their cognate domains. *FASEB J.* 14, 231–241.
- Laparie, M., Larvor, V., Frenot, Y., Renault, D., 2011. Starvation resistance and effects of diet on energy reserves in a predatory ground beetle (*Merizodus soledadinus*; Carabidae) invading the Kerguelen Islands. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 161, 122–129.

- Lehtonen, K.K., 1994. Metabolic effects of short-term starvation on the benthic amphipod *Pontoporeia affinis* Lindström from the northern Baltic Sea. *J. Exp. Mar. Biol. Ecol.* 176, 269–283.
- Leroy, D., Haubruge, E., De Pauw, E., Thome, J., Francis, F., 2010. Development of ecotoxicoproteomics on the freshwater amphipod *Gammarus pulex*: identification of PCB biomarkers in glycolysis and glutamate pathways. *Ecotoxicol. Environ. Saf.* 73, 343–352.
- Lin, C.Y., Viant, M.R., Tjeerdema, R.S., 2006. Metabolomics: methodologies and applications in the environmental sciences. *J. Pest Sci.* 31, 245–251.
- Liu, J., Yeo, H.C., Övervik-Douki, E., Hagen, T., Doniger, S.J., Chu, D.W., Brooks, G.A., Ames, B.N., 2000. Chronically and acutely exercised rats: biomarkers of oxidative stress and endogenous antioxidants. *J. Appl. Physiol.* 89, 21–28.
- Lorenz, M.W., Gade, G., 2009. Hormonal regulation of energy metabolism in insects as a driving force for performance. *Integr. Comp. Biol.* 49, 380–392.
- Lozano, S.J., Scharold, J.V., Nalepa, T.F., 2001. Recent declines in benthic macroinvertebrate densities in Lake Ontario. *Can. J. Fish. Aquat. Sci.* 58, 518–529.
- Maity, S., Jannash, A., Adamec, J., Gribskov, M., Nalepa, T., Höök, T., Sepúlveda, M.S., 2012. Metabolite profiles in starved *Diporeia* spp. using liquid chromatography-mass spectrometry (LC-MS) based metabolomics. *J. Crustacean Biol.*
- Marzolf, G.R., 1965. Substrate relations of the burrowing amphipod *Pontoporeia affinis* in Lake Michigan. *Ecology* 46, 580–592.
- Matés, J.M., Perez-Gomez, C., de Castro, I.N., Asenjo, M., Marquez, J., 2002. Glutamine and its relationship with intracellular redox status, oxidative stress and cell proliferation/death. *Int. J. Biochem. Cell Biol.* 34, 439–458.
- Mente, E., Coutteau, P., Houlihan, D., Davidson, I., Sorgeloos, P., 2002. Protein turnover, amino acid profile and amino acid flux in juvenile shrimp *Litopenaeus vannamei*: effects of dietary protein source. *J. Exp. Biol.* 205, 3107–3122.
- Mente, E., Davidson, I., Karapanagiotidis, I.T., Fountoulaki, E., Nengas, I., 2010. Amino acid analysis in the shore crab *Carcinus maenas* (Decapoda: Brachyura). *J. Crustacean Biol.* 30, 643–650.
- Meyers, S., 2000. Use of neurotransmitter precursors for treatment of depression. *Alt. Med. Rev.* 5, 64–71.
- Michaelis, E.K., 1998. Molecular biology of glutamate receptors in the central nervous system and their role in excitotoxicity, oxidative stress and aging. *Prog. Neurobiol.* 54, 369–415.
- Mohler, R.E., Dombek, K.M., Hoggard, J.C., Young, E.T., Synovec, R.E., 2006. Comprehensive two-dimensional gas chromatography time-of-flight mass spectrometry analysis of metabolites in fermenting and respiring yeast cells. *Anal. Chem.* 78, 2700–2709.
- Nalepa, T., 1998. Dramatic changes in benthic macroinvertebrate populations in southern Lake Michigan. *Aquat. Nuis. Sp.* 4, 1–2.
- Nalepa, T.F., Fanslow, D.L., Messick, G., 2005. Characteristics and potential causes of declining *Diporeia* spp. populations in southern Lake Michigan and Saginaw Bay, Lake Huron. *Great Lakes Fish Commun. Tech. Rep.* 66, 157–188.
- Netea, M.G., Brown, G.D., Kullberg, B.J., Gow, N.A.R., 2008. An integrated model of the recognition of *Candida albicans* by the innate immune system. *Nat. Rev. Microbiol.* 6, 67–78.
- Nicholson, J.K., Connelly, J., Lindon, J.C., Holmes, E., 2002. Metabonomics: a platform for studying drug toxicity and gene function. *Nat. Rev. Drug Discovery* 1, 153–161.
- Oh, C., Huang, X., Regnier, F.E., Buck, C., Zhang, X., 2008. Comprehensive two-dimensional gas chromatography/time-of-flight mass spectrometry peak sorting algorithm. *J. Chromatogr. A* 1179, 205–215.
- Pongsuwan, W., Bamba, T., Harada, K., Yonetani, T., Kobayashi, A., Fukusaki, E., 2008. High-throughput technique for comprehensive analysis of Japanese green tea quality assessment using ultra-performance liquid chromatography with time-of-flight mass spectrometry (UPLC/TOF MS). *J. Agric. Food Chem.* 56, 10705–10708.
- Prestwich, G.D., Collins, M.S., 1982. Chemical defense secretions of the termite soldiers of *Acorhinotermes* and *Rhinotermes* (Isoptera, Rhinotermitinae). *J. Chem. Ecol.* 8, 147–161.
- Ralston Hooper, K., Adamec, J., Jannash, A., Mollenhauer, R., Ochoa Acuña, H., Sepúlveda, M., 2011. Use of GC×GC/TOF MS and LC/TOF MS for metabolomic analysis of *Hyalella azteca* chronically exposed to atrazine and its primary metabolite, desethylatrazine. *J. Appl. Toxicol.* 31, 399–410.
- Ralston-Hooper, K., Hopf, A., Oh, C., Zhang, X., Adamec, J., Sepúlveda, M.S., 2008. Development of GC×GC/TOF-MS metabolomics for use in ecotoxicological studies with invertebrates. *Aquat. Toxicol.* 88, 48–52.
- Roesijadi, G., 1981. The significance of low molecular weight, metallothionein-like proteins in marine invertebrates: current status. *Mar. Environ. Res.* 4, 167–179.
- Rosa, R., Nunes, M.L., 2003. Seasonal changes in nucleic acids, amino acids and protein content in juvenile Norway lobster (*Nephrops norvegicus*). *Mar. Biol.* 143, 565–572.
- Rosa, R., Nunes, M.L., 2005. Seasonal patterns of nucleic acid concentrations and amino acid profiles of *Parapenaeus longirostris* (Crustacea, Decapoda): relation to growth and nutritional condition. *Hydrobiologia* 537, 207–216.
- Rosas, C., Cuzon, G., Gaxiola, G., Le Priol, Y., Pascual, C., Rossignol, J., Contreras, F., Sanchez, A., Van Wormhoudt, A., 2001. Metabolism and growth of juveniles of *Litopenaeus vannamei*: effect of salinity and dietary carbohydrate levels. *J. Exp. Mar. Biol. Ecol.* 259, 1–22.
- Rosenstiel, P., Philipp, E.E.R., Schreiber, S., Bosch, T.C.G., 2009. Evolution and function of innate immune receptors: insights from marine invertebrates. *Innate Immun.* 1, 291–300.
- Sahly, H., Keisari, Y., Crouch, E., Sharon, N., Ofek, I., 2008. Recognition of bacterial surface polysaccharides by lectins of the innate immune system and its contribution to defense against infection: the case of pulmonary pathogens. *Infect. Immun.* 76, 1322–1332.
- Samuelsson, L.M., Larsson, D.G.J., 2008. Contributions from metabolomics to fish research. *Mol. Biosyst.* 4, 974–979.
- Scaraffia, P., Wells, M., 2003. Proline can be utilized as an energy substrate during flight of *Aedes aegypti* females. *J. Insect Physiol.* 49, 591–601.
- Schock, T.B., Stancyk, D.A., Thibodeaux, L., Burnett, K.G., Burnett, L.E., Boroujerdi, A.F.B., Bearden, D.W., 2010. Metabolomic analysis of Atlantic blue crab, *Callinectes sapidus*, hemolymph following oxidative stress. *Metabolomics* 6, 250–262.
- Sepúlveda, M.S., Ralston-Hopper, K.J., Sanchez, B.C., Hopf-Jannash, A., Baker, S.D., Diaz, N., Adamec, J., 2011. Use of proteomic and metabolomics techniques in ecotoxicological research. In: Casciano, D.A., Sahu, S.C. (Eds.), *Handbook of Systems Toxicology*. John Wiley and Sons Ltd, Chichester, UK, pp. 227–254.
- Sharon, N., 2007. Lectins: carbohydrate-specific reagents and biological recognition molecules. *J. Biol. Chem.* 282, 2753–2764.
- Shellie, R.A., Welthagen, W., Zrostliková, J., Spranger, J., Ristow, M., Fiehn, O., Zimmermann, R., 2005. Statistical methods for comparing comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry results: metabolomic analysis of mouse tissue extracts. *J. Chromatogr. A* 1086, 83–90.
- Tanguy, A., Boutet, I., Moraga, D., 2005. Molecular characterization of the glutamine synthetase gene in the Pacific oyster *Crassostrea gigas*: expression study in response to xenobiotic exposure and developmental stage. *Biochim. Biophys. Acta, Gene Struct. Expression* 1681, 116–125.
- Van Ruitenbeek, P., Sambeth, A., Vermeeren, A., Young, S., Riedel, W., 2009. Effects of L-histidine depletion and L-tyrosine/L-phenylalanine depletion on sensory and motor processes in healthy volunteers. *Br. J. Pharmacol.* 157, 92–103.
- Vazquez, L., Alpuche, J., Maldonado, G., Agundis, C., Pereyra-Morales, A., Zenteno, E., 2009. Review: Immunity mechanisms in crustaceans. *Innate Immun.* 15, 179–188.
- Viant, M.R., 2008. Recent developments in environmental metabolomics. *Mol. Biosyst.* 4, 980–986.
- Wiklund, A., Sundelin, B., Rosa, R., 2008. Population decline of amphipod *Monoporeia affinis* in Northern Europe: consequence of food shortage and competition? *J. Exp. Mar. Biol. Ecol.* 367, 81–90.
- Wishart, D., 2005. Metabolomics: the principles and potential applications to transplantation. *Am. J. Transplant.* 5, 2814–2820.
- Yaron, A., Naider, F., Scharpe, S., 1993. Proline-dependent structural and biological properties of peptides and proteins. *Crit. Rev. Biochem. Mol. Biol.* 28, 31–81.
- Zarrinpar, A., Bhattacharyya, R.P., Lim, W.A., 2003. The structure and function of proline recognition domains. *Sci. STKE* 179, 1–10.
- Zhao, Y.Y., Takahashi, M., Gu, J.G., Miyoshi, E., Matsumoto, A., Kitazume, S., Taniguchi, N., 2008. Functional roles of N-glycans in cell signaling and cell adhesion in cancer. *Cancer Sci.* 99, 1304–1310.