

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

2-2005

Colonization of Gnotobiotic Piglets by a *luxS* Mutant Strain of *Escherichia coli* O157:H7

Dianna M. Jordan
Iowa State University

Vanessa Sperandio
University of Maryland School of Medicine

James B. Kaper
University of Maryland School of Medicine

Evelyn A. Dean-Nystrom
Pre-Harvest Food Safety and Enteric Diseases Research Unit, evelyn.nystrom@ars.usda.gov

Harley W. Moon
Iowa State University

Follow this and additional works at: <https://digitalcommons.unl.edu/usdaarsfacpub>

 Part of the [Agricultural Science Commons](#)

Jordan, Dianna M.; Sperandio, Vanessa; Kaper, James B.; Dean-Nystrom, Evelyn A.; and Moon, Harley W., "Colonization of Gnotobiotic Piglets by a *luxS* Mutant Strain of *Escherichia coli* O157:H7" (2005).
Publications from USDA-ARS / UNL Faculty. 935.
<https://digitalcommons.unl.edu/usdaarsfacpub/935>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Colonization of Gnotobiotic Piglets by a *luxS* Mutant Strain of *Escherichia coli* O157:H7

Dianna M. Jordan,^{1*} Vanessa Sperandio,^{2†} James B. Kaper,²
Evelyn A. Dean-Nystrom,³ and Harley W. Moon¹

Department of Veterinary Pathology, Iowa State University,¹ and Pre-Harvest Food Safety and Enteric Disease Research Unit, National Animal Disease Center, Agricultural Research Service, U.S. Department of Agriculture,³ Ames, Iowa, and Center for Vaccine Development and Department of Microbiology and Immunology, University of Maryland School of Medicine, Baltimore, Maryland²

Received 12 July 2004/Returned for modification 20 August 2004/Accepted 7 October 2004

Gnotobiotic piglets inoculated with *Escherichia coli* O157:H7, its *luxS* mutant derivative, or nonpathogenic *E. coli* were evaluated for attaching and effacing lesions. Although no differences in clinical symptoms were seen between pigs inoculated with the parent and those inoculated with the *luxS* mutant, the *luxS* mutant-inoculated pigs had a lower frequency of attaching and effacing lesions in the spiral colon than parent strain-inoculated pigs.

Quorum sensing (QS) is a bacterial cell-to-cell signaling mechanism used by a wide variety of bacterial species for recognition of the surrounding microbial community. The *luxS* QS system is responsible for production of a signaling compound called autoinducer 2 (AI-2) that can control gene expression in a wide variety of gram-positive and gram-negative species (10). In *Escherichia coli* O157:H7, the LuxS enzyme contributes to the production of another signaling compound, AI-3, which is involved in the QS regulation of some enterohemorrhagic *E. coli* (EHEC) virulence genes (11, 13). The AI-3 signaling compound is also produced by the human resident intestinal flora and hypothesized to be involved in signaling to EHEC that it has reached the large intestine of the host (4, 13). This QS signaling cascade activates transcription of the genes within the locus of enterocyte effacement (LEE) pathogenicity island, which encode a type III secretion system, bacterial effectors, and the bacterial adhesin intimin, essential for formation of the attaching and effacing (A/E) lesions in enterocytes (6, 11, 13). A *luxS* mutation in EHEC results in decreased transcription of the LEE region genes and a deficiency in type III secretion in broth cultures (11). Due to the influence of LuxS on the synthesis of AI-3, which modulates EHEC colonization factors, it is hypothesized that LuxS facilitates intestinal colonization of gnotobiotic pigs by *E. coli* O157:H7. The objective of this study was to determine if mutation of the *luxS* gene would impair the ability of *E. coli* O157:H7 to colonize gnotobiotic piglets.

Germfree piglets, 24 h old, were orally inoculated with 10⁵ CFU of bacteria. Eleven pigs from four litters were inoculated with the parent strain, a spontaneous streptomycin- and nali-

dix acid-resistant derivative of *E. coli* O157:H7 86-24 (14). Fourteen pigs from five litters were inoculated with strain VS97, a *luxS* mutant derived from the parent strain by methods described by Sperandio et al. (12). Six pigs from two litters were inoculated with nonpathogenic *E. coli* strain 123 (NPE 123) (7) to serve as controls.

Strains were grown as overnight cultures in tryptic soy broth with appropriate antibiotics. The parent strain was grown with 50 µg of streptomycin/ml and 10 µg of nalidixic acid/ml, the *luxS* mutant was grown with 10 µg of tetracycline/ml, and NPE 123 was grown without antibiotics. Inocula were prepared as previously described (1, 2, 9).

Pigs inoculated with the parent strain and VS97 were necropsied when neurologic signs were demonstrated or at 7 days postinoculation (p.i.). Control pigs were necropsied at 4 days p.i., the average time of onset of clinical signs for pigs inoculated with the pathogenic strains. At necropsy, sections of ilea, ceca, spiral colons, cerebrums, cerebellums, and brain stems were collected in neutral buffered formalin for histopathology, processed, and stained with hematoxylin and eosin. A specific horseradish peroxidase immunohistochemical stain with the primary antibody targeting the O157 antigen was used for the intestinal sections (3). Cross-sections from each intestinal segment were examined microscopically in a blind fashion and scored for A/E lesions with intimately associated bacteria as a measure of colonization (5, 8). Multiple sections of spiral colons from five pigs that were inoculated with the parent strain and the *luxS* mutant were prepared and evaluated by transmission electron microscopy.

Contingency analysis for ordinal data, followed by pairwise testing, was done for the A/E scores. JMP (version 5.0.1a; SAS Institute, Inc.) was the statistical program utilized for analysis; a *P* value of <0.05 was used as the level of significance for all evaluations.

Pigs from groups inoculated with the parent strain or its *luxS* mutant derivative showed clinical signs of ataxia, head-press-

* Corresponding author. Present address: Veterinary Diagnostic and Production Animal Medicine, Iowa State University, Ames, IA 50011. Phone: (515) 294-1950. Fax: (515) 294-3564. E-mail: dmjordan@iastate.edu.

† Present address: Department of Microbiology, University of Texas Southwestern Medical Center, Dallas, TX 75390-9048.




FIG. 1. A/E lesion scores in tissues of gnotobiotic piglets inoculated with nonpathogenic *E. coli* (NPE), *E. coli* O157:H7 strain VS97 (*luxS* mutant), or 86-24 (Parent). The A/E lesion scores of the formalin-fixed intestinal sections were as follows: 0, no A/E lesions with discernable associated bacteria visible throughout the sections; 0.5, infrequent A/E lesions identified only with the immunohistochemical stain; 1, discernable lesions on fewer than 10% of the enterocytes of the sections; 2, discernable lesions on more than 10% but fewer than 50% of the enterocytes; 3, lesions present on 50% or more of the enterocytes. Contingency analysis detected statistically significant differences between the scores for the *luxS* mutant-inoculated tissues and those for the parent-inoculated tissues from the spiral colons ($P < 0.03$).

ing, and/or recumbency due to the systemic effects of Shiga toxins (15). Eleven of 14 pigs inoculated with the *luxS* mutant developed neurologic signs between 3.5 and 7 days p.i. Nine of 11 pigs inoculated with the parent strain developed neurologic signs between 3 and 7 days p.i. No clinical signs were observed in the control pigs necropsied at 4 days p.i.

Upon necropsy, 10 of 14 *luxS* mutant-inoculated pigs and all 11 parent strain-inoculated pigs had mesocolonic edema. No gross enteric lesions were present in the NPE 123-inoculated control pigs.

Histologically, the A/E lesions in the ilea, ceca, and spiral colons from the *luxS* mutant-inoculated pigs were qualitatively similar to lesions from the pigs inoculated with the parent strain. The distributions of the A/E lesion scores are summarized in Fig. 1. The parent strain-inoculated group had significantly higher ($P = 0.0025$) A/E scores (median, 2; range, 1 to 3) for the spiral colon than did the *luxS* mutant-inoculated pigs (median, 1; range, 0 to 3). For the spiral colon sections, both the parent strain- and the *luxS* mutant-inoculated pigs had significantly higher ($P < 0.0001$) A/E lesion scores than did the NPE 123-inoculated pigs. There was no significant difference ($P > 0.5$) between the cecal A/E lesion scores for the parent strain (median, 2; range, 0 to 3) and those for the *luxS* mutant-inoculated pigs (median, 1; range, 0 to 3); these two groups had significantly higher cecal A/E lesion scores than did the NPE 123-inoculated pigs ($P < 0.002$). There were no A/E lesions in any intestinal segments from the pigs inoculated with NPE 123. There were no significant differences in A/E lesion scores for the ileal sections from pigs inoculated with any of the strains. Classical A/E lesions (8) were identified with electron microscopy in the spiral colons from all the parent strain- and *luxS*

TABLE 1. Clinical and microscopic findings for gnotobiotic piglets inoculated with *E. coli* O157:H7 strain 86-24, VS97, or NPE 123

Strain used for inoculation	No. of samples positive for indicated finding/no. of pigs inoculated					
	Clinical signs	Tissue colonization			Brain histopathology	
		Ileum	Cecum	Spiral colon	Microhemorrhage	Vessel necrosis
Parent	9/11	3/6	10/11	11/11	2/11	8/11
<i>luxS</i> mutant	11/14	2/6	13/14	14/14	6/14	9/14
NPE 123	0/6	0/6	0/6	0/6	0/6	0/6

mutant-inoculated pigs examined. Microhemorrhages and necrotic vessels were detected in the brains of pigs inoculated with the parent strain or its *luxS* derivative; results are shown in Table 1. There was no statistically significant difference between the scores of these two groups of pigs for neurological lesions. Pigs inoculated with NPE 123 had no neurologic lesions.

We hypothesized that LuxS was a critical enzyme in the production of autoinducer signals that are important for the expression of virulence factors, such as those required for A/E lesion production by *E. coli* O157:H7 in vitro (11, 12). This study demonstrated that A/E lesions in spiral colons occurred less frequently in gnotobiotic piglets inoculated with the *luxS* mutant strain than in those inoculated with the parent strain. There were no significant differences in the frequencies of lesions in the ceca and ilea. Mutation in *luxS* did not eliminate the ability of *E. coli* O157:H7 to produce A/E lesions or to cause clinical disease in piglets due to the effects of Shiga toxin. These results were consistent with a recent report by Sperandio et al. (13) demonstrating that the eukaryotic hormones epinephrine and norepinephrine activate the type III secretion system, *LEE* transcription, and, consequently, A/E lesion formation by the *luxS* mutant. Therefore, the host's epinephrine and norepinephrine may be substituting for the AI-3 bacterial signaling (13). In this study, the presence of these hormones or other host factors in the intestines of the gnotobiotic piglets presumably may have signaled to the bacteria to express the *LEE* region genes for the development of A/E lesions; therefore, compelling differences were not demonstrated in vivo as they were in vitro with the *luxS* mutant of *E. coli* O157:H7.

This work was supported by NIH grant 1R01DK58957, awarded to J.B.K.

We thank Lawayne Nusz and the NADC animal caretaker, Roger Spaete, for their assistance; Judy Stasko for transmission electron microscopy assistance; Richard Evans for statistical consultation; and, especially, Sheridan Booher for technical assistance in bacteriology and animal handling.

REFERENCES

1. Booher, S. L., N. A. Cornick, and H. W. Moon. 2002. Persistence of *Escherichia coli* O157:H7 in experimentally infected swine. *Vet. Microbiol.* **89**:69-81.
2. Cornick, N. A., S. L. Booher, T. A. Casey, and H. W. Moon. 2000. Persistent colonization of sheep by *Escherichia coli* O157:H7 and other *E. coli* pathotypes. *Appl. Environ. Microbiol.* **66**:4926-4934.
3. Dean-Nystrom, E. A., B. T. Bosworth, W. C. Cray, Jr., and H. W. Moon. 1997. Pathogenicity of *Escherichia coli* O157:H7 in the intestines of neonatal calves. *Infect. Immun.* **65**:1842-1848.
4. Falcao, J. P., F. Sharp, and V. Sperandio. 2004. Cell-to-cell signaling in intestinal pathogens. *Curr. Issues Intest. Microbiol.* **5**:9-17.
5. Jordan, D. M., N. Cornick, A. G. Torres, E. A. Dean-Nystrom, J. B. Kaper,

- and H. W. Moon. 2004. Long polar fimbriae contribute to colonization by *Escherichia coli* O157:H7 in vivo. *Infect. Immun.* **72**:6168–6171.
6. McDaniel, T. K., K. G. Jarvis, M. S. Donnenberg, and J. B. Kaper. 1995. A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proc. Natl. Acad. Sci. USA* **92**:1664–1668.
 7. Moon, H. W., and S. C. Whipp. 1971. Systems for testing the enteropathogenicity of *Escherichia coli*. *Ann. N. Y. Acad. Sci.* **176**:197–211.
 8. Moon, H. W., S. C. Whipp, R. A. Argenzio, M. M. Levine, and R. A. Giannella. 1983. Attaching and effacing activities of rabbit and human enteropathogenic *Escherichia coli* in pig and rabbit intestines. *Infect. Immun.* **41**:1340–1351.
 9. Sarmiento, J. I., T. A. Casey, and H. W. Moon. 1988. Postweaning diarrhea in swine: experimental model of enterotoxigenic *Escherichia coli* infection. *Am. J. Vet. Res.* **49**:1154–1159.
 10. Schauder, S., K. Shokat, M. G. Surette, and B. L. Bassler. 2001. The LuxS family of bacterial autoinducers: biosynthesis of a novel quorum-sensing signal molecule. *Mol. Microbiol.* **41**:463–476.
 11. Sperandio, V., J. L. Mellies, W. Nguyen, S. Shin, and J. B. Kaper. 1999. Quorum sensing controls expression of the type III secretion gene transcription and protein secretion in enterohemorrhagic and enteropathogenic *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **96**:15196–15201.
 12. Sperandio, V., A. G. Torres, J. A. Girón, and J. B. Kaper. 2001. Quorum sensing is a global regulatory mechanism in enterohemorrhagic *Escherichia coli* O157:H7. *J. Bacteriol.* **183**:5187–5197.
 13. Sperandio, V., A. G. Torres, B. Jarvis, J. P. Nataro, and J. B. Kaper. 2003. Bacteria-host communication: the language of hormones. *Proc. Natl. Acad. Sci. USA* **100**:8951–8956.
 14. Tarr, P. I., M. A. Neill, C. R. Clausen, J. W. Newland, R. J. Neill, and S. L. Moseley. 1989. Genotypic variation in pathogenic *Escherichia coli* O157:H7 isolated from patients in Washington, 1984–1987. *J. Infect. Dis.* **159**:344–347.
 15. Tzipori, S., H. Karch, K. I. Wachsmuth, R. M. Robins-Browne, A. D. O'Brien, H. Lior, M. L. Cohen, J. Smithers, and M. M. Levine. 1987. Role of a 60-megadalton plasmid and Shiga-like toxins in the pathogenesis of infection caused by enterohemorrhagic *Escherichia coli* O157:H7 in gnotobiotic piglets. *Infect. Immun.* **55**:3117–3125.

Editor: J. B. Bliska