University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

Historical Publications in Weed Science and Weed Technology

Agronomy and Horticulture Department

4-24-1987

INSECT, PLANT DISEASE, & WEED SCIENCE NEWS [No. 87-6] [April 24, 1987]

Fred W. Roeth University of Nebraska-Lincoln, fwroeth41@gmail.com

Alex Martin University of Nebraska - Lincoln, amartin2@unl.edu

Follow this and additional works at: https://digitalcommons.unl.edu/weedscihist

Roeth, Fred W. and Martin, Alex, "INSECT, PLANT DISEASE, & WEED SCIENCE NEWS [No. 87-6] [April 24, 1987]" (1987). *Historical Publications in Weed Science and Weed Technology*. 7. https://digitalcommons.unl.edu/weedscihist/7

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Publications in Weed Science and Weed Technology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

COOPERATIVE EXTENSION SERVICE

INSTITUTE OF AGRICULTURE AND NATURAL RESOURCES



INSECT PLANT DISEASE WEED SCIENCE

NEWS

DEPARTMENT OF AGRONOMY (WEED SCIENCE) UNIVERSITY OF NEBRASKA-LINCOLN, EAST CAMPUS 68583-0915 PHONE 472-1555 87-6 April 24, 1987

In This Issue:

-Musk Thistle Control

-Book Announcement: SUSTAINABLE AGRICULTURE

-1987 Weed Science Tour

-David Mortensen Joins Staff at Lincoln

Musk Thistle Control

As the weather warms and the grass greens, musk and assorted other pasture thistles are set to compete, bolt, flower, and scatter seeds across the land. Springing from an innocent-looking flat rosette in late April to a gigantic seed factory in late June, this same scene repeats itself yearly. A robust musk thistle can produce 20,000 seeds. Last summer and fall had ideal weather for musk thistle seed production, germination, and rosette growth. With abundant root reserves, the thistles will be vigorous in 1987. The only effective way to control them chemically is by spraying the rosettes. Once the center stalk elongates, musk thistle has a much higher tolerance for herbicides and is not easily killed. So apply either 2,4-D at 1 1/2-2 qt/A, Banvel + 2,4-D at 0.5 + 1 qt/A, or Tordon 22K at 0.5 pt/A within the next two weeks. Normally May 1 to May 10 are the critical dates for musk thistle bolting across Nebraska from southeast to northwest.

Be especially careful of sensitive, non-target plants in the vicinity of the spray area and avoid drift onto these species. A11 of the thistle herbicides will injure trees, shrubs, flowers, garden crops, ornamentals, alfalfa, and other broadleaf crops. The surest way to avoid plant injury is to spray as soon as possible and to use good judgment regarding wind speed, wind direction, and spray application.

Even though musk thistle is a biennial it is an annual problem, so give some consideration to a fall treatment for controlling the 1988 crop. Fall application is effective, less risky to many sensitive plants, and usually more convenient for the applicator.

Book Announcement: SUSTAINABLE AGRICULTURE

A new book, "Sustainable Agriculture...Wise and Profitable Use of Our Resources in Nebraska", is now available from the Department of Agronomy at the University of Nebraska-Lincoln. The book highlights practices which help producers cut costs by reducing fine-tuning inputs, particularly fertilizer and pesticide

(Continued)



UNIVERSITY OF NEBRASKA-LINCOLN, COOPERATING WITH THE COUNTIES AND THE U.S. DEPARTMENT OF AGRICULTURE The Cooperative Extension Service provides information and educational programs to all people without regard to race, color, national origin, sex or handicap.

applications to those levels considered appropriate (and necessary). There are a number of articles on non-chemical approaches to pest control and to meeting a part of the fertility needs of crops with rotations and legumes.

<u>Sustainable Agriculture</u> is available at cost from the Department of Agronomy at the address shown below. Each book costs \$7.00 to cover printing and mailing.

> Sustainable Agriculture Book Department of Agronomy 279 Plant Sciences University of Nebraska Lincoln, NE 68583-0910

1987 Weed Science Tour

The itinerary has been set for the 1987 Weed Tour. We will be completing the entire tour during the same week this year. The proposed itinerary is as follows:

June 22	, 12:30 p.m. 6:30 p.m.	Concord - Northeast REC Lincoln - 84th and Havelock Avenue
June 23	, 10:00 a.m. 12:00 Noon	Clay Center - South Central REC Adjourn
June 24	, 8:00 a.m. 3:00 p.m. (MDT)	North Platte - West Central REC Sidney - High Plains Station
June 25	, 8:00 a.m. (MDT)	Scottsbluff - Panhandle REC
	11:00 a.m. (MDT)	Adjourn

David Mortensen Joins Staff at Lincoln

Dr. David Mortensen has joined the Agronomy Department at Lincoln as an Assistant Professor. His responsibilities will be both Research and Teaching with research emphasis on weed control in conservation tillage systems. Dave is a native of New York and recently completed his Ph.D. under Dr. Harold Coble at North Carolina State University. Welcome aboard, Dave!

Fred Rock

Fred W. Roeth Extension Weeds Specialist

alex Martin

Alex R. Martin Extension Weeds Specialist