

Epidemiology and Infection

<http://journals.cambridge.org/HYG>

Additional services for *Epidemiology and Infection*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Pathogen burden and cortisol profiles over the day

A. STEPTOE, Å. GYLFE, A. SHAMAEI-TOUSI, S. BERGSTROM and B. HENDERSON

Epidemiology and Infection / Volume 137 / Issue 12 / December 2009, pp 1816 - 1824
DOI: 10.1017/S0950268809002738, Published online: 19 May 2009

Link to this article: http://journals.cambridge.org/abstract_S0950268809002738

How to cite this article:

A. STEPTOE, Å. GYLFE, A. SHAMAEI-TOUSI, S. BERGSTROM and B. HENDERSON (2009). Pathogen burden and cortisol profiles over the day. *Epidemiology and Infection*, 137, pp 1816-1824 doi:10.1017/S0950268809002738

Request Permissions : [Click here](#)

Pathogen burden and cortisol profiles over the day

A. STEPTOE¹*, Å. GYLFE², A. SHAMAEI-TOUSI³, S. BERGSTROM⁴
AND B. HENDERSON⁵

¹ Department of Epidemiology and Public Health, University College London, UK

² Department of Chemistry, Umeå University, Umeå, Sweden

³ Helperby Therapeutics Limited, London, UK

⁴ Department of Molecular Biology, Umeå University, Umeå, Sweden

⁵ Division of Microbial Diseases, UCL Eastman Dental Institute, London, UK

(Accepted 13 April 2009; first published online 19 May 2009)

SUMMARY

Hypothalamic–pituitary–adrenocortical (HPA) regulation in adults is influenced by early psychosocial adversity, but the role of infectious disease history is poorly understood. We studied the association between cumulative pathogen burden and cortisol profile over the day in a sample of 317 healthy men and women aged 51–72 years. Cumulative pathogen burden was defined as positive serostatus for *Chlamydia pneumoniae*, cytomegalovirus (CMV) and herpes simplex virus 1 (HSV-1). Salivary cortisol was sampled repeatedly over the day. The cortisol slope was defined as the decrease across the day and evening. Age, gender, grade of employment, body mass index, smoking status, self-rated health, cardiovascular medication, depressed mood and time of waking were included as covariates. The pathogen burden averaged 1.76 (s.d. = 0.92). The cortisol slope was inversely associated with pathogen burden after controlling for covariates. When individual pathogens were studied, only CMV was associated with flatter cortisol rhythms in isolation. We conclude that pathogen burden is independently associated with flatter cortisol slopes over the day, and may contribute to disturbed neuroendocrine regulation.

Key words: Cortisol awakening response, *Chlamydia pneumoniae*, cytomegalovirus, herpes simplex virus 1, psychosocial adversity.

INTRODUCTION

There is abundant evidence that early childhood adversity influences adult health outcomes, with factors such as low childhood socioeconomic status (SES), poverty, and poor or overcrowded housing being associated with increased risk of cardiovascular, respiratory and psychiatric illness [1–3]. These effects

may be mediated in part through stress processes, since childhood abuse, family discord and negative life events increase risk for later ill health [4–6], and tend to cluster in children of lower SES [7]. Childhood stressors also impact on biological processes relevant to adult disease risk [8, 9]. Extensive research both in animals and humans has documented the impact of early psychosocial adversity on neuroendocrine function in adult life, with effects on cortisol regulation and stress reactivity [10, 11]. Childhood SES is associated with aberrant profiles of cortisol over the day independently of adult SES [12]. Stress-induced

* Address for correspondence: Professor A. Steptoe, Department of Epidemiology and Public Health, University College London, 1-19 Torrington Place, London WC1E 6BT, UK.
(Email: a.steptoe@ucl.ac.uk)

neuroendocrine dysregulation has been proposed as a key process linking early life experience with adult risk of physical disease and psychopathology [13–15]. Cortisol has a pronounced diurnal rhythm, with high levels early in the morning, declining throughout the day to a nadir in the late evening and early hours of sleep [16]. The full diurnal profile requires repeated blood sampling, usually in clinical or laboratory settings, but the introduction of salivary cortisol measurements has permitted the profile over waking hours to be assessed in large samples in everyday life [17]. Different aspects of the diurnal profile appear to be associated with SES and other psychosocial factors, notably the cortisol awakening response (CAR; the rise over the first 30–45 min after waking), the overall cortisol output over the day, and the slope of cortisol decline over the day [17, 18].

Another pathway through which early life adversity may influence adult disease risk is exposure to common pathogens. Thus seropositivity for cytomegalovirus (CMV), *Chlamydia pneumoniae* (*C. pneumoniae*), herpes simplex virus 1 (HSV-1), and *Helicobacter pylori* is greater in individuals of low SES defined by education, poverty or occupational status, and with household crowding [19–25]. These pathogens are typically acquired in childhood [26], when they may or may not be symptomatic [27–29]. Reactivation of specific pathogens such as CMV and HSV-1 may be relevant to disease risk and cognitive decline in older age [30, 31]. In addition, cumulative pathogen burden, defined by positive serostatus for a range of pathogens, has been associated with cardiovascular risk factors and with coronary heart disease (CHD) in case-control and longitudinal cohort studies [32–35].

The relationship between pathogen burden and cortisol profile in adults has not been extensively investigated. Animal studies have demonstrated that early-life exposure to endotoxin alters later-life hypothalamic–pituitary–adrenocortical (HPA) function, with raised corticosterone output over the day [36]. Glucocorticoids are known to enhance the growth of *C. pneumoniae* in cell cultures [37] and the replication of human CMV in embryonic cells [38]. Additionally, early childhood exposure to upper respiratory infection was negatively correlated with cortisol sampled in the clinic and in the late evening in the Barry Caerphilly Growth cohort [39]. In this study, we hypothesized that cumulative pathogen burden, defined by positive serostatus for *C. pneumoniae*, CMV and HSV-1, influences cortisol regulation

in adult life. We measured salivary cortisol over the day and evening, specifically focusing on total cortisol output, the CAR, and the slope of cortisol decline over the day. A blunted CAR has been associated with cardiovascular risk factors [40], while a flattened cortisol profile over the day is related to subclinical coronary artery disease [41] and abdominal adiposity [42]. Other factors such as age, gender, low SES, body mass, smoking, depression, health status, and time of waking have been related to cortisol profiles over the day, and were therefore taken into account in the analyses.

METHODS

Participants

Data were analysed from 361 members (237 men, 124 women) aged 51–72 years from the Whitehall II epidemiological cohort [43]. Data were collected from consecutive attendees at a medical screening session where we completed assessments of serostatus for *C. pneumoniae*, CMV, and HSV-1, as detailed elsewhere [20]. All participants were in good health, with no indications of current infection. In total, 378 participants carried out salivary cortisol assessments over the day and evening, of whom 17 were excluded because they were taking either oral or inhaled corticosteroid medication. The individuals taking part in this substudy did not differ from the remaining healthy participants in the medical screening in terms of age, gender and SES. None of the participants were shift-workers. The study was approved by the University College London/University College London Hospitals Medical Ethics Committee, and all participants gave signed consent.

MEASURES

Participants underwent a physical examination during which fasting blood samples were obtained. Height and weight were measured, and body mass index (BMI) was computed. Current or most recent occupational grade in the civil service was used to index SES, and participants were divided into higher, intermediate and lower grade groups. Self-rated health was measured with a single item ('In general, would you say your health is: excellent, very good, good, fair, or poor') that has consistently been shown to predict adverse health outcomes [44]. Depressed mood was measured using the Center for Epidemiologic

Studies Depression (CES-D) scale. Current medication was also assessed.

Salivary sampling with salivettes (Sarstedt, UK) was explained and practised during the health screening. Participants were instructed to collect saliva immediately after waking, 30 min later, and then at 2½ h, 8 h and 12 h after waking, with a final sample being obtained just before bedtime. They were instructed to record the time of sample collection, take samples as soon as they woke, avoid caffeine and acidic drinks in the first 30 min, and not brush teeth or eat or drink anything for 15 min before a sample collection. Time of waking was recorded along with the actual times the saliva samples were collected. Participants stored salivettes in domestic refrigerators before posting them back to the laboratory. Saliva sampling took place within 7–10 days of the collection of blood samples used for the analysis of seropositivity.

Laboratory methods

Serostatus for *C. pneumoniae*, CMV, and HSV-1 was assayed at Umeå University using methods described previously [45]. Briefly, *C. pneumoniae* IgG was measured with the *Chlamydia* microtitre indirect immunofluorescence test (Focus Technologies, USA) using a serum dilution of 1/16. Each serum was tested against antigen from *C. pneumoniae* and a control antigen. Two control sera were used, one seropositive and one seronegative. The results were read manually by fluorescent microscopy by two independent investigators and sera with a specific apple green fluorescence were considered seropositive. Sera reacting with the control antigen were not possible to interpret and therefore excluded from the analyses. All procedures and interpretation were performed according to the manufacturer's instructions. The antigen for the CMV IgG ELISA was derived from CMV-strain Ad169 cultured in human fibroblasts. The antibody activity was expressed in arbitrary units (AU); a value of ≥ 10 AU was considered positive for presence of IgG and < 5 AU as negative. A grey zone area was defined in the range between 5–9 AU, but participants who fell into this range were classified conservatively as seronegative. The antigen for the HSV-1 IgG ELISA was a local HSV-1 strain cultured in GMK-cells. Both tests followed the protocols and materials used for clinical tests accredited by SWEDAC (Swedish Board for Accreditation and Confirmatory Assessment), and external international quality

assurance programmes from Labquality and the UK National External Quality Assessment Service validate the tests several times a year. The sensitivity of detecting these latent viral infections is estimated to be almost 100%. The specificity is also very high since each serum sample is controlled for unspecific reactivity by a control antigen. The high and low reacting control sera used have been tested at two independent accredited laboratories according to standard accreditation regulations.

Salivary samples were sent to the Technical University Dresden for the analysis of cortisol. The salivettes were centrifuged at 3000 rpm for 5 min, resulting in a clear supernatant of low viscosity. Salivary cortisol levels were measured using a commercial immunoassay with chemiluminescence detection (CLIA; IBL-Hamburg, Germany). The lower limit of the assay was 0.44 nmol/l, and intra- and inter-assay coefficients of variance were $< 8\%$.

Statistical analysis

Delays between waking in the morning and taking the 'waking' cortisol sample can lead to inaccurate waking values and a diminished CAR [46]. We excluded 44 individuals who reported a delay of more than 15 min between waking and taking sample 1, leaving 317 in the analysis. There were no differences in age, gender, seropositivity, self-rated health or other factors between those who were included and excluded from the final analysis. Cumulative pathogen burden was defined as the number of pathogens (*C. pneumoniae*, CMV, HSV-1) for which the individual was seropositive, so could range from 0 to 3. The associations between pathogen burden and age, gender, SES (grade of employment), BMI, smoking status, self-rated health, depression and time of waking were assessed using analysis of variance for continuous measures and χ^2 tests and logistic regression for categorical variables. The most common medications were related to cardiovascular health, including antihypertensives and statins. The use of cardiovascular medication was therefore included as an additional covariate. Total cortisol output over the day was computed using area under the curve methods involving all six samples [47]. The slope or rate of decline in cortisol over the day was computed by calculating regression slopes across all data-points except the 30-min post-waking sample (sample 2), and was expressed as nmol/l per h. The CAR was assessed as the difference between samples 1 and 2. The cortisol

Table 1. *Pathogen burden*

Pathogen	Serostatus, <i>N</i> (%)	
	Positive	Negative
<i>Chlamydia pneumoniae</i> (<i>n</i> =317)	200 (64.7%)	109 (35.3%)
Cytomegalovirus (<i>n</i> =315)	165 (52.4%)	150 (47.6%)
Herpes simplex virus 1 (<i>n</i> =317)	193 (60.9%)	124 (39.1%)
Total pathogen burden (<i>n</i> =317) mean (s.d.)	1.76 (0.92)	

output, cortisol slope over the day and CAR were normally distributed, while levels on waking and bedtime were skewed, and therefore log-transformed before analysis. The relationship of cortisol with burden of infection was assessed with analysis of covariance, adjusting for age, gender, SES, BMI, smoking status, self-rated health, depression and time of waking. Results are presented as means \pm standard deviations.

RESULTS

The 317 participants in the analysis included 205 men and 112 women aged 61.0 ± 5.7 years. Serological status for the three pathogens is summarized in Table 1. The pathogen burden was 0 in 9.8%, 1 in 27.8%, 2 in 39.1% and 3 in 23.3%, with a mean of 1.76 ± 0.92 . Pathogen burden did not differ between men and women, but was positively related to age ($P=0.004$). As previously described, participants of lower SES tended to have a greater pathogen burden ($P=0.016$) [20]. Pathogen burden was positively associated with BMI, but was not related to smoking, self-rated health, cardiovascular medication, depressed mood or time of waking in the morning (Table 2).

Figure 1 illustrates the profile of salivary cortisol over the study. Participants woke at 6:41 on average, and took the 30-min post-waking sample at $7:12 \pm 56$ min. Subsequent measures were taken at the expected intervals. Cortisol increased by 6.88 ± 10.4 nmol/l between waking and 30 min later, falling over the remainder of the day. Cortisol output over the day averaged 7.87 ± 2.87 nmol/l, while the rate of cortisol decline over the day averaged 1.204 ± 0.89 nmol/l per h.

There was no relationship between pathogen burden and total cortisol output over the day (Table 3).

This was also true when the three individual pathogens were tested. However, there was a significant linear association between pathogen burden and cortisol decline over the day after adjustment for age, gender, grade of employment, BMI, smoking status, self-rated health, cardiovascular medication, depression and time of waking ($P=0.010$). As can be seen in Figure 2, the rate of cortisol decrease over the day was greatest in participants with zero pathogen burden, and least in those with the highest pathogen burden. This indicates that the cortisol rhythm over the day was flatter in participants with greater pathogen burden, independently of covariates associated with pathogen exposure and cortisol output. When we analysed relationships with individual pathogens, a flatter cortisol slope was associated with seropositivity for CMV ($P<0.001$), but not the other pathogens in isolation. Results were unchanged when waist/hip ratio was included as an additional covariate.

The flatter cortisol decline over the day in participants with a greater pathogen burden was driven primarily by differences in cortisol levels on waking. Separate analysis of the waking values showed a significant association with pathogen burden after adjustment for age, gender, SES, BMI, smoking status, self-rated health, cardiovascular medication, depression and time of waking ($P=0.031$). Cortisol levels on waking were higher in participants with zero pathogen burden scores. By contrast, pathogen burden was not related to cortisol at bedtime ($P=0.63$), although values were lowest in participants with zero pathogen burden.

There was no association between pathogen burden and CAR ($P=0.58$). The CAR averaged 6.76 ± 10.1 , 7.34 ± 10.3 , 6.18 ± 10.1 and 8.58 ± 11.0 nmol/l in participants with burden scores of 0, 1, 2 and 3 respectively, after adjustment for covariates.

DISCUSSION

Cumulative pathogen burden has been postulated to be a risk factor for CHD, contributing to atherosclerosis [20, 32, 33]. Exposure to pathogens is greater in lower SES groups and in individuals who experience poverty and adversity in early life. Since early stress and adversity are related to HPA regulation, we reasoned that older adults with greater pathogen burden scores might show disturbances of cortisol output. It should be emphasized that these analyses of associations between cortisol output over the day

Table 2. Participant characteristics associated with pathogen burden

	Pathogen burden				P trend
	0 (n = 31)	1 (n = 88)	2 (n = 124)	3 (n = 74)	
Age (years)	59.1 (5.3)	60.5 (5.5)	60.9 (6.0)	62.5 (5.2)	0.004
Gender (male), n (%)	18 (58.1%)	61 (69.3%)	75 (60.5%)	51 (68.9%)	0.69
Grade of employment					
Higher	6 (20.0%)	29 (33.0%)	28 (22.6%)	14 (18.9%)	0.016
Intermediate	16 (53.3%)	47 (53.4%)	59 (47.6%)	36 (48.6%)	
Lower	8 (26.7%)	12 (13.6%)	37 (29.8%)	24 (32.4%)	
Body mass index	26.4 (5.8)	25.4 (4.1)	26.5 (4.3)	27.9 (4.2)	0.032
Current smokers, n (%)	2 (6.5%)	8 (9.2%)	11 (8.9%)	8 (10.8%)	0.51
Self-rated health (fair or poor), n (%)	6 (21.4%)	7 (8.2%)	14 (12.2%)	10 (14.7%)	0.43
Cardiovascular medication, n (%)	3 (9.7%)	5 (5.7%)	15 (12.1%)	12 (16.2%)	0.068
CES-D depression	8.80 (9.3)	8.73 (8.2)	8.52 (7.8)	8.14 (7.9)	0.95
Time of waking in the morning (h:min)	6:33 (38)	6:42 (52)	6:42 (59)	6:39 (57)	0.96

Results expressed as means (standard deviation) and n (percent).

Table 3. Salivary cortisol in relation to pathogen burden

Time (h:min)	Pathogen burden			
	0	1	2	3
Waking (6:41 ± 0:55)	21.29 ± 1.66	18.29 ± 0.89	17.95 ± 0.76	17.26 ± 1.01
Wake + 30 min (7:13 ± 0:56)	28.36 ± 2.11	25.59 ± 1.13	23.93 ± 0.96	25.63 ± 1.30
Wake + 2.5 h (9:17 ± 0:58)	7.79 ± 1.07	10.75 ± 0.60	9.74 ± 0.51	10.47 ± 0.66
Wake + 8 h (14:51 ± 1:05)	5.99 ± 0.84	6.73 ± 0.47	6.78 ± 0.40	7.59 ± 0.52
Wake + 12 h (18:52 ± 1:04)	4.34 ± 0.63	3.60 ± 0.36	3.19 ± 0.30	3.44 ± 0.39
Bedtime (23:10 ± 0:57)	2.47 ± 0.61	3.12 ± 0.34	2.77 ± 0.29	3.06 ± 0.38

The times of sampling are presented as means ± standard deviation. Cortisol results are presented expressed as means ± standard error, adjusted for age, gender, grade of employment, body mass index, smoking status, self-rated health, cardiovascular medication, depression and time of waking.




Fig. 1. Mean salivary cortisol at the six assessment points over the day. Error bars are standard error of the mean (S.E.M.).

and cumulative pathogen burden are preliminary, since only three pathogens contributed to the cumulative burden index, and salivary cortisol was

measured over only a single day. Nevertheless, we found that pathogen burden was associated with a flatter cortisol rhythm over the day, independently of factors that potentially influence cortisol including age, gender, current SES, smoking, BMI, depression, self-rated health, and time of waking in the morning. The flatter cortisol slope was driven primarily by lower cortisol values on waking in participants with greater pathogen burden, rather than raised evening values. Flatter cortisol slopes appear to be associated with increased health risk, and have been related in other studies to coronary disease and abdominal adiposity [41, 42]. A longitudinal study of patients with breast cancer found that flatter cortisol slopes predicted early mortality, independently of covariates [48]. Additionally, flatter cortisol slopes are associated with some aspects of psychological stress and coping; e.g. men and women with marital problems have




Fig. 2. Mean rate of salivary cortisol decrease over the day in participants with infectious burden scores of 0, 1, 2 or 3. The difference was significant after adjustment for age, gender, grade of employment, body mass index, smoking status, self-rated health, cardiovascular medication, depressive symptoms and time of waking in the morning. Error bars are S.E.M.

flatter cortisol slopes [49, 50], while individuals who utilize adaptive coping styles have steeper slopes [51]. By contrast, there was no association with peak cortisol, suggesting that the different components of the cortisol diurnal rhythm have varying associations with health-related biological processes. Previous studies have shown limited associations between cortisol slopes, peak levels, and average output [52].

The mechanisms underlying the association between pathogen burden and cortisol rhythm are not certain. The effect appeared to be driven primarily by CMV seropositivity, although the significant trend across pathogen burden categories indicates that *C. pneumoniae* and HSV-1 also contributed. There are four broad possibilities. First, there could be a direct effect of pathogens on HPA function. Infection with these common pathogens is likely to have taken place early in life [19, 26, 53]. CMV is a potent immunomodulator [54], and is believed to contribute to T-cell immunosenescence [55]. Viral infections are associated with concomitant activation of the HPA, and cytokines released in response to viral infection induce corticotrophin-releasing factor (CRF), have direct effects on adrenocorticotrophic hormone (ACTH) release from the anterior pituitary, and promote glucocorticoid secretion from the adrenal glands [56]. Levels of mRNA in the paraventricular nucleus of the hypothalamus and the peptides corticotrophin-releasing hormone and arginine vasopressin in the median eminence are raised by endotoxin treatment in early life, and may lead to resetting of adult HPA function [57]. HSV-1 may cause down-regulation of glucocorticoid receptors in the hippocampus, thereby

augmenting HPA activation [58]. These responses could lead to chronic HPA dysregulation expressed as a flattened diurnal cortisol profile, although direct mechanistic evidence is currently lacking.

A second possibility is that disturbed cortisol profiles may have preceded infection and increased susceptibility. Even though cortisol was assessed in middle-age whereas infection was probably acquired early in life, the temporal relationship between infection and cortisol cannot be determined in this cross-sectional design. However, administration of glucocorticoids to human cell cultures enhances the growth and replication of CMV and *C. pneumoniae* [37, 38]. Third, exposure to infection in childhood and psychosocial adversity may coexist. Acquisition of a variety of pathogens early in life is greater in children from low SES backgrounds [19, 22, 59], and living in poorer, crowded residences may increase risk of family conflict and maltreatment [3]. These factors can in turn stimulate HPA dysregulation that is manifested in adult life [13, 14]. If this is the case, then infectious burden could be a marker of stressful experiences in early life that lead to HPA dysfunction, and have no direct role. Interestingly, other population studies have shown differential associations between childhood SES and different measures of cortisol rhythm in adult life, consistent with the current findings [12].

A fourth possibility is that the association is due to stress-related increases in antibody titres for latent infections. There is a growing literature relating antibody titres with psychological stress [60, 61], and evidence that cortisol is involved in the up-regulation of viral replication [62–64]. It has recently been shown that HSV-1 antibody levels are heightened in adolescents who experienced severe stress or abuse as children [65]. It is theoretically possible that our results were related to reactivation of pathogens, increasing the likelihood that antibody levels exceed the threshold for seropositivity tests. However, this is unlikely for two reasons. First, the detection thresholds in the tests for serostatus were set low. Second, we found no association between cortisol and the levels of CMV or HSV-1 antibodies in analyses limited to individuals who had positive serostatus (results not presented).

This study has several limitations. The sample was weighted towards men, as is also the case in the larger Whitehall study from which participants were drawn [43]. We have no evidence about when seropositivity for these infectious agents was acquired. We only tested three pathogens, and other organisms

are relevant to pathologies such as CHD, notably periodontal infectious agents [66]. Multiple days of cortisol monitoring would have been desirable, since measures from a single day may not be representative [67]. Despite adjusting statistically for the principle influences on cortisol regulation, unmeasured factors correlated with infection exposure may have affected cortisol output. We were not able to measure ACTH in our study, and this would have been valuable for the understanding of cortisol dynamics. Nor was it possible to conclude whether overall exposure to cortisol throughout the complete diurnal cycle differed between pathogen burden groups, since values were not recorded in the night. Nevertheless, this study on a large well characterized population suggests that cumulative pathogen burden is positively related to cortisol dysregulation. Since early life psychosocial adversity frequently coincides with pathogen exposure, this association may be relevant to the processes through which life experience affects HPA function in health and disease.

ACKNOWLEDGEMENTS

This research was supported by the British Heart Foundation (grant PG/03/029). Västerbotten County Council provided financial support for Å.G. We thank the men and women in the Whitehall II cohort and the Whitehall study team for their participation.

DECLARATION OF INTEREST

None.

REFERENCES

1. Kuh D, Ben-Shlomo Y. *A Life Course Approach to Chronic Disease Epidemiology*, 2nd edn. Oxford: Oxford University Press, 2004.
2. Galobardes B, et al. Systematic review of the influence of childhood socioeconomic circumstances on risk for cardiovascular disease in adulthood. *Annals of Epidemiology* 2006; **16**: 91–104.
3. Poulton R, et al. Association between children's experience of socioeconomic disadvantage and adult health: a life-course study. *Lancet* 2002; **360**: 1640–1645.
4. Dong M, et al. Adverse childhood experiences and self-reported liver disease: new insights into the causal pathway. *Archives of Internal Medicine* 2003; **163**: 1949–1956.
5. Dong M, et al. Insights into causal pathways for ischemic heart disease: adverse childhood experiences study. *Circulation* 2004; **110**: 1761–1766.
6. Evans GW, Kim P. Childhood poverty and health: cumulative risk exposure and stress dysregulation. *Psychological Science* 2007; **18**: 953–957.
7. Taylor SE, et al. Health psychology: what is an unhealthy environment and how does it get under the skin? *Annual Review of Psychology* 1997; **48**: 411–447.
8. Taylor SE, et al. Relationship of early life stress and psychological functioning to adult C-reactive protein in the coronary artery risk development in young adults study. *Biological Psychiatry* 2006; **60**: 819–824.
9. Danese A, et al. Childhood maltreatment predicts adult inflammation in a life-course study. *Proceedings of the National Academy of Sciences USA* 2007; **104**: 1319–1324.
10. Elzinga BM, et al. Diminished cortisol responses to psychosocial stress associated with lifetime adverse events a study among healthy young subjects. *Psychoneuroendocrinology* 2008; **33**: 227–237.
11. Heim C, et al. Pituitary-adrenal and autonomic responses to stress in women after sexual and physical abuse in childhood. *Journal of the American Medical Association* 2000; **284**: 592–597.
12. Li L, et al. Life-time socio-economic position and cortisol patterns in mid-life. *Psychoneuroendocrinology* 2007; **32**: 824–833.
13. Gunnar M, Quevedo K. The neurobiology of stress and development. *Annual Review of Psychology* 2007; **58**: 145–173.
14. Luecken LJ, Lemery KS. Early caregiving and physiological stress responses. *Clinical Psychology Review* 2004; **24**: 171–191.
15. Belmaker RH, Agam G. Major depressive disorder. *New England Journal of Medicine* 2008; **358**: 55–68.
16. Van Cauter E, et al. Effects of gender and age on the levels and circadian rhythmicity of plasma cortisol. *Journal of Clinical Endocrinology and Metabolism* 1996; **81**: 2468–2473.
17. Kirschbaum C, Hellhammer D. Salivary cortisol. In: Fink G, ed. *Encyclopedia of Stress*, 2nd edn, vol. 3. Oxford: Academic Press, 2007, pp. 405–409.
18. Chida Y, Steptoe A. Cortisol awakening response and psychosocial factors: a systematic review and meta-analysis. *Biological Psychology* 2009; **80**: 265–278.
19. Dowd JB, et al. Socioeconomic disparities in the seroprevalence of cytomegalovirus infection in the US population: NHANES III. *Epidemiology and Infection* 2009; **137**: 58–65.
20. Steptoe A, et al. Socioeconomic status, pathogen burden and cardiovascular disease risk. *Heart* 2007; **93**: 1567–1570.
21. Ekesbo R, et al. Combined seropositivity for H. pylori and C. pneumoniae is associated with age, obesity and social factors. *Journal of Cardiovascular Risk* 2000; **7**: 191–195.
22. Xu F, et al. Seroprevalence of herpes simplex virus type 1 in children in the United States. *Journal of Pediatrics* 2007; **151**: 374–377.
23. Aiello AE, et al. Socioeconomic and psychosocial gradients in cardiovascular pathogen burden and immune

- response: the multi-ethnic study of atherosclerosis. *Brain, Behavior and Immunity* 2008 (in press).
24. **Simanek AM, et al.** Persistent pathogens linking socioeconomic position and cardiovascular disease in the US. *International Journal of Epidemiology* 2008 (in press).
 25. **Zajacova A, et al.** Socioeconomic and race/ethnic patterns in persistent infection burden among U.S. adults. *Journals of Gerontology. A: Biological Science and Medical Science* 2009; **64**: 272–279.
 26. **Dowd JB, et al.** Early origins of health disparities: Burden of infection, health, and socioeconomic status in U.S. children. *Social Science and Medicine* 2009; **68**: 699–707.
 27. **Smith JS, Robinson NJ.** Age-specific prevalence of infection with herpes simplex virus types 2 and 1: a global review. *Journal of Infectious Disease* 2002; **186** (Suppl. 1): S3–28.
 28. **McQuillan GM, et al.** Racial and ethnic differences in the seroprevalence of 6 infectious diseases in the United States: data from NHANES III, 1988–1994. *American Journal of Public Health* 2004; **94**: 1952–1958.
 29. **Malaty HM, et al.** Age at acquisition of *Helicobacter pylori* infection: a follow-up study from infancy to adulthood. *Lancet* 2002; **359**: 931–935.
 30. **Aiello AE, et al.** The influence of latent viral infection on rate of cognitive decline over 4 years. *Journal of the American Geriatric Society* 2006; **54**: 1046–1054.
 31. **Sorlie PD, et al.** A prospective study of cytomegalovirus, herpes simplex virus 1, and coronary heart disease: the atherosclerosis risk in communities (ARIC) study. *Archives of Internal Medicine* 2000; **160**: 2027–2032.
 32. **Espinola-Klein C, et al.** Impact of infectious burden on extent and long-term prognosis of atherosclerosis. *Circulation* 2002; **105**: 15–21.
 33. **Zhu J, et al.** Prospective study of pathogen burden and risk of myocardial infarction or death. *Circulation* 2001; **103**: 45–51.
 34. **Georges JL, et al.** Impact of pathogen burden in patients with coronary artery disease in relation to systemic inflammation and variation in genes encoding cytokines. *American Journal of Cardiology* 2003; **92**: 515–521.
 35. **Kiechl S, et al.** Chronic infections and the risk of carotid atherosclerosis: prospective results from a large population study. *Circulation* 2001; **103**: 1064–1070.
 36. **Shanks N, et al.** Early-life exposure to endotoxin alters hypothalamic-pituitary-adrenal function and predisposition to inflammation. *Proceedings of the National Academy of Sciences USA* 2000; **97**: 5645–5650.
 37. **Komura H, et al.** Effects of antiasthma drugs on the growth of *Chlamydia pneumoniae* in HEp-2 cells. *Journal of Infection and Chemotherapy* 2003; **9**: 160–164.
 38. **Tanaka J, et al.** Enhanced replication of human cytomegalovirus in human fibroblasts treated with dexamethasone. *Journal of General Virology* 1984; **65**: 1759–1767.
 39. **Vedhara K, et al.** Relationship of early childhood illness with adult cortisol in the Barry Caerphilly Growth (BCG) cohort. *Psychoneuroendocrinology* 2007; **32**: 865–873.
 40. **Wirtz PH, et al.** Evidence for altered hypothalamus-pituitary-adrenal axis functioning in systemic hypertension: blunted cortisol response to awakening and lower negative feedback sensitivity. *Psychoneuroendocrinology* 2007; **32**: 430–436.
 41. **Matthews K, et al.** Diurnal cortisol decline is related to coronary calcification: CARDIA study. *Psychosomatic Medicine* 2006; **68**: 657–661.
 42. **Rosmond R, et al.** Stress-related cortisol secretion in men: relationships with abdominal obesity and endocrine, metabolic and hemodynamic abnormalities. *Journal of Clinical Endocrinology and Metabolism* 1998; **83**: 1853–1859.
 43. **Marmot MG, et al.** Health inequalities among British civil servants: the Whitehall II study. *Lancet* 1991; **337**: 1387–1393.
 44. **Idler EL, Benyamini Y.** Self-rated health and mortality: a review of twenty-seven community studies. *Journal of Health and Social Behavior* 1997; **38**: 21–37.
 45. **Steptoe A, et al.** Presence of human heat shock protein (Hsp) 60 is associated with decreased seropositivity to pathogens. Is circulating Hsp60 protective against infection? *Clinical and Vaccine Immunology* 2007; **14**: 204–207.
 46. **Dockray S, et al.** The cortisol awakening response in relation to objective and subjective measures of waking in the morning. *Psychoneuroendocrinology* 2008; **33**: 77–82.
 47. **Pruessner JC, et al.** Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change. *Psychoneuroendocrinology* 2003; **28**: 916–931.
 48. **Sephton SE, et al.** Diurnal cortisol rhythm as a predictor of breast cancer survival. *Journal of the National Cancer Institute* 2000; **92**: 994–1000.
 49. **Adam EK, Gunnar MR.** Relationship functioning and home and work demands predict individual differences in diurnal cortisol patterns in women. *Psychoneuroendocrinology* 2001; **26**: 189–208.
 50. **Barnett RC, et al.** Marital-role quality and stress-related psychological indicators. *Annals of Behavioral Medicine* 2005; **30**: 36–43.
 51. **O'Donnell K, et al.** Psychological coping styles and cortisol over the day in healthy older adults. *Psychoneuroendocrinology* 2008; **33**: 601–611.
 52. **Stone AA, et al.** Individual differences in the diurnal cycle of salivary free cortisol: a replication of flattened cycles for some individuals. *Psychoneuroendocrinology* 2001; **26**: 295–306.
 53. **Xu F, et al.** Trends in herpes simplex virus type 1 and type 2 seroprevalence in the United States. *Journal of the American Medical Association* 2006; **296**: 964–973.
 54. **Mocarski Jr. ES.** Immunomodulation by cytomegaloviruses: manipulative strategies beyond evasion. *Trends in Microbiology* 2002; **10**: 332–339.

55. **Koch S, et al.** Cytomegalovirus infection: a driving force in human T cell immunosenescence. *Annals of the New York Academy of Sciences* 2007; **1114**: 23–35.
56. **Silverman MN, et al.** Immune modulation of the hypothalamic-pituitary-adrenal (HPA) axis during viral infection. *Viral Immunology* 2005; **18**: 41–78.
57. **Shanks N, et al.** Neonatal endotoxin exposure alters the development of the hypothalamic-pituitary-adrenal axis: early illness and later responsivity to stress. *Journal of Neuroscience* 1995; **15**: 376–384.
58. **Bener D, et al.** Glucocorticoid resistance following herpes simplex-1 infection: role of hippocampal glucocorticoid receptors. *Neuroendocrinology* 2007; **85**: 207–215.
59. **Dowd JB, et al.** Socioeconomic gradients in immune response to latent infection. *American Journal of Epidemiology* 2008; **167**: 112–120.
60. **Glaser R, et al.** Stress-related activation of Epstein-Barr virus. *Brain, Behavior and Immunity* 1991; **5**: 219–232.
61. **Phillips AC, et al.** Cytomegalovirus is associated with depression and anxiety in older adults. *Brain, Behavior and Immunity* 2008; **22**: 52–55.
62. **Glaser R, et al.** Hormonal modulation of Epstein-Barr virus replication. *Neuroendocrinology* 1995; **62**: 356–361.
63. **Cacioppo J, et al.** Autonomic and glucocorticoid associations with the steady-state expression of latent Epstein-Barr virus. *Hormones and Behavior* 2002; **42**: 32–41.
64. **Sainz B, et al.** Stress-associated immunomodulation and herpes simplex virus infections. *Medical Hypotheses* 2001; **56**: 348–356.
65. **Shirtcliff EA, et al.** Early childhood stress is associated with elevated antibody levels to herpes simplex virus type 1. *Proceedings of the National Academy of Sciences USA* 2009; **106**: 2963–2967.
66. **Spahr A, et al.** Periodontal infections and coronary heart disease: role of periodontal bacteria and importance of total pathogen burden in the Coronary Event and Periodontal Disease (CORODONT) study. *Archives of Internal Medicine* 2006; **166**: 554–559.
67. **Hellhammer J, et al.** Several daily measurements are necessary to reliably assess the cortisol rise after awakening: state- and trait components. *Psychoneuroendocrinology* 2007; **32**: 80–86.