

1 **Molecular evidence of Pleistocene bidirectional faunal exchange between**
2 **Europe and the Near East: the case of the bicolored shrew (*Crocidura***
3 ***leucodon*, Soricidae)**

4
5 **Sylvain Dubey, * Jean-François Cosson,‡ Vladimir Vohralík, † Boris Kryštufek, § Ebru**
6 **Diker¶ & Peter Vogel ***

7
8 * *Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland*

9 ‡ *Centre de Biologie et de Gestion des Populations (CBGP), INRA-EFPA, Campus*

10 *International de Baillarguet, CS 30016, 34988 Montferrier / Lez cedex, France*

11 † *Department of Zoology, Charles University, Viničná 7, CZ-128 44 Prague 2, Czech*

12 *Republic*

13 § *Science and Research Centre of Koper, University of Primorska, Garibaldijeva 18, SI-6000*

14 *Koper, Slovenia*

15 ¶ *Department of Biology, Faculty of Art and Sciences, Trakya University, Edirne, Turkey*

16
17 Running title: Phylogeography of the bicolored shrew

18
19 Correspondence:

20 Sylvain Dubey, Département d'Ecologie et d'Evolution, Bâtiment du Biophore, Université de
21 Lausanne, 1015 Lausanne, Switzerland

22 Phone: +41-21-692-4163; Fax: +41-21-692-4165; e-mail: sylvain.dubey@unil.ch

1 **Abstract**

2 We sequenced 1077 bp of the mitochondrial cytochrome *b* gene and 511 bp of the nuclear
3 Apolipoprotein B gene in bicolored shrew (*Crocidura leucodon*, Soricidae) populations
4 ranging from France to Georgia. The aims of the study were to identify the main genetic
5 clades within this species and the influence of Pleistocene climatic variations on the
6 respective clades. The mitochondrial analyses revealed a European clade distributed from
7 France eastwards to northwestern Turkey and a Near East clade distributed from Georgia to
8 Romania; the two clades separated during the Middle Pleistocene. We clearly identified a
9 population expansion after a bottleneck for the European clade based on mitochondrial and
10 nuclear sequencing data; this expansion was not observed for the eastern clade. We
11 hypothesise that the western population was confined to a small Italo-Balkan refugium,
12 whereas the eastern population subsisted in several refugia along the southern coast of the
13 Black Sea.

14 *Keywords:* *Crocidura leucodon*; Cytochrome *b* gene; Apolipoprotein B gene;
15 Phylogeography.

16

17

18

1 **Introduction**

2 The impact of Pleistocene climatic fluctuations on European biota is well documented.
3 Studies of mitochondrial DNA markers reveal general patterns involving southern refugia and
4 northern recolonisation routes, as a modification of genetic diversity of terrestrial biota
5 throughout the Holarctic (Hewitt, 2000, 2004a,b). Populations were isolated in different glacial
6 refugia by barriers such as mountains and seas, leading to various recolonisation patterns
7 (e.g., Taberlet *et al.*, 1998; Hewitt, 1999). Widely accepted refugia include the Iberian
8 Peninsula in the west and the Italo-Balkan region in the east (e.g., Thorpe, 1984; Ferris *et*
9 *al.*, 1993, 1998; Dumolin-Lapegue *et al.*, 1997; Santucci *et al.*, 1998; Taberlet *et al.*, 1998;
10 Dubey *et al.*, 2006); however, a number of phylogeographical studies of various taxa have
11 revealed an additional pattern of colonisation of Europe by populations originating from
12 eastern areas such as the Caucasus, southern Urals, and western Asia (Cooper *et al.*, 1995;
13 Bilton *et al.*, 1998; Nesbo *et al.*, 1999; Palme & Vendramin, 2002; Seddon *et al.*, 2002;
14 Michaux *et al.*, 2004; Culling *et al.*, 2006; Dubey *et al.*, 2006). This pattern of post-glacial
15 recolonisation appears to be more common than previously suspected and affected probably a
16 large range of taxa.

17 With regard to population dynamics, few studies have detailed the impact of
18 Pleistocene climatic oscillations, e.g., the dating of population expansions and identifying
19 differences in patterns among clades of different geographic origin (Michaux *et al.*, 2004;
20 Brändli *et al.*, 2005; Culling *et al.*, 2006; Dubey *et al.*, 2006; Dubey *et al.*, in press a; Koch *et*
21 *al.*, 2006; Marmi *et al.*, 2006; Vörös *et al.*, 2006). In the same way, the majority of previous
22 studies are based solely on mitochondrial DNA, whereas the inclusion of uniparentally and
23 biparentally inherited markers present in nuclear genomes can enhance our understanding of
24 population history (Brändli *et al.*, 2005; Dubey *et al.*, 2006).

1 In the present study, we focused on the bicolored shrew, *Crocidura leucodon*
2 (Hermann, 1780), which is a strictly western Eurasian species, distributed from France to
3 central and southern Europe to Turkey and Georgia. This species seems to be an interesting
4 candidate for a phylogeographic study, as Vogel *et al.* (2003), based on only two samples,
5 from Switzerland and Turkey, noted a substantial Kimura 2-paramters genetic distance of
6 3.8% between them for the Cytochrome *b* gene. In addition, Poulakakis *et al.* (2005) analysed
7 three Greek samples (from Peloponnesus and Lesvos) obtained from owl pellets, based on the
8 same gene, promoting such material as usable for phylogeographic studies, without giving
9 any information concerning the distances between haplotypes. Nevertheless, the intraspecific
10 genetic distances were considered as low.

11 We tested (i) in which way the Bosphorus strait has isolated the European and Near
12 East populations, and (ii) in which way the Pleistocene climatic fluctuations have had
13 different impacts on the western and eastern populations, such as bottleneck effects. To
14 resolve these questions, we analysed sequence data from mitochondrial and nuclear markers
15 (Cytochrome *b* and Apolipoprotein B) and considered the molecular clock.

16 17 **Material and methods**

18 **Sampling**

19 We analysed 60 samples of *Crocidura leucodon* collected from France to Georgia
20 (Fig. 1 and Table 1), three other Eurasian crocidurine taxa, and as an out-group a soricine,
21 *Sorex minutus*. This set of samples (Table 1) included material from the collections of
22 Lausanne (IZEA), Switzerland; Prague (DZCU), Czech Republic; and Ljubljana (PMS),
23 Slovenia. Some sequences were taken from Ohdachi *et al.* (2004), Fontanillas *et al.* (2005)
24 and Dubey *et al.* (2006). We also used samples from Poulakakis *et al.* (2005) that were treated
25 separately.

26 **DNA extraction and amplification of Cytochrome *b* and Apolipoprotein B genes**

1 Samples (livers) from IZEA collection were first frozen in the field in liquid nitrogen
2 and kept for several years at -70°C before being stored in ethanol until DNA extraction.
3 Samples from the other collections were directly stored in ethanol. DNA extraction was
4 carried out using the QIA Amp DNA Mini Kit (Qiagen). Double-stranded DNA
5 amplifications of the mitochondrial cytochrome *b* gene (*cyt-b*) were performed with the
6 primer pairs L14724/H15149, C1/C2, C3/H15915, and L14724/H15915 (Irwin *et al.*, 1991;
7 Dubey *et al.*, 2006). Amplification of the Apolipoprotein B (*ApoB*) nuclear genes was
8 performed using the primer pairs ApoBf/ApoBr (Dubey *et al.*, in press b). Amplification
9 conditions for the *ApoB* consisted of 40 cycles of 45 s denaturation at 94°C , 45 s annealing at
10 50°C , and 90 s extension at 72°C ; for the *cyt-b*, the conditions of Dubey *et al.* (2006) were
11 used.

12 PCR products were checked on a 1% agarose gel and then purified using the QIAquick
13 PCR Purification Kit (QIAGEN) following the manufacturer's instructions. DNA sequencing
14 was performed in a total volume of 10 μl containing 1-3 μl of amplified PCR product, 1 μl of
15 10 μM primer, and 4 μl of ABI PRISMTM Dye Terminator 1 (Perkin-Elmer). Sequence
16 reactions were visualised on an ABI 3100 genetic analyser (Applied Biosystems, USA).

17 **Phylogenetic methods**

18 Nucleotide sequences of *cyt-b* and *ApoB* genes were edited with Sequence Navigator
19 (Parker, 1997) and manually aligned. Two methods of phylogenetic analyses were carried out
20 for *cyt-b*, using PAUP*version 4.0b10 PPC (Swofford, 1998). Tests were conducted on the
21 complete fragment (1077 bp), all codon positions were used, and trees were rooted using
22 sequences of *Sorex minutus* (DQ630379). A Neighbour Joining (NJ) tree was constructed
23 using the general time reversible (GTR; Rodriguez *et al.*, 1990) genetic distance, which was
24 selected previously using Modeltest 3.06 according to the protocol of Posada & Crandall
25 (1998). The Parsimony analyses (MP) were performed using the following options: heuristic

1 search, stepwise-addition of sequences, 10 replicates of random addition of taxa, and TBR
2 branch swapping (Swofford, 1998); all codon positions were equally weighted. Bootstrap
3 support values were obtained with 1000 pseudo-replicates and 10 random replicates of
4 stepwise-addition sequences. Fast maximum likelihood (ML) heuristic searches and bootstrap
5 analyses (1000 replicates) were performed using PHYML (Guindon & Gascuel, 2003) with a
6 GTR model.

7 **Nucleotide diversity and genetic structure**

8 The nucleotide diversities (π) of *cyt-b* and *ApoB* and the population genetic structure
9 for *cyt-b* (analysis of molecular variance, AMOVA) were estimated using Arlequin version
10 2.0 Software (Schneider, 2000). AMOVA was performed at two different hierarchical levels:
11 among clades and within clades.

12 **Molecular clock**

13 The molecular clock hypothesis was tested following the method of Posada and
14 Crandall (1998). Estimation of divergence time from the molecular data was performed
15 according to the calibration developed for *cyt-b* in Soricidae by Fumagalli *et al.* (1999), based
16 on an estimate of 20 Myr for the split between the Soricinae and the Crocidurinae and on the
17 number of third-position transversions observed. Nevertheless, it could not be used directly
18 because of the low number of third-position transversions observed at a specific level within
19 the family. Thus, we used the calibration of Dubey *et al.* (2006), based on the previous study
20 that allowed a better estimated, due to the use of a ML distance including all the codon
21 positions with a divergence rate of 0.057 ML distance/Myr (95% CI: 0.044-0.070).

22 **Isolation with migration**

23 We used the “isolation with migration” (or IM, Nielsen and Wakeley 2001) model to
24 date the divergence between populations of both side of the Marmara Sea within each clade.
25 As suggested by Hey (2007), we started the simulations using a burn-in of 100,000 steps

1 followed by a ½ hour run with maximum values for theta, m and t arbitrary set to 10 and the –
2 J1 run option. From there, appropriate values for priors were chosen. In each following runs,
3 plots of parameter trend lines were systematically consulted for assessing how well the
4 Markov chain was exploring the parameter space. Convergence by the Markov chain
5 simulations was assessed by monitoring three independent chains, and by assessing the
6 autocorrelation of parameter values over the course of each run. Individual simulations were
7 run for 10 million updates or more. For each of the demographic parameters, we recorded the
8 marginal density. The peaks of the resulting distributions were taken as estimates of the
9 parameters and the 90% highest posterior density (HPD) interval was taken for the credibility
10 intervals. Estimates with IM are scaled by the overall neutral mutation rate per gene per
11 generation. We assigned an inheritance scalar of 0.25 as usual for mtDNA and assumed 1
12 year per generation. To convert parameter estimates to time scale units, we used the average
13 rate of evolution for the *cyt-b* gene in *Crocidura* species from Dubey et al. (2006),
14 corresponding to $3.07E^{-5}$ (95% CI: $2.37E^{-5}$ - $3.77E^{-5}$) mutation events/locus/year.

15 **Expansion time**

16 To test the hypothesis of recent population growth from a low-diversity founder
17 population within the different clades, several tests were performed for *cyt-b*. We used three
18 methods implemented in Arlequin version 2.0 (Schneider, 2000). The first method, Fu's
19 (1997) F_S statistic, tests the probability of having no fewer than the number of observed
20 alleles in the sample given that θ (heterozygosity per sites) = π . This statistic tends to be
21 negative when there is an excess of recent mutations (or rare alleles). The second method,
22 Tajima's (1989) D statistic, tests the null hypothesis that two estimates of the neutral mutation
23 parameter, one derived from the average number of pairwise nucleotide differences and the
24 other based on the number of segregating sites in the sample, are equal. In the third test,
25 pairwise mismatch distributions among individuals were plotted and tested for goodness-of-fit

1 to a model of sudden expansion using parametric bootstrapping with 1000 replicates
2 (Schneider & Excoffier, 1999). Expansion time after the bottleneck was estimated from the
3 mismatch distribution (τ) (Rogers, 1995) and uncorrected distances (p). Evolutionary rate for
4 uncorrected (p) distance was estimated using the molecular clock developed by Fumagalli *et*
5 *al.* (1999) with a divergence rate of 0.061 uncorrected distance (p)/Myr (95% CI: 0.054-
6 0.069).

7

8 **Results**

9 **Cytochrome *b* gene**

10 The 60 *C. leucodon* samples showed 41 different haplotypes of 1077 bp and contained
11 366 variable sites, of which 218 were parsimony-informative. No insertions or deletions were
12 observed. As the three phylogenetic methods gave identical arrangements of the main
13 branches, the relationship between haplotypes is given only for the ML analysis in Fig. 2.

14 The *Crocidura leucodon* samples formed a monophyletic unit (all bootstrap values of
15 100%), well differentiated from the other Eurasian species (Fig. 2). Within this unit, two
16 major and strongly supported clades were found (all bootstrap values $\geq 80\%$).

17 Clade I (western clade) included European samples from Czech Republic, France, Germany,
18 Greece (Lesvos), Italy, Serbia, Slovakia, Slovenia, Switzerland, and samples from
19 northwestern Anatolia, Turkey. Clade II (eastern clade) included the European samples from
20 Bulgaria and Romania and the samples from Georgia and Anatolia (with the exception of
21 westernmost Anatolia).

22 AMOVA showed that the majority of mtDNA variation (84.09%) is distributed
23 between the two clades of *C. leucodon*; only a small percentage of this variation (15.91%) is
24 observed within clades. Mean pair-wise GTR distance between clades is 3.94%. The mean

1 pairwise GTR distances within clades and nucleotide diversities were 0.65% and 0.057 for the
2 western clade, respectively, and 0.71% and 0.059 for the eastern clade.

3 Three additional sequences from Lesvos, Stymfalia, and Parnitha (Greece) described
4 by Poulakakis *et al.* (2005) belong to the western clade, but were not included in our
5 phylogenetic analyses. The analysis of these data (AY452166, AY452176, AY452165)
6 revealed an unexpected result. The mean GTR distance between our samples and the
7 Poulakakis *et al.* samples from Lesvos was 2.8%. In comparing the sequences, we observed a
8 large number of mutations at the beginning and end of their *cyt-b* sequence. When the last 13
9 bp of their sequence were deleted, the mean GTR distance fell to 0.89%, and this was further
10 reduced to 0.42% by deleting the first 48 bp of their sequence (see Table 2 for the details of
11 the mutations observed at the beginning and the end of the sequences of Poulakakis *et al.*). In
12 addition, the mean GTR distance between the Poulakakis *et al.* samples from Lesvos and
13 those from continental Greece was 2.15%, whereas the mean distance between our samples
14 from Lesvos and the Poulakakis *et al.* samples was 6.22%; this value is still much greater than
15 the mean GTR distance between all of our sequences from this clade (0.65%) and the mean
16 GTR distance between the two major clades (western and eastern) obtained in our analyses
17 (3.94%, Fig. 2).

18 A likelihood ratio test led to the acceptance of the molecular clock hypothesis for the
19 whole sample ($df = 43$, $\chi^2 = 53.44$, $P = 0.13$). – Ln Likelihood values with and without the
20 molecular clock assumption are 4169.79 and 4143.07, respectively, for the best trees. On the
21 basis of the calibration of Fumagalli *et al.* (1999) and Dubey *et al.* (2006), we estimated the
22 divergence time between the western and eastern clade to be 0.691 Myr (95% CI: 0.510-
23 0.980; Fig. 2).

24 The IM model has six demographic parameters but we were particularly interested in
25 three of them: t , the time of population splitting (in generation) in the past, and m_1 and m_2 ,

1 the gene-flow rates per gene copy per generation. Once priors were correctly defined, we ran
2 three runs for each clade. The different runs revealed unambiguous marginal posterior
3 probability distributions of the parameters in each clade. The migration parameters revealed a
4 peak at the lower limit of resolution in both directions (from East to West and vice versa) in
5 both clades, as expected given that the Marmara Sea is a nonnegotiable barrier to dispersal
6 once field with water (during interglacials). We hereafter interpret the locations of these peaks
7 as being at zero and then simulated three more runs for each clade fixing m_1 and m_2 to zero
8 for estimating the time of the splits. The marginal posterior probability distribution of t
9 revealed a sharp peak at 0.525 and 1.785 for the Western and the Eastern clade respectively
10 (Fig. 3). When converted to a scale of years, the divergence times between both sides of the
11 Marmara Sea were estimated to be 17,100 yr (90% HPD interval: 1,700-40,600) for the
12 Western clade, and 58,200 yr (90% HPD interval: 16,100-109,000) for the Eastern one.

13 We observed a non-significant P -value for the mismatch distribution test of goodness-
14 of-fit for the western clade (*Harpending's Raggedness index* = 0.02, P = 0.28) and a
15 significant P -value for the eastern clade (*Harpending's Raggedness index* = 0.12, P = 0.03;
16 Table 2). The frequency of the mean pairwise difference between haplotypes showed a bell-
17 shaped distribution for the western clade, contrasting with the more complex distribution
18 obtained for the eastern clade (Fig. 1). Fu's F_S statistics and Tajima's D were significant (F_S =
19 -10.68 , P < 0.001; D = -1.48 , P = 0.04; Table 3) for the western clade but not significant for
20 the eastern (F_S = -2.02 , P = 0.16; D = -0.67 , P = 0.26); consequently, we inferred a scenario
21 of expansion for the western population and non-expansion for the eastern population.

22 The timing of expansions was estimated from the mismatch distribution according to
23 the method proposed by Rogers (1995; Fig. 1B, C). The τ value for the western population
24 was 6.71 (95% CI: 4.19-8.39). Assuming no saturation of uncorrected distances (p), as shown
25 in Dubey *et al.* (2006), distance was 0.0612 per million years (95% CI: 0.054-0.069). With a

1 generation time of 1 year, the population expansion time was estimated to be 51,900 years
2 (95% CI: 28,100-72,100).

3 **Apolipoprotein B gene**

4 The 44 analysed samples showed four different *ApoB* alleles of 511 bp, named A1 to
5 A4 (Genbank accession: EF011555-EF011558), all the alleles differing from each others by
6 only one mutation at the position 62, 96 and 144 of our sequence alignment. Consequently,
7 the different alleles of heterozygous samples were easily determined, as only one mutation
8 was observed between alleles. The western mitochondrial clade samples were all homozygous
9 for the allele A1 (Table 1). In contrast, thirteen A1 homozygotes, two A1/A2 compound
10 heterozygotes, one A1/A3 compound heterozygote, one A1/A4 compound heterozygote, and
11 one A4 homozygote were found in the eastern mitochondrial clade. Consequently, nucleotide
12 diversities were 0.0000 for the western clade and 0.0006 for the eastern clade.

13

14 **Discussion**

15 **Quality of samples**

16 The integration of our *cyt-b* sequences with those of Ohdachi *et al.* (2004), Fontanillas *et al.*
17 (2005) and Dubey *et al.* (2006), analysed from preserved tissues, showed expected genetic
18 distances between samples. In contrast, the material of *C. leucodon* from owl pellets analysed
19 by Poulakakis *et al.* (2005) showed GTR distances with our *C. leucodon* samples much higher
20 than expected, even between samples from the same island (Lesvos; mean GTR distance:
21 2.8%). A distance that fell to 0.42% by deleting the first and the last bp of their sequence from
22 Lesvos. Moreover, no non-synonymous mutations were observed in the *cyt-b* gene for our *C.*
23 *leucodon* samples, whereas mutations at the end of the Poulakakis *et al.* (2005) sequences
24 resulted in two amino-acid substitutions; thus, these mutations appear to be highly suspect.
25 These mutations may in fact be artefacts of DNA sequencing as a result of the poor template

1 quality obtained from the owl pellets (Taberlet and Fumagalli, 1996; Waits and Paetkau,
2 2005). This reinforces the value of repeating the DNA extraction and/or the analysis for each
3 non-invasive sample several times in order to validate the sequencing result (Waits and
4 Paetkau, 2005). Consequently, the affirmation of Poulakakis *et al.* (2005), concerning the use
5 of their samples for phylogenetic and phylogeographical studies of small mammals, has to be
6 questioned.

7 **Biogeography**

8 Based on our mitochondrial phylogenetic analyses, *Crocidura leucodon* populations
9 are divided in two main mitochondrial clades (mean GTR distance between clades: 3.94%;
10 Fig. 1A & 2). The first clade (western clade) includes samples from western and central
11 Europe, and unexpectedly, two samples from Lesvos Island and three from northwestern
12 Anatolia, to which the Greek samples (Lesvos, Parnitha, and Stymfalia) from Poulakakis *et*
13 *al.* (2005; results not shown) should be added. The second clade (eastern clade) includes
14 samples from Bulgaria, Romania, Anatolia (with the exception of westernmost Anatolia) and
15 Georgia. These two clades could represent the chromosomal differences observed between the
16 Georgian samples and those from Czech Republic and Lesvos that were detected in the
17 karyotype analyses undertaken by Biltueva *et al.* (2001).

18 The separation between these two mitochondrial lineages of *C. leucodon* occurred in
19 the Middle Pleistocene, 0.691 Myr (95% CI: 0.510-0.980), the period immediately following
20 the Günz glacial events (790,000-950,000 years BP). This suggests the isolation of
21 populations by submergence of the Bosphorus Strait (between the Black Sea and the Marmara
22 Sea) with increasing sea levels following the glacial period.

23 Sedimentological and palaeontological evidence reveals that the Bosphorus Strait has
24 alternatively submerged and emerged since the Middle Pleistocene before being completely
25 submerged from the Mid-Late Holocene (Kerey *et al.*, 2004). Consequently, the fact that we

1 observed a lack of clear structure between samples situated on either side of the Bosphorus
2 Strait indicates that south-eastern Europe was probably colonised by the eastern
3 mitochondrial lineages during a recent land bridge connection between Europe and the Near
4 East (Late Pleistocene). A hypothesis confirmed by the migration analyses that estimated this
5 event of vicariance to the Upper Pleistocene (58,200 years BP; 90% HPD: 16,100 to
6 109,000). Based on similar results, the colonisation of western Anatolia by the western
7 mitochondrial lineage occurred during the same period (17,100 years BP; 90% HPD: 1,700 to
8 40,600; Late Pleistocene-Holocene). Thus, this strait appears to be a permeable biogeographic
9 barrier for *C. leucodon*. This permeability has already been demonstrated in classical
10 zoogeography for various species (e.g., Hosey, 1982) on the basis of unidirectional migrations
11 from Europe to Anatolia (Hosey, 1982; Filippucci & Simson, 1996; Kryštufek, 2002).
12 However, our study demonstrate for the first time a Late Pleistocene-Holocene bidirectional
13 exchange of two different conspecific lineages between Europe and the Near East, whereas
14 previous studies have only revealed unidirectional colonisation.

15 The last climatic fluctuations of the Upper Pleistocene (126,000-11,500 years BP;
16 Ogg, 2004) had contrasting impacts on the western and eastern mitochondrial clades. For the
17 western clade, the bell-shaped curve of the mismatch distributions (Fig. 1B) of the *cyt-b* gene
18 indicates an expansion following a bottleneck 50,800 years ago (95% CI: 28,100-72,100).
19 This finding reveals that the last glacial maximum of the Pleistocene (22,000 years BP) had a
20 moderate impact on these small mammal populations. Conversely, no sign of expansion was
21 detected for the eastern mitochondrial clade of *C. leucodon*, which is consistent with the
22 absence of nuclear polymorphism in the western clade (π : 0.0000), whereas four alleles are
23 present in the eastern (π : 0.0006).

24 Thus, the European population appears to have been reduced to a small number of
25 individuals during the last glaciations, probably confined to within a small Italo-Balkan

1 refugium. Conversely, the eastern clade appears to have persisted with a greater population
2 size and probably in several refugia around the Black Sea. This pattern has also been
3 suggested for mammals and plants (Michaux *et al.*, 2004; Heuertz *et al.*, 2006; Kučera *et al.*,
4 2006), based on a higher genetic diversity observed in Anatolian populations compared to
5 European ones. Moreover, these results are also supported by palynological data that indicate
6 that open wooded cover in this area persisted through the full glacial condition of the
7 Pleistocene (Tarasov *et al.*, 2000).

8 **Conclusion**

9 Based on molecular data, we confirmed that the Bosphorus Strait was a permeable
10 barrier. This finding has been suggested previously by the colonisation of western Anatolia by
11 European populations of the lesser white-toothed shrew in the Lower Pleistocene (*C.*
12 *suaveolens*; Dubey *et al.*, 2006) and more recently by the hedgehog (Filippucci & Simson,
13 1996; Kryštufek, 2002). Moreover, we provide evidence that European populations of the
14 bicolored shrew may be the source of recent West Anatolian populations.

15 We also found a marked difference in population history between the two divergent
16 mitochondrial lineages of bicolored shrew, suggesting that Pleistocene climatic variations
17 have more strongly reduced the genetic diversity of the European population than that of the
18 Near East population. We hypothesise that this could be a general pattern for fauna and flora,
19 as other studies comparing these geographical areas have reported similar differences
20 (Michaux *et al.*, 2004; Heuertz *et al.*, 2006; Kučera *et al.*, 2006).

21

22 **Acknowledgments**

23 We thank N. Di Marco for laboratory facilities, C. Ohayon for scientific and linguistic advice
24 and M. Fontaine for noticeable advice in using IM software. We are grateful to P.

25 Bokenhagen, P. Brunnet-Lecompte, H. Brünner, F. Catzeflis, P. Müller, R. Parapanov, C.

1 Vogel, and M. Zaitsev for tissues samples, and to A. Andreeva, S. Uzunov (Bulgaria), C.
2 Kurtonur (Turkey), D. Murariu (Romania) for integrating us (PV) in local fieldwork for
3 trapping shrews. The study was partly sponsored by GACR (grant 206/05/2334) and MSMT
4 CR (project 0021620828).

5

6 **References**

- 7 Bilton, D.T., Mirol, P.M., Mascheretti, S. Fredga, K., Zima, J. & Searle J.B. 1998.
8 Mediterranean Europe as an area of endemism for small mammals rather than a source
9 for northwards postglacial colonization. *Proc. R. Soc. Lond., Ser. B Biol. Sci.* **265**:
10 1219-1226.
- 11 Biltueva, L.S., Rogatcheva, M.B., Perelman, P.L., Borodin P.M., Oda, S.I., Koyasu, K.,
12 Harada, M., Zima, J. & Graphodatsky, A.S. 2001. Chromosomal phylogeny of certain
13 shrews of the genera *Crocidura* and *Suncus* (Insectivora). *J. Zool. Syst. Evol. Res.* **39**:
14 69-76.
- 15 Brändli, L., Handley, L.J., Vogel, P. & Perrin, N. 2005. Evolutionary history of the greater
16 white-toothed shrew (*Crocidura russula*) inferred from analysis of mtDNA, Y and X
17 chromosome markers. *Mol. Phylogenet. Evol.* **37**: 832-844.
- 18 Cooper, S.J.B., Ibrahim, K.M. & Hewitt, G.M. 1995. Postglacial expansion and genome
19 subdivision in the European Grasshopper *Chorthippus parallelus*. *Mol. Ecol.* **4**: 49-60.
- 20 Culling, M.A., Janko, K., Boron, A., Vasil'ev, V.P., Cote, I.M. & Hewitt, G.M. 2006.
21 European colonization by the spined loach (*Cobitis taenia*) from Ponto-Caspian refugia
22 based on mitochondrial DNA variation. *Mol. Ecol.* **15**: 173-190.
- 23 Dubey, S., Cosson, J.-F., Magnanou, E, Vohralík, V., Benda, P., Frynta, D., Hutterer, R.,
24 Vogel, V & Vogel, P. Mediterranean populations of the lesser white-toothed shrew

- 1 (*Crocidura suaveolens* group): an unexpected puzzle of Pleistocene survivors and
2 prehistoric introductions. *Mol. Ecol., in press a.*
- 3 Dubey, S., Salamin, N., Ohdachi, S.D., Barriere, P. & Vogel, P. Molecular phylogenetics of
4 shrews (Mammalia: Soricidae) reveal timing of transcontinental colonizations. *Mol.*
5 *Phylogenet. Evol., in press b.*
- 6 Dubey, S., Zaitsev, M., Cosson, J.-F., Abdulkadier, A. & Vogel, P. 2006. Pliocene and
7 Pleistocene diversification and multiple refugia in a Eurasian shrew (*Crocidura*
8 *suaveolens* group). *Mol. Phylogenet. Evol.* **38**: 635-647.
- 9 Dumolin-Lapegue, S., Demesure, B., Fineschi, S., LeCorre, V. & Petit, R.J. 1997.
10 Phylogeographic structure of white oaks throughout the European continent. *Genetics*
11 **146**: 1475-1487.
- 12 Ferris, C., Oliver, R.P., Davy, A.J. & Hewitt, G.M. 1993. Native oak chloroplasts reveal an
13 ancient divide across Europe. *Mol. Ecol.* **2**: 337-344.
- 14 Ferris, C., King, R.A., Vainola, R. & Hewitt, G.M. 1998. Chloroplast DNA recognizes three
15 refugial sources of European oaks and suggests independent eastern and western
16 immigrations to Finland. *Heredity* **80**: 584-593.
- 17 Filippucci, M.G. & Simson, S. 1996. Allozyme variation and divergence in Erinaceidae
18 (Mammalia, Insectivora). *Israel Journal of Zoology* **42**: 335-345.
- 19 Fontanillas, P., Depraz, A., Giorgi, M.S. & Perrin, N. 2005. Nonshivering thermogenesis
20 capacity associated to mitochondrial DNA haplotypes and gender in the greater white-
21 toothed shrew, *Crocidura russula*. *Mol. Ecol.* **14**: 661-670.
- 22 Fu, Y.X. 1997. Statistical tests of neutrality of mutations against population growth,
23 hitchhiking and background selection. *Genetics* **147**: 915-925.

- 1 Fumagalli, L., Taberlet, P., Stewart, D.T., Gielly, L., Hausser, J. & Vogel, P. 1999. Molecular
2 phylogeny and evolution of *Sorex* shrews (Soricidae: Insectivora) inferred from
3 mitochondrial DNA sequence data. *Mol. Phylogenet. Evol.* **11**: 222-235.
- 4 Guindon, S. & Gascuel, O. 2003. A simple, fast, and accurate algorithm to estimate large
5 phylogenies by maximum likelihood. *Syst. Biol.* **52**: 692-704.
- 6 Heuertz, M., Carnevale, S., Fineschi, S., Sebastiani, F., Hausman, F., Paule, L. & Vendramin,
7 G.G. 2006. Chloroplast DNA phylogeography of European ashes, *Fraxinus* sp.
8 (Oleaceae): roles of hybridization and life history traits. *Mol. Ecol.* **15**: 2131-2140.
- 9 Hewitt, G.M. 1999. Post-glacial re-colonization of European biota. *Biol. J. Linn. Soc.* **68**: 87-
10 112.
- 11 Hewitt, G.M. 2000. The genetic legacy of the quaternary ice ages. *Nature* **405**: 907-913.
- 12 Hewitt, G.M. 2004a. Genetic consequences of climatic oscillations in the
13 Quaternary. *Phil. Trans. R. Soc. Lond. Ser. B.* **359**: 183–195.
- 14 Hewitt, G.M. 2004b. The structure of biodiversity – insights from molecular phylogeography.
15 *Front. Zool.* **1**: 4.
- 16 Hey, J. 2007. Using the IM program, IM Documentation.
17 [http://lifesci.rutgers.edu/~heylab/ProgramsandData/Programs/IM/Using_IM_3_5_200](http://lifesci.rutgers.edu/~heylab/ProgramsandData/Programs/IM/Using_IM_3_5_2007.pdf)
18 [7.pdf](http://lifesci.rutgers.edu/~heylab/ProgramsandData/Programs/IM/Using_IM_3_5_2007.pdf).
- 19 Hosey, G.R. 1982. The Bosphorus land-bridge and mammal distributions in Asia Minor and
20 the Balkans. *Säugetierkd. Mitt.* **30**: 53-62.
- 21 Irwin, D.M., Kocher, T.D. & Wilson, A.C. 1991. Evolution of the cytochrome *b* gene of
22 Mammals. *J. Mol. Evol.* **32**: 128-144.
- 23 Kerey, I.E., Meric, E., Kelling, G., Brenner, R.A. & Dogan, A.U. 2004. Black Sea-Marmara
24 Sea Quaternary connections: new data from the Bosphorus, Istanbul, Turkey.
25 *Paleogeogr. Paleocl. Paleoecol.* **204**: 277-295.

- 1 Koch M.A., Kiefer C., Ehrich D., Vogel, J., Brochmann, C., & Mummenhoff, K. 2006. Three
2 times out of Asia Minor: the phylogeography of *Arabis alpina* L. (Brassicaceae). *Mol.*
3 *Ecol.* **15**: 825-839.
- 4 Kučera, J., Lihová, J. & Marhold, K. 2006. Taxonomy and phylogeography of *Cardamine*
5 *impatiens* and *C. pectinata* (Brassicaceae). *Bot. J. Linn. Soc.* **152**: 169-195.
- 6 Kryštufek, B. 2002. Cranial variability in the eastern hedgehog *Erinaceus concolor*
7 (Mammalia: Insectivora). *J. Zool. (London)* **258**: 365-373.
- 8 Marmi, J., Lopez-Giraldez, F., MacDonald, D.W., Calafell, F., Zholnerovskaya, E. &
9 Domingo-Roura, X. 2006. Mitochondrial DNA reveals a strong phylogeographic
10 structure in the badger across Eurasia. *Mol. Ecol.* **15**: 1007-1020.
- 11 Michaux, J.R., Libois, R., Paradis, E. & Filippucci, M.G. 2004. Phylogeographic history of
12 the yellow-necked fieldmouse (*Apodemus flavicollis*) in Europe and in the Near and
13 Middle East. *Mol. Phylogenet. Evol.* **32**: 788-798.
- 14 Nesbo, C.L., Fosshem, T., Vollestad, L.A. & Jakobsen, K.S. 1999. Genetic divergence and
15 phylogeographic relationships among European perch (*Perca fluviatilis*) populations
16 reflect glacial refugia and postglacial colonization. *Mol. Ecol.* **8**: 1387-1404.
- 17 Nielsen, R., & J. Wakeley. 2001. Distinguishing migration from isolation: a Markov chain
18 Monte Carlo approach. *Genetics*, **158**: 885–896.
- 19 Ohdachi, S.D., Iwasa, M.A., Nesterenko, V.A. Abe, H., Masuda, R. & Haberl, W. 2004.
20 Molecular phylogenetics of *Crocidura* shrews (Insectivora) in east and central Asia. *J.*
21 *Mammal*, **85**: 396-403.
- 22 Ogg, J.G. 2004. Status of divisions of the International Geologic Time Scale. *Lethaia* **37**:
23 183–199.
- 24 Palme, A.E. & Vendramin, G.G. 2002. Chloroplast DNA variation, postglacial recolonization
25 and hybridization in hazel, *Corylus avellana*. *Mol. Ecol.* **11**: 1769-1780.

- 1 Parker, S.R., 1997. Sequence Navigator. Multiple sequence alignment software. *Methods*
2 *Mol. Biol.* **70**: 145-154.
- 3 Posada, D. & Crandall, K.A. 1998. MODELTEST: testing the model of DNA substitution.
4 *Bioinformatics* **14**: 817-818.
- 5 Poulakakis, N., Lymberakis, P., Paragamian, K. & Mylonas, M. 2005. Isolation and
6 amplification of shrew DNA from barn owl pellets. *Biol. J. Linn. Soc.* **85**: 331-340.
- 7 Rodriguez, R., Oliver, J.L., Marin, A. & Medina, J.R., 1990. The general
8 stochastic model of nucleotide substitution. *J. Theor. Bio.* **142**: 485–501.
- 9 Rogers, A.R., 1995. Genetic-evidence for a Pleistocene population explosion. *Evolution* **49**:
10 608-615.
- 11 Santucci, F., Emerson, B.C. & Hewitt, G.M., 1998. Mitochondrial DNA phylogeography of
12 European hedgehogs. *Mol. Ecol.* **7**: 1163-1172.
- 13 Seddon, J.M., Santucci, F., Reeve, N. & Hewitt, G.M. 2002. Caucasus Mountains divide
14 postulated postglacial colonization routes in the white-breasted hedgehog, *Erinaceus*
15 *concolor*. *J. Evol. Biol.* **15**: 463-467.
- 16 Schneider, I. 2000. Software programs for DNA sequence analysis. *Genet. Eng. News* **20**: 8.
- 17 Schneider, S. & Excoffier, L. 1999. Estimation of past demographic parameters from the
18 distribution of pairwise differences when the mutation rates vary among sites:
19 Application to human mitochondrial DNA. *Genetics* **152**: 1079-1089.
- 20 Swofford, D.L. 1998. Paup*. Phylogenetic Analysis Using Parsimony (* and other Methods).
21 Version 4.0b1. Sinauer Associates, Sunderland, MA.
- 22 Taberlet, P. & Fumagalli, L. 1996. Owl pellets as a source of DNA for genetics studies of
23 small mammals. *Mol. Ecol.* **5**: 301–305.
- 24 Taberlet, P., Fumagalli, L., Wust-Saucy, A.G. & Cosson, J.F. 1998. Comparative
25 phylogeography and postglacial colonization routes in Europe. *Mol. Ecol.* **7**: 453-464.

- 1 Tajima, F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA
2 polymorphism. *Genetics* **123**: 585-595.
- 3 Tarasov, P.E., Volkova, V.S., Webb, T. Guiot, J., Andreev, A.A., Bezusko, L-G., Bezusko,
4 T.V., Bykova, G.V., Dorofeyuk, N.I., Kvavadze, E.V., Osipova, I.M., Panova, N.K. &
5 Sevastyanov, D.V. 2000. Last glacial maximum biomes reconstructed from pollen and
6 plant macrofossil data from northern Eurasia. *J. Biogeogr.* **27**: 609–620.
- 7 Thorpe, R.S. 1984. Primary and secondary transition zones in speciation and population
8 differentiation – a phylogenetic analysis of range expansion. *Evolution* **38**: 233-243.
- 9 Vogel, P., Cosson, J.F. & Jurado, L.F.L. 2003. Taxonomic status and origin of the shrews
10 (Soricidae) from the Canary Islands inferred from a mtDNA comparison with the
11 European *Crocidura* species. *Mol. Phylogenet. Evol.* **27**: 271-282.
- 12 Vörös, J., Alcobendas, M., Martinez-Solano, I. & Garcia-Paris, M. 2006. Evolution of
13 *Bombina bombina* and *Bombina variegata* (Anura: Discoglossidae) in the Carpathian
14 Basin: A history of repeated mt-DNA introgression across species. *Mol. Phylogenet.*
15 *Evol.* **38**: 705-718.
- 16 Waits, L.P. & Paetkau, D. 2005. Noninvasive genetic sampling tools for wildlife biologists: a
17 review of applications and recommendations for accurate data collection. *J. Wildl.*
18 *Manage.* **69**: 1419–1433.
- 19

1 **Fig. 1** (A) Location of analysed samples, tentative distribution of mitochondrial clades (grey
2 areas), allelic frequencies of nuclear genes (cheese), and nucleotide diversity of mitochondrial
3 and nuclear gene; (B, C) Observed (solid line) and expected (dotted-line) mismatch
4 distributions for a sudden expansion of the western and eastern clades, respectively.

5
6 **Fig. 2** Phylogeny of the 1077bp *cyt-b* fragment analysed with maximum likelihood, using the
7 GTR model of substitution. Values in branches are indices of support for the major branches
8 for maximum likelihood (ML), maximum parsimony (MP) and distance (NJ) analyses
9 (percentage of 1000 replications). Codes are as in Table 1.

10

11 **Fig. 3** Marginal posterior probability (p) distribution of the divergence time parameter (t) in
12 year, from the isolation with migration model, between populations situated western and
13 eastern of the Bosphorus Strait within the Western and Eastern Clade, and estimated time for
14 the splits with the 90% highest posterior density interval (HPD).

15

1 **Table 1** Details of samples used in this study: Species, samples location, collection, location
2 on map (Fig. 1), *cyt-b* Id. Code, accession number of *cyt-b* sequences, and *ApoB* alleles.
3 Abbreviations of countries are: Bulgaria (BG), Czech Republic (CZ), Germany (DE), France
4 (FR), Georgia (GE), Greece (GR), Hungary (HU), Italy (IT), Russia (RU), Serbia (SE),
5 Slovak Republic (SK), Slovenia (SI), Switzerland (CH), and Turkey (TR).

Species	Samples location	Coll. code	Number on the map	<i>Cyt-b</i> Id. code	Accession (<i>cyt-b</i>) and <i>ApoB</i> allele
<i>Sorex minutus</i>	Champmartin, CH	IZEA 7622	/	/	DQ630379/-
<i>Crocidura suaveolens</i>	Fulophasa, HU	Dubey et al. (2006)	/	/	AY843451/-
<i>Crocidura russula</i>	Vaud, CH	Fontanillas et al. (2005)	/	/	AY769264/-
<i>Crocidura lasiura</i>	Ussurisk, RU	Ohdachi et al. (2004)	/	/	AB077071/-
<i>Crocidura leucodon</i>	Visp, CH	IZEA 7553	1	CH1	DQ994744/A1
<i>Crocidura leucodon</i>	Réchy, CH	IZEA 7552	1	CH2	DQ994745/A1
<i>Crocidura leucodon</i>	Brigerbad, CH	IZEA 2951	1	CH3	DQ994747/-
<i>Crocidura leucodon</i>	Raron, CH	IZEA 7532	1	CH4	DQ994746/-
<i>Crocidura leucodon</i>	Réchy, CH	IZEA 5590	1	CH9	DQ994794/-
<i>Crocidura leucodon</i>	Brigerbad, CH	IZEA 7526	1	/	-/A1
<i>Crocidura leucodon</i>	Gordevio, CH	IZEA 5965	2	CH5	DQ994748/-
<i>Crocidura leucodon</i>	Gordevio, CH	IZEA 5963	2	CH6	DQ994749/A1
<i>Crocidura leucodon</i>	Quartino, CH	IZEA 5455	2	CH7	DQ994750/A1
<i>Crocidura leucodon</i>	Grison, CH	IZEA 9013	3	CH8	DQ994785/A1
<i>Crocidura leucodon</i>	Karlsruhe, DE	IZEA 5441	4	DE5	DQ994795/A1
<i>Crocidura leucodon</i>	Rendsburg, DE	IZEA 7835	5	DE1	DQ994761/-
<i>Crocidura leucodon</i>	Rendsburg, DE	IZEA 7836	5	DE2	DQ994762/-
<i>Crocidura leucodon</i>	Rendsburg, DE	IZEA 7837	5	DE3	DQ994763/A1
<i>Crocidura leucodon</i>	Rendsburg, DE	IZEA 7838	5	DE4	DQ994764/-
<i>Crocidura leucodon</i>	Mignouillard, FR	IZEA 9006	6	FR4	DQ994780/A1
<i>Crocidura leucodon</i>	Mignouillard, FR	IZEA 9002	6	FR5	DQ994781/A1
<i>Crocidura leucodon</i>	Chapelle, FR	IZEA 9003	7	FR2	DQ994779/A1
<i>Crocidura leucodon</i>	Chapelle, FR	IZEA 9005	7	FR3	DQ994782/A1
<i>Crocidura leucodon</i>	St-Martin, FR	IZEA 9008	8	FR1	DQ994783/A1
<i>Crocidura leucodon</i>	St-Etienne, FR	IZEA 9009	9	FR6	DQ994784/A1
<i>Crocidura leucodon</i>	Vercelli, IT	IZEA 7948	10	IT1	DQ994767/-
<i>Crocidura leucodon</i>	Vercelli, IT	IZEA 7946	10	IT2	DQ994768/-
<i>Crocidura leucodon</i>	Vercelli, IT	IZEA 7952	10	IT3	DQ994769/-
<i>Crocidura leucodon</i>	Serramazoni, IT	IZEA 5662	11	IT4	DQ994770/A1
<i>Crocidura leucodon</i>	Serramazoni, IT	IZEA 5649	11	IT5	DQ994771/-
<i>Crocidura leucodon</i>	Serramazoni, IT	IZEA 5663	11	IT6	DQ994772/A1
<i>Crocidura leucodon</i>	Piacenza, IT	IZEA 7517	12	IT7	DQ994773/A1
<i>Crocidura leucodon</i>	Bratislava, SK	IZEA 5728	13	SK1	DQ994765/-
<i>Crocidura leucodon</i>	Bratislava, SK	IZEA 5723	13	SK2	DQ994766/A1
<i>Crocidura leucodon</i>	Lesbos, GR	IZEA 4153	14	GR1	DQ994777/A1
<i>Crocidura leucodon</i>	Lesbos, GR	IZEA 3929	14	GR2	DQ994778/A1
<i>Crocidura leucodon</i>	Lesbos, GR	Poulakakis et al., 2005	14	-	AY452165/-
<i>Crocidura leucodon</i>	Burgas, BG	IZEA 8059	15	BG1	DQ994774/A1/A2
<i>Crocidura leucodon</i>	Burgas, BG	IZEA 8060	15	BG2	DQ994775/A1
<i>Crocidura leucodon</i>	Burgas, BG	IZEA 8063	15	BG3	DQ994776/A1
<i>Crocidura leucodon</i>	Burgas, BG	IZEA 8059	15	-	-/A1/A2
<i>Crocidura leucodon</i>	Alazani, GE	IZEA 23629	16	GE1	DQ994756/A1/A4
<i>Crocidura leucodon</i>	Alazani, GE	IZEA 23635	16	GE2	DQ994757/A4
<i>Crocidura leucodon</i>	Alazani, GE	IZEA 23613	16	GE3	DQ994758/A1
<i>Crocidura leucodon</i>	Dusheti, GE	IZEA 2880	17	GE4	DQ994760/A1/A3
<i>Crocidura leucodon</i>	Cakalli, TR	IZEA 6064	18	TR1	DQ994751/A1

<i>Crocidura leucodon</i>	Rize, TR	IZEA 6045	19	TR2	DQ994752/-
<i>Crocidura leucodon</i>	Rize, TR	IZEA 6049	19	TR3	DQ994753/A1
<i>Crocidura leucodon</i>	Rize, TR	IZEA 6047	19	TR4	DQ994754/A1
<i>Crocidura leucodon</i>	Rize, TR	IZEA 6046	19	TR5	DQ994755/A1
<i>Crocidura leucodon</i>	Altindere, TR	IZEA 6039	20	TR6	DQ994759/A1
<i>Crocidura leucodon</i>	Altindere, TR	IZEA 6038	20	TR7	DQ994796/-
<i>Crocidura leucodon</i>	Altindere, TR	IZEA 6044	20	-	-/A1
<i>Crocidura leucodon</i>	Yellibeli, Kar., TR	DZCU TU-1179	23	TR8	DQ994787/A1
<i>Crocidura leucodon</i>	Çiğlikara, Ant., TR	DZCUTU-1195	24	TR9	DQ994786/A1
<i>Crocidura leucodon</i>	Karlovy Vary, CZ	DZCU 01400	25	-	-/A1
<i>Crocidura leucodon</i>	Karlovy Vary, CZ	DZCU 01399	25	CZ1	DQ994788/A1
<i>Crocidura leucodon</i>	Karlovy Vary, CZ	DZCU 01398	25	CZ2	DQ994789/A1
<i>Crocidura leucodon</i>	Mt. Cer, SE	PMS 7391	26	SE1	DQ994790/A1
<i>Crocidura leucodon</i>	Cataloi, RO	IZEA 8160	27	RO1	DQ994791/A1
<i>Crocidura leucodon</i>	Slobozia, RO	IZEA 8170	28	RO2	DQ994792/A1
<i>Crocidura leucodon</i>	Vrhnik, SI	PMS Slo 1	29	SI1	DQ994793/A1
<i>Crocidura leucodon</i>	Katranci-Biga, TR	2003.131	30	TR10	EF417543/A1
<i>Crocidura leucodon</i>	Özbek, İzmir, TR	2003.199	31	TR11	EF417544 /A1
<i>Crocidura leucodon</i>	Terzialan-Çan., TR	2003.217	32	TR12	EF417545/A1
<i>Crocidura leucodon</i>	Parnitha, GR	Poulakakis et al., 2005	21	-	AY452166/-
<i>Crocidura leucodon</i>	Stymfalia, GR	Poulakakis et al., 2005	22	-	AY452176/-

1

2

1 **Table 2** Example of mutations observed between some of our samples and the samples of
2 Poulakakis *et al.* (2005; *, Lesvos, AY452165; Parnitha, AY452166; Stymfalia, AY452176).
3 Only the mutations including the first 107 bp and the last 52 bp of the Poulakakis sequences
4 are shown. The samples SK2, IT5, TR10, GR1 and GR2 (this study) belong to the western
5 mitochondrial clade, and the samples TR6 and GE1 from the eastern mitochondrial clade.

Samples	Example of mutations observed between sequences																		
SK2, Bratislava	TC	TGC	TTA	ATT	GCA	CAA	ATC	CTA	ACA	GGA	TTA	TTC	CTA	GCC	ATA	CAC	TAT	ACA	[53]
IT5, Serramazoni	[53]
TR10, Katranci	[53]
GR2, LesvosT	[53]
GR1, Lesvos	[53]
*Lesvos	NN	NNN	NNN	NNN	NNNTT	[53]
*Parnitha	NN	NNN	NNN	NNN	NNNGT	[53]
*StymfaliaT	C..	..T	[53]
TR6, Altindere	T..	C..	...	T..	[53]
GE1, Alazani	T..	C..	[53]
SK2, Bratislava	TCT	GAT	ACT	ACA	ACA	GCT	TTC	TCC	TCC	GTA	ACC	CAT	ATT	TGC	CGA	GAT	GTA	AAT	[107]
IT5, Serramazoni	..A	[107]
TR10, Katranci	..A	[107]
GR2, Lesvos	..A	[107]
GR1, Lesvos	..A	[107]
*Lesvos	..A	[107]
*Parnitha	..AC	[107]
*Stymfalia	..ACT	[107]
TR6, Altindere	..AA	[107]
GE1, Alazani	..AA	[107]
SK2, Bratislava	TA	GGA	TAT	GTT	CTT	CCC	TGA	GGT	CAA	ATA	TCA	TTT	TGA	GGT	GCA	ACA	GTA	AT	[332]
IT5, Serramazoni	[332]
TR10, Katranci	[332]
GR2, Lesvos	[332]
GR1, Lesvos	[332]
*LesvosACG	..T	G..	NNN	NN	[332]
*ParnithaACG	..T	G..	NNN	NN	[332]
*StymfaliaACG	..T	G..	AGT	T.	[332]
TR6, AltindereG	[332]
GE1, AlazaniG	[332]


6
7
8
9
10
11
12
13

1 **Table 3** Number of samples (Ns) and number of haplotypes (Nh) within the western and
 2 eastern mitochondrial bicolored shrew clades, genetic diversity, nucleotide diversity, mean
 3 pairwise differences between haplotypes, Goodness-of-fit test probability, Fu's F_s test
 4 probability, Tajima's D test probability, estimated τ value and expansion time for western
 5 clade.

Clade (Ns, Nh)	Nucleotide div./Mean pairwise diff.	Goodness-of-fit test		Fu's F_s test		Tajima's test		τ	Expansion time
		Rag. Index	P	F_s	P	Tajima's D	P		
Western (34, 23)	0.0057/6.21	0.02	= 0.28	-10.68	<0.001	-1.48	= 0.04	6.80 (95% CI: 4.19-8.39)	51,900 y BP (95% CI: 28,100-72,100)
Eastern (16, 11)	0.0059/6.31	0.12	= 0.03	-2.02	= 0.16	-0.67	= 0.26	7.47 (95% CI: 3.71-10.85)	/

6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

1



2

3

4

5

6

7

8

9

10

11

12

13

14

15


16

17

18


19

1



2
3
4
5
6
7
8
9
10

1



2