

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

US Army Research

U.S. Department of Defense

2013

Genome Sequence of *Staphylococcus pseudintermedius* Strain E140, an ST71 European-Associated Methicillin-Resistant Isolate

Arshnee Moodley

University of Copenhagen, asm@sund.ku.dk

Matthew C. Riley

U.S. Army Medical Service Corps

Stephen A. Kania

University of Tennessee - Knoxville

Luca Guardabassi

University of Copenhagen

Follow this and additional works at: <https://digitalcommons.unl.edu/usarmyresearch>

Moodley, Arshnee; Riley, Matthew C.; Kania, Stephen A.; and Guardabassi, Luca, "Genome Sequence of *Staphylococcus pseudintermedius* Strain E140, an ST71 European-Associated Methicillin-Resistant Isolate" (2013). *US Army Research*. 243.

<https://digitalcommons.unl.edu/usarmyresearch/243>

This Article is brought to you for free and open access by the U.S. Department of Defense at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in US Army Research by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Genome Sequence of *Staphylococcus pseudintermedius* Strain E140, an ST71 European-Associated Methicillin-Resistant Isolate

Arshnee Moodley,^a Matthew C. Riley,^{b,c} Stephen A. Kania,^b Luca Guardabassi^a

Department of Veterinary Disease Biology, Faculty of Health and Medical Sciences, University of Copenhagen, Frederiksberg C, Copenhagen, Denmark^a; Department of Biomedical and Diagnostic Sciences, University of Tennessee, Knoxville, Tennessee, USA^b; U.S. Army Medical Service Corps^c

We report the first genome sequence of the methicillin-resistant *Staphylococcus pseudintermedius* (MRSP) strain E140, isolated from a canine bite wound infection in Denmark. This strain represents the dominant clonal lineage associated with canine MRSP infections in Europe.

Received 14 December 2012 Accepted 5 February 2013 Published 7 March 2013

Citation Moodley A, Riley MC, Kania SA, Guardabassi L. 2013. Genome sequence of *Staphylococcus pseudintermedius* strain E140, an ST71 European-associated methicillin-resistant isolate. *Genome Announc.* 1(2):e00207-12. doi:10.1128/genomeA.00207-12.

Copyright © 2013 Moodley et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Arshnee Moodley, asm@sund.ku.dk.

Staphylococcus pseudintermedius is a commensal Gram-positive species residing on the skin and mucosae of dogs, and it is the most common bacterial pathogen associated with canine infections, predominantly skin infections. Methicillin-resistant *S. pseudintermedius* (MRSP) was first reported in Europe in 2006 (1). The prevalent clone circulating in Europe belongs to sequence type (ST) 71, which displays resistance to all antibiotics routinely used in small animal practice (1–3). This clonal type has also been reported in Canada and the United States, although less frequently (2). MRSP ST71 can be transmitted to veterinarians and dog owners (4, 5), and human infection has been reported as a result of exposure to a colonized dog (6).

The genome of an MRSP ST71 strain (E140) isolated from a canine bite wound infection in Denmark was sequenced using three platforms: (i) a Roche GS FLX sequencer generating 133,947 reads, with an average length of 569 bp; (ii) an Illumina paired-end (PE) technology (500-bp library) generating 300 Mb data and 3,333,334 paired reads; and (iii) an Ion Torrent 318 generating 3,752,077 reads with an average length of 190 bp. *De novo* genome assembly of the various read sets was performed using Geneious and DNASTar SeqMan NGen and produced (i) Roche 454 data, 86 contigs of $\geq 1,000$ bp with a mean length of 34,063 bp; (ii) Illumina PE data, 26 contigs of $\geq 1,000$ bp with a mean length of 106,456 bp; and (iii) Ion Torrent data, 73 contigs of $\geq 1,000$ bp with a mean length of 94,909 bp. In addition, a high-resolution AflII whole-genome restriction map of E140 was generated using an Argus whole-genome mapping system (Opgen, Inc.). The whole-genome map was used to align contigs from different assemblies and to verify final genome assembly. Automated annotation of genes was performed on the RAST (Rapid Annotation using Subsystem Technology) server (7).

The draft genome of MRSP E140 has a size of 2,769,487 bp, which is ~150 to 200 kb larger than the two published methicillin susceptible *S. pseudintermedius* genomes of strain ED99 (8) and strain HKU10-03 (9). This shift in genome size is in part due to the presence of a novel ~30-kb prophage, as seen in the sequencing

contigs and verified in the optical map. E140 has a GC content of 38.0% and has 2,678 coding sequences, 5 ribosomal operons, and 68 tRNAs. Methicillin resistance is attributed to the presence of the staphylococcal cassette chromosome *mec* element (SCC*mec*) type II-III associated with ST71 (2, 10).

The discovery of the genome sequence of this MRSP will facilitate future studies to understand the rapid spread, fitness, and pathogenesis of this multidrug-resistant, dominant clonal lineage, which causes infections in Europe and in North America. In addition, the genome sequence information will assist investigators in tracking the evolution of methicillin and multidrug resistance in this species as more MRSP genomes become available.

Nucleotide sequence accession number. The draft genome sequence of *S. pseudintermedius* E140 has been deposited in the GenBank database with the accession number [ANOI01000001](https://www.ncbi.nlm.nih.gov/nuccore/ANOI01000001).

ACKNOWLEDGMENTS

Funding was received from the Fund for Scientific Studies in Small Animals (University of Copenhagen) and the American Kennel Club Canine Health Foundation (project 01421).

REFERENCES

1. Loeffler A, Linek M, Moodley A, Guardabassi L, Sung JM, Winkler M, Weiss R, Lloyd DH. 2007. First report of multiresistant, *mecA*-positive *Staphylococcus intermedius* in Europe: 12 cases from a veterinary dermatology referral clinic in Germany. *Vet. Dermatol.* 18:412–421.
2. Perreten V, Kadlec K, Schwarz S, Grönlund Andersson U, Finn M, Greko C, Moodley A, Kania SA, Frank LA, Bemis DA, Franco A, Iurescia M, Battisti A, Duim B, Wagenaar JA, van Duijkeren E, Weese JS, Fitzgerald JR, Rossano A, Guardabassi L. 2010. Clonal spread of methicillin-resistant *Staphylococcus pseudintermedius* in Europe and North America: an international multicentre study. *J. Antimicrob. Chemother.* 65:1145–1154.
3. Vanni M, Tognetti R, Pretti C, Crema F, Soldani G, Meucci V, Intorre L. 2009. Antimicrobial susceptibility of *Staphylococcus intermedius* and *Staphylococcus schleiferi* isolated from dogs. *Res. Vet. Sci.* 87:192–195.
4. Paul NC, Moodley A, Ghibaudo G, Guardabassi L. 2011. Carriage of

- methicillin-resistant *Staphylococcus pseudintermedius* in small animal veterinarians: indirect evidence of zoonotic transmission. *Zoonoses Public Health* 58:533–539.
- van Duijkeren E, Kamphuis M, van der Mije IC, Laarhoven LM, Duim B, Wagenaar JA, Houwers DJ. 2011. Transmission of methicillin-resistant *Staphylococcus pseudintermedius* between infected dogs and cats and contact pets, humans and the environment in households and veterinary clinics. *Vet. Microbiol.* 150:338–343.
 - Stegmann R, Burnens A, Maranta CA, Perreten V. 2010. Human infection associated with methicillin-resistant *Staphylococcus pseudintermedius* ST71. *J. Antimicrob. Chemother.* 65:2047–2048.
 - Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75.
 - Ben Zakour NL, Bannoehr J, van den Broek AH, Thoday KL, Fitzgerald JR. 2011. Complete genome sequence of the canine pathogen *Staphylococcus pseudintermedius*. *J. Bacteriol.* 193:2363–2364.
 - Tse H, Tsoi HW, Leung SP, Urquhart IJ, Lau SK, Woo PC, Yuen KY. 2011. Complete genome sequence of the veterinary pathogen *Staphylococcus pseudintermedius* strain HKU10-03, isolated in a case of canine pyoderma. *J. Bacteriol.* 193:1783–1784.
 - Descloux S, Rossano A, Perreten V. 2008. Characterization of new staphylococcal cassette chromosome mec (SCCmec) and topoisomerase genes in fluoroquinolone- and methicillin-resistant *Staphylococcus pseudintermedius*. *J. Clin. Microbiol.* 46:1818–1823.