



# RNA

A PUBLICATION OF THE RNA SOCIETY

## The role of the putative 3' end processing endonuclease Ysh1p in mRNA and snoRNA synthesis

Monika Garas, Bernhard Dichtl and Walter Keller

*RNA* 2008 14: 2671-2684 originally published online October 29, 2008  
Access the most recent version at doi:[10.1261/rna.1293008](https://doi.org/10.1261/rna.1293008)

---

### References

This article cites 71 articles, 30 of which can be accessed free at:  
<http://rnajournal.cshlp.org/content/14/12/2671.full.html#ref-list-1>

### Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#)

---

---

To subscribe to *RNA* go to:  
<http://rnajournal.cshlp.org/subscriptions/>

---

# The role of the putative 3' end processing endonuclease Ysh1p in mRNA and snoRNA synthesis

MONIKA GARAS,<sup>1</sup> BERNHARD DICHTL,<sup>2</sup> and WALTER KELLER<sup>1</sup>

<sup>1</sup>Biozentrum, University of Basel, CH-4056 Basel, Switzerland

<sup>2</sup>Institute of Molecular Biology, University of Zürich, CH-8057 Zürich, Switzerland

## ABSTRACT

Pre-mRNA 3' end formation is tightly linked to upstream and downstream events of eukaryotic mRNA synthesis. The two-step reaction involves endonucleolytic cleavage of the primary transcript followed by poly(A) addition to the upstream cleavage product. To further characterize the putative 3' end processing endonuclease Ysh1p/Brr5p, we isolated and analyzed a number of new temperature- and cold-sensitive mutant alleles. We show that Ysh1p plays a crucial role in 3' end formation and in RNA polymerase II (RNAP II) transcription termination on mRNA genes. In addition, we observed a range of additional functional deficiencies in *ysh1* mutant strains, which were partially allele-specific. Interestingly, snoRNA 3' end formation and RNAP II termination were defective on specific snoRNAs in the cold-sensitive *ysh1-12* strain. Moreover, we observed the accumulation of several mRNAs including the *NRD1* transcript in this mutant. We provide evidence that *NRD1* autoregulation is associated with endonucleolytic cleavage and that this process may involve Ysh1p. In addition, the *ysh1-12* strain displayed defects in RNA splicing indicating that a functional link may exist between intron removal and 3' end formation in yeast. These observations suggest that Ysh1p has multiple roles in RNA synthesis and processing.

**Keywords:** 3' end formation; 3' endonuclease; RNA polymerase II transcription termination; snoRNA; Nrd1 autoregulation

## INTRODUCTION

RNA polymerase II (RNAP II) transcribes protein-encoding genes and a subset of noncoding small nuclear RNAs (snRNA) and small nucleolar RNAs (snoRNA). In order to be biologically functional, all primary transcripts of RNAP II need to undergo extensive processing. For mRNAs, this includes capping at the 5' end, removal of introns by splicing, and 3' end processing.

Pre-mRNA 3' end processing is initiated by endonucleolytic cleavage at the poly(A) site, followed by polyadenylation of the upstream cleavage product. In yeast, 3' end processing is performed by a large complex of proteins, including cleavage and polyadenylation factor (CPF), cleavage factor IA (CF IA), and cleavage factor IB (CF IB) (Zhao et al. 1999). Ysh1p/Brr5p is associated with CPF and was first identified by sequence homology with mammalian CPSF73 (Chanfreau et al. 1996; Jenny et al. 1996). It was recognized early on that the protein carries a

highly conserved  $\beta$ -lactamase fold commonly found in metal-dependent hydrolytic enzymes (Aravind 1999; Daiyasu et al. 2001; Callebaut et al. 2002). Although this suggested that catalytic activity was associated with Ysh1p, convincing structural and biochemical evidence for endonucleolytic activity has emerged only recently (Ryan et al. 2004; Wickens and Gonzalez 2004; Mandel et al. 2006).

Cleavage and polyadenylation factors are required for termination of RNAP II transcription (Connelly and Manley 1988; Proudfoot 2004; Buratowski 2005). Two general models have been proposed to explain the mechanism of termination (Proudfoot et al. 2002; Luo and Bentley 2004; Bentley 2005; Buratowski 2005). In the "anti-terminator" model, RNAP II complex undergoes conformational changes in response to the emerging terminator sequences on RNA either by recruiting termination factors and/or by displacing positive elongation factors (Logan et al. 1987; Orozco et al. 2002; Zhang et al. 2005; Zhang and Gilmour 2006). This renders the RNAP II complex termination competent and leads to the loss of processivity and gradual termination. The "torpedo" model proposes that RNAP II transcription termination is triggered by poly(A) site cleavage and subsequent degradation of the 3' downstream RNA by Rat1p 5'-3' exonuclease (Connelly and

**Reprint requests to:** Walter Keller, Biozentrum, University of Basel, Klingelbergstrasse 70, CH-4056 Basel, Switzerland; e-mail: [walter.keller@unibas.ch](mailto:walter.keller@unibas.ch); fax: +41 61 267 2079.

Article published online ahead of print. Article and publication date are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.1293008>.

Manley 1988; Kim et al. 2004; West et al. 2004). Experimental support for either model has been provided in several eukaryotic experimental systems (Gilmour and Fan 2008), and a unified model of termination has been proposed that combines mechanistic predictions of both models (Luo et al. 2006).


All snRNAs and most snoRNAs in yeast are synthesized independently by RNAP II. The 3' ends of snoRNA transcripts are produced by 3'-5' exonucleolytic trimming that follows either endonucleolytic cleavage or RNAP II termination (Chanfreau et al. 1998; Allmang et al. 1999; van Hoof et al. 2000; Butler 2002). Numerous protein factors were reported to be essential for transcription termination on snoRNA genes, including the RNAP II subunits Rpb3p and Rpb11p, the sequence-specific RNA-binding proteins Nab3p and Nrd1p, the Sen1p helicase, the CTD kinase Ctk1p, and the RNAP II-associated Paf1 complex (Ursic et al. 1997; Conrad et al. 2000; Steinmetz et al. 2001, 2006a; Sheldon et al. 2005). Furthermore, several subunits of the 3' end processing complexes CF IA (Pcf11p, Rna15p, and Rna14p) and CPF (Pta1p, Pti1p, Ref2p, Ssu72p, Swd2p, and Glc7p) were implicated in snoRNA termination (Fatica et al. 2000; Morlando et al. 2002; Dheur et al. 2003; Ganem et al. 2003; Nedea et al. 2003, 2008; Steinmetz and Brow 2003; Cheng et al. 2004; Dichtl et al. 2004; Kim et al. 2006).

In this study, we isolated a number of conditional *ysh1* mutant alleles and characterized them for defects in mRNA and snoRNA synthesis. We found that all analyzed *ysh1* mutants were deficient in pre-mRNA 3' end formation and RNAP II transcription termination on mRNA genes. Moreover, a cold-sensitive *ysh1* mutant strain displayed distinct defects in snoRNA 3' end formation, termination on snoRNA genes, and RNA splicing. We also provide evidence that implies endonucleolytic cleavage and functional Ysh1p in the process of regulated premature termination at the *NRD1* locus. Altogether, this study underscores the central role of the 3' end processing endonuclease Ysh1p in cellular RNA metabolism.

## RESULTS

### Ysh1p is required for 3' end cleavage and polyadenylation of pre-mRNA in vitro

To address the cellular role of the putative 3' endonuclease Ysh1p, we initially produced point mutations within the  $\beta$ -lactamase consensus signature motif  $H_{68}X_{69}H_{70}X_{71}D_{72}H_{73}$ , which is located at the amino-terminal end of the protein and which contributes to formation of the catalytic core of the protein (Fig. 1C; Aravind 1999; Mandel et al. 2006). We found that alanine substitutions of the highly conserved histidines  $H_{68}$  and  $H_{70}$  or of aspartate  $D_{72}$  resulted in lethality (Fig. 1A). This observation underscored the functional significance of this motif but hampered further



**FIGURE 1.** Isolation of conditional *ysh1* mutants. (A) The conserved  $\beta$ -lactamase signature  $H_{68}X_{69}H_{70}X_{71}D_{72}H_{73}$  is essential for cell viability. Plasmid shuffling was used to test the requirement of  $H_{68}$ ,  $H_{70}$ , and  $D_{72}$  residues of the  $\beta$ -lactamase consensus motif for cell viability. The LM109 strain with a disrupted chromosomal *YSH1* gene and carrying *YSH1-URA3* plasmid was transformed with the *ADE2*-plasmid bearing either the wild-type *YSH1* gene or its mutant versions  $H_{68}A$ ,  $H_{70}A$ , and  $D_{72}A$ , followed by counterselection on 5'-FOA plates. (B) Tenfold serial dilutions of cultures of wild-type and *ysh1* mutant strains were spotted onto YPAD medium and incubated for 3 d at 23°–37°C or 5 d at 15°C. (C) Schematic representation of *ysh1* mutant sequences underlying the respective temperature- or cold-sensitive phenotypes. (Shaded boxes) The  $\beta$ -lactamase,  $\beta$ -CASP, and C-terminal domains and the conserved  $H_{68}F_{69}H_{70}L_{71}D_{72}H_{73}$  signature are marked approximately.


functional analyses. Therefore, we isolated temperature-sensitive (ts) and cold-sensitive (cs) alleles of the *YSH1/BRR5* gene using random mutagenesis (Fig. 1B; see Materials and Methods). Of the mutant strains chosen for further analysis, three were ts, with *ysh1-32* being lethal at 33°C, and *ysh1-13* and *ysh1-15* being lethal at 37°C; in contrast, the cs *ysh1-12* strain ceased growth at 15°C (Fig. 1B). Amino acid changes within mutant Ysh1p proteins were

checked by DNA sequencing and found to localize predominantly to the nonconserved C-terminus (Fig. 1C); this part of the protein is involved in both catalytic steps of 3' end processing and mediates the interaction with Pta1p (Zhelkovsky et al. 2006).

Initially, we tested the *ysh1* mutant strains for cleavage and polyadenylation in vitro (Fig. 2A,B). Total cell extracts

were produced from wild-type and mutant strains (*ysh1-12*, *ysh1-13*, *ysh1-15*, *ysh1-32*) and assayed on a synthetic *CYC1* substrate for cleavage and on pre-cleaved *CYC1-pre* RNA for polyadenylation. Comparing the effect of the mutations to wild-type, we observed that both cleavage and polyadenylation were reduced in all the *ysh1* mutant strains, with the defects being more pronounced at the restrictive temperatures (Fig. 2A,B).

To confirm that the observed defects in pre-mRNA 3' end formation were due to inactive CPF (the factor that contains Ysh1p), we aimed to reconstitute its activity by addition of purified wild-type CPF to the *ysh1-32* extract (Fig. 2C,D). We found that CPF partially rescued the cleavage defect of the mutant extract at both permissive and restrictive temperatures (Fig. 2C, lanes 4,9). Likewise, specific polyadenylation activity of the *ysh1-32* extract was restored by addition of CPF (Fig. 2D, lanes 4,9). CPF alone was not able to cleave the *CYC1* RNA (Fig. 2C, lanes 5,10) and gave unspecific hyperadenylation of the *CYC1-pre* RNA substrate as previously observed (Fig. 2D, lanes 5,10; Preker et al. 1997). Taken together, the above results underscore an important role for Ysh1p in both catalytic steps of pre-mRNA 3' end formation.




**FIGURE 2.** Ysh1p is required for cleavage and polyadenylation of pre-mRNA in vitro. (A, upper panel) In vitro cleavage and (lower panel) polyadenylation assays with protein extracts prepared from wild-type and *ysh1* temperature-sensitive strains as indicated. Input shows a control where no protein was added. Positions of substrate RNAs, 5' and 3' end cleavage products and polyadenylation products bands are shown. HpaII-digested pBR322 fragments were 5' end labeled and served as markers. Internally [ $^{32}$ P]-labeled substrate RNAs, *CYC1* for the cleavage assay and *CYC1-pre* for specific polyadenylation, were used. Reactions were performed either at 30°C (lanes 1–6) or at 37°C (lanes 7–12). Extracts were preincubated at restrictive temperatures for 10 min prior to assaying. (B) As in A, except reactions were performed either at the permissive temperature 30°C (lanes 1–3) or at the non-permissive temperatures 17°C (lanes 4–6) for cleavage, and at 15°C for polyadenylation. (C,D) Reconstitution of specific cleavage and polyadenylation activities of the *ysh1-32* extract in vitro. (C) Cleavage and (D) polyadenylation assays were performed essentially as in A, with protein extracts prepared from wild-type and the *ysh1-32* strain as indicated. The *ysh1-32* extract was combined with purified CPF to restore the specific 3' end processing activity. Twice more CPF was used for the reconstitution of polyadenylation than for the reconstitution of cleavage.

### Distinct effects of mutations within Ysh1p on mRNA abundance

Next, we performed Northern blotting to analyze the steady-state levels of mRNAs in *ysh1* mutant strains before and after shift to restrictive temperatures. We found that several mRNAs (*ASC1*, *ASN1*, *PGK1*, *RPS16A*) were significantly reduced in *ysh1-32* strains following growth at 37°C (Fig. 3A, lanes 6–8); similar phenotypes were observed for *ysh1-13* and *ysh1-15* mutant strains (data not shown). The well-characterized *rna15-1* strain was used as a control and showed the expected strong reduction of mRNAs after 1-h shift to 37°C (Fig. 3A, lane 4; Minvielle-Sebastia et al. 1994). Unexpectedly, the reverse effect was observed in *ysh1-12* cells, where the levels of all tested mRNAs were relatively stable or even increased upon shift to non-permissive temperature (Fig. 3A, lanes 12–14); this included *NAB2* mRNA, which increased up to 13-fold. Interestingly, Nab2p is believed to autoregulate the levels of its own mRNA in a process that requires the nuclear exosome component Rrp6p (Roth et al. 2005), and our results suggest a role for Ysh1p in this process (see Discussion).

To evaluate the global phenotypes of our mutants, we analyzed the length distribution of cellular poly(A). The *ts ysh1-32*, *ysh1-13*, and *ysh1-15* strains showed a marked reduction of poly(A) tail length after growth at 37°C (Fig. 3B, left panel; data not shown). In contrast, the poly(A) tail length distribution remained unchanged following shift to the restrictive temperature for *ysh1-12* (Fig. 3B, right panel). Thus, global poly(A) length reflected the trends of





**FIGURE 3.** Ysh1p is required for normal mRNA accumulation and poly(A) site selection. (A) Northern blot analysis of mRNA steady-state levels in *ysh1* strains. Wild-type, *ysh1-32*, and *rna15-1* strains were grown in YPD at 25°C and shifted for 1, 3, and 6 h to 37°C. The cold-sensitive strain *ysh1-12* was grown in YPD at 30°C and shifted for 1, 3, and 6 h to 15°C. Total RNA was extracted from wild-type and mutant *ysh1* cells and separated on formaldehyde/1.2% agarose gels. Filters were developed with either random-primed labeled DNA probes or end-labeled oligonucleotides directed against RNA species indicated on the left. 18S rRNA and SCR1 served as loading controls. (B) Poly(A) tails shorten in the *ysh1-32* strain after shift to restrictive temperature. 3' End labeling of poly(A) tails with total RNA extracted from (left panel) wild-type and *ysh1-32* or (right panel) wild-type and *ysh1-12* mutant strains after growth at permissive temperatures (23°C or 30°C, respectively), and after shift for 1, 3, and 6 h to respective restrictive temperatures (37°C or 15°C). Poly(A) tail length (in nucleotides) is indicated on the left. (C) Western blot analysis of wild-type, *ysh1-32*, and *ysh1-12* mutant extracts prepared from cells grown analogously to A. Equal amounts of total protein were loaded in each lane. Blots were probed with antibodies directed against the proteins indicated on the left. (D) Northern blot analysis of total RNAs extracted from wild-type and mutant *ysh1* cells. The *ysh1-13*, *ysh1-15*, *ysh1-32*, and *rna15-1* strains were grown in YPD at 23°C and shifted for 1, 3, and 6 h to 37°C. *ysh1-12* was grown at 30°C and shifted for 1, 3, and 6 h to 15°C. The analysis was performed essentially as in A. 18S rRNA served as a loading control. The positions of ACT1 poly(A) sites I–V are labeled. The scheme below the panel represents the relative positions of the different ACT1 poly(A) sites. (E) Analysis of ACT1 poly(A) site usage in wild-type, *ysh1-13*, and *ysh1-15* cells. Total RNAs extracted from strains grown as described in E were treated with RNase H and oligonucleotides ACT1-RNase H and oligo(dT), and analyzed by polyacrylamide Northern blotting with a probe specific for the 3' end of ACT1 mRNA. Positions of the poly(A) sites I–V are indicated on the right. RNA levels were quantified by PhosphorImager scanning (Molecular Dynamics). The ratios of poly(A) site I versus site V usage for each lane are indicated at the bottom.

mRNA levels observed in the Northern blot analyses for both the *ysh1* ts strains (reduced mRNA) and the *ysh1-12* cs strain (unchanged or increased mRNA).

To test whether the observed phenotypes were due to underaccumulation of Ysh1p or other subunits of the 3' end processing complex, we carried out Western blot analyses (Fig. 3C). We observed reduced levels of Ysh1p and Ssu72p in the *ysh1-32* strain after 1-h growth at 37°C (Fig. 3C, lanes 4–6), whereas other CPF components (Pfs2p, Pap1p, Fip1p) and Rna15p remained largely unchanged. Interestingly, the amounts of all analyzed

proteins remained relatively stable in the *ysh1-12* strain during a 6-h shift to 15°C (Fig. 3C, lanes 9–12). These results support the view that the ts and cs *ysh1* mutations caused distinct functional deficiencies (see Discussion).

### Ysh1p is required for correct poly(A) site recognition of ACT1 pre-mRNA

Several subunits of the CPF complex including Yhh1p and Ydh1p were previously shown to be required for poly(A)

site recognition (Dichtl et al. 2002b; Kyburz et al. 2003). The well-characterized 3'-UTR of the *ACT1* gene includes five alternative polyadenylation sites and is therefore useful to examine poly(A) site selection in 3' end processing mutants (see scheme in Fig. 3D; Mandart and Parker 1995). Low-resolution Northern blot analysis revealed a change in the *ACT1* hybridization pattern in *ysh1* mutants compared to wild-type (Fig. 3D). To corroborate this phenotype, we further analyzed RNA obtained from two mutant strains (*ysh1-13* and *ysh1-15*) on high-resolution Northern blots following RNase H treatment (Mandart and Parker 1995). In these experiments, total RNA was targeted by RNase H and oligonucleotides complementary to the 3' end of *ACT1* mRNA and to the poly(A) tail allowing the resolution of the five poly(A) sites (Fig. 3E). Quantification showed that the most proximal poly(A) site I was preferentially used in wild-type cells, as expected; downstream sites II–V were utilized four to seven times less frequently. In contrast, *ysh1* mutant strains displayed a change in poly(A) site usage toward the minor site V, both at permissive and restrictive temperatures. Taken together, these results demonstrate the involvement of Ysh1p in correct poly(A) site recognition.

### Ysh1p is required for RNAP II transcription termination on mRNA genes

To address the role of Ysh1p in RNAP II transcription termination, we carried out transcriptional run-on analysis (TRO) on a plasmid-borne *CYC1* gene, which is placed under the control of the *GAL1/10* promoter (Birse et al. 1998). Correct termination produces run-on signals over probes P1–P3, whereas defective termination results in increased signals also over probes P4–P6 (see scheme in Fig. 4A). TRO was performed with wild-type and *ysh1-12* cells following growth in galactose-containing medium at permissive and restrictive conditions. Quantification of slot hybridizations showed that the *ysh1-12* mutant displayed an approximately twofold to threefold increased run-on signal over probes P3–P6 (Fig. 4A,B). Similar results were obtained with *ysh1-32* and *ysh1-15* mutants (Fig. 4C; data not shown).


It has previously been shown that defective termination can result in production of polycistronic transcripts (Connelly and Manley 1988). In line with this notion, we observed read-through RNAs in *ysh1-12* cells at restrictive temperature. For example, the *YSH1* transcript extended into the downstream gene *DBP9* and *ADH1* mRNA extended into the downstream gene *YOL087C* (Fig. 4E). The read-through transcripts were not detectable in the *ysh1* mutant strains (*ysh1-13*, *ysh1-15*, *ysh1-32*) (data not shown), which is presumably resulting from a reduced stability of read-through RNAs in these strains. Taken together, the above results support a role for Ysh1p in termination of RNAP II transcription on mRNA genes.

### snoRNA 3' end formation is impaired in the *ysh1-12* strain

To address the question whether Ysh1p is required for the 3' end processing of snoRNA, we performed primer extension analysis on total RNA; in these experiments, the appearance of an extension signal is an indication of a defect in 3' end formation of these noncoding RNAs. In the *ysh1-12* strain, we detected accumulation of 3' extended transcripts of snR13, snR33, snR39B, snR46, snR47, and snR128 snoRNAs (Fig. 5A, lanes 3–6). In contrast, no extended RNAs were observed for the *ysh1-32* strain, when snR13, snR33, and snR47 were analyzed (data not shown), suggesting that the snoRNA processing defects in the *ysh1* strains are allele-specific. However, we were unable to detect any defects in the 3' end formation of several other snoRNAs such as snR3, snR50, snR71, and snR45 neither for *ysh1-12* nor for the positive control strains *nrđ1-102* and *sen1-1* (data not shown); this could be due to limitations of the primer extension procedure.

Next, we used transcription run-on analysis to measure RNAP II termination efficiency on the endogenous *SNR3* locus (Steinmetz et al. 2001). Two probes were used, one spanning the *SNR3* gene and its terminator sequences (probe A), and one downstream from this position (probe B) (Fig. 5B). These single-stranded DNAs do not hybridize with mRNA derived from the downstream ORF *YJR129c*; however, we cannot exclude the possibility that they could hybridize to antisense transcripts that may originate from this gene locus. Interestingly, *ysh1-12* mutant cells revealed up to sixfold higher density of transcribing RNAP II over the downstream probe compared to wild type (Fig. 5B,C), indicating defective termination on this transcription unit. In contrast, termination on the *SNR3* gene was unaffected in the *ysh1-32* strain (data not shown), which is consistent with the primer extension analysis above.

Conditional depletion of Ysh1p in the *brr5-td* degon strain was reported not to affect termination at snoRNA genes, as determined by RNAP II chromatin immunoprecipitation (ChIP) experiments (Kim et al. 2006). As these data conflicted with our observations, we included the *brr5-td* strain in our analysis. The degon-Ysh1p levels were significantly reduced after 1-h growth at 37°C, and almost completely diminished after 3 h (Fig. 5D). In contrast, the levels of Fip1p, Ssu72p, and Rna15p remained unchanged following depletion of Ysh1p. We compared snoRNA 3' end formation in *brr5-td*, its isogenic wild type (FY23), and *ysh1-12* strains by primer extension analysis as described above. 3'-Extended forms of the tested snoRNAs (snR13, snR33, snR39B, and snR47) were readily detected in the *ysh1-12* strain (Fig. 5E) but not following depletion of Ysh1p. These observations indicated that defects in snoRNA 3' end formation were apparent only with distinct *ysh1* mutant strains on a subset of snoRNA transcription units (see Discussion).




**FIGURE 4.** Mutations in *YSH1* impair correct RNAP II transcription termination on mRNA genes in vivo. (A) Slot hybridizations of run-on transcripts obtained after transcriptional run-on analysis (TRO) (Birise et al. 1998). The scheme represents the pGAL-*CYC1* gene construct that was used for TRO analysis. The order of M13 probes spanning the *CYC1* gene (P1–P6) and poly(A) site (position 506) are depicted. Wild-type and *ysh1-12* cells transformed with pUGCYC1 plasmid were grown in synthetic medium lacking uracil and containing 2% galactose under permissive growth conditions (30°C) and after shift for 1 h to restrictive temperature (15°C). P1–P6 represent probes complementary to *CYC1* transcripts as indicated on the scheme. Empty M13 was used as a background hybridization control. Hybridization of RNAP III transcripts to the tRNA probe is shown. (B) Quantitative analysis of transcriptional run-on profiles for the *ysh1-12* mutant. Values obtained by PhosphorImager scanning (Molecular Dynamics) were corrected by subtraction of the M13 background signal and normalized to the value of P1, which was fixed at 100%. Results shown in the diagram represent the average value of three independent experiments. (C) Quantitative analysis of transcriptional run-on profiles for the *ysh1-32* strain. Values were obtained as described in B. (D,E) Read-through mRNA transcripts accumulate in the *ysh1-12* strain in vivo. Northern blot analysis of total RNA extracted from wild-type and *ysh1-12* cells (lanes 1–3,7–9) grown in YPD medium at permissive temperature 30°C and (lanes 4–6,10–12) followed by a shift for 1, 3 or 6 h to 15°C. RNAs were resolved as described in Figure 3A. The filter was developed with random-primed labeled probe directed against (D) *YSH1* or (E) *ADH1*, then subsequently washed and probed for the RNA of the downstream gene (*DBP9* or *YOL087C*, respectively), as indicated. (\*) Unidentified intermediate-sized transcripts. Probing to the RNAP III transcript *SCR1* served as a loading control. (Bottom) Schematic representation of the (D) *YSH1-DBP9* and the (E) *ADH1-YOL087C* gene loci.

### Putative role for Ysh1p in mediating premature termination within the *NRD1* gene

Nrd1p controls its own expression through regulated premature termination, and levels of *NRD1* mRNA are increased in *nrd1*, *nab3*, and *sen1* mutant strains (Steinmetz et al. 2001; Arigo et al. 2006). Interestingly, we observed increased levels of *NRD1* mRNA in *ysh1-12* cells at the non-permissive temperature (Fig. 6A), which resulted in increased amounts of Nrd1p protein (Fig. 6B). Such a phenotype was not observed with the three *ysh1* ts mutant strains (data not shown). Since a similar increase in Nrd1p

protein was observed in the *nrd1-102* strain, this indicated that Ysh1p is involved in regulating Nrd1p levels. Moreover, some *NRD1* transcripts extended into the downstream gene *MRPL17*, resulting in a dicistronic *NRD1-MRPL17* transcript (Fig. 6A, lanes 4–6). We also observed an ~1-kb RNA that corresponded to 3'-truncated *NRD1* RNA in the wild-type strain as previously reported (Arigo et al. 2006). We detected this RNA with a probe directed against the 5' end of *NRD1* mRNA but not with a probe directed against the 3' end (data not shown). Interestingly, we found that this RNA was absent in *ysh1-12* (Fig. 6A).





**FIGURE 5.** Mutations in *ysh1-12* impair snoRNA 3' end formation and transcription termination. (A) 3' end extended snoRNA transcripts are produced in the *ysh1-12* mutant strain. Primer extension (PE) analysis of extended transcripts of several snoRNAs in the wild-type and *ysh1-12* strains was carried out with radioactively labeled oligonucleotides complementary to sequences located downstream from the mature 3' ends of the indicated snoRNAs. An oligonucleotide complementary to the mature U3 snoRNA was used as control. (B) Transcription termination on the *SNR3* snoRNA gene is impaired in *ysh1-12*. Slot hybridizations of run-on transcripts were obtained after TRO. (Top) The scheme represents the *SNR3-YJR129c* genomic locus and location of the TRO probes A and B. Note that *SNR3* and *YJR129c* are transcribed from opposite strands of the DNA, as indicated by the arrows. (Lanes 1,2) Wild-type and *ysh1-12* cells were grown in YPD medium containing 2% glucose under permissive growth conditions (30°C) and (lanes 3,4) after the shift to restrictive temperature (15°C) for 1 h. M13 slots are single-stranded phagemids with no insert and were used as a background hybridization control. The *ACT1* probe served as a positive control for hybridization. (C) Quantitative analysis of transcriptional run-on profiles for the *ysh1-12* mutant. Values obtained by PhosphorImager scanning (Molecular Dynamics) were corrected by subtraction of the M13 background signal and normalized to the value of probe A, which was set at 100%. Results shown in the diagram represent the average value of three independent experiments. (D) Western blot analysis of isogenic wild-type (FY23) and *brr5-td* degon strain extracts prepared from cells grown at 25°C and shifted for 1, 2, or 3 h to 37°C. Equal amounts of total protein were loaded in each lane. The blot was probed with antibody directed against Ysh1p and subsequently re-probed with antibodies directed against other CPF subunits, as indicated. (E) Primer extension analysis of extended transcripts of several snoRNAs in the wild-type FY23, *brr5-td* (both shifted for 2 h to 37°C), and *ysh1-12* (shifted for 6 h to 15°C) strains was carried out with radioactively labeled oligonucleotides complementary to sequences located downstream from the mature 3' ends of the indicated snoRNAs. (Lanes 1,3) Levels of the RNAP III transcript U6 served as a loading control and showed that lower amounts of total RNA extracted from the FY23 and *brr5-td* strains were used in this experiment.


The above observations on the *ysh1-12* phenotype suggested that Ysh1p might be involved in the regulation of *NRD1* expression. We speculated that Ysh1p could facilitate premature termination through endonucleolytic cleavage of the *NRD1* transcript. Direct evidence for occurrence of cleavage within *NRD1* would be the identification of downstream 3' cutoff products. Since such intermediates are predicted to be rapidly degraded by 5'-3' exonucleases, we analyzed total RNA extracted from the *rat1-1*, *xrn1Δ* double-mutant strain (Henry et al. 1994).

Primer extension analysis was carried out with oligonucleotide D, which is complementary to sequences located downstream from the predicted premature termination region (see scheme in Fig. 6C). Most interestingly, a primer extension stop was obtained that corresponded in size to a predicted downstream 3' cutoff product (Fig. 6C, lane 4). An equivalent primer extension stop was not detected with wild-type RNA, where a higher molecular weight band was detectable instead (Fig. 6C, lane 3); this stop most likely represented the 5' end of the *NRD1* mRNA. As a control, we used oligonucleotide U, which is complementary to the 5' end of the *NRD1* transcript and which gave primer extension stops that corresponded to the 5' end of *NRD1* mRNA, as expected (Fig. 6C, lanes 1,2). These data are consistent with the proposal that premature termination within the *NRD1* gene involves endonucleolytic cleavage of the primary transcript and that the putative endonuclease Ysh1p may provide this activity.

### Splicing is impaired in the *ysh1-12* strain

The *brr5-1* allele of *YSH1* originated from a screen for factors defective in RNA splicing (Noble and Guthrie 1996). To determine whether our *ysh1* mutant strains were defective in splicing, we analyzed *SNR17A* and *SNR17B* pre-snoRNAs using primer extension analysis (Noble and Guthrie 1996). We found that the *ysh1-12* strain exhibited a gradual accumulation of unspliced precursors derived from both *U3* genes following shift to nonpermissive temperature (Fig. 7A, lanes 3–6). The *rse1-1* mutant strain was analyzed as control and found to exhibit a similar splicing defect, as expected (Chen et al. 1998; data now shown). In contrast, the *ysh1-32* ts strain was fully competent in splicing (data not shown). Moreover, Northern blot analysis of intron-containing *ACT1* and *CYH2* transcripts in the *ysh1-12* strain detected extended species that were likely to represent unspliced pre-mRNAs (Fig. 7B). This was confirmed by primer extension analysis with an oligonucleotide complementary to the *ACT1* exon 2 (data not shown). The above results suggest a function for Ysh1p in pre-mRNA





**FIGURE 6.** Involvement of Ysh1p in the autoregulation of *NRD1* mRNA levels. (A) Increased levels of *NRD1* mRNA accumulate in the *ysh1-12* mutant strain. Northern blot analysis of *NRD1* mRNA in wild-type and *ysh1-12* strains was performed as in Figure 3A, with probes directed against the 5' end of *NRD1* mRNA, or against *MRPL17*, as indicated on the left. (Top) The scheme represents the *NRD1-MRPL17* gene loci; transcripts originating from the *NRD1* transcription start are indicated. Hybridization to the RNAP III transcript *SCR1* served as a loading control. (B) The amount of Nrd1 protein increases in *ysh1-12* cells. Western blot analysis of wild-type and mutant extracts prepared from cells grown in YPD at 25°C and shifted for 3 h to restrictive temperatures (15°C or 37°C, respectively). Equal amounts of total protein were loaded in each lane. Blots were probed with antibodies directed against the proteins indicated on the left. Act1p levels served as control for equal loading. (C) *NRD1* premature termination occurs via an endonucleolytic cleavage of the *NRD1* pre-mRNA. (Upper part) Schematic representation of the *NRD1* gene locus. Positions of the oligonucleotides U (+300 nt) and D (+1300 nt; relative to the ORF start) used in the primer extension analyses, the full-length and prematurely terminated *NRD1* mRNA transcripts, and of the putative 3' end cutoff product are shown. (Below, right panel) Primer extension analysis of total RNA extracted from wild-type (*W303*) and *rat1-1, xrn1Δ* strains (grown for 2 h at 37°C), carried out with radioactively labeled oligonucleotide D complementary to sequences located downstream from the predicted premature termination region. (Left panel) Oligonucleotide U complementary to the 5' end of *NRD1* transcript was used as an internal control.


and pre-snoRNA splicing and may reflect a functional coupling of 3' end processing and splicing events.

## DISCUSSION

In this study, we investigated the role of Ysh1p in mRNA and snoRNA metabolism. Our results are consistent with the view that Ysh1p plays an essential role in 3' end formation, probably through catalysis of endonucleolytic cleavage of the pre-mRNA. In addition to this, we provide evidence that Ysh1p is involved in termination of RNAP II both at mRNA and snoRNA genes, that it may contribute to the accumulation of distinct mRNAs independent of its role in 3' end formation, and that it may act in pre-mRNA splicing.

A striking outcome of our study is that distinct sets of phenotypes were associated with specific groups of *ysh1* mutant strains. Table 1 summarizes a broad range of

phenotypic observations, which were either resulting from our work or were previously described by others. Based on biochemical phenotypes two groups, class I and class II, can be formed by the ts strains (*ysh1-13*, *ysh1-15*, *ysh1-32*, and *brr5-td*) and the cs strains (*ysh1-12* and *brr5-1*), respectively. Extracts prepared from strains of both classes displayed defects in 3' end processing in vitro. The only exception is the *brr5-1* extract, which was reported not to have a clear cleavage defect (Chanfreau et al. 1996). Moreover, all analyzed ts and cs alleles of *YSH1* showed an aberrant pattern of *ACT1* poly(A) site usage. These observations are consistent with the proposal that Ysh1p indeed represents the 3' endonuclease. However, defects in 3' end cleavage and polyadenylation are generally associated with reduced mRNA stability. Consistently, class I mutants show under-accumulation of several mRNAs and a reduction of the global poly(A) tail length distribution. As this, in turn, leads to reduced protein levels of certain



**FIGURE 7.** The *ysh1-12* mutant is defective in splicing. (A) Primer extension analysis of spliced and unspliced *U3* transcripts in wild-type and *ysh1-12* mutant strains grown in YPD for up to 6 h at 15°C. Analysis was carried out with a radioactively labeled oligonucleotide complementary to sequences located in *U3* exon 2. HpaII-digested pBR322 fragments were 5' end labeled and served as markers. (Right) Bands corresponding to pre-*U3A*, pre-*U3B*, and mature *U3* snoRNA are indicated. (B) Northern blot analysis of total RNA extracted from wild-type and mutant *ysh1-12* cells, performed as in Figure 3A. Filters were developed with random-primed labeled probes directed against RNA transcripts as indicated on the left.

factors (such as, for example, Ssu72p), it cannot be excluded that defects observed in this class of strains are indirect. In stark contrast, class II mutations did not cause a general loss of mRNA levels. Moreover, phenotypes in snoRNA 3' end formation and termination, in the control of *NRD1* mRNA accumulation, and in RNA splicing were observed in the *ysh1-12* cs mutant strain. Notably, these phenotypes were not seen with class I mutants. Such striking differences of mutant phenotypes cannot easily be explained by a simple loss-of-function model and thus they complicate the mechanistic interpretation of the role of Ysh1p in different pathways of cellular RNA synthesis. Consistent with this idea, it was suggested previously that cold-sensitive phenotypes can unravel physiologic features that may be resistant to mutation to temperature sensitivity (Moir et al. 1982; Noble and Guthrie 1996).

In the torpedo model of RNAP II termination, cleavage at the poly(A) site is an obligatory step to provide an entry point for the Rat1/Xrn2 exonuclease, which contributes to termination through exonucleolytic degradation of the RNA fragment that remains associated with the polymerase following 3' end cleavage (Kim et al. 2004; West et al. 2004, 2008). Recently, this model has been challenged by the observation that although Rat1p associates with 3' end factors and may contribute to the assembly of the processing machinery, degradation of the downstream RNA appears not to be essential for termination (Luo et al. 2006). Interestingly, it was also shown that degradation of the 3' cleavage fragment is stimulated when coupled to 3' end cleavage in mammalian cells (Kaneko et al. 2007). Because it was demonstrated previously that exonuclease activity is required for Rat1p termination activity (Kim et al. 2004), the role of Rat1p in this process remains paradoxical (Rosonina et al. 2006). We found that class I and class II *ysh1* mutants were defective in transcription termination when assayed on the *CYC1* gene via transcriptional run-on. These results suggest that Ysh1p, the endonuclease acting in the 3' end processing of pre-mRNA transcripts, is also essential for RNAP II termination. Unlike other 3' end processing factors previously shown to be involved in termination, Ysh1p interacts neither with the CTD of RNAP II nor with RNA (Kyburz et al. 2003; Sadowski et al. 2003; M. Garas, B. Dichtl, and W. Keller, unpubl.). We, therefore, predict that it is the endonucleolytic cleavage that contributes mechanistically to transcription termination, possibly by providing the entry site for exonucleases. Moreover, transcriptional read-through products were detected in the class II *ysh1-12* and *brr5-1* strains but not in the class I strains. As both classes of mutants were defective in 3' end cleavage and in RNAP II termination, it seems that aberrant transcripts were somehow stabilized in class II mutants, pointing to a possible role of Ysh1p in RNA quality control.

The mechanism of 3' end processing and transcription termination on independently transcribed snoRNA genes remains unclear. Several lines of evidence support the notion that the canonical 3' end processing complex is involved in pre-snoRNA 3' end formation and that an endonucleolytic cleavage step may be associated with this process. Specific cleavage sites were identified on *SNR13* and *SNR47* snoRNAs as primer extension stops in *rat1-1*  $\Delta$ *xrn1* double-mutant cells (Fatica et al. 2000). Several subunits of CF IA and CPF complexes are important for snoRNA termination (see Introduction), and ChIP experiments clearly show that poly(A) factors and Rat1p exonuclease are recruited to snoRNA genes (Kim et al. 2006). ChIP analysis of elongating RNAP II suggested that degron-mediated depletion of Ysh1p did not affect snoRNA transcription termination (Kim et al. 2006); however, these experiments did not reveal whether or not cleavage occurs at the snoRNA 3' ends. In our analyses, the

**TABLE 1.** *ysh1/brr5* phenotypes

Allele	Class I				Class II	
	<i>ysh1-13</i>	<i>ysh1-15</i>	<i>ysh1-32</i>	<i>brr5-td</i>	<i>ysh1-12</i>	<i>brr5-1</i>
Growth phenotype	ts <sup>a</sup>	ts <sup>a</sup>	ts <sup>a</sup>	ts-degron <sup>b</sup>	cs <sup>a</sup>	cs <sup>c</sup>
Cleavage defect in vitro	Yes <sup>a</sup>	Yes <sup>a</sup>	Yes <sup>a</sup>	Yes <sup>b</sup>	Yes <sup>a</sup>	No <sup>d</sup>
Polyadenylation defect in vitro	Yes <sup>a</sup>	Yes <sup>a</sup>	Yes <sup>a</sup>	Yes <sup>b</sup>	Yes <sup>a</sup>	Yes <sup>d</sup>
mRNA stability	Unstable <sup>a</sup>	Unstable <sup>a</sup>	Unstable <sup>a</sup>		Stable <sup>a</sup>	
<i>ACT1</i> poly(A) site selection defect	Yes <sup>a</sup>	Yes <sup>a</sup>	Yes <sup>a</sup>		Yes <sup>a</sup>	
Termination defect on mRNA genes	ND	Yes <sup>a</sup>	Yes <sup>a</sup>	Yes <sup>e</sup>	Yes <sup>a</sup>	No <sup>b</sup>
Read-through mRNA transcripts	No <sup>a</sup>	No <sup>a</sup>	No <sup>a</sup>		Yes <sup>a</sup>	Yes <sup>d</sup>
<i>NRD1</i> mRNA up-regulation	No <sup>a</sup>	No <sup>a</sup>	No <sup>a</sup>		Yes <sup>a</sup>	
Termination defect on snoRNA genes	ND	ND	No <sup>a</sup>	No <sup>e</sup>	Yes <sup>a</sup>	
3' End extended snoRNA transcripts	ND	ND	No <sup>a</sup>		Yes <sup>a</sup>	
Splicing defect	ND	ND	No <sup>a</sup>		Yes <sup>a</sup>	Yes <sup>c</sup>

(ND) Not determined.

<sup>a</sup>This study.<sup>b</sup>Zhelkovsky et al. (2006).<sup>c</sup>Noble and Guthrie (1996).<sup>d</sup>Chanfreau et al. (1996).<sup>e</sup>Kim et al. (2006).

*brr5-td* strain did not accumulate 3' extended snoRNA transcripts following depletion of Ysh1p, suggesting that the protein may not play an essential role in this process. Nevertheless, our analysis of the class II *ysh1-12* allele demonstrated that Ysh1p has a function in snoRNA 3' end formation and transcription termination. One potential explanation for this apparent paradox apart from other possible indirect effects could be that for reasons not known, the 3' extended snoRNA transcripts are stabilized in the *ysh1-12* cs mutant, but not in the *ysh1* ts/td strains. Canonical 3' end processing factors and Rat1p are recruited to genes transcribed by RNAP II regardless of whether they encode mRNA or snoRNA (Kim et al. 2006). It is possible that an early stage of termination is similar for both types of transcripts and that in a later phase of termination a combination of factors used by RNAP II for 3' end formation may vary depending on the specific snoRNA gene (Kim et al. 2006). A speculative scenario therefore is that the phenotype associated with the cs *ysh1-12* strain might unravel an early stage of the snoRNA transcription termination that requires a certain function of Ysh1p/CPF.

Whereas a mechanistic understanding of RNAP II termination at the end of transcription units is emerging, there is little information available on regulated premature termination within a transcription unit. Indeed, the case of *NRD1* autoregulation is one of the best documented examples (Arigo et al. 2006). It has been found that the RNA-binding proteins Nrd1p and Nab3p recognize sequence elements within the *NRD1* mRNA to trigger RNAP II termination prematurely, i.e., before the "regular" poly(A)-dependent terminator in the 3'-UTR is transcribed (Arigo et al. 2006). In agreement with these observations, the early *NRD1* terminator resembles Nrd1p/Nab3p-

dependent snoRNA terminators (Steinmetz et al. 2001; Arigo et al. 2006). It was shown that this autoregulation involves the exosome and the TRAMP complex (Arigo et al. 2006), but it remains unclear whether endonucleolytic cleavage and possibly a torpedo-like termination mechanism may play a role as well. We observed elevated levels of the full-length *NRD1* mRNA and of the Nrd1p protein in the *ysh1-12* strain, suggesting a negative role for Ysh1p in the regulation of *NRD1* expression. Therefore it is possible that the increased amounts of the Nrd1p protein cause the snoRNA formation defect in the *ysh1-12* mutant due to an imbalance in the stoichiometry of the components that form the Nrd1 complex. As Nrd1p bound to Nab3p and Sen1p associates with the RNAP II CTD, the excess free Nrd1 could displace this complex from the CTD, resulting in termination defects. Importantly, we were also able to demonstrate the existence of a *NRD1* 3' cutoff product in *rat1-1*,  $\Delta xrn1$  cells, which is consistent with a mechanism involving endonucleolytic cleavage. These results imply a role for Ysh1p in *NRD1* premature termination and thus for regulation of *NRD1* mRNA levels, possibly via its cleavage activity. Interestingly, it was previously suggested that the Nrd1p/Nab3p/Sen1p pathway acts primarily on transcripts that are 500 nucleotides (nt) or shorter (Steinmetz et al. 2001, 2006b). The observation that the prematurely terminated *NRD1* RNA is at least 1000 nt long is consistent with the involvement of Ysh1p, which is a component of CPF that preferentially mediates termination of longer transcripts. In addition to Ssu72p (Ganem et al. 2003; Steinmetz and Brow 2003) and Hrp1p (Kuehner and Brow 2008), Ysh1p may therefore represent another example of a 3' end processing factor involved in regulated premature RNAP II termination.

Interestingly, Northern blot analyses showed that the levels of several other mRNAs were elevated in the *ysh1-12* strain as well. Among the messages accumulating in the *ysh1-12* mutant, we identified *NAB2*, which is believed to autoregulate the levels of its own mRNA in a process that requires the nuclear exosome component Rrp6p (Roth et al. 2005). This post-transcriptional mechanism involves a sequence of 26 adenosines (A<sub>26</sub>) in the *NAB2* 3'-UTR, which represses *NAB2* 3' end formation and subsequently directs the transcript for degradation by Rrp6p (Roth et al. 2005). The increased concentration of *NAB2* mRNA in the *ysh1-12* strain suggests a contribution of Ysh1p to the autogenic control of *NAB2* levels. Interestingly, we found that *YSH1* genetically interacts with the nuclear exosome subunit Rrp6p, as deletions of *RRP6* suppressed the *cs* growth phenotype of *ysh1-12* cells (data not shown). However, further investigations of the involvement of Ysh1p in *NAB2* autoregulation are required. Our results indicate that Ysh1p directly or indirectly influences the nuclear turnover of mRNAs. Class II mutations in *ysh1* alleles could inhibit correct mRNP assembly, which in turn might lead to impaired mRNA export and retention of the mRNA in the nucleus. An alternative and more intriguing possibility would be that the endonucleolytic cleavage activity of Ysh1p directly contributes to RNA surveillance. In this case, we would predict that also class I mutations result in some up-regulation of *NAB2* and/or *NRD1* mRNAs; however, such phenotypes were clearly not observed in these mutant strains. Taken together, our results support the idea that Ysh1p is involved in the negative regulation of certain mRNAs and that it may contribute to RNA surveillance.

Our analyses of the *ysh1-12* allele revealed a splicing defect that correlates well with the phenotype observed previously

for the *cs brr5-1* allele (Chanfreau et al. 1996; Noble and Guthrie 1996). However, the molecular basis of these defects remains unclear. snRNA synthesis appeared normal in *ysh1* mutant strains (data not shown), and this excludes the possibility that snRNA depletion underlies the defect. We cannot rule out the possibility that secondary effects resulting from under-accumulation of critical splicing components contribute to the phenotype. Nevertheless, the possibility remains that Ysh1p acts in the coupling of splicing and 3' end processing. Evidence for such coupling is ample in mammalian cells but so far was not reported in yeast. Mechanisms of coupling include physical interactions between components of the 3' end formation and splicing machineries, e.g., an interaction between CPSF and the U2 snRNP (Kyburz et al. 2006), and between U2AF 65 and CF I<sub>m</sub> (Millevoi et al. 2006). Interestingly, Rse1p, the yeast homolog of SF3b130 and a component of U2 snRNP, genetically interacts with different subunits of CPF (A. Kyburz and W. Keller, unpubl.). Future experiments will have to address the question whether there is, indeed, a functional interdependence of splicing and poly(A) addition in yeast.

## MATERIALS AND METHODS

### Yeast growth and strain construction

The *S. cerevisiae* strains used in this study and their genotypes are listed in Table 2. Construction and isolation of temperature-sensitive *ysh1* alleles was done with an error-prone PCR and in vivo gap-repair approach as described previously (Dichtl et al. 2002b); details are available upon request. Manipulations and growth of *S. cerevisiae* were carried out by established procedures. Typically, for temperature shift experiments, yeast cultures were grown to midexponential phase at 23°C (*ts* strains) or 30°C (*cs*

**TABLE 2.** Yeast strains used in this study

Strain	Genotype	Reference
W303-1A	<i>MATa ade2 leu2 ura3 trp1-1 his3</i>	Thomas and Rothstein (1989)
LM109	<i>MATa ura3-1 trp1Δ ade2-1 leu2-3,112 his3-11,15 TRP1::ysh1 [YSH1-URA1-CEN]</i>	Jenny et al. (1996)
YMG44 ( <i>YSH1</i> )	<i>MATa ura3-1 trp1Δ ade2-1 leu2-3,112 his3-11,15 TRP1::ysh1 [YSH-HIS3-CEN]</i>	This study
YMG17 ( <i>ysh1-12</i> )	<i>MATa ura3-1 trp1Δ ade2-1 leu2-3,112 his3-11,15 TRP1::ysh1 [ysh1-12-HIS3-CEN]</i>	This study
YMG29 ( <i>ysh1-32</i> )	<i>MATa ura3-1 trp1Δ ade2-1 leu2-3,112 his3-11,15 TRP1::ysh1 [ysh1-32-HIS3-CEN]</i>	This study
YMG16 ( <i>ysh1-13</i> )	<i>MATa ura3-1 trp1Δ ade2-1 leu2-3,112 his3-11,15 TRP1::ysh1 [ysh1-13-HIS3-CEN]</i>	This study
YMG27 ( <i>ysh1-15</i> )	<i>MATa ura3-1 trp1Δ ade2-1 leu2-3,112 his3-11,15 TRP1::ysh1 [ysh1-15-HIS3-CEN]</i>	This study
YJC1166 ( <i>nrd1-102</i> )	<i>MATα ura3Δ0 his3Δ1 leu2Δ0 met15Δ0 nrd1-102</i>	Steinmetz et al. (2001)
FY23	<i>MATa, ura3-52, leu2Δ1, trp1Δ63</i>	Winston et al. (1995)
YAZ212 ( <i>brr5-td</i> )	<i>MATa ura3-52 leu2Δ1 trp1Δ63 brr5::DHFR<sup>ts</sup>-BRR5</i>	Zhelkovsky et al. (2006)
<i>rna15-1</i>	<i>MATa ura3-1 trp1-1 ade2-1 leu2-3,112 his3-11,15 rna15-1</i>	Minvielle-Sebastia et al. (1994)
D174 ( <i>rat1-1 xrn1Δ</i> )	<i>MATa ade2-1 xrn1::URA3 rat1-1</i>	Henry et al. (1994)



strains), and transferred to a water bath at 37°C or 15°C, respectively. Yeast cells were grown in rich YPD medium (2% glucose, 2% bacto-tryptone, 1% yeast extract) or in synthetic drop-out medium (2% glucose or galactose, respectively, 0.67% yeast nitrogen base, 1× amino acids) as indicated in the figure legends. For the droplet test, strains were grown overnight and diluted to an OD<sub>600</sub> of 0.1, 0.01, 0.001, and 0.0001. Five microliters of each dilution were spotted on five different YPAD plates and incubated at 15°C, 23°C, 30°C, 33°C, and 37°C.

### RNA analyses

Yeast total RNA was extracted with the hot phenol method. Northern analyses and RNase H experiments were carried out as described (Dichtl et al. 2002b, 2004). Oligonucleotides used as probes in Northern analyses were routinely labeled with 30 μCi of [ $\gamma$ -<sup>32</sup>P]ATP and T4 polynucleotide kinase (New England Biolabs). Random-primed Northern probes were as published previously (Dichtl et al. 2002a,b, 2004) or were obtained using the following oligonucleotides:

ASN1 (ASN1-For, 5'-ATGTGTGGTATTTTCGCCCGC; ASN1-Rev, 5'-GAATCGTATACGTGGCCCCGG);  
 DBP9 (DBP9-For, 5'-AGCTTGAAGGCGAGCCTG; DBP9-Rev, 5'-TCATTTGAAGTTCTTCAACG);  
 RPB1 (RPB1-For, 5'-CAGAGGCTAAAAAGAAAGTTTTG; RPB1-Rev, 5'-GATCTATGTGGAAAAGTTTGTG);  
 SCRI (SCRI-For, 5'-AGGCTGTAATGGCTTCTGGTGGG; SCRI-Rev, 5'-TATGGTTCAGGACACACTCCATCCC);  
 SEN1 (SEN1-For, 5'-CTCAATACGTCGCGAGGCTGAGGC; SEN1-Rev, 5'-GAAAGAACAGTTGGTGGTAGTTG);  
 YOL087C (YOL087C-For, 5'-GTCCTGATAATGTACACGATGG; YOL087C-Rev, 5'-GTTTCATGGTGATGATGATGATGG); and  
 YSH1 (YSH1-For, 5'-GTCAATGGTATCAAATTTACGGC; YSH1-Rev, 5'-GGTTATTTCCGGATTATTTATGG).

Total poly(A) tail labeling was performed as described previously (Minvielle-Sebastia et al. 1998). The poly(A) tails were separated on a 10% denaturing polyacrylamide/8.3 M urea/TBE gel; images were generated by a PhosphorImager (STORM). Transcriptional run-on (TRO) analyses on the *CYCI* gene with the multi-copy plasmid pUGCYC were performed according to Birse et al. (1998). TRO analysis on the endogenous *SNR3* gene was done as described previously (Steinmetz et al. 2001) with the probes A and B. Primer extension analysis (PE) was carried out as described previously (Beltrame and Tollervey 1992) with SuperScript III reverse transcriptase (Invitrogen) and 6 μg of total RNA. Extension primers were as published (Steinmetz et al. 2001) and additionally as the following:

SNR46 (5'-ATCGACCAGCTCTTTAGCATCC);  
 U3exon2 (5'-GAGCCACTGAATCCAACCTGGTTGAT);  
 U6 (5'-GGGGAAGTCTGATCATCTCTGTATTG);  
 NRD1 U (5'-CGTGTATTATGGCATGGGCACAAG); and  
 NRD1 D (5'-CAGCGTCCGTGAGCCTGTGCATAG).

### Extract preparation and in vitro cleavage and polyadenylation assays

3' End processing extracts were made following the procedure described previously (Ohnacker et al. 2000). Cleavage and poly-

adenylation assays were carried out according to Minvielle-Sebastia et al. (1994). When cleavage only was assayed, EDTA replaced magnesium acetate (MgAc), and ATP was omitted. Internally [<sup>32</sup>P]-labeled *CYCI* and *CYCI-pre* RNA substrates were produced by in vitro run-off transcription. For reactions at restrictive temperatures, both the reaction mix and protein extracts were preincubated separately for 10 min at the respective restrictive temperatures, combined, and assayed for 1 h. Each of the processing extracts was prepared independently at least three times and subsequently tested for cleavage and polyadenylation activity in vitro in order to confirm the specific processing defect. CPF was obtained by affinity purification as described previously (Ohnacker et al. 2000).

### Western blotting analysis

Proteins were separated on 10% SDS polyacrylamide gels, transferred to nitrocellulose membranes, and incubated according to standard procedures with the antibodies indicated in the legends to the figures. The sources of the antibodies used in this work were the following: anti-Ysh1p (Jenny et al. 1996); anti-Ssu72p (Dichtl et al. 2002a); anti-Rna15p (Minvielle-Sebastia et al. 1994); anti-Fip1p and anti-Pap1p (Preker et al. 1995); anti-Pfs2p (Ohnacker et al. 2000); anti-Nrd1p (Steinmetz and Brow 1998); anti-Act1p (Chemicon International). Peroxidase-conjugated swine anti-rabbit and rabbit anti-mouse immunoglobulins (DAKO) served as secondary antibodies for detection of the primary antibodies.

### ACKNOWLEDGMENTS

We thank Drs. Claire Moore and Jeffrey Corden for generous gifts of yeast strains, Drs. David Brow and Eric Steinmetz for anti-Nrd1p antibody, and Dr. Bertrand Paguet for purified CPF. We also thank the reviewers for constructive suggestions. We are grateful to Drs. Stepanka Vanacova and Bertrand Paguet for comments and discussions on the project and for improving the manuscript. W.K. was supported by the University of Basel, the Swiss National Science Fund, and, as part of the European Science Foundation EUROCORES Programme EuroDYNA, from funds of the EC Sixth Framework Programme, under contract ERAS-CT-2003-980409. B.D. was supported by the Swiss National Science Fund (Grant no. PP00A-102941).

Received July 26, 2008; accepted September 15, 2008.

### REFERENCES

- Allmang, C., Kufel, J., Chanfreau, G., Mitchell, P., Petfalski, E., and Tollervey, D. 1999. Functions of the exosome in rRNA, snoRNA and snRNA synthesis. *EMBO J.* **18**: 5399–5410.
- Aravind, L. 1999. An evolutionary classification of the metallo-β-lactamase fold proteins. *In Silico Biol.* **1**: 69–91.
- Arigo, J.T., Carroll, K.L., Ames, J.M., and Corden, J.L. 2006. Regulation of yeast NRD1 expression by premature transcription termination. *Mol. Cell* **21**: 641–651.
- Beltrame, M. and Tollervey, D. 1992. Identification and functional analysis of two U3 binding sites on yeast pre-ribosomal RNA. *EMBO J.* **11**: 1531–1542.
- Bentley, D.L. 2005. Rules of engagement: Co-transcriptional recruitment of pre-mRNA processing factors. *Curr. Opin. Cell Biol.* **17**: 251–256.

- Birse, C.E., Minvielle-Sebastia, L., Lee, B.A., Keller, W., and Proudfoot, N.J. 1998. Coupling termination of transcription to messenger RNA maturation in yeast. *Science* **280**: 298–301.
- Buratowski, S. 2005. Connections between mRNA 3' end processing and transcription termination. *Curr. Opin. Cell Biol.* **17**: 257–261.
- Butler, J.S. 2002. The yin and yang of the exosome. *Trends Cell Biol.* **12**: 90–96.
- Callebaut, I., Moshous, D., Mornon, J.P., and de Villartay, J.P. 2002. Metallo- $\beta$ -lactamase fold within nucleic acids processing enzymes: The  $\beta$ -CASP family. *Nucleic Acids Res.* **30**: 3592–3601.
- Chanfreau, G., Noble, S.M., and Guthrie, C. 1996. Essential yeast protein with unexpected similarity to subunits of mammalian cleavage and polyadenylation specificity factor (CPSF). *Science* **274**: 1511–1514.
- Chanfreau, G., Legrain, P., and Jacquier, A. 1998. Yeast RNase III as a key processing enzyme in small nucleolar RNAs metabolism. *J. Mol. Biol.* **284**: 975–988.
- Chen, E.J., Frand, A.R., Chitouras, E., and Kaiser, C.A. 1998. A link between secretion and pre-mRNA processing defects in *Saccharomyces cerevisiae* and the identification of a novel splicing gene, RSE1. *Mol. Cell. Biol.* **18**: 7139–7146.
- Cheng, H., He, X., and Moore, C. 2004. The essential WD repeat protein Swd2 has dual functions in RNA polymerase II transcription termination and lysine 4 methylation of histone H3. *Mol. Cell. Biol.* **24**: 2932–2943.
- Connelly, S. and Manley, J.L. 1988. A functional mRNA polyadenylation signal is required for transcription termination by RNA polymerase II. *Genes & Dev.* **2**: 440–452.
- Conrad, N.K., Wilson, S.M., Steinmetz, E.J., Patturajan, M., Brow, D.A., Swanson, M.S., and Corden, J.L. 2000. A yeast heterogeneous nuclear ribonucleoprotein complex associated with RNA polymerase II. *Genetics* **154**: 557–571.
- Daiyasu, H., Osaka, K., Ishino, Y., and Toh, H. 2001. Expansion of the zinc metallo-hydrolase family of the  $\beta$ -lactamase fold. *FEBS Lett.* **503**: 1–6.
- Dheur, S., Vo, L.T.A., Voisinet-Hakil, F., Minet, M., Schmitter, J.-M., Lacroute, F., Wyers, F., and Minvielle-Sebastia, L. 2003. Pti1p and Ref2p found in association with the mRNA 3' end formation complex direct snoRNA maturation. *EMBO J.* **22**: 2831–2840.
- Dichtl, B., Blank, D., Ohnacker, M., Friedlein, A., Roeder, D., Langen, H., and Keller, W. 2002a. A role for SSU72 in balancing RNA polymerase II transcription elongation and termination. *Mol. Cell* **10**: 1139–1150.
- Dichtl, B., Blank, D., Sadowski, M., Hübner, W., Weiser, S., and Keller, W. 2002b. Yhh1p/Cft1p directly links poly(A) site recognition and RNA polymerase II transcription termination. *EMBO J.* **21**: 4125–4135.
- Dichtl, B., Aasland, R., and Keller, W. 2004. Functions for *S. cerevisiae* Swd2p in 3' end formation of specific mRNAs and snoRNAs and global histone 3 lysine 4 methylation. *RNA* **10**: 965–977.
- Fatica, A., Morlando, M., and Bozzoni, I. 2000. Yeast snoRNA accumulation relies on a cleavage-dependent/polyadenylation-independent 3'-processing apparatus. *EMBO J.* **19**: 6218–6229.
- Ganem, C., Devaux, F., Torchet, C., Jacq, C., Quevillon-Cheruel, S., Labesse, G., Facca, C., and Faye, G. 2003. Ssu72 is a phosphatase essential for transcription termination of snoRNAs and specific mRNAs in yeast. *EMBO J.* **22**: 1588–1598.
- Gilmour, D.S. and Fan, R. 2008. Derailing the locomotive: Transcription termination. *J. Biol. Chem.* **283**: 661–664.
- Henry, Y., Wood, H., Morrissey, J.P., Petfalski, E., Kearsley, S., and Tollervey, D. 1994. The 5' end of yeast 5.8S rRNA is generated by exonucleases from an upstream cleavage site. *EMBO J.* **13**: 2452–2463.
- Jenny, A., Minvielle-Sebastia, L., Preker, P.J., and Keller, W. 1996. Sequence similarity between the 73-kilodalton protein of mammalian CPSF and a subunit of yeast polyadenylation factor I. *Science* **274**: 1514–1517.
- Kaneko, S., Rozenblatt-Rosen, O., Meyerson, M., and Manley, J.L. 2007. The multifunctional protein p54nrb/PSF recruits the exonuclease XRN2 to facilitate pre-mRNA 3' processing and transcription termination. *Genes & Dev.* **21**: 1779–1789.
- Kim, M., Krogan, N.J., Vasiljeva, L., Rando, O.J., Nedeá, E., Greenblatt, J.F., and Buratowski, S. 2004. The yeast Rat1 exonuclease promotes transcription termination by RNA polymerase II. *Nature* **432**: 517–522.
- Kim, M., Vasiljeva, L., Rando, O.J., Zhelkovsky, A., Moore, C., and Buratowski, S. 2006. Distinct pathways for snoRNA and mRNA termination. *Mol. Cell* **24**: 723–734.
- Kuehner, J.N. and Brow, D.A. 2008. Regulation of a eukaryotic gene by GTP-dependent start site selection and transcription attenuation. *Mol. Cell* **31**: 201–211.
- Kyburz, A., Sadowski, M., Dichtl, B., and Keller, W. 2003. The role of the yeast cleavage and polyadenylation factor subunit Ydh1p/Cft2p in pre-mRNA 3' end formation. *Nucleic Acids Res.* **31**: 3936–3945.
- Kyburz, A., Friedlein, A., Langen, H., and Keller, W. 2006. Direct interactions between subunits of CPSF and the U2 snRNP contribute to the coupling of pre-mRNA 3' end processing and splicing. *Mol. Cell* **23**: 195–205.
- Logan, J., Falck-Pedersen, E., Darnell Jr., J.E., and Shenk, T. 1987. A poly(A) addition site and a downstream termination region are required for efficient cessation of transcription by RNA polymerase II in the mouse  $\beta$  maj-globin gene. *Proc. Natl. Acad. Sci.* **84**: 8306–8310.
- Luo, W. and Bentley, D. 2004. A ribonucleolytic rat torpedoes RNA polymerase II. *Cell* **119**: 911–914.
- Luo, W., Johnson, A.W., and Bentley, D.L. 2006. The role of Rat1 in coupling mRNA 3' end processing to transcription termination: Implications for a unified allosteric-torpedo model. *Genes & Dev.* **20**: 954–965.
- Mandart, E. and Parker, R. 1995. Effects of mutations in the *Saccharomyces cerevisiae* RNA14, RNA15, and PAPI1 genes on polyadenylation in vivo. *Mol. Cell. Biol.* **15**: 6979–6986.
- Mandel, C.R., Kaneko, S., Zhang, H., Gebauer, D., Vethantham, V., Manley, J.L., and Tong, L. 2006. Polyadenylation factor CPSF-73 is the pre-mRNA 3' end processing endonuclease. *Nature* **444**: 953–956.
- Millevoi, S., Loulergue, C., Dettwiler, S., Karaa, S.Z., Keller, W., Antoniou, M., and Vagner, S. 2006. An interaction between U2AF 65 and CF I<sub>m</sub> links the splicing and 3' end processing machineries. *EMBO J.* **25**: 4854–4864.
- Minvielle-Sebastia, L., Preker, P.J., and Keller, W. 1994. RNA14 and RNA15 proteins as components of a yeast pre-mRNA 3' end processing factor. *Science* **266**: 1702–1705.
- Minvielle-Sebastia, L., Beyer, K., Krecic, A.M., Hector, R.E., Swanson, M.S., and Keller, W. 1998. Control of cleavage site selection during mRNA 3' end formation by a yeast hnRNP. *EMBO J.* **17**: 7454–7468.
- Moir, D., Stewart, S.E., Osmond, B.C., and Botstein, D. 1982. Cold-sensitive cell-division-cycle mutants of yeast: Isolation, properties, and pseudoreversion studies. *Genetics* **100**: 547–563.
- Morlando, M., Greco, P., Dichtl, B., Fatica, A., Keller, W., and Bozzoni, I. 2002. Functional analysis of yeast snoRNA and snRNA 3' end formation mediated by uncoupling of cleavage and polyadenylation. *Mol. Cell. Biol.* **22**: 1379–1389.
- Nedeá, E., He, X., Kim, M., Pootoolal, J., Zhong, G., Canadien, V., Hughes, T., Buratowski, S., Moore, C.L., and Greenblatt, J. 2003. Organization and function of APT, a subcomplex of the yeast cleavage and polyadenylation factor involved in the formation of mRNA and small nucleolar RNA 3' ends. *J. Biol. Chem.* **278**: 33000–33010.
- Nedeá, E., Nalbant, D., Xia, D., Theoharis, N.T., Suter, B., Richardson, C.J., Tatchell, K., Kislinger, T., Greenblatt, J.F., and Nagy, P.L. 2008. The Glc7 phosphatase subunit of the cleavage and polyadenylation factor is essential for transcription termination on snoRNA genes. *Mol. Cell* **29**: 577–587.
- Noble, S.M. and Guthrie, C. 1996. Identification of novel genes required for yeast pre-mRNA splicing by means of cold-sensitive mutations. *Genetics* **143**: 67–80.

- Ohnacker, M., Barabino, S.M., Preker, P.J., and Keller, W. 2000. The WD-repeat protein Pfs2p bridges two essential factors within the yeast pre-mRNA 3' end processing complex. *EMBO J.* **19**: 37–47.
- Orozco, I.J., Kim, S.J., and Martinson, H.G. 2002. The poly(A) signal, without the assistance of any downstream element, directs RNA polymerase II to pause in vivo and then to release stochastically from the template. *J. Biol. Chem.* **277**: 42899–42911.
- Preker, P.J., Lingner, J., Minvielle-Sebastia, L., and Keller, W. 1995. The FIP1 gene encodes a component of a yeast pre-mRNA polyadenylation factor that directly interacts with poly(A) polymerase. *Cell* **81**: 379–389.
- Preker, P.J., Ohnacker, M., Minvielle-Sebastia, L., and Keller, W. 1997. A multisubunit 3' end processing factor from yeast containing poly(A) polymerase and homologues of the subunits of mammalian cleavage and polyadenylation specificity factor. *EMBO J.* **16**: 4727–4737.
- Proudfoot, N. 2004. New perspectives on connecting messenger RNA 3' end formation to transcription. *Curr. Opin. Cell Biol.* **16**: 272–278.
- Proudfoot, N.J., Furger, A., and Dye, M.J. 2002. Integrating mRNA processing with transcription. *Cell* **108**: 501–512.
- Rosonina, E., Kaneko, S., and Manley, J.L. 2006. Terminating the transcript: Breaking up is hard to do. *Genes & Dev.* **20**: 1050–1056.
- Roth, K.M., Wolf, M.K., Rossi, M., and Butler, J.S. 2005. The nuclear exosome contributes to autogenous control of NAB2 mRNA levels. *Mol. Cell. Biol.* **25**: 1577–1585.
- Ryan, K., Calvo, O., and Manley, J.L. 2004. Evidence that polyadenylation factor CPSF-73 is the mRNA 3' processing endonuclease. *RNA* **10**: 565–573.
- Sadowski, M., Dichtl, B., Hübner, W., and Keller, W. 2003. Independent functions of yeast Pcf11p in pre-mRNA 3' end processing and in transcription termination. *EMBO J.* **22**: 2167–2177.
- Sheldon, K.E., Mauger, D.M., and Arndt, K.M. 2005. A requirement for the *Saccharomyces cerevisiae* Paf1 complex in snoRNA 3' end formation. *Mol. Cell* **20**: 225–236.
- Steinmetz, E.J. and Brow, D.A. 1998. Control of pre-mRNA accumulation by the essential yeast protein Nrd1 requires high-affinity transcript binding and a domain implicated in RNA polymerase II association. *Proc. Natl. Acad. Sci.* **95**: 6699–6704.
- Steinmetz, E.J. and Brow, D.A. 2003. Ssu72 protein mediates both poly(A)-coupled and poly(A)-independent termination of RNA polymerase II transcription. *Mol. Cell. Biol.* **23**: 6339–6349.
- Steinmetz, E.J., Conrad, N.K., Brow, D.A., and Corden, J.L. 2001. RNA-binding protein Nrd1 directs poly(A)-independent 3' end formation of RNA polymerase II transcripts. *Nature* **413**: 327–331.
- Steinmetz, E.J., Ng, S.B., Cloute, J.P., and Brow, D.A. 2006a. *Cis-* and *trans-*acting determinants of transcription termination by yeast RNA polymerase II. *Mol. Cell. Biol.* **26**: 2688–2696.
- Steinmetz, E.J., Warren, C.L., Kuehner, J.N., Panbehi, B., Ansari, A.Z., and Brow, D.A. 2006b. Genome-wide distribution of yeast RNA polymerase II and its control by Sen1 helicase. *Mol. Cell* **24**: 735–746.
- Thomas, B.J. and Rothstein, R. 1989. Elevated recombination rates in transcriptionally active DNA. *Cell* **56**: 619–630.
- Ursic, D., Himmel, K.L., Gurley, K.A., Webb, F., and Culbertson, M.R. 1997. The yeast SEN1 gene is required for the processing of diverse RNA classes. *Nucleic Acids Res.* **25**: 4778–4785.
- van Hoof, A., Lennertz, P., and Parker, R. 2000. Yeast exosome mutants accumulate 3'-extended polyadenylated forms of U4 small nuclear RNA and small nucleolar RNAs. *Mol. Cell. Biol.* **20**: 441–452.
- West, S., Gromak, N., and Proudfoot, N.J. 2004. Human 5' → 3' exonuclease Xrn2 promotes transcription termination at co-transcriptional cleavage sites. *Nature* **432**: 522–525.
- West, S., Proudfoot, N.J., and Dye, M.J. 2008. Molecular dissection of mammalian RNA polymerase II transcriptional termination. *Mol. Cell* **29**: 600–610.
- Wickens, M. and Gonzalez, T.N. 2004. Knives, accomplices, and RNA. *Science* **306**: 1299–1300.
- Winston, F., Dollard, C., and Ricupero-Hovasse, S.L. 1995. Construction of a set of convenient *Saccharomyces cerevisiae* strains that are isogenic to S288C. *Yeast* **11**: 53–55.
- Zhang, Z. and Gilmour, D.S. 2006. Pcf11 is a termination factor in *Drosophila* that dismantles the elongation complex by bridging the CTD of RNA polymerase II to the nascent transcript. *Mol. Cell* **21**: 65–74.
- Zhang, Z., Fu, J., and Gilmour, D.S. 2005. CTD-dependent dismantling of the RNA polymerase II elongation complex by the pre-mRNA 3' end processing factor, Pcf11. *Genes & Dev.* **19**: 1572–1580.
- Zhao, J., Hyman, L., and Moore, C. 1999. Formation of mRNA 3' ends in eukaryotes: Mechanism, regulation, and interrelationships with other steps in mRNA synthesis. *Microbiol. Mol. Biol. Rev.* **63**: 405–445.
- Zhelkovsky, A., Tacahashi, Y., Nasser, T., He, X., Sterzer, U., Jensen, T.H., Domdey, H., and Moore, C. 2006. The role of the Brr5/Ysh1 C-terminal domain and its homolog Syc1 in mRNA 3' end processing in *Saccharomyces cerevisiae*. *RNA* **12**: 435–445.