



Letter to the Editor: Backbone resonance assignment of the 298 amino acid catalytic domain of protein tyrosine phosphatase 1B (PTP1B)

Sebastian Meier^a, Yu-Chin Li^b, James Koehn^b, Isidoros Vlattas^c, James Wareing^c, Wolfgang Jahnke^d, Lawrence P. Wennogle^c & Stephan Grzesiek^{a,*}

^aDivision of Structural Biology and Division of Biophysical Chemistry, Biozentrum, University of Basel, 4056 Basel, Switzerland; ^bCore Technology Department and ^cNovartis Institute for Biomedical Research, Novartis Pharmaceuticals Corporation, 556 Morris Avenue, Summit, NJ 07901-1398, U.S.A.; ^dCore Technologies, Novartis Pharma AG, CH-4002 Basel, Switzerland

Received 26 July 2002; Accepted 13 August 2002

Key words: diabetes, heteronuclear NMR, obesity

Biological context

Protein activation via tyrosine phosphorylation by tyrosine kinases and the concomitant deactivation/dephosphorylation by protein tyrosine phosphatases (PTPs) plays a central role in cellular signaling. Tyrosine phosphorylation regulates cell homeostasis by the control of gene-transcription, cell-cycle progression, cell growth, cell metabolism, immune response as well as programmed cell death (Walton et al., 1993; Zhang, 2002). Due to the low level of cellular phosphotyrosine these pathways and the respective enzymes were identified only rather recently: the first tyrosine kinase src by Hunter et al. (1980), the first protein tyrosine phosphatases by Nelson and Branton (1984). Receptor tyrosine kinases include the insulin receptor and a large variety of other growth factor receptors. Examples of PTPs include cdc25A responsible for cell cycle progression, CD45 activating B- and T-cells, the Vaccinia virus minimal variant VH1, YOP 2b required for the virulence of the pathogenic *Yersinia* genus of bacteria, and PTP1B, which is a potential target for diabetes and obesity treatment.

PTP1B can dephosphorylate a number of tyrosine kinases *in vitro* and in cell culture at the endoplasmic reticulum (ER) (Haj et al., 2002). *In vivo* specificity of this phosphatase for insulin receptor dephosphorylation was suggested by the finding that the active site mutant C215S copurifies with insulin receptors from eucaryotic cell lines (Seely et al., 1996; Bandy-

opadhyay et al., 1997). Association with the activated insulin receptor is mediated by three tyrosine residues in the catalytic domain that become phosphorylated upon insulin treatment (Bandyopadhyay et al., 1997). The importance of PTP1B in the regulation of glucose metabolism was underlined by the finding that homozygous PTP1B knockout in mice leads to strongly increased blood glucose clearance and hypersensitivity to insulin (Elchebly et al., 1999). Additionally, homo- and heterozygous knockouts proved to decrease weight gain by 50% on high fat diet. For these reasons, PTP1B appears as an attractive target for specific inhibitors that would counteract type II diabetes and obesity.

Full length human PTP1B consists of 435 residues. It encompasses the highly conserved 298 amino acid, N-terminal catalytic domain (PTP1B^{cat}), a proline rich domain and a C-terminal hydrophobic anchor that targets the protein to the membrane of the ER (Frangioni et al., 1992; Haj et al., 2002). The crystal structure of the PTP1B^{cat} construct 1-321 (Barford et al., 1994) shows that this domain consists of 12 β -strands and 8 α -helices. The catalytic residue C215 is located near the surface in a crevice between β 12 and α 4. Here we report the NMR resonance assignments of the conserved 298 amino acid PTP1B^{cat} domain in complex with the peptidic Novartis inhibitor CGS35385.

Methods and experiments

Human PTP-1B (1-298) was cloned from a human hippocampal cDNA library (Clontech) using PCR and placed into a pET 19-b vector (Novagen). Re-

*To whom correspondence should be addressed. E-mail: Stephan.Grzesiek@unibas.ch

