

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31

A Novel Capture Compound for the Identification and Analysis of Cyclic di-GMP Binding Proteins

Jutta Nesper¹⁾, Alberto Reinders¹⁾, Timo Glatter^{*1)}, Alexander Schmidt*, and Urs Jenal

Affiliations:

Biozentrum of the University of Basel, Klingelbergstrasse 50, CH-4054 Basel,
Switzerland

*Proteomics Core Facility, Biozentrum of the University of Basel, Klingelbergstrasse
50, CH-4054 Basel, Switzerland

1) These authors contributed equally to this work

For correspondence: urs.jenal@unibas.ch

Keywords: *Caulobacter crescentus*; *Pseudomonas aeruginosa*; *Salmonella enterica* serovar *typhimurium*; c-di-GMP effector; EAL GGDEF; Capture Compound mass spectrometry

1 **Abstract**

2 The second messenger cyclic di-GMP is a near-ubiquitous signaling molecule that
3 globally alters bacterial cell physiology to promote biofilm formation and community
4 behavior. Much progress was made in recent years towards the identification and
5 characterization of diguanylate cyclases and phosphodiesterases, enzymes involved
6 in the synthesis and degradation of this signaling compound. In contrast, our
7 knowledge of the nature and mechanistic details of c-di-GMP effector proteins lags
8 behind, primarily because effective tools for their specific enrichment and rapid
9 analysis are missing. In this report we demonstrate that a novel tri-functional c-di-
10 GMP-specific Capture Compound (cdG-CC) can be effectively used to identify and
11 validate c-di-GMP binding proteins. The cdG-CC was able to specifically and
12 efficiently pull-down *bona fide* c-di-GMP effector proteins. Furthermore, in
13 combination with mass spectrometry (CCMS), this technology robustly identified a
14 substantial fraction of the known c-di-GMP signaling components directly from cell
15 extracts of different model organisms. Finally, we applied the CCMS technique to
16 profile c-di-GMP binding proteins of *Salmonella enterica serovar typhimurium*. Our
17 studies establish CCMS as a powerful and versatile tool to identify and analyze
18 components of the cellular c-di-GMP pathway in a wide range of different organisms.
19

1 Cyclic di-GMP is a ubiquitous second messenger regulating growth and behavior of a
2 wide range of gram-positive and gram-negative bacteria. In particular, c-di-GMP
3 mediates the switch between single cell behavior and a community life style called
4 biofilm, which is often associated with chronic infections of bacterial pathogens [1].
5 Major components of the regulatory network are the GGDEF and EAL domains that
6 are widespread in bacteria and catalyze c-di-GMP synthesis and degradation,
7 respectively [2]. The list of cellular processes controlled by c-di-GMP is ever-
8 increasing and includes the biosynthesis and secretion of surface adhesins and
9 exopolysaccharide (EPS) matrix components, different forms of cellular motility, long-
10 term survival and response to environmental stress, synthesis of secondary
11 metabolites, regulated proteolysis and cell cycle progression, delivery of anti-
12 bacterial toxins, intracellular growth and the production of virulence factors in a
13 range of animal and plant pathogens [2-4]. Despite of this wide variety of cellular
14 functions that are modulated by c-di-GMP, the list of effector proteins has remained
15 relatively sparse [2-4]. These include PilZ, small switch-like domains that undergo
16 conformational change upon binding c-di-GMP. In addition, several members of the
17 CRP/FNR and response regulator superfamilies of transcription regulators were
18 shown to specifically bind c-di-GMP. Finally, a subgroup of GGDEF and EAL domains
19 was recognized as c-di-GMP effector proteins adopting their novel functionality
20 through the combined loss of catalytic activity and exploitation of their allosteric and
21 active site binding pockets, respectively. While most known effector proteins were
22 discovered through an “educated guess” approach that was based on their
23 functional linkage to c-di-GMP mediated cellular processes, unbiased screening for
24 novel effectors was hampered primarily by the lack of reliable and effective
25 biochemical tools for their enrichment and isolation. Only one global recent study
26 used a chemical proteomics approach to identify c-di-GMP binding proteins in *P.*
27 *aeruginosa* [5]. Here we introduce a novel tri-functional capture molecule (cdG-CC)
28 as an effective tool to identify specific c-di-GMP binding proteins directly from a
29 complex mixture of macromolecules. The compound is based on a chemical scaffold
30 harboring specificity, reactivity, and sorting properties (Fig. 1A) [6, 7].
31 To evaluate the potential of this compound for the specific enrichment and isolation
32 of c-di-GMP binding proteins we first tested if the cdG-CC is able to pull down known

1 c-di-GMP effector molecules directly from a solution with purified protein. We used
2 the well-studied PilZ domain protein DgrA from *C. crescentus* [8]. The binding affinity
3 of wild-type DgrA for c-di-GMP is in the low nanomolar range. In contrast, the
4 binding mutant DgrAW75A has a K_d of 6.2 μM and a DgrA mutant lacking
5 coordinating Arg residues in the N-terminus (RR11AA) fails to bind c-di-GMP
6 completely [8]. 0.5 μM purified wild-type and mutant hexahistidine-tagged proteins
7 (Fig. 1B) were incubated with 10 μM cdG-CC, UV cross-linked for 4 min and pulled
8 out with streptavidin coated magnetic beads. While DgrA was readily captured in the
9 absence of any other ligand, the addition of a 100x excess of c-di-GMP abolished
10 cdG-CC almost entirely (Fig. 1B). In contrast, an equal molar excess of GTP did not
11 interfere with the cdG-CC pull down (Fig. 1B) arguing that cdG-CC binding to DgrA is
12 highly specific. In agreement with this, the cdG-CC only inefficiently pulled-down the
13 DgrAW75A mutant form and completely failed to bind the DgrARR11AA mutant (Fig.
14 1B). These data are fully consistent with the binding affinities of the different DgrA
15 forms for c-di-GMP and strongly argue that cdG-CC is able to specifically capture
16 DgrA. Moreover, these experiments demonstrate that the cdG-CC is a valuable
17 diagnostic tool to verify candidate c-di-GMP binding proteins.

18 These results encouraged us to probe if the capture compound can be applied for
19 the selective enrichment of c-di-GMP binding proteins from a more complex mixture
20 of proteins. To test this, we captured soluble c-di-GMP binding proteins from cell
21 extracts and probed immunoblots with PopA specific antibodies. PopA is a GGDEF
22 effector protein that regulates cell cycle progression in *C. crescentus* in response to a
23 cellular upshift of c-di-GMP during the G1-S phase transition [9, 10]. To bind c-di-
24 GMP PopA utilizes a conserved and well-defined binding pocket, which, in related
25 catalytic GGDEF domains, is used as an allosteric I-site for product inhibition of the
26 diguanylate cyclase (DGC) activity [11, 12]. As shown in Fig. 1C, PopA with a known
27 K_d for c-di-GMP of 2.5 μM , was readily captured from *C. crescentus* cell lysates using
28 3 μM cdG-CC and a total of 400 μg soluble protein. Pull down of PopA was inhibited
29 in the presence of a large excess of c-di-GMP (1mM), while GTP did not interfere
30 with cdG-CC binding (Fig. 1C), arguing that the cdG-CC interaction with PopA is highly
31 specific. Likewise, no PopA was bound to the cdG-CC when using a strain expressing
32 a PopA mutant that lacks the highly conserved Arg residue of the canonical RxxD I-

1 site binding motif (R357G) [9] (Fig. 1C). This indicated that the cdG-CC can enrich c-
2 di-GMP binding proteins directly from whole cell extracts in a highly specific manner
3 and that this compound is suited for a global isolation procedure of c-di-GMP binding
4 proteins. Therefore we combined capture experiments with the analysis of isolated
5 proteins by LC-MS/MS (CCMS, [6, 7]). When applying 10, 5 or 2.5 μ M cdG-CC with
6 400 μ g of soluble *C. crescentus* proteins, nine of eleven proteins predicted to contain
7 either a PilZ, GGDEF or EAL domain (Table 1A and Supplementary Table S1A) were
8 significantly enriched as compared to the competition control based on the spectral
9 counts of the identified peptides. In addition to the analysis of soluble proteins we
10 also aimed at evaluating the efficiency of CCMS for the enrichment of c-di-GMP
11 binding proteins from membrane fractions. Although the numbers of spectral counts
12 were lower as compared to proteins from the soluble fraction, three of the five
13 integral membrane proteins predicted to bind c-di-GMP were identified when 400
14 μ g DDM solubilized membrane proteins and 10 μ M cdG-CC were used for CCMS
15 (Table 1A and Supplementary Table S1A). Only four of the known components of the
16 *C. crescentus* c-di-GMP network were not identified by CCMS. Two of these are
17 integral membrane proteins with several predicted membrane spanning domains in
18 their N-terminal regions (CC0740, CC0896). It is possible that they were not
19 solubilized by the detergent used or not detected by LC-MS/MS. Another possibility is
20 that they are not expressed, as it might be the case for CC3094 and CC3148.

21 To expand these studies to a different organism, we also tried to isolate known c-di-
22 GMP binding proteins from *P. aeruginosa* PA01. With over 40 GGDEF and EAL
23 domain and eight PilZ domain proteins annotated, the complexity of the c-di-GMP
24 signaling network in this organism is much higher than in *C. crescentus*. Using 10 μ M,
25 7.5 μ M, 5 μ M, 2.5 μ M and 1.25 μ M cdG-CC and 350 μ g of soluble protein extract,
26 several of these proteins were unambiguously identified by CCMS (Table 1B and
27 Supplementary Table S1B). This includes four PilZ domain proteins, four GGDEF and
28 two composite GGDEF-EAL domain proteins. The fraction of the *P. aeruginosa*
29 proteins predicted to bind c-di-GMP that were isolated by CCMS is substantially
30 lower (10 of 28 predicted soluble proteins) as compared to *C. crescentus* but is
31 comparable to a recent study using sepharose-coupled c-di-GMP [5]. The

1 discrepancy between *C. crescentus* and *P. aeruginosa* might be related to the fact
2 that many of these proteins are engaged in the transition between a motile,
3 planktonic and a sessile, surface attached life-style. Since *C. crescentus* has
4 integrated this developmental transition into its reproductive cycle [13], most of the
5 components of the c-di-GMP network orchestrating this switch, need to be
6 expressed in cells growing in liquid cultures [10]. In contrast, *P. aeruginosa* adapts its
7 expression program upon surface colonization in response to environmental cues
8 [14]. It is thus possible that several components of the *P. aeruginosa* core machinery
9 required to adapt to surface communities are simply not expressed in logarithmically
10 growing cultures in liquid media. To test this, it would be interesting to repeat these
11 experiments with cell extracts harvested from surface grown *P. aeruginosa*
12 communities.

13 To evaluate whether cdG-CC based CCMS can be used to profile an entire cellular
14 network of c-di-GMP binding proteins we used *S. typhimurium*. This organism was
15 chosen to perform an unbiased CCMS experiment because its c-di-GMP network
16 appears to be of lower complexity as compared to other bacterial species [15]. In
17 contrast to the CCMS experiments with *C. crescentus* and *P. aeruginosa*, in which
18 spectral counts were extracted for known c-di-GMP binders, we tested whether
19 CCMS is capable to enrich for c-di-GMP binding proteins upon MS1 label-free
20 quantification. CCMS experiments were performed in triplicates with 350 µg soluble
21 whole cell proteins using 7.5 µM cdG-CC and competition controls with an excess of
22 c-di-GMP (1 mM) (Supplementary Table S2A). Following mass spectrometry analysis
23 and label-free quantification (Supplementary Table S2B) significant differences in
24 protein enrichment between cdG-CC experiments and control samples with
25 competing c-di-GMP were visualized in a volcano plot. The graph shows a significant
26 enrichment (>2 fold) of 36 proteins as compared to the control with a q-value <0.05
27 (Supplementary Table S2C and Fig. 2). Among the enriched proteins is the PilZ
28 domain protein YcgR (Fig. 2) [16]. Many of the identified components that were not
29 previously associated with the c-di-GMP network were metabolic proteins and
30 proteins involved in fatty acid and LPS biosynthesis (Fig. 2). Such proteins are of
31 great interest in the light of switching between a virulent planktonic and a surface
32 attached persistent lifestyle. However, since none of these proteins show homology

1 to known c-di-GMP binding proteins, they first need to be validated to specifically
2 bind c-di-GMP and be part of the c-di-GMP signaling network. In addition also some
3 background proteins are enriched. They are mostly binders of other nucleotides or
4 proteins that are known to be highly abundant, such as five tRNA-related proteins or
5 the chaperone GroEL (Supplementary Table S2C).
6

1 **Acknowledgements**

2 We thank Samuel Steiner for help in optimization of Capture Compound binding
3 assays. This work was supported by the Swiss National Science Foundation (SNF)
4 Sinergia grant CRSII3_127433.

5

6

1 **References:**

- 2 [1] Furukawa S, Kuchma SL, O'Toole GA. Keeping their options open: Acute versus
3 persistent infections. *J Bacteriol.* 2006;188:1211-7.
- 4 [2] Schirmer T, Jenal U. Structural and mechanistic determinants of c-di-GMP
5 signalling. *Nat Rev Microbiol.* 2009;7:724-35.
- 6 [3] Hengge R. Cyclic-di-GMP reaches out into the bacterial RNA world. *Sci Signal.*
7 2010;3:pe44.
- 8 [4] Sondermann H, Shikuma NJ, Yildiz FH. You've come a long way: C-di-GMP
9 signaling. *Curr Opin Microbiol.* 2012.
- 10 [5] Duvel J, Bertinetti D, Moller S, Schwede F, Morr M, Wissing J, et al. A chemical
11 proteomics approach to identify c-di-GMP binding proteins in *Pseudomonas*
12 *aeruginosa*. *J Microbiol Methods.* 2012;88:229-36.
- 13 [6] Koster H, Little DP, Luan P, Muller R, Siddiqi SM, Marappan S, et al. Capture
14 compound mass spectrometry: A technology for the investigation of small molecule
15 protein interactions. *Assay Drug Dev Technol.* 2007;5:381-90.
- 16 [7] Lenz T, Fischer JJ, Dreger M. Probing small molecule-protein interactions: A new
17 perspective for functional proteomics. *J Proteomics.* 2011.
- 18 [8] Christen M, Christen B, Allan MG, Folcher M, Jenö P, Grzesiek S, et al. DgrA is a
19 member of a new family of cyclic diguanosine monophosphate receptors and
20 controls flagellar motor function in *Caulobacter crescentus*. *Proc Natl Acad Sci U S A.*
21 2007;104:4112-7.
- 22 [9] Duerig A, Abel S, Folcher M, Nicollier M, Schwede T, Amiot N, et al. Second
23 messenger-mediated spatiotemporal control of protein degradation regulates
24 bacterial cell cycle progression. *Genes Dev.* 2009;23:93-104.
- 25 [10] Abel S, Chien P, Wassmann P, Schirmer T, Kaefer V, Laub MT, et al. Regulatory
26 cohesion of cell cycle and cell differentiation through interlinked phosphorylation
27 and second messenger networks. *Mol Cell.* 2011;43:550-60.
- 28 [11] Christen B, Christen M, Paul R, Schmid F, Folcher M, Jenö P, et al. Allosteric
29 control of cyclic di-GMP signaling. *J Biol Chem.* 2006;281:32015-24.
- 30 [12] Wassmann P, Chan C, Paul R, Beck A, Heerklotz H, Jenal U, et al. Structure of
31 BeF₃⁻-modified response regulator PleD: Implications for diguanylate cyclase
32 activation, catalysis, and feedback inhibition. *Structure.* 2007;15:915-27.

- 1 [13] Curtis PD, Brun YV. Getting in the Loop: Regulation of development in
2 *Caulobacter crescentus*. Microbiology and Molecular Biology Reviews. 2010;74:13-41.
- 3 [14] Kuchma SL, O'Toole GA. Surface-induced and biofilm-induced changes in gene
4 expression. Curr Opin Biotechnol. 2000;11:429-33.
- 5 [15] Solano C, Garcia B, Latasa C, Toledo-Arana A, Zorraquino V, Valle J, et al. Genetic
6 reductionist approach for dissecting individual roles of GGDEF proteins within the c-
7 di-GMP signaling network in *Salmonella*. Proc Natl Acad Sci U S A. 2009;106:7997-
8 8002.
- 9 [16] Ryjenkov DA, Simm R, Romling U, Gomelsky M. The PilZ domain is a receptor for
10 the second messenger c-di-GMP: The PilZ domain protein YcgR controls motility in
11 enterobacteria. J Biol Chem. 2006;281:30310-4.
- 12

1 **Figure legends:**

2 **Fig. 1** - The cdG-CC can specifically pull down bona fide c-di-GMP binding proteins
3 from purified and from crude cell extracts. A) Chemical structure of cdG-CC. C-di-
4 GMP provides the selectivity for capturing proteins. The photo-reactivity group
5 allows irreversible nitren formation between the compound and the captured
6 protein. Biotin as sorting function allows the isolation of the compound by binding to
7 streptavidin coated magnetic beads. B) Immunoblot of purified and captured DgrA
8 (wt), DgrA W75A and DgrA RR11AA. In all capture experiments 10 μ M cdG-CC was
9 present, in competition experiments the proteins were preincubated with a 100 x
10 excess of c-di-GMP (cdG) or GTP. All proteins were His-tagged and detected using
11 anti-His antibodies. C) Immunoblot of PopA. PopA (wt) or the PopA I-site mutant
12 R357G was expressed in NA1000 Δ popA (lanes marked with 'proteins'; note that this
13 I-site mutant is less abundant in the cell). PopA was captured in the presence of 3
14 μ M cdG-CC. Addition of 1 mM c-di-GMP but not 1 mM GTP prevented the binding to
15 the cdG-CC. In contrast the PopA I-site mutant could not be captured at all. PopA
16 was detected using anti-PopA antibodies.

17

18 **Fig. 2** - Volcanoplot based visualization of proteins significantly enriched by CCMS of
19 *S. typhimurium*. Following capturing, LC-MS/MS analysis and label-free quantification,
20 log₂-intensity ratio of all detected peptide features between capturing and
21 competition experiment were calculated and plotted versus values derived from
22 significance analysis. Proteins within the significance thresholds for q-values <0.05
23 and intensity ratios >2-fold are indicated in a box. Experiments in triplicate were
24 performed in the presence of 7.5 μ M cdG-CC and with 1 mM c-di-GMP added to the
25 competition reactions.

26

1 Table 1 - Identified known c-di-GMP binding proteins.

2 A) *C. crescentus*

| Protein Name | ID | Domain architecture | CCMS experiment/CCMS competition ¹⁾ | | |
|--------------------------|------------|------------------------|--|-------|------|
| | | | No spectral counts of identified peptides | | |
| Soluble fraction: | | | | | |
| Experiment ²⁾ | | | 1 | 2 | 3 |
| PopA | CCNA_01918 | GGDEF | 20/1 | 25/3 | 15/5 |
| PleD | CCNA_02546 | Rec-Rec-GGDEF | 18/0 | 21/2 | 13/5 |
| DgcA | CCNA_03394 | GGDEF | 5/0 | 3/0 | 1/0 |
| DgcB | CCNA_01926 | GGDEF | 12/1 | 16/3 | 10/2 |
| CC0655 | CCNA_00692 | PAC-GGDEF-EAL | 8/0 | 18/0 | 19/0 |
| CC1086 | CCNA_01140 | PAS-EAL | 8/0 | 5/0 | 4/0 |
| PdeA | CCNA_03507 | GGDEF-EAL | 3/0 | 5/0 | 6/0 |
| DgrA | CCNA_01671 | PilZ | 5/0 | 6/3 | 6/1 |
| DgrB | CCNA_03268 | PilZ | | 2/0 | |
| Membrane fraction: | | | | | |
| Experiment ³⁾ | | | | | |
| TipF | CCNA_00747 | EAL | 3/0 | 23/13 | |
| PdeB | CCNA_00089 | 3x(MHYT)-PAS-GGDEF-EAL | 3/0 | 6/0 | |
| CC0857 | CCNA_00900 | CHASE4-GGDEF-EAL | | 2/0 | |

3

4 1) All competition experiments were performed in the presence of 1 mM c-di-GMP.

5 2) 3 independent experiments are indicated using 10, 5 or 2.5 μ M cdG-CC
6 respectively.

7 3) 2 independent experiments are indicated using 10 μ M cdG-CC.

8

1 B) Soluble proteins of *P. aeruginosa*

| Protein Name | ID | Domain architecture | CCMS experiment/CCMS competition ¹⁾ | | | | |
|-----------------------------|--------|---------------------|--|------|-----|------|-----|
| | | | No of spectral counts of identified peptides | | | | |
| Experiment No ⁴⁾ | | | 1 | 2 | 3 | 4 | 5 |
| FimX | PA4959 | PAS-GDSIF-EVL | 9/0 | 10/0 | 9/0 | 8/0 | 7/0 |
| | PA3353 | PilZ | 5/0 | 6/0 | 6/0 | 10/0 | 7/0 |
| WspR | PA3702 | Rec-GGEEF | 2/0 | 2/0 | 2/0 | | |
| | PA0012 | PilZ | | | 2/0 | 1/0 | |
| | PA0169 | GGEEF | 2/0 | 2/0 | 2/0 | | |
| | PA4843 | Rec-Rec-GGEEF | | 2/0 | 2/0 | 5/0 | 3/0 |
| | PA2989 | PilZ | 1/0 | 1/0 | 1/0 | 1/0 | 1/0 |
| | PA0290 | PAS-Rec-GGEEF | 1/0 | | | 2/0 | 2/0 |
| | PA2567 | GAF-SPTRF-EAL | | 1/0 | 1/0 | | 1/0 |
| | PA4608 | PilZ | 1/0 | 2/0 | 2/0 | 2/0 | 1/0 |


2

3 4) Experiment 1 was performed with 10 μ M cdG-CC, experiment 2 with 7.5 μ M cdG-CC,
 4 experiment 3 with 2.5 μ M cdG-CC, experiment 4 with 2.5 μ M cdG-CC and experiment 5
 5 with 1.25 μ M cdG-CC.


6

Figure 1:

A



B



C





Fig. 2

[Click here to download high resolution image](#)

Figure 2:



Materials & Methods

[Click here to download Supplementary material: SupplementaryMaMe.docx](#)

Table S1

[Click here to download Supplementary material: SupplementaryTable S1.xlsx](#)

Table S2

[Click here to download Supplementary material: SupplementaryTable S2.xlsx](#)