



FGF signalling through Fgfr2 isoform IIIb regulates adrenal cortex development

Leonardo Guasti^a, W.C. Candy Sze^a, Tristan McKay^a, Richard Grose^b, Peter J. King^{a,*}

^a Centre for Endocrinology, William Harvey Research Institute, Queen Mary University of London, London, UK

^b Tumour Biology, Barts Cancer Institute, Queen Mary University of London, London, UK

ARTICLE INFO

Article history:

Available online 29 January 2013

Keywords:

Adrenal
Development
FGF
FGFR2
Shh

ABSTRACT

Developmental signalling pathways are implicated in the formation and maintenance of the adrenal gland, but their roles are currently not well defined. In recent years it has emerged that Sonic hedgehog (Shh) and Wnt/ β catenin signalling are crucial for the growth and development of the adrenal cortex. Here we demonstrate that Fibroblast growth factor receptor (Fgfr) 2 isoforms IIIb and IIIc are expressed mainly in the adrenal subcapsule during embryogenesis and that specific deletion of the Fgfr2 IIIb isoform impairs adrenal development, causing reduced adrenal growth and impaired expression of SF1 and steroidogenic enzymes. The hypoplastic adrenals also have thicker, disorganised capsules which retain *Gli1* expression but no longer express *Dlk1*. Fgfr2 ligands were detected in both the capsule and the cortex, suggesting the importance of signalling between the capsule and the cortex in adrenal development.


© 2013 Elsevier Ireland Ltd. Open access under [CC BY license](http://creativecommons.org/licenses/by/3.0/).

1. Introduction

The adrenal cortex is the primary site of steroid synthesis, producing glucocorticoids under the control of the hypothalamic–pituitary–adrenal (HPA) axis and mineralocorticoids under the control of the renin–angiotensin system (RAS). Adrenocortical development originates in a group of cells lying between the urogenital ridge and the dorsal aorta, forming the precursors of the adrenal glands and the gonads, the adrenogonadal primordium (AGP) (Hatano et al., 1996). At around embryonic day e9.0 in the mouse these cells begin to express the steroidogenic transcription factor Steroidogenic Factor-1 (SF1), which is essential for both adrenal and gonadal development (Luo et al., 1994). By e12.0, cells of the dorsal medial AGP form the adrenocortical primordium, which is invaded by migrating

adrenal from e12.5 throughout life (King et al., 2009; Ching and Vilain, 2009; Huang et al., 2010). The Shh signal is transduced by capsule cells, some of which delaminate into the cortex, lose their responsiveness to Shh, and become steroidogenic (King et al., 2009; Laufer et al., 2012). Deletion of *Shh* from steroidogenic tissues causes adrenal hypoplasia, and interestingly the adrenal capsule is markedly thinner. Similar targeted deletion of β *catenin*, which like Shh is expressed just underneath the capsule in the murine adrenal cortex from e12.5, also causes adrenal hypoplasia, demonstrating that canonical Wnt signalling is critical for adrenal development.

Like Shh and Wnts, the pathways activated by the binding of Fibroblast Growth Factors (FGFs) to their receptors are of crucial importance for many developmental processes as well as in disease (Gazit and Itsk, 2001; Turner and Gross, 2010). Four signalling

brought to you by  CORE

provided by St George's Online Research Archive

are transmembrane immunoglobulin (Ig) domains. FGFRs 1–3 are alternatively spliced in the third Ig domain such that an invariant exon IIIa is spliced to either exon IIIb or IIIc to produce IIIb or IIIc isoforms. Embryos with a global *Fgfr2 IIIb* deletion have hypoplastic adrenals (Revest et al., 2001), and deletion of both isoforms of FGFR2 from steroidogenic tissue recapitulates this phenotype (Kim et al., 2007), implying that the receptor is both expressed in the adrenal cortex and necessary for development. However, neither the expression pattern of FGFR2 isoforms, nor the adrenal phenotypes of the knockouts, has been investigated further.

Here we demonstrate that Fgfr2 IIIb and IIIc isoforms are expressed underneath the embryonic adrenal capsule, in a similar expression pattern to Shh and β catenin. *Fgfr2 IIIb* null mouse

and similar papers at core.ac.uk

time in a poorly understood manner. Zonation occurs around birth (for reviews see (Kim et al., 2009; Laufer et al., 2012). Over the last few years signalling pathways involved in adrenal development have begun to be elucidated. We and others have shown that sonic hedgehog (Shh) is expressed in relatively undifferentiated steroidogenic cells in the subcapsular region of the mouse

Abbreviations: Shh, sonic hedgehog; FGFR, fibroblast growth factor receptor.

* Corresponding author. Address: Centre for Endocrinology, William Harvey Research Institute, Barts and the London, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, UK. Tel.: +44 (0)20 78826237; fax: +44 (0)20 78826197.

E-mail address: p.j.king@qmul.ac.uk (P.J. King).

embryos have hypoplastic adrenals with impaired steroidogenic differentiation, and a larger and less defined capsule. However, the expression of the Shh target gene *Gli1* in capsular cells persisted in *Fgfr2 IIIb* null adrenals. Instead, the capsular expression of *Dlk1* (also known as *Pref-1*), a member of the Notch family of EGF repeat-containing proteins (for review see (Sul, 2009)), was strongly and specifically reduced in the *Fgfr2 IIIb* knock-out. Finally, we performed RT-PCR analysis of FGFs in laser capture dissected mouse embryonic adrenal tissues, and detected *Fgf1* in the adrenal capsule and *Fgf2* and *Fgf9* in the adrenal cortex.

2. Methods

2.1. Mice

Mice were housed in rooms with controlled light and temperature and treated under the Home Office Animals (Scientific Procedures) Act 1986.

We used Cre-Lox transgenics to generate *Fgfr2 IIIb* null embryos, taking advantage of a $K5^{cre}$ construct which, in female mice, is mis-expressed in the oocyte in addition to basal keratinocytes, resulting in global deletion of *Fgfr2 IIIb* (Grose et al., 2007; Ramirez et al., 2004). Briefly, *Fgfr2 IIIb*^{fllox/fllox} male mice were crossed with female *keratin K5^{cre}/+*; *Fgfr2 IIIb*^{fllox/+} mice. Wild type and $K5^{cre/+}$ females were sacrificed by CO₂ asphyxiation at day 15.5 post-fertilisation and embryos were snap frozen in liquid nitrogen and stored at –80 °C. Sagittal sections were obtained on a freezing-microtome (Leica GM 1510 S or Leica 3050 S) at 12 μm thickness, and mounted on Superfrost Plus slides (VWR). Sections were stained with hematoxylin and eosin using standard procedures.

2.2. Non-radioactive in situ hybridization (NR-ISH)

RNA was extracted from adult adrenal tissues using an RNeasy Mini kit (Qiagen), and cDNA was prepared from random-primed total RNA using Maloney Murine Leukemia Virus-Reverse Transcriptase (MMLV-RT, Promega). *Fgfr2*, *Fgfr2 IIIb*, *Fgfr2 IIIc* and *Dlk1* cDNAs were amplified (PCR primer sequences are listed in Supplemental) and cloned into pGEM T-easy vectors (Promega). The vectors were sequenced and linearised with the appropriate restriction enzymes. The mouse *Gli1* probe (King et al., 2009) and *Sf1* probe (Guasti et al., 2011) have been described previously.

Digoxigenin (DIG)-labelled cRNA probes synthesis and NR-ISH was performed as described (Guasti et al., 2011).

2.3. Immunostaining

Immunostaining of sections and image acquisition and processing was performed according to (Guasti et al., 2011). Anti-Cyp11b1 (1:20 dilution), anti-P450 Side Chain Cleavage (Sc, 1:1000 dilution, Millipore AB1244), anti-PCNA (F2, sc25280, Santa Cruz Biotechnology, 1:1000 dilution) and anti-Cleaved caspase 3 (Cell Signaling, D175, 1:500 dilution) were used in this study. For immunohistochemical comparative analysis (as well as for *Gli1*, *Sf1* and *Dlk1* NR-ISH), all steps from fixation and sectioning to staining and image acquisition were performed in parallel in wild-type and knock-out mice.

2.4. Laser capture microdissection

Sections of e15.5 mouse embryos were cut at 12 μm thickness and mounted onto polyethylene naphthalate-membrane slides (Life Technologies). Sections were fixed and stained with cresyl violet using the LCM staining kit (Ambion) according to the manufacturer's instructions. Sections were allowed to dry at room

temperature for 5 min and immediately used for laser capture microdissection using a PALM MicroBeam (Zeiss). Adrenal capsule, adrenal cortex and pooled tissues (brain, liver, heart, skin, gut, and lungs) were collected in RLT lysis buffer (RNeasy kit, Qiagen) and RNA was extracted according to the manufacturer's instructions. cDNA was synthesised using a Sensiscript RT kit (Qiagen) according to the manufacturer's instructions.

3. Results

3.1. Expression pattern of FGFR2 isoforms

We investigated the expression of *Fgfr2* isoforms *IIIb* and *IIIc* in e15.5 wild type (WT) mouse adrenals by non-radioactive *in situ* hybridization (NR-ISH) using isoform-specific probes (Fig. 1A–F): both *IIIb* and *IIIc* isoforms were expressed underneath the capsule in cells that do not express *Cyp11b1*. This is the same expression pattern we observed for Shh in developing and adult mouse (King et al., 2009) and rat (Guasti et al., 2011) adrenals. A probe recognising both isoforms gave similar results; however, upon longer exposure, some capsular *Fgfr2* expression could be detected with this probe (Fig. S1).

3.2. *Fgfr2 IIIb* null embryos have hypoplastic adrenals

As *Fgfr2 IIIb* null embryos die at birth, we analysed e15.5 embryos. The adrenals in the null embryos are clearly smaller and have a markedly thicker capsule containing rounded rather than the typically elongated cells (Fig. 1G–L), and a less well-defined boundary with the cortex (arrows in Fig. 1L). The expression of *Fgfr2 IIIc* did not seem to be affected by *IIIb* deletion (Fig. S1). To investigate the mechanisms for the hypoplasticity of the gland we studied markers of apoptosis and proliferation. While apoptotic cells (detected with an anti-cleaved caspase 3 antibody) were rare and mainly localised in the medulla in both WT and knock-out (KO) adrenals (Fig. S2), analysis of cell proliferation with an anti-PCNA antibody showed that, whereas the number of PCNA-positive cortical cells immediately adjacent to the capsule did not differ between WT and KO mice, PCNA positive nuclei were clearly present in the capsule of the KO, but absent in the WT, and there were fewer proliferative cells in the inner cortex of the KO adrenal compared to WT adrenals.

3.3. *Fgfr2 IIIb* null adrenals have reduced steroidogenic enzyme expression

Next, the expression of the pan-steroidogenic marker *Cyp11a1* (Sc) and the ZF marker *Cyp11b1* was compared between WT and knockout (KO) adrenals using specific antibodies (Fig. 2). The expression of both enzymes was noticeably reduced in KO embryos, with fewer cells expressing the enzymes and at a lower level. NR-ISH analysis also showed that *Sf1* mRNA expression was reduced in KO adrenals (Fig. 2I and J).

3.4. Analysis of gene expression in the capsule of *Fgfr2 IIIb* null adrenals

As signalling through *Fgfr2 IIIb* is required to establish Shh expression in the developing limb bud (Revest et al., 2001), it was therefore possible that the defects in adrenal development described above are a consequence of perturbed Shh signalling. Shh activity at the capsule was assessed by *Gli1* NR-ISH (Fig. 3). *Gli1* expression was readily detected in the *Fgfr2 IIIb* KO adrenal capsule, and was apparently expressed at higher levels than in the wild type mouse, with more clusters of *Gli1* positive cells observed. To investigate the capsule phenotype further, we also studied the




Fig. 1. (A–F) NR-ISH analysis of e15.5 mouse adrenal glands using *Fgfr2 IIIb* (B and C) and *IIIc* (E and F) specific probes combined with Cyp11b1 immunolocalization. *Fgfr2* positive cells are labelled red, Cyp11b1 cells are labelled green and cell nuclei are labelled blue (Cap. = capsule, Med. = medulla). (G–L) Hematoxylin and eosin staining of wild-type (+/+, G–I) and *fgfr2 IIIb* knock out (-/-, J–L) adrenals. (M, N) Immunohistochemical analysis of PCNA expression in wild-type (+/+, M) and *fgfr2 IIIb* knock-out (-/-, N) adrenals. Arrows in N point to PCNA-positive cells in the adrenal capsule. Scale bars = F, 50 μ m (applies to A–F); J, 150 μ m (applies to G and J); L, 50 μ m (applies to H, I, K, and L); M and N = 100 μ m. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

expression of *Dlk1*, another marker of the mouse adrenal capsule (Beuschlein et al., 2002), which is also expressed in the medulla. Interestingly, *Dlk1* expression was completely lost from the

adrenal capsule but retained in the medulla, which is comparable in size to that of the wild type adrenal, indicating that medulla formation is unaffected.




Fig. 2. Expression of *Scc* (A–D), *Cyp11b1* (E–H) and *Sf1* in e15.5 WT (A, B, E, F, and I) and *Fgfr2 IIIb* KO (C, D, G, H, and J) adrenals. Scale bar = 200 μ m (A, C, E, and G); 50 μ m (B, D, F, H, I, and J). Med. = medulla, Cx. = cortex.

3.5. *Fgf* expression in mouse embryo adrenals

Finally, we performed laser capture microdissection on e15.5 WT mouse adrenal tissues, collecting tissues from both the capsule

and the cortex. RT-PCR performed on RNA isolated from these tissues, using primers against *FGF* isoforms, detected expression of *Fgf1*, in the cortex, and *Fgf2* and 9, in the capsule (Fig. 4). The detection of *Shh* expression in the cortical fraction and the preferential




Fig. 3. NR-ISH of WT (+/+) and *Fgfr2 IIIb* KO (-/-) e15.5 mouse adrenals using *Gli1* (A and B) and *Dlk1* (C–F) riboprobes. Arrows in A and B indicate clusters of *Gli1* positive cells in the capsule. Capsule cells in A, B, E and F are between the brackets. Scale bars = 100 μ m. Med. = medulla.




Fig. 4. RT-PCR analysis of *Fgfs* and *Fgfrs* expression in laser capture microdissected e15.5 adrenal capsule and cortex. Control cDNA was made from pooled tissues including lung, heart, liver, bone and gut.

expression of its receptor *Ptch1* in the capsule (King et al., 2009) indicated the accuracy of the microdissection. Both *Fgfr2 IIIb*, albeit at lower levels than in the cortex, and *IIIc*, were amplified in the capsule, in keeping with the NR-ISH experiments (see Fig. 1 and Fig. S1).

4. Discussion

Germline knockout of the *IIIb* exon of the *Fgfr2* gene results in mice that die at birth from multiple developmental defects, identifying *Fgfr2 IIIb* as a critical mediator of organogenesis (De

Moerlooze et al., 2000; Revest et al., 2001). We have shown here that adrenal development is also affected, with a detectable but smaller gland at e15.5 in which expression of the steroidogenic markers *Sf1*, *Sccl* and *Cyp11b1* is reduced, compared to a wild type adrenal. Analysis of apoptosis and proliferation markers demonstrates that the hypoplasia is likely a consequence of a reduction of inner cortical cell proliferation. In *Shh* null adrenals, apoptosis and proliferation indices are largely normal in the cortex and the defect in growth is believed to be a consequence of the very thin capsule, from which mesenchymal cells differentiate to populate the steroidogenic cortex in the wild type organ. β *catenin* KO, however, does result in reduced cortical proliferation.

Fgf and *Shh* pathways interact in several developmental settings including the limb bud (Laufer et al., 1994), the cerebellum (Fogarty et al., 2007) and the lung (Chuang et al., 2003), and *Shh* has been shown to be under both positive and negative regulation by FGFs. In the developing limb, for example, *Fgf10* signalling to the apical ectodermal ridge through *Fgfr2 IIIb* is required to initiate *Shh* expression in the zone of polarising activity in the posterior limb bud and thus establish the anterior–posterior limb axis (Revest et al., 2001), and expression of *Shh* is restricted to the posterior limb bud by inhibitory *Fgf* signalling in the anterior limb to control limb patterning (Mao et al., 2009; Zhang et al., 2009). The maintained expression of the *Shh* target gene *Gli1* that we observe in the adrenal capsule demonstrates that FGF signalling through *Fgfr2 IIIb* is not required for *Shh* signalling in this setting, and may act to restrict it, given the apparent increased *Gli1* expression in the *Fgfr2 IIIb* KO adrenal capsule. *Fgf* and *Wnt*/ β *catenin*

pathways also interact, with Fgf10 activating β catenin signalling in developing skin (Reddy et al., 2001), mammary tissue (Veltmaat et al., 2006) and liver (Berg et al., 2007; Mavila et al., 2012), and Fgfs 1, 4 and 8 activating β catenin signalling in murine embryonic liver (Sekhon et al., 2004). We have not examined β catenin expression in the null adrenal but note that the phenotypes of the Fgfr2 IIIb and β catenin KO adrenals are similar, with impaired development, reduced expression of steroidogenic markers and reduced cortical cell proliferation evident (Kim et al., 2008). However, it does not appear that β catenin KO causes a capsule phenotype as we see here, so we do not expect that this aspect of the phenotype is a consequence of impaired Wnt signalling.

The two receptor isoforms are apparently predominantly expressed in the same cells in the cortex, although expression was detected in the adrenal capsule at lower levels. As the adrenal gland is present, signalling through Fgfr2 IIIb is not necessary for the formation of the adrenal primordium but, like Shh and β catenin (Kim et al., 2008), this receptor is required for the subsequent growth of the gland. The Fgfr2 IIIb KO examined here is a global knockout and it is therefore possible that the observed effects on adrenal development may be secondary to those on other organ systems, although no effect on the gonads was observed, indicating that the adrenal hypoplasia was not secondary to effects on the adrenogonadal primordium (Revest et al., 2001). Conditional deletion of both isoforms of Fgfr2 from steroidogenic tissue also leads to adrenal hypoplasia (Kim et al., 2007) and global hemizygous deletion of exon IIIc of Fgfr2 does not appear to have any gross effect on adrenal development, unlike on the development of other visceral organs (Hajihosseini et al., 2001). Taken together, these observations suggest that the effect of loss of Fgfr2 IIIb on the adrenal is direct, although formal proof would require the generation of adrenal specific knockouts.

Fgfr2 IIIb is activated by five known ligands, Fgfs 1, 3, 7, 10 and 22 (Zhang et al., 2006), but of these we could only detect Fgf1, which is expressed in the cortex. Fgfr2 IIIc is also expressed in the adrenal cortex, and Fgfs 2 and 9 (Lavine et al., 2005) activate this isoform, both being detected in the capsule. Fgf2 acts as a mitogen for adrenocortical cells in culture and in regeneration experiments (Crickard et al., 1981; Lepique et al., 2004; Chu et al., 2009). However, the normal adrenal phenotype observed on hemizygous Fgfr2 IIIc deletion suggests that signalling through this receptor may not be necessary during development (Hajihosseini et al., 2001). Given the detection, albeit at low levels, of receptor expression in the adrenal capsule, it is not possible to determine whether Fgf1 is acting on the IIIb isoform receptor in the cortex or in the capsule, or both. It will be interesting to determine the relative importance of signalling from the cortex to the capsule (Fgf1/Fgfr2 IIIb), from the capsule to the cortex (Fgfs 2 and 9/Fgfr2 IIIc) or within the capsule (Fgfs 2 and 9/Fgfr2 IIIc) or cortex (Fgf1/Fgfr2 IIIb), for adrenal growth and development.

The thicker and disorganised adrenal capsule phenotype we observe is similar to that seen in *melanocortin 2 receptor* (*Mc2r*, *ACTH receptor*) KO adrenals (Chida et al., 2007), perhaps indicating an interaction between the FGF and ACTH signalling pathways, and this warrants further investigation. However, as both *Mc2r* and *Pomc* (Karpac et al., 2005) KO adrenals are otherwise normal at birth, we do not feel that impaired ACTH signalling is a likely mechanism for the effects on the growth and differentiation of the gland we see here. The observation that *Dlk1* expression is specifically lost from the adrenal capsule in the null adrenals is fascinating and suggests that this may be the cause of the capsule phenotype. Enucleation experiments in rats, in which the inner mass of the adrenal is removed and the cortex regenerates from the capsule and adherent subcapsular cells, have demonstrated that *Dlk1* expression is immediately downregulated and not re-established until the zonation of the cortex has been restored

(Halder et al., 1998). This led to the suggestion that *Dlk1* is a negative regulator of adrenal differentiation, analogous to its role in adipogenesis (Smas and Sul, 1993). Unusually, however, the expression pattern of *Dlk1* differs between rats and mice, with it being specifically expressed in subcapsular cells in the rat (Halder et al., 1998), rather than in the capsule (Beuschlein et al., 2002), perhaps making comparisons of the role of the protein in the two species invalid. Nevertheless, it is possible that *Dlk1* negatively regulates differentiation of the Gli1-positive capsular stem cells in a similar manner and its downregulation is in response to the impaired recruitment of these cells into the cortex caused by the Fgfr2 IIIb KO, perhaps as a consequence of the effects on the growth and organisation of the capsule.

These data highlight the importance of FGF signalling in adrenal development. Shh expression is not greatly perturbed by the loss of FGF signalling, so presumably other, as yet unidentified, mechanisms are involved in the control of adrenal development and growth of the capsule by signalling through Fgfr2 IIIb. Determining these mechanisms is an important goal for understanding adrenal development.

Acknowledgements

We are very grateful to Dr. C. Gomez-Sanchez (University of Mississippi Medical Center, Jackson, Mississippi, USA) for the anti-Cyp11b1 antibody.

This work was supported by Grants from the Medical Research Council (PJK), the Wellcome Trust (CS) and an Early Career Grant from the Society for Endocrinology (LG).

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.mce.2013.01.014>.

References

- Berg, T., Rountree, C.B., Lee, L., Estrada, J., Sala, F.G., Choe, A., Veltmaat, J.M., De Langhe, S., Lee, R., Tsukamoto, H., Crooks, G.M., Bellusci, S., Wang, K.S., 2007. Fibroblast growth factor 10 is critical for liver growth during embryogenesis and controls hepatoblast survival via beta-catenin activation. *Hepatology* 46, 1187–1197.
- Beuschlein, F., Mutch, C., Bavers, D.L., Ulrich-Lai, Y.M., Engeland, W.C., Keegan, C., Hammer, G.D., 2002. Steroidogenic factor-1 is essential for compensatory adrenal growth following unilateral adrenalectomy. *Endocrinology* 143, 3122–3135.
- Chida, D., Nakagawa, S., Nagai, S., Sagara, H., Katsumata, H., Imaki, T., Suzuki, H., Mitani, F., Ogishima, T., Shimizu, C., Kotaki, H., Kakuta, S., Sudo, K., Koike, T., Kubo, M., Iwakura, Y., 2007. Melanocortin 2 receptor is required for adrenal gland development, steroidogenesis, and neonatal gluconeogenesis. *Proc. Natl. Acad. Sci. USA* 104, 18205–18210.
- Ching, S., Vilain, E., 2009. Targeted disruption of Sonic Hedgehog in the mouse adrenal leads to adrenocortical hypoplasia. *Genesis* 47, 628–637.
- Chu, Y., Ho, W.J., Dunn, J.C., 2009. Basic fibroblast growth factor delivery enhances adrenal cortical cellular regeneration. *Tissue Eng. Part A* 15, 2093–2101.
- Chuang, P.T., Kawcak, T., McMahon, A.P., 2003. Feedback control of mammalian Hedgehog signaling by the Hedgehog-binding protein, Hip1, modulates Fgf signaling during branching morphogenesis of the lung. *Genes Dev.* 17, 342–347.
- Crickard, K., III, C.R., Jaffe, R.B., 1981. Control of proliferation of human fetal adrenal cells in vitro. *J. Clin. Endocrinol. Metab.* 53, 790–796.
- De Moerloose, L., Spencer-Dene, B., Revest, J.M., Hajihosseini, M., Rosewell, I., Dickson, C., 2000. An important role for the IIIb isoform of fibroblast growth factor receptor 2 (FGFR2) in mesenchymal-epithelial signalling during mouse organogenesis. *Development* 127, 483–492.
- Fogarty, M.P., Emmenegger, B.A., Gräsfelder, L.L., Oliver, T.G., Wechsler-Reya, R.J., 2007. Fibroblast growth factor blocks Sonic hedgehog signaling in neuronal precursors and tumor cells. *Proc. Natl. Acad. Sci. USA* 104, 2973–2978.
- Grose, R., Fantl, V., Werner, S., Chionni, A.M., Jarosz, M., Rudling, R., Cross, B., Hart, I.R., Dickson, C., 2007. The role of fibroblast growth factor receptor 2b in skin homeostasis and cancer development. *EMBO J.* 26, 1268–1278.
- Guasti, L., Paul, A., Laufer, E., King, P., 2011. Localization of Sonic hedgehog secreting and receiving cells in the developing and adult rat adrenal cortex. *Mol. Cell. Endocrinol.* 336, 117–122.

- Hajihosseini, M.K., Wilson, S., De Moerloozee, L., Dickson, C., 2001. A splicing switch and gain-of-function mutation in *FgfR2-IIIc* hemizygotes causes Apert/Pfeiffer-syndrome-like phenotypes. *Proc. Natl. Acad. Sci. USA* 98, 3855–3860.
- Halder, S.K., Takemori, H., Hatano, O., Nonaka, Y., Wada, A., Okamoto, M., 1998. Cloning of a membrane-spanning protein with epidermal growth factor-like repeat motifs from adrenal glomerulosa cells. *Endocrinology* 139, 3316–3328.
- Hatano, O., Takakusu, A., Nomura, M., Morohashi, K., 1996. Identical origin of adrenal cortex and gonad revealed by expression profiles of Ad4BP/SF-1. *Genes Cells* 1, 663–671.
- Huang, C.C., Miyagawa, S., Matsumaru, D., Parker, K.L., Yao, H.H., 2010. Progenitor cell expansion and organ size of mouse adrenal is regulated by sonic hedgehog. *Endocrinology* 151, 1119–1128.
- Karpac, J., Ostwald, D., Bui, S., Hunnewell, P., Shankar, M., Hochgeschwender, U., 2005. Development, maintenance, and function of the adrenal gland in early postnatal proopiomelanocortin-null mutant mice. *Endocrinology* 146, 2555–2562.
- Kim, A.C., Barlaskar, F.M., Heaton, J.H., Else, T., Kelly, V.R., Krill, K.T., Scheys, J.O., Simon, D.P., Trovato, A., Yang, W.H., Hammer, G.D., 2009. In search of adrenocortical stem and progenitor cells. *Endocr. Rev.* 30, 241–263.
- Kim, A.C., Reuter, A.L., Zubair, M., Else, T., Serecky, K., Bingham, N.C., Lavery, G.G., Parker, K.L., Hammer, G.D., 2008. Targeted disruption of beta-catenin in Sf1-expressing cells impairs development and maintenance of the adrenal cortex. *Development* 135, 2593–2602.
- Kim, Y., Bingham, N., Sekido, R., Parker, K.L., Lovell-Badge, R., Capel, B., 2007. Fibroblast growth factor receptor 2 regulates proliferation and Sertoli differentiation during male sex determination. *Proc. Natl. Acad. Sci. USA* 104, 16558–16563.
- King, P., Paul, A., Laufer, E., 2009. Shh signaling regulates adrenocortical development and identifies progenitors of steroidogenic lineages. *Proc Natl Acad Sci USA* 106, 21185–21190.
- Laufer, E., Kesper, D., Vortkamp, A., King, P., 2012. Sonic hedgehog signaling during adrenal development. *Mol. Cell. Endocrinol.* 351, 19–27.
- Laufer, E., Nelson, C.E., Johnson, R.L., Morgan, B.A., Tabin, C., 1994. Sonic hedgehog and Fgf-4 act through a signaling cascade and feedback loop to integrate growth and patterning of the developing limb bud. *Cell* 79, 993–1003.
- Lavine, K.J., Yu, K., White, A.C., Zhang, X., Smith, C., Partanen, J., Ornitz, D.M., 2005. Endocardial and epicardial derived FGF signals regulate myocardial proliferation and differentiation in vivo. *Dev. Cell* 8, 85–95.
- Lepique, A.P., Moraes, M.S., Rocha, K.M., Eichler, C.B., Hajj, G.N., Schwindt, T.T., Armelin, H.A., 2004. C-Myc protein is stabilized by fibroblast growth factor 2 and destabilized by ACTH to control cell cycle in mouse Y1 adrenocortical cells. *J. Mol. Endocrinol.* 33, 623–638.
- Luo, X., Ikeda, Y., Parker, K.L., 1994. A cell-specific nuclear receptor is essential for adrenal and gonadal development and sexual differentiation. *Cell* 77, 481–490.
- Mao, J., McGlinn, E., Huang, P., Tabin, C.J., McMahon, A.P., 2009. Fgf-dependent *Etv4/5* activity is required for posterior restriction of Sonic Hedgehog and promoting outgrowth of the vertebrate limb. *Dev. Cell* 16, 600–606.
- Mavila, N., James, D., Utley, S., Cu, N., Coblens, O., Mak, K., Rountree, C.B., Kahn, M., Wang, K.S., 2012. Fibroblast growth factor receptor-mediated activation of AKT-beta-catenin-cbp pathway regulates survival and proliferation of murine hepatoblasts and hepatic tumor initiating stem cells. *PLoS ONE* 7, e50401.
- Ornitz, D.M., Itoh, N., 2001. Fibroblast growth factors. *Genome Biol.* 2, Reviews 3005.
- Ramirez, A., Page, A., Gandarillas, A., Zanet, J., Pibre, S., Vidal, M., Tusell, L., Genesca, A., Whitaker, D.A., Melton, D.W., Jorcano, J.L., 2004. A keratin K5Cre transgenic line appropriate for tissue-specific or generalized Cre-mediated recombination. *Genesis* 39, 52–57.
- Reddy, S., Andl, T., Bagasra, A., Lu, M.M., Epstein, D.J., Morrisey, E.E., Millar, S.E., 2001. Characterization of Wnt gene expression in developing and postnatal hair follicles and identification of Wnt5a as a target of Sonic hedgehog in hair follicle morphogenesis. *Mech. Dev.* 107, 69–82.
- Revest, J.M., Spencer-Dene, B., Kerr, K., De Moerloozee, L., Rosewell, I., Dickson, C., 2001. Fibroblast growth factor receptor 2-IIIb acts upstream of Shh and Fgf4 and is required for limb bud maintenance but not for the induction of Fgf8, Fgf10, Msx1, or Bmp4. *Dev. Biol.* 231, 47–62.
- Sekhon, S.S., Tan, X., Micsenyi, A., Bowen, W.C., Monga, S.P., 2004. Fibroblast growth factor enriches the embryonic liver cultures for hepatic progenitors. *Am. J. Pathol.* 164, 2229–2240.
- Smas, C.M., Sul, H.S., 1993. Pref-1, a protein containing EGF-like repeats, inhibits adipocyte differentiation. *Cell* 73, 725–734.
- Sul, H.S., 2009. Minireview: Pref-1: role in adipogenesis and mesenchymal cell fate. *Mol. Endocrinol.* 23, 1717–1725.
- Turner, N., Grose, R., 2010. Fibroblast growth factor signalling: from development to cancer. *Nature Rev. Cancer* 10, 116–129.
- Veltmaat, J.M., Relaix, F., Le, L.T., Kratochwil, K., Sala, F.G., Van Veelen, W., Rice, R., Spencer-Dene, B., Mailloux, A.A., Rice, D.P., Thiery, J.P., Bellusci, S., 2006. Gli3-mediated somitic Fgf10 expression gradients are required for the induction and patterning of mammary epithelium along the embryonic axes. *Development* 133, 2325–2335.
- Zhang, X., Ibrahim, O.A., Olsen, S.K., Umemori, H., Mohammadi, M., Ornitz, D.M., 2006. Receptor specificity of the fibroblast growth factor family. The complete mammalian FGF family. *J. Biol. Chem.* 281, 15694–15700.
- Zhang, Z., Verheyden, J.M., Hassell, J.A., Sun, X., 2009. FGF-regulated *Etv* genes are essential for repressing Shh expression in mouse limb buds. *Dev. Cell* 16, 607–613.