

POSTER PRESENTATION

Open Access

ErbB2/HER2-specific NK cells for adoptive cancer immunotherapy

Congcong Zhang¹, Michael Burger², Kurt Schönfeld¹, Sabrina Genßler¹, Christiane Sahn¹, Sonja Naundorf³, Marcus Odendahl⁴, Paulina Nowakowska⁵, Torsten Tonn⁴, Manuel Grez¹, Joachim P Steinbach², Winfried S Wels^{1*}

From Society for Immunotherapy of Cancer 28th Annual Meeting
National Harbor, MD, USA. 8-10 November 2013

Significant progress has been made over the last decade towards realizing the potential of natural killer (NK) cells for cancer immunotherapy. NK cells can respond rapidly to transformed and stressed cells, and have the intrinsic potential to extravasate and reach their targets in almost all body tissues. In addition to donor-derived primary NK cells, also continuously expanding cytotoxic cell lines such as NK-92 are being considered for adoptive cancer immunotherapy. High cytotoxicity of NK-92 has previously been shown against malignant cells of hematologic origin in preclinical studies, and general safety of infusion of NK-92 cells has been established in phase I clinical trials. To enhance their therapeutic utility, we genetically modified NK-92 cells to express chimeric antigen receptors (CAR) specific for tumor-associated surface antigens. Such CAR were composed of a tumor-specific scFv antibody fragment fused via hinge and transmembrane domains to intracellular signaling moieties such as CD3 zeta chain, or composite fusion molecules also containing a costimulatory protein domain in addition to CD3 zeta. For development towards clinical applications, here a codon-optimized second generation CAR was constructed that consists of an ErbB2-specific scFv antibody domain fused via a linker to a composite CD28-CD3 zeta signaling domain. GMP-compliant protocols for vector production, lentiviral transduction and expansion of a genetically modified NK-92 single cell clone (NK-92/5.28.z) were established. Functional analysis of NK-92/5.28.z cells revealed high and stable CAR expression, selective cytotoxicity against ErbB2-expressing but otherwise NK-resistant tumor cells of different origins in vitro, as well as homing to ErbB2-expressing tumors in vivo. Furthermore, antigen

specificity and selective cytotoxicity of these cells were retained in vivo, resulting in antitumoral activity against subcutaneous and intracranial glioblastoma xenografts in NSG mice. Ongoing work now focuses on the development of these cells for adoptive immunotherapy of ErbB2-positive glioblastoma.

Authors' details

¹Georg-Speyer-Haus, Frankfurt, Germany. ²Institute for Neurooncology, University Hospital, University of Frankfurt, Frankfurt, Germany. ³EUFETS GmbH, Idar-Oberstein, Germany. ⁴German Red Cross Blood Service North-East, Dresden, Germany. ⁵German Red Cross Blood Service Baden-Württemberg - Hessen, Frankfurt, Germany.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P38

Cite this article as: Zhang et al.: ErbB2/HER2-specific NK cells for adoptive cancer immunotherapy. *Journal for ImmunoTherapy of Cancer* 2013 **1**(Suppl 1):P38.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Georg-Speyer-Haus, Frankfurt, Germany
Full list of author information is available at the end of the article