

COMPLEX HADAMARD MATRICES AND THE SPECTRAL SET CONJECTURE

MIHAIL N. KOLOUNTZAKIS & MÁTÉ MATOLCSI

ABSTRACT. By analyzing the connection between complex Hadamard matrices and spectral sets we prove the direction “spectral \Rightarrow tile” of the Spectral Set Conjecture for all sets A of size $|A| \leq 5$ in any finite Abelian group. This result is then extended to the infinite grid \mathbb{Z}^d for any dimension d , and finally to \mathbb{R}^d .

It was pointed out recently in [16] that the corresponding statement fails for $|A| = 6$ in the group \mathbb{Z}_3^5 , and this observation quickly led to the failure of the Spectral Set Conjecture in \mathbb{R}^5 [16], and subsequently in \mathbb{R}^4 [13]. In the second part of this note we reduce this dimension further, showing that the direction “spectral \Rightarrow tile” of the Spectral Set Conjecture is false already in dimension 3.

In a computational search for counterexamples in lower dimension (one and two) one needs, at the very least, to be able to decide efficiently if a set is a tile (in, say, a cyclic group) and if it is spectral. Such efficient procedures are lacking however and we make a few comments for the computational complexity of some related problems.

2000 Mathematics Subject Classification. Primary 52C22, Secondary 20K01, 42B99.

Keywords and phrases. *Spectral sets, complex Hadamard matrices, translational tiles, Spectral Set Conjecture*

1. INTRODUCTION

Let G be a locally compact Abelian group and $\Omega \subseteq G$ be a bounded open set. We call Ω *spectral* if there exists a set Λ of continuous characters of G which forms an orthogonal basis of the space $L^2(\Omega)$ (with respect to Haar measure). Such a set Λ is called a *spectrum* of Ω .

The set Ω tiles G by translation if there exists a set $T \subseteq G$ of translates, called a *tiling complement* of Ω , such that $\sum_{t \in T} \chi_\Omega(x - t) = 1$, for almost all (with respect to the Haar measure) $x \in G$. Here χ_Ω denotes the indicator function of Ω .

The *Spectral Set Conjecture* (originally proposed in $G = \mathbb{R}^d$ by Fuglede, [2]) states that Ω is a tile in G if and only if it is spectral.

This conjecture has attracted considerable attention over the last decade. Many positive results were obtained in special cases (i.e., under various restrictions on Ω, G , and T ; see [2, 5, 6, 7, 10, 11, 12, 14]), until Tao [16] recently proved that the direction “spectral \Rightarrow tile” does not hold in dimension 5 and higher. Tao’s result was supplemented by a result of Matolcsi [13], which allowed the dimension of the counterexample to be reduced to 4. In the second part of this paper we combine

Date: October 2004.

Partially supported by European Commission IHP Network HARP (Harmonic Analysis and Related Problems), Contract Number: HPRN-CT-2001-00273 - HARP.

the ideas of [16] and [13] in order to reduce the dimension to 3. We also remark that the “tile \Rightarrow spectral” direction was recently disproved in [9] in dimension 5 and higher. The existing counterexamples do not exclude the possibility that the conjecture may still be true in low dimensions (in particular, in dimension 1), or under some natural restrictions (such as Ω being a convex set in Euclidean space).

The results of this paper concern the direction “spectral \Rightarrow tile” of the Spectral Set Conjecture in finite Abelian groups and \mathbb{Z}^d and \mathbb{R}^d . In Section 2 we prove the validity of this direction of the conjecture for sets of at most 5 elements (and the corresponding union of unit cubes in the case of \mathbb{R}^d). On the other hand, in Section 3 we show a particular example of a 6-element set in $G = \mathbb{Z}_8^3$ which is spectral but does not tile G by translation. This allows us to improve the results of [16] and [13], by producing a counterexample in \mathbb{R}^3 . Finally in Section 4 we discuss the algorithmic aspect of when a given set tiles a finite cyclic group or is spectral in that group. Such questions arise naturally and, unfortunately, with no satisfying answer, when one searches for counterexamples to the Spectral Set Conjecture in cyclic groups (any such counterexample would extend to a counterexample in \mathbb{R}).

2. SPECTRAL SETS OF SMALL SIZE

We recall the following notations and definitions:

A $k \times k$ complex matrix H is called a (complex) *Hadamard matrix* if all entries of H have absolute value 1, and $HH^* = kI$ (where I denotes the identity matrix). This means that the rows (and also the columns) of H form an orthogonal basis of \mathbb{C}^k . A *log-Hadamard matrix* is any real square matrix $(h_{i,j})_{i,j=1}^k$ such that the matrix $(e^{2\pi i h_{i,j}})_{i,j=1}^k$ is Hadamard.

Based on the ideas of [16] and [13] we will make use of the connection between complex Hadamard matrices and spectral sets to prove the following:

Theorem 2.1. *Let G be any finite Abelian group and $A \subset G$ a spectral set in G with $|A| \leq 5$. Then A tiles G .*

Proof. We begin the proof by recalling a general property of spectral sets and tiles. Namely, if $H \leq G$ is a subgroup and $A \subset H$, then A is spectral (resp. a tile) in G if and only if A is spectral (resp. a tile) in H . The only non-trivial statement here is that if A is spectral in H then it is also spectral in G . This follows directly from the fact that each character of H can be extended, not necessarily uniquely, to a character of G . To see this, we need to show that the natural (restriction) homomorphism $\phi : \hat{G} \rightarrow \hat{H}$ is surjective. Any element in the kernel of ϕ is constant on each coset of H , therefore it can be identified with a character of the factor group G/H . Then, the surjectivity of ϕ follows from comparing the number of elements of the groups $G, H, G/H$.

It is a well known fact that for any finite Abelian group G we may choose natural numbers N, d such that $G \leq \mathbb{Z}_N^d$. By the remark above, it is enough to prove the statement for groups of the type $G = \mathbb{Z}_N^d$. This observation makes the proof technically simpler.

The essential part of the proof relies on the fact that we have a full characterization of complex Hadamard matrices up to order 5.

We identify the elements of G and \hat{G} , the group of characters of G , with d -dimensional column- and row-vectors, respectively. Let $A \subset G = \mathbb{Z}_N^d$, with $|A| = k \leq 5$. We regard A as a $d \times k$ matrix with integer coefficients. If $L \subset \hat{G}$ is a spectrum of A (regarded as a $d \times k$ matrix), then $H := \frac{1}{N}L \cdot A$ is log-Hadamard (where the matrix multiplication can be taken mod N). Multiplication by the matrix L defines a homomorphism from G to \mathbb{Z}_N^k , and the images of the elements of A are given by the columns c_j of $L \cdot A$ ($0 \leq j \leq k-1$). By Proposition 2.3 in [13] it is enough to prove that the vectors c_j tile \mathbb{Z}_N^k .

In the cases $k = 1, 2, 3, 5$ this follows immediately from the uniqueness (up to natural equivalence) of complex Hadamard matrices of order k . This uniqueness is trivial for $k = 1, 2, 3$, while the case $k = 5$ is settled in [4]. Indeed, we can assume without loss of generality that $0 \in A$ and $0 \in L$ (due to the trivial translation invariance of the notion of spectrality and spectrum), and this already implies that the matrix $H = h_{m,j}$ is (after a permutation of columns) given by $h_{m,j} = \frac{1}{k}mj$ ($0 \leq m, j \leq k-1$). It follows that N is a multiple of k , say $N = Mk$, and the column vectors c_j are given as $c_j = (0, jM, 2jM, \dots, (k-1)jM)^T$. In order to see that these vectors tile \mathbb{Z}_N^d we invoke Proposition 2.3 in [13] once again. Let $V := (0, 1, 0, \dots, 0)$, and consider the mod N product $V \cdot L \cdot A = (0, M, 2M, \dots, (k-1)M)$. It is obvious that the set $\{0, M, 2M, \dots, (k-1)M\}$ tiles \mathbb{Z}_N , and therefore the columns c_j tile \mathbb{Z}_N^k .

The case $k = 4$ is settled in a similar manner, although we have no uniqueness of Hadamard matrices in this case. The general form of a 4×4 complex Hadamard matrix is given (see e.g. [4], Proposition 2.1) by the parametrization

$$U = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & e^{2\pi i \phi} & -e^{2\pi i \phi} \\ 1 & -1 & -e^{2\pi i \phi} & e^{2\pi i \phi} \end{pmatrix}.$$

Due to the presence of -1's it follows that N must be a multiple of 2, say $N = 2M$. The matrix LA is then given by

$$LA = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & M & M \\ 0 & M & M+t & t \\ 0 & M & t & M+t \end{pmatrix}.$$

To see that the columns of this matrix tile \mathbb{Z}_N^4 , consider the matrix

$$V_2 := \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Then

$$V_2 LA = \begin{pmatrix} 0 & 0 & M & M \\ 0 & M & M+t & t \end{pmatrix}.$$

It is trivial to check that the columns of this matrix tile \mathbb{Z}_N^2 , and, by Proposition 2.3 in [13], this implies that the columns of LA tile \mathbb{Z}_N^4 .

□

Next, we extend the previous result to the infinite grid \mathbb{Z}^d . First, we need to establish the rationality of the spectra in the cases considered.

Proposition 2.2. *Let $A \subset \mathbb{Z}^d$ be a spectral set with $|A| \leq 5$. Then A admits a rational spectrum.*

Proof. Note that we do not claim that *all* spectra of A must be rational, but only that the spectrum can be chosen rational.

The proof is an easy argument from linear algebra. Let us first consider the case $|A| = 5$ (the cases $|A| = 1, 2, 3$ are settled the same way, while $|A| = 4$ will require some extra considerations). Let $L \subset \mathcal{T}^d$ denote a spectrum of A . We may assume that $0 \in A$ and $0 \in L$. Then, after a permutation of elements of A , we have

$$LA = \frac{1}{5} \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 3 & 4 \\ 0 & 2 & 4 & 1 & 3 \\ 0 & 3 & 1 & 4 & 2 \\ 0 & 4 & 3 & 2 & 1 \end{pmatrix} \pmod{1}.$$

Let $l_{m,j}$ denote the elements of L . Considering, for example, the second row of L we see that there exist integers $z_{2,1}, \dots, z_{2,5}$ such that

$$(l_{2,1}, \dots, l_{2,d}) \cdot A = \left(z_{2,1}, z_{2,2} + \frac{1}{5}, \dots, z_{2,5} + \frac{4}{5} \right).$$

Regarding this equation as a set of linear equations with variables $l_{2,1}, \dots, l_{2,d}$ we see that if there exists a solution, then the solution can be chosen rational. The same argument holds for the other rows of L .

We now turn to the case $|A| = 4$. Then, for some $q \in [0, 1]$ we have

$$LA = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & \frac{1}{2} & \frac{1}{2} + q & q \\ 0 & \frac{1}{2} & q & \frac{1}{2} + q \end{pmatrix} \pmod{1}.$$

If q is rational then the previous argument applies. If q is irrational we need some additional considerations. Applying the previous argument we see that the first two rows of L can be chosen rational even in this case (they do not depend on q). It is also clear that the fourth row of L can be chosen as the sum of the second and third rows. Consider therefore the third row only. For some integers $z_{3,1}, \dots, z_{3,4}$ we have

$$(l_{3,1}, \dots, l_{3,d}) \cdot A = \left(z_{3,1}, z_{3,2} + \frac{1}{2}, z_{3,3} + \frac{1}{2} + q, z_{3,4} + q \right)$$

Regarding this equation as a set of linear equations with variables $l_{2,1}, \dots, l_{2,d}, q$ we see that if there exists a solution then all the variables (including q) can be chosen rational. \square

Now, we are in position to prove the analogue of Theorem 2.1 in \mathbb{Z}^d .

Theorem 2.3. *Let $A \subset \mathbb{Z}^d$ be a spectral set in \mathbb{Z}^d with $|A| \leq 5$. Then A tiles \mathbb{Z}^d .*

Proof. By Proposition 2.2 we can choose the spectrum of A rational. This means that A is already spectral in some finite group $G = \mathbb{Z}_{n_1} \times \dots \times \mathbb{Z}_{n_d}$. By Theorem 2.1 we conclude that A tiles G , and therefore it tiles \mathbb{Z}^d . \square

Finally, we formulate the corresponding statement in \mathbb{R}^d .

Theorem 2.4. *Let A be a set in \mathbb{Z}^d with $|A| \leq 5$. Let $\Omega := A + [0, 1]^d$ be the corresponding union of unit cubes. If Ω is spectral then it tiles \mathbb{R}^d .*

Proof. By Theorem 4.2 in [9] the set Ω is spectral in \mathbb{R}^d if and only if A is spectral in \mathbb{Z}^d . Therefore, Theorem 2.3 implies the result. \square

3. THE SPECTRAL SET CONJECTURE IS FALSE IN DIMENSION 3

In the previous section we saw how a characterization of Hadamard matrices (up to order 5) leads to an understanding of tiling properties of spectral sets. The situation changes drastically when we consider Hadamard matrices of order 6. On the one hand side we do not seem to have a full characterization of complex Hadamard matrices of order 6 or greater. On the other hand we do possess particular examples (and even a description [1] of some parametric families of 6×6 Hadamard matrices) which show that the statement of Theorem 2.1 does not hold for sets of size $|A| = 6$. Such an example was exploited in [16] (and subsequently in [13]) to disprove Fuglede’s conjecture in \mathbb{Z}_3^5 (resp. \mathbb{Z}_3^4) and then transfer the counterexample to \mathbb{Z}^5 and \mathbb{R}^5 (resp. \mathbb{Z}^4 and \mathbb{R}^4).

In this section we improve the results mentioned above by showing that the direction “spectral \Rightarrow tile” of the Spectral Set Conjecture fails already in dimension 3. We use a combination of the ideas of [16] and [13].

Our starting point is the following numerical example of a particular 6×6 log-Hadamard matrix. Let

$$H := \frac{1}{8} \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 4 & 2 & 6 & 6 & 2 \\ 0 & 2 & 4 & 1 & 5 & 6 \\ 0 & 6 & 3 & 4 & 2 & 7 \\ 0 & 6 & 7 & 2 & 4 & 3 \\ 0 & 2 & 6 & 5 & 1 & 4 \end{pmatrix}.$$

It is easy to check that H is log-Hadamard (this example corresponds to a particular element of the one-parameter family of 6×6 Hadamard matrices given in [1, p. 5357]).

Next, we observe that the “mod 8 rank” of $8H$ is 3. This means that there exist integer matrices A and L of size 3×6 and 6×3 , respectively, such that $8H = LA \pmod{8}$. A possible example of such a decomposition is the following:

$$A := \begin{pmatrix} 0 & 2 & 4 & 1 & 5 & 6 \\ 0 & 6 & 3 & 4 & 2 & 7 \\ 0 & 6 & 7 & 2 & 4 & 3 \end{pmatrix} \quad \text{and} \quad L := \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 7 & 1 & 1 \end{pmatrix}.$$

This means that the set (of the columns of) A is spectral in the group $G = \mathbb{Z}_8^3$. However, $|A| = 6$, therefore A cannot tile G due to obvious divisibility reasons.

We can transfer this counterexample to \mathbb{Z}^3 and \mathbb{R}^3 in the same way as described in [16] and [13]. Namely, the set $A_n := A + 8[0, n]^3$ is spectral in \mathbb{Z}^3 for every n , but it does not tile \mathbb{Z}^3 for large enough values of n (see Propositions 2.1 and 2.5 in [13]). Finally, for such a large value n , the set $\Omega = A_n + [0, 1]^3$ is a finite union of

unit cubes, which is spectral in \mathbb{R}^3 but does not tile \mathbb{R}^3 (cf. the proof of Theorem 3.1 in [13]).

We see that the counterexample is simply based on the existence of a particular log-Hadamard matrix with prescribed properties (namely, to get the non-divisibility condition the size (6) must not divide the common denominator of the entries (8); the dimension (3) is then determined by the 'rank' of the matrix, i.e. the smallest possible decomposition). In principle, this would seem to be the way to search for counterexamples in dimensions 2 and 1.

However, due to the lack of characterization of Hadamard matrices of order greater than 5, we do not know of any methodical way of *how* to produce Hadamard matrices with the desired properties. The particular example above was simply found in the literature by looking through the existing (fairly scattered) results on complex Hadamard matrices.

It seems that in order to prove or disprove this direction of the Spectral Set Conjecture in dimensions 1 and 2, some new ideas will be needed. Let us also mention that the direction "tile \Rightarrow spectral" has recently been disproved [9] in dimension 5 and higher, but remains open in lower dimensions.

4. ALGORITHMIC QUESTIONS

In searching for counterexamples to the Spectral Set Conjecture using a computer one is immediately faced with the following problem: it appears that no efficient algorithm is known to decide if a given subset E of some group, say the cyclic group \mathbb{Z}_n , is a tile. The customary definition of efficiency for such an algorithm demands that it should run in time polynomial in n . No efficient algorithm appears to be known also in the case of deciding if a given set is spectral or not. The obvious algorithms, namely to test all possible tiling complements and all possible spectra, are clearly exponential in n .

Both questions would easily be solvable given an efficient algorithm to decide the following question.

PROBLEM DIFF:

- INPUT: We are given a positive integer n and sets $E, D \subseteq \mathbb{Z}_n$
- OUTPUT: Maximal k such that there exists $A \subseteq E$, with $|A| = k$, and $A - A \subseteq D$.

Indeed, assuming one could solve the above problem in polynomial time in n , one could use this postulated procedure to decide if $T \subseteq \mathbb{Z}_n$ tiles \mathbb{Z}_n by using $E = \mathbb{Z}_n$ and $D = (T - T)^c \cup \{0\}$. If $x = n/|T|$ is an integer and the answer to this query is equal to x then T tiles \mathbb{Z}_n with translation set equal to the set A in the description of problem DIFF, otherwise T is not a tile.

To decide spectrality of $S \subseteq \mathbb{Z}_n$ in polynomial time, first compute the Fourier Transform of χ_S , its zero set Z and invoke the procedure for problem DIFF with $E = D = \{0\} \cup Z$. If the answer k is equal to $|S|$ then, and only then, S is spectral in \mathbb{Z}_n and one of its spectra is the set A in the description of problem DIFF.

It is easily seen that DIFF is solvable in polynomial time if and only if the corresponding decision problem, described below, is solvable in polynomial time.

PROBLEM DIFF':

- INPUT: We are given positive integers k and n and sets $E, D \subseteq \mathbb{Z}_n$
- OUTPUT: YES if there exists $A \subseteq E$, with $|A| = k$, and $A - A \subseteq D$, and NO otherwise.

We show now that problem DIFF' is NP-complete [3]. We remark however that this does not prove that the corresponding tiling and spectrality decision problems are NP-complete.

Theorem 4.1. *The decision problem DIFF' is NP-complete.*

Proof. The following problem (decision version of the maximum independent set optimization problem) is well known [3] to be NP-complete:

PROBLEM IND:

- INPUT: We are given a simple graph on n vertices and a positive integer k .
- OUTPUT: YES if there exists a set of vertices A none of which is connected via an edge to another vertex in A (the set A is then called an independent set) and such that $|A| = k$.

To prove that DIFF' is NP-complete it suffices to give a polynomial time algorithm to solve the arbitrary instance of IND using a black-box algorithm that solves DIFF' in unit time.

Let $V = \{1, \dots, n\}$ be the set of vertices of a given graph G . The first step is to define an efficiently computable function $\phi : V \rightarrow \mathbb{Z}_m$, with m bounded by a polynomial in n , and a set $A \subseteq \mathbb{Z}_m$ such that

$$(1) \quad i, j \in V \text{ are connected in } G \Leftrightarrow \phi(i) - \phi(j) \in A.$$

For this we note that one can easily construct such a map ϕ and set A using the greedy method, as long as m is large compared to n . Indeed, choose $\phi(1) = 0$, and, having defined $\phi(1), \dots, \phi(r)$, $r < n$, define $\phi(r + 1)$ to be the minimum $\nu \in \{1, \dots, m - 1\}$ such that

$$\nu \notin \{\phi(i) + \phi(j) - \phi(l) : 1 \leq i, j, l \leq r\}.$$

(It is easy to see that this procedure can be carried out if m is large enough, namely $m \geq Cn^3$.) This ensures that all differences $\phi(i) - \phi(j)$, $1 \leq i \neq j \leq n$ are distinct.

Next we define

$$A = \{\phi(i) - \phi(j) : i \text{ and } j \text{ are connected in } G\}.$$

To decide if G has an independent set of size k it is enough to use the algorithm for DIFF' for \mathbb{Z}_m with the set $E = \phi(\{1, \dots, n\})$ and A as constructed above. \square

REFERENCES

- [1] P. Dita, Some results on the parametrization of complex Hadamard matrices, J. Phys. A: Math. Gen. **37** (2004), no. 20, 5355-5374.
- [2] B. Fuglede, Commuting self-adjoint partial differential operators and a group theoretic problem, J. Funct. Anal. **16** (1974), 101-121.
- [3] M.R. Garey, D.S. Johnson, *Computers and Intractability, A Guide to the Theory of NP-Completeness*, Freeman, New York, 1979.

- [4] U. Haagerup, Orthogonal maximal abelian *-subalgebras of the $n \times n$ matrices and cyclic n -roots, in *Operator algebras and quantum field theory* (Rome, 1996), 296–322, Internat. Press, Cambridge, MA, 1997.
- [5] A. Iosevich, N. Katz, T. Tao, Convex bodies with a point of curvature do not admit exponential bases, *Amer. J. Math.* **123** (2001), 115-120.
- [6] A. Iosevich, N. Katz, T. Tao, The Fuglede spectral conjecture holds for convex bodies in the plane, *Math. Res. Letters* **10** (2003), 559-570.
- [7] M.N. Kolountzakis, Non-symmetric convex domains have no basis of exponentials, *Ill. J. Math.* **44** (2000), no. 3, 542-550.
- [8] M.N. Kolountzakis, The study of translational tiling with Fourier Analysis, Proceedings of the Workshop on Fourier Analysis and Convexity, Università di Milano-Bicocca, 2001, **to appear**.
- [9] M.N. Kolountzakis, M. Matolcsi, Tiles with no spectra, preprint.
- [10] S. Konyagin, I. Laba, Spectra of certain types of polynomials and tiling of integers with translates of finite sets, *J. Number Th.* **103** (2003), no. 2, 267-280.
- [11] J.C. Lagarias and Y. Wang, Spectral sets and factorizations of finite Abelian groups, *J. Func. Anal.* **145** (1997), 73-98.
- [12] J.C. Lagarias, S.Szabó, Universal spectra and Tijdeman’s conjecture on factorization of cyclic groups, *J. Fourier Anal. Appl.* **7** (2001), no. 1, 63-70.
- [13] M. Matolcsi, Fuglede’s conjecture fails in dimension 4, **preprint**.
- [14] S. Pedersen, Y. Wang, Universal spectra, universal tiling sets and the spectral set conjecture, *Math. Scand.* **88** (2001), no. 2, 246-256.
- [15] W. Rudin, *Fourier analysis on groups*, Reprint of the 1962 original. Wiley Classics Library. A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1990.
- [16] T. Tao, Fuglede’s conjecture is false in 5 and higher dimensions, *Math. Res. Letters*, to appear.

M.K.: SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, 686 CHERRY ST NW, ATLANTA, GA 30332, USA, AND DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CRETE, KNOSSOS AVE., GR-714 09, IRAKLIO, GREECE

E-mail address: kolount@member.ams.org

M.M.: ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES POB 127 H-1364 BUDAPEST, HUNGARY.

E-mail address: matomate@renyi.hu