

FUGLEDE'S CONJECTURE FAILS IN DIMENSION 4

MÁTÉ MATOLCSI

ABSTRACT. In this note we give an example of a set $\Omega \subset \mathbb{R}^4$ such that $L^2(\Omega)$ admits an orthonormal basis of exponentials $\{\frac{1}{|\Omega|^{1/2}}e^{2\pi i\langle x, \xi \rangle}\}_{\xi \in \Lambda}$ for some set $\Lambda \subset \mathbb{R}^4$, but which does not tile \mathbb{R}^4 by translations. This improves Tao's recent 5-dimensional example, and shows that one direction of Fuglede's conjecture fails already in dimension 4. Some common properties of translational tiles and spectral sets are also proved.

2000 Mathematics Subject Classification. Primary 42B99, Secondary 20K01.

Keywords and phrases. *Translational tiles, spectral sets, Fuglede's conjecture, Hadamard matrices.*

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^d$ be a Lebesgue measurable set of finite non-zero measure. A set $\Lambda \subset \mathbb{R}^d$ is said to be a *spectrum* of Ω if the functions $\{\frac{1}{|\Omega|^{1/2}}e^{2\pi i\langle x, \xi \rangle}\}_{\xi \in \Lambda}$ form an orthonormal basis of $L^2(\Omega)$ (here $|\Omega|$ denotes the Lebesgue measure of Ω). If such Λ exists, Ω is said to be *spectral*. The set Ω is said to be a (translational) *tile* if it is possible to tile \mathbb{R}^d with a family of translates $\{t + \Omega : t \in \Sigma\}$ of Ω (ignoring overlaps and gaps of measure zero).

Tiles and spectral sets are connected by Fuglede's famous

Conjecture 1.1. [1] *A set $\Omega \subset \mathbb{R}^d$ of finite, non-zero Lebesgue measure is a tile if and only if it is spectral.*

Fuglede [1] proved the conjecture in the special case when the spectrum Λ or the translation set Σ is assumed to be a lattice. The general case of the conjecture was open for nearly 30 years, until last year Tao [13] showed an example to disprove one direction of the conjecture in 5 and higher dimensions. Namely, he gave an example of a spectral set in \mathbb{R}^5 which is not a tile. The aim of this note is to modify Tao's example and prove that the conjecture fails already in dimension 4. We emphasize that Propositions 2.1 and 2.5 are only new in the generality formulated below, and credit should be given to [13], where special

cases are proved with the same idea. The essentially new results of the paper are contained in Propositions 2.2, 2.3 and Theorem 3.1.

We will restrict our attention to sets of the form of finite union of unit cubes placed at points with integer coordinates, i.e.

$$(1) \quad \Omega = \cup_{t \in \Gamma} \{t + [0, 1)^d\},$$

where $\Gamma \subset \mathbb{Z}^d$ is finite (this means, essentially, that we will be considering the tiling and spectral properties of sets in \mathbb{Z}^d). Besides giving a counterexample in dimension 4, we also prove some interesting properties that are shared by spectral sets and tiles of the form (1).

To assess our present knowledge concerning this topic it is interesting to note that Fuglede's conjecture was recently proved in 2 dimensions for *convex* bodies in [5]. Even for finite union of intervals the conjecture is still open in dimension 1. Some partial results concerning this case appeared in [10], [11]. In arbitrary dimensions certain classes of non-tiling sets are already known to admit no spectrum, either. Results in this direction are contained in [4], [7] and [8]. For further recent results related to Fuglede's conjecture see [3], [12] [2] and [6]. The recent survey [9] gives an overview not only of the Fuglede conjecture but also of some recent results in the theory of translational tilings.

Note, also, that one direction of the conjecture is still open in all dimensions, namely we have no examples of tiles which do not admit a spectrum.

2. COMMON PROPERTIES OF TILING AND SPECTRAL SETS

In this section we examine some natural properties of spectral sets and tiles of the form (1).

We will need the following notations and definitions.

The family of spectral sets (resp. tiles) of the form (1) will be denoted by \mathcal{S}^d (resp. \mathcal{T}^d).

A $k \times k$ complex matrix H is called a *Hadamard matrix* if all entries of H have absolute value 1, and $HH^* = kI$ (where I denotes the identity matrix). This means that the rows (and also the columns) of H form an orthogonal basis of \mathbb{C}^k . A *log-Hadamard matrix* is any real square matrix $(h_{i,j})_{i,j=1}^k$ such that the matrix $(e^{2\pi i h_{i,j}})_{i,j=1}^k$ is Hadamard.

For a given finite set $\{t_1, \dots, t_k\} \subset \mathbb{Z}^d$ let T denote the $d \times k$ matrix with j th column t_j . (With a slight abuse of notation we will denote the set itself by T as well.) The set $T = \{t_1, \dots, t_k\}$ is called *spectral* if there exists a $k \times d$ real matrix L such that LT is log-Hadamard (in the usual terminology of harmonic analysis it is also required that the entries of L belong to the interval $[0, 1)$ but the definition above will be more

convenient for present purposes). The (rows of the) matrix L is the (not necessarily unique) *spectrum* of T . The set $T = \{t_1, \dots, t_k\} \subset \mathbb{Z}^d$ is called a *tile* if it is possible to tile \mathbb{Z}^d with a family of translates of T . The family of spectral sets (resp. tiles) of \mathbb{Z}^d will be denoted by \mathcal{S}_0^d (resp. \mathcal{T}_0^d).

The set $T = \{t_1, \dots, t_k\} \subset \mathbb{Z}^d$ is called *m-spectral* for some positive integer m if it admits a spectrum L with all entries being multiples of $1/m$ (which, essentially, means that T is spectral in the group \mathbb{Z}_m^d). The set T is called an *m-tile* if it is possible to tile the cube $[0, m)^d$ with translates of $T \bmod m$. (This means that T tiles the group \mathbb{Z}_m^d .) The family of *m-spectral* sets (resp. *m-tiles*) of \mathbb{Z}^d will be denoted by \mathcal{S}_m^d (resp. \mathcal{T}_m^d).

It is obvious that $\mathcal{S}_m^d \subset \mathcal{S}_0^d$ and $\mathcal{T}_m^d \subset \mathcal{T}_0^d$, and it is also clear that if $T \in \mathcal{T}_0^d$ then $T + [0, 1)^d \in \mathcal{T}^d$. It is also well-known that if $T \in \mathcal{S}_0^d$ with spectrum $L = \{l_1, \dots, l_k\}$ then $\Omega = T + [0, 1)^d \in \mathcal{S}^d$ with spectrum $\Lambda = L + \mathbb{Z}^d$ (the orthogonality of the functions $e^{2\pi i \langle \lambda, x \rangle}$ is an easy calculation, while the completeness can be seen by checking the Parseval identity for functions of the form $e^{2\pi i \langle s, x \rangle}$ and noting that the linear span of these functions is dense in $L^2(\Omega)$).

Tao's example [13] shows that $\mathcal{S}_0^d \neq \mathcal{T}_0^d$ and $\mathcal{S}^d \neq \mathcal{T}^d$ for $d \geq 5$, therefore Fuglede's conjecture is false, in general. Nevertheless, it is interesting to see that \mathcal{S}_0^d and \mathcal{T}_0^d share certain properties. Namely, it is reasonable to look for 'natural' properties of \mathcal{T}_0^d and check whether the same holds for \mathcal{S}_0^d , and vice versa. The first result in this direction establishes the 'composition property' for \mathcal{S}_0^d and \mathcal{T}_0^d . This property is already used in Tao's example (in less generality, but the idea is the same).

Proposition 2.1. *Assume that the set $T = \{t_1, \dots, t_k\} \subset \mathbb{Z}^d$ is an m -tile (resp. m -spectral) and the set $S = \{s_1, \dots, s_r\} \subset \mathbb{Z}^d$ is an n -tile (resp. n -spectral). Then the set $\Gamma := T + mS$ is an mn -tile (resp. mn -spectral).*

Proof. This property is 'natural' for tiles, therefore we prove only the spectral part of the proposition.

Let $L := \{\frac{l_1}{m}, \dots, \frac{l_k}{m}\}$ be the spectrum of T , and $Q := \{\frac{q_1}{n}, \dots, \frac{q_r}{n}\}$ be the spectrum of S . We claim that the spectrum of Γ is $\Lambda = L + \frac{Q}{m}$.

The cardinality of Λ and Γ is the same (mn), therefore it is enough to check orthogonality of the functions $e^{2\pi i \langle \lambda, x \rangle}$.

To see this write $\lambda_{i_1} = \frac{l_{i_1}}{m} + \frac{q_{i_1}}{nm}$, $\lambda_{i_2} = \frac{l_{i_2}}{m} + \frac{q_{i_2}}{nm}$, and

$$(2) \quad \sum_{x \in \Gamma} e^{2\pi i \langle \lambda_{i_1} - \lambda_{i_2}, x \rangle} = \sum_{t \in T} \sum_{s \in S} e^{2\pi i \langle \frac{l_{i_1} - l_{i_2}}{m} + \frac{q_{i_1} - q_{i_2}}{nm}, t + ms \rangle}.$$

Notice that if $q_{i_1} \neq q_{i_2}$ then the second summation gives 0 for all fixed $t \in T$ (because of the spectral property of Q with respect to S), therefore the whole sum is 0. On the other hand, if $q_{i_1} = q_{i_2}$ then the second summation gives $r e^{2\pi i \langle \frac{l_{i_1} - l_{i_2}}{m}, t \rangle}$, and the whole sum equals 0 because of the spectral property of L with respect to T . \square

The next two propositions are not directly used in our 4-dimensional counterexample below, but they are of independent interest. First we examine a 'natural' ascendance property of spectral sets.

Proposition 2.2. *Assume that for a given set $T = \{t_1, \dots, t_k\} \subset \mathbb{Z}^d$ there exists a $d_1 \times d$ matrix L_1 with integer entries for which the columns of $L_1 T$ form a spectral (resp. m -spectral) set T_1 in \mathbb{Z}^{d_1} . Then T itself is spectral (resp. m -spectral).*

Proof. If L denotes a spectrum of T_1 then LL_1 is clearly a spectrum of T , because $(LL_1)T = L(L_1T)$. \square

It may be a little surprising that the same property holds also for tiles.

Proposition 2.3. *Assume that for a given set $T = \{t_1, \dots, t_k\} \subset \mathbb{Z}^d$ there exists a $d_1 \times d$ matrix L_1 with integer entries for which the columns of $L_1 T$ form a tile (resp. m -tile) T_1 in \mathbb{Z}^{d_1} . Then T itself is a tile (resp. m -tile).*

Proof. We prove the statement for tiles, and remark that the case of m -tiles is settled exactly the same way.

Note first that the elements of T_1 must be different from each other because of the tiling condition. Let Σ_1 denote the translation set corresponding to T_1 , i.e. $\Sigma_1 + T_1 = \mathbb{Z}^{d_1}$. Define $\Sigma := L_1^{-1}[\Sigma_1] = \{\sigma \in \mathbb{Z}^d : L_1 \sigma \in \Sigma_1\}$. We claim that Σ is a good translation set for T .

To see this, we check first that the translated copies of T are disjoint. Assume therefore that $\sigma_1 + t_i = \sigma_2 + t_j$ for some $\sigma_1, \sigma_2 \in \Sigma$ and $t_i, t_j \in T$. Applying the transformation L_1 we get $L_1 \sigma_1 + L_1 t_i = L_1 \sigma_2 + L_1 t_j$, and in view of the tiling condition on T_1 we conclude that $L_1 t_i = L_1 t_j$. This implies $t_i = t_j$, and hence $\sigma_1 = \sigma_2$.

Next we check that the translates of T cover the whole of \mathbb{Z}^d . Let $z \in \mathbb{Z}^d$ arbitrary, and let $z^1 = L_1 z \in \mathbb{Z}^{d_1}$. Then, by assumption, $z^1 = \sigma^1 + t^1$ with some $\sigma^1 \in \Sigma_1$ and $t^1 \in T_1$. Take the inverse image

of t^1 , i.e. the element $t_i \in T$ for which $L_1 t_i = t^1$. Then $\sigma := z - t_i$ is contained in Σ because $L_1(z - t_i) = z^1 - t^1 = \sigma^1 \in \Sigma_1$. \square

Let us mention an interesting consequence of this proposition.

Corollary 2.4. *Let $T = \{t_1, \dots, t_k\} \subset \mathbb{Z}^d$ be a linearly independent set of vectors (over \mathbb{R}). Then T is a tile in \mathbb{Z}^d .*

Proof. By the assumption of linear independence we can delete $d - k$ rows of the matrix T , so that the remaining k -dimensional columns $PT = \{Pt_1, \dots, Pt_k\}$ span \mathbb{R}^k . (The deletion of rows corresponds to multiplication by a $k \times d$ matrix P with entries 1's and 0's.) In view of Proposition 2.3, it is enough to show that PT is a tile in \mathbb{Z}^k .

Consider the column vectors $\underline{k} = (0, 1, \dots, k-1)^*$ (the $*$ in the exponent denotes transposition) and $\underline{j} := (\det PT) \cdot PT^{*-1} \underline{k}$. Notice that the matrix $(\det PT) \cdot PT^{*-1}$ (and hence the vector j) has integer entries, and the 1-dimensional set $\underline{j}^* \cdot PT = (\det PT) \underline{k}^*$ obviously tiles \mathbb{Z} . Hence, Proposition 2.3 applies, and PT is a tile in \mathbb{Z}^k . \square

Finally, we examine a property of 'non-tiling' sets which played an essential role in the counterexample of [13] (we give a slightly generalized version here).

Proposition 2.5. *Assume that the set $T = \{t_1, \dots, t_k\} \subset [0, m]^d$ is not an m -tile. Then for large n the set $S_n := T + (m \cdot [0, n]^d)$ is not a tile.*

Proof. The proof proceeds along the same lines as in [13], and we include it only for completeness.

Assume, for contradiction, that S_n tiles \mathbb{Z}^d with some translation set Σ . Take a cube $C_l = [0, l]^d$, where l is much larger than n . Let $\Sigma_l := \{\sigma \in \Sigma : (\sigma + S_n) \cap C_l \neq \emptyset\}$. Note that $\#S_n = kn^d$. We have $\#\Sigma_l \leq \frac{(l+2mn)^d}{kn^d}$, because all Σ_l -translates of S_n are contained in the cube $(-mn, l + mn)^d$.

Let A denote the annulus $A := [-m, mn+m]^d - [m, mn-m]^d$. Then $\#A (\approx 4dm(mn)^{d-1}) \leq 5dm(mn)^{d-1}$, if n is large enough compared to m . Hence, $\Sigma_l + A$ cannot cover the cube $C_{l-m} := [0, l-m]^d$ because $(\#\Sigma_l)(\#A) \leq (l+2mn)^d \left(\frac{5dm(mn)^{d-1}}{kn^d} \right) < (l-m)^d$, if the numbers n, l are chosen so that n is sufficiently large compared to m , and l is sufficiently large compared to n .

Take a point $x \in C_{l-m}$ not covered by $\Sigma_l + A$. Consider the cube $C_m^x := x + [0, m]^d$. This cube is fully inside C_l , therefore if any translate $\sigma + S_n$ intersects C_m^x then σ necessarily belongs to Σ_l . The point x is not covered by the annulus $\sigma + A$, therefore C_m^x is contained in the cube $\sigma + [0, mn]^d$. Let S denote the set $T + m \cdot \mathbb{Z}^d$. In view of what has been

said, we have $(\sigma + S_n) \cap C_m^x = (\sigma + S) \cap C_m^x = (x + [0, m]^d) \cap (\sigma + T + m\mathbb{Z}^d)$. The mod m reduction of this set is exactly the translate $\sigma + T \pmod{m}$. Hence, the tiling of the cube C_m^x by Σ -translates of S_n contradicts the assumption that T is not an m -tile. \square

It would be interesting to see whether the corresponding result holds also for spectral sets.

3. COUNTEREXAMPLE IN DIMENSION 4

We can now show that one direction of Fuglede's conjecture fails already in dimension 4.

Theorem 3.1. *For any $d \geq 4$ there exists a finite union of unit cubes $\Omega = \cup_{j=1}^k \{s_j + [0, 1]^d\}$ in \mathbb{R}^d , such that Ω is spectral but not a tile.*

Proof. Take the matrix

$$K := \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 2 & 2 \\ 0 & 1 & 0 & 2 & 2 & 1 \\ 0 & 1 & 2 & 0 & 1 & 2 \\ 0 & 2 & 2 & 1 & 0 & 1 \\ 0 & 2 & 1 & 2 & 1 & 0 \end{pmatrix}$$

and observe that $\frac{1}{3}K$ is log-Hadamard (and corresponds to the Hadamard matrix H given in [13]).

Note that K is of rank 4 mod 3, therefore it is possible to make a mod 3 decomposition $K = L \cdot T$, where L is a 6×4 and T is a 4×6 matrix. We give a specific mod 3 decomposition for convenience:

$$L := \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 2 \\ 1 & 0 & 2 & 2 \\ 1 & 2 & 0 & 1 \\ 2 & 2 & 1 & 0 \\ 2 & 1 & 2 & 1 \end{pmatrix} \quad \text{and} \quad T := \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 2 \\ 0 & 0 & 1 & 0 & 0 & 2 \\ 0 & 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 0 & 1 & 2 \end{pmatrix}.$$

Note, however, that infinitely many decompositions exist, and each decomposition corresponds to a counterexample as explained below. (We also remark that the original example of [13], although not explicitly stated in the paper, corresponds to the decomposition $K = K \cdot I$, where I denotes the identity matrix.)

Forgetting about mod 3 calculations and regarding L and T as real matrices we see that $K' := \frac{1}{3}L \cdot T$ is a log-Hadamard matrix. Therefore

the set $T \in \mathbb{Z}^4$ above is 3-spectral, but obviously not a 3-tile because the number of elements of T (i.e. 6) does not divide 3^4 . Therefore, for large n , the set $S := T + 3[0, n]^4$ is $3n$ -spectral, but not a tile in view of Propositions 2.1 and 2.5.

This means that the set $\Omega := \cup_{s \in S} \{s + [0, 1]^4\}$ is spectral in \mathbb{R}^4 (this fact was already mentioned in the discussion preceding Proposition 2.1). The fact that Ω does not tile \mathbb{R}^4 follows as in Proposition 2.5. Namely, in a large cube C_l we find a point x which is not covered by any translate of the annulus A , and note that any translate of Ω is either disjoint from the cube $x + [0, 3]^4$ or it covers exactly 6 unit volumes of it. Therefore $x + [0, 3]^4$ cannot be covered by translates of Ω because of obvious divisibility reasons.

□

REFERENCES

- [1] B. Fuglede, *Commuting self-adjoint partial differential operators and a group theoretic problem*, J. Funct. Anal. **16**(1974), 101-121.
- [2] B. Fuglede, *Orthogonal exponentials on the ball*, Expo. Math. **19**(2001), 267-272.
- [3] A. Iosevich, S. Pedersen, *Spectral and tiling properties of the unit cube*, Inter. Math. Res. Notices **16**(1998), 819-828.
- [4] A. Iosevich, N. Katz, T. Tao, *Convex bodies with a point of curvature do not have Fourier bases*, Amer. J. Math. **123**(2001), 115-120.
- [5] A. Iosevich, N. Katz, T. Tao, *The Fuglede conjecture holds for convex planar domains*, Math. Res. Letters, **10**(2003), 559-570.
- [6] A. Iosevich, M. Rudnev, *A combinatorial approach to orthogonal exponentials*, Inter. Math. Res. Notices **49**(2003)1-12.
- [7] M. Kolountzakis, *Non-symmetric convex domains have no basis of exponentials*, Illinois J. Math. **44**(2000), 542-550.
- [8] M. Kolountzakis, M. Papadimitrakakis, *A class of non-convex polytopes which admit no orthogonal basis of exponentials*, Illinois J. Math. **46**(2002), 4, 1227-1232.
- [9] M. Kolountzakis, *The study of translational tiling with Fourier analysis*, Proceedings of the Milano Conference on Fourier Analysis and Convexity, to appear.
- [10] I. Laba, *Fuglede's conjecture for a union of two intervals*, Proc. Amer. Math. Soc. **129**(2001), 2965-2972.
- [11] I. Laba, *The spectral set conjecture and multiplicative properties of roots of polynomials*, J. London Math. Soc. **65**(2002), 661-671.
- [12] J.C. Lagarias, P.W. Shor, Y. Wang, *Orthonormal bases for exponentials for the n -cube*, Duke Math. J. **103**(2000), 1, 25-37.
- [13] T. Tao, *Fuglede's conjecture is false in 5 and higher dimensions*, Math. Res. Letters, to appear.

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES POB 127 H-1364 BUDAPEST, HUNGARY TEL: (+361) 483-8302, FAX: (+361) 483-8333

E-mail address: matomate@renyi.hu