

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/115410>

Please be advised that this information was generated on 2017-12-05 and may be subject to change.



## Never-depressed females with a family history of depression demonstrate affective bias

Iris van Oostrom<sup>a,b,\*</sup>, Barbara Franke<sup>a,b,c</sup>, Alejandro Arias Vasquez<sup>a,b,c</sup>, Mike Rinck<sup>d</sup>,  
Indira Tendolkar<sup>a,b,e</sup>, Maaike Verhagen<sup>a,d</sup>, Annemarie van der Meij<sup>a</sup>, Jan K. Buitelaar<sup>b,f</sup>,  
Joost G.E. Janzing<sup>a</sup>

<sup>a</sup> Department of Psychiatry, Radboud University Nijmegen Medical Centre, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands

<sup>b</sup> Donders Institute for Brain, Cognition and Behavior, Radboud University Nijmegen Medical Centre, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands

<sup>c</sup> Department of Human Genetics, Radboud University Nijmegen Medical Centre, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands

<sup>d</sup> Behavioural Science Institute, Radboud University Nijmegen, P.O. Box 9104, 6500 HE Nijmegen, The Netherlands

<sup>e</sup> LVR Klinikum Essen, Faculty of Medicine, University of Essen-Duisburg, Virchowstrasse 174, 45147 Essen, Germany

<sup>f</sup> Department of Cognitive Neuroscience, Radboud University Nijmegen Medical Centre, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands

### ARTICLE INFO

#### Article history:

Received 23 February 2012

Received in revised form

9 July 2012

Accepted 5 August 2012

#### Keywords:

Emotional Stroop

Endophenotype

Affective processing bias

Major depressive disorder

Family history

Neuropsychology

### ABSTRACT

According to cognitive theories of depression, individuals susceptible to depression attend selectively to negative information. The purpose of the study was to examine if such an affective processing bias is present in never-depressed individuals with a family history of major depressive disorder (MDD). Formerly depressed female patients having at least one first-degree relative with a history of MDD ( $n=23$ ), their never-depressed female siblings ( $n=21$ ) and never-depressed female controls ( $n=21$ ) performed a conventional and an emotional Stroop task using negative, positive and neutral words. A significant effect was found of group on negative processing bias; post hoc comparisons indicated that never-depressed siblings showed a larger negative processing bias than never-depressed controls. No significant differences were observed in positive bias or conventional interference between the three groups. Our findings suggest that never-depressed females with a family history of depression, like depressed patients, have more difficulties to inhibit negative material and to direct their attention towards task-specific material. This adds to the existing evidence that affective processing bias is a trait characteristic that contributes to the onset of depression and that could be a useful endophenotype for MDD.

© 2012 Elsevier Ireland Ltd. Open access under the [Elsevier OA license](http://creativecommons.org/licenses/by/3.0/).

### 1. Introduction

Major depressive disorder (MDD) causes significant impairment and results in a high disease burden and increased mortality risk. The genetic contribution to MDD is estimated to be in range of 31–42 % (Sullivan et al., 2000). Family members of depressed patients have a two-fold increased risk to develop MDD (Levinson, 2005). The search for specific genes for depression has, however, remained largely unsuccessful, potentially due to the high heterogeneity of MDD and to complex gene–environment interactions in the etiology (Cannon and Keller, 2006). The intermediate phenotype approach (Gottesman and Gould, 2003) promises more successful genetic analyses by studying apparently simpler and less heterogeneous constructs that are linked to clinically observable mental

disorders. Besides facilitating genetic analysis, endophenotypes can improve our understanding of the etiology of psychiatric disorders. Gottesman and Gould (2003) stated that endophenotypes for psychiatric disorders must fulfill several criteria in order to be useful. Endophenotypes should be heritable traits that are associated with the disorder, should co-segregate with the disorder within families and should be found in non-affected family members at a higher rate than in the general population.

Biased information processing in attention, memory and interpretation is proposed to be one of the central cognitive dysfunctions found in MDD (Gotlib and Joormann, 2010) and fulfills several of the endophenotype criteria formulated by Gottesman and Gould (2003). Recent cognitive theories of depression posit that depression is caused and maintained by affective processing bias and by deficits in cognitive control when processing negative information (Gotlib and Joormann, 2010). Reduced cognitive control and depressogenic schemas are assumed to engender impaired attentional inhibitory control over negative elaborative processes, such as rumination

\* Corresponding author at: Department of Psychiatry, Radboud University Nijmegen Medical Centre, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands. Tel.: +31 24 361 35 13; fax: +31 24 366 82 83.

E-mail address: [i.vanoostrom@psy.umcn.nl](mailto:i.vanoostrom@psy.umcn.nl) (I. van Oostrom).

(De Raedt and Koster, 2010). Due to this impaired inhibitory control, depressed patients attend selectively to negative information and also store and retrieve more negative information in memory, resulting in negative thinking, rumination and low mood. Affective processing bias has indeed been frequently demonstrated in depressed patients (Beck, 2008) and in remitted patients (Teasdale and Dent, 1987). Evidence is now emerging that affective processing bias may be heritable, since it has been associated with several variants in candidate genes for depression, like the 5-HTTLPR, COMT Val108/158Met (Hayden et al., 2008; Williams et al., 2010; Thomason et al., 2010; Perez-Edgar et al., 2010) and BDNF Val66Met (Van Oostrom et al., 2012). Affective processing bias is probably a trait characteristic; it has been observed in patients in remission (Bhagwagar et al., 2004; Joormann and Gotlib, 2007; Ramel et al., 2007) as well as in highly neurotic, never-depressed individuals (Chan et al., 2007), who are at increased risk of depression. There is however also evidence that affective processing bias is activated by mood state and stress and is also state-dependent (Bower, 1981).

To date, knowledge about the presence of affective processing bias in never-depressed family members of MDD patients is limited. Affective processing bias has been studied in never-depressed children of depressed parents (Jaenicke et al., 1987; Joormann et al., 2007; Taylor and Ingram, 1999). These studies have generally reported biased information processing in high risk children, but not always (Mannie et al., 2007). Children of depressed parents also demonstrate increased amygdala and nucleus accumbens activation to fearful faces during unconstrained viewing of emotional expressions on faces (Monk et al., 2008). Children have however not yet reached the peak age of onset in MDD, that is between 20 and 50 years of age (Sadock and Sadock, 2007). Although studying affective processing bias in adult first degree relatives of MDD patients is relevant to determine whether affective bias can be considered a valid endophenotype for depression, it has only been studied in adult first degree relatives of depressed patients in two fMRI studies only (Amico et al., 2012; Lisiecka et al., 2012). First degree relatives demonstrated activation of the right Heschl's gyrus during performance in an emotional dot probe task as well as increased activation of the right middle cingulate cortex and left caudate nucleus during inhibition of negative pictures. To the best of our knowledge, no behavioral studies have been conducted to date.

This study aimed at examining affective processing bias in remitted depressed patients with a family history of depression, in their non-affected female siblings and in never-depressed controls selected from the general population. Only female participants were included in view of the female preponderance in depression that may suggest sex-specific etiologic pathways (Kendler et al., 2002). We hypothesized affective processing bias to be present in remitted depressed patients and never-depressed female siblings at a higher rate than in a matched control group. Adopting the emotional Stroop paradigm to assess affective processing bias, we expected remitted depressed patients and non-affected female siblings to show longer color naming latencies when naming the color of negative words than neutral or positive words (Williams et al., 1996). Furthermore, we expected affective processing bias and depression to co-segregate within families and hypothesized that remitted depressed patients would demonstrate stronger affective processing bias, and thus longer color naming latencies when naming the color of negative words, as compared to their non-affected female siblings. Finally, we expected no interference effects in general selective attention in remitted depressed patients or their female siblings using the traditional Stroop paradigm, since studies using neuropsychological tasks have found little empirical support for cognitive deficits in remitted depressed patients (Gotlib and Joormann, 2010).

## 2. Methods

### 2.1. Participants

Women aged 18–65 years from European Caucasian origin were invited to participate in a clinical study of the genetics of MDD, GenMood (Verhagen et al., 2008), through psychiatric treatment settings and advertisements. All participants were screened before inclusion for current and lifetime diagnosis of MDD and other psychiatric diagnoses and for family history of depression.

Formerly depressed female patients ( $n=23$ ) had a history of MDD, at least one first-degree relative with a history of formerly diagnosed and treated MDD, MDD in remission at enrollment and no history of alcohol abuse, manic or psychotic episodes. Never-depressed female siblings ( $n=21$ ) were sisters of the participants in the patient group, without either a current or lifetime history of depression. Never-depressed siblings who participated in the study thus had a sibling and another first degree relative with a lifetime history of MDD. Unaffected female controls ( $n=21$ ) had no personal or family history of current or lifetime psychiatric disorder. They were matched for age and education to resemble the patient and family member groups as closely as possible. The study was approved by the Dutch central medical ethics review board.

### 2.2. Materials

#### 2.2.1. Diagnostic instruments

The Dutch version of the Composite International Diagnostic Interview (CIDI), version 2.1, was used to examine depressive and other psychiatric diagnoses. The following CIDI sections were used: lifetime presence of mood disorders, anxiety disorders, schizophrenia and other non-affective psychotic disorders, alcohol abuse and dependence. The family history of depression was examined using an adapted version of the Family Interview for Genetic Studies (Maxwell, 1992) and the Family History Research Diagnostic Criteria (Endicott et al., 1975) following the method described for the GenMood project (Verhagen et al., 2008).

#### 2.2.2. Stroop tasks

The *emotional Stroop task* (Williams et al., 1996) was administered to assess processing bias for negative and positive information. Words were generated with negative, positive and neutral content and were rated for valence by psychology students. Subsequently, 10 words per affective category were selected based on the valence ratings and matched for word frequency and length (see Table 1). The task consisted of three blocks with 100 presentations each. One block contained the selected negative words, one block the positive words and one block the neutral words. Words appeared one by one in the middle of a computer screen for 1997 ms. Participants were asked to name the ink color of the words as fast as possible by speaking it into a microphone. Reaction times (RTs) were registered using computer software (Voogd, 2005). The experimenter used a button box to register incorrect or invalid answers (like "uhh").

The *conventional Stroop task* was used to assess impairments in attention, especially cognitive interference from automatic responses, and consisted of a computerized version of the original Stroop (1935) task. The task consisted of three blocks with 100 items each, that appeared one by one in the middle of a computer screen for 1997 ms. The first block consisted of color words printed in black ink; participants were asked to read aloud the words as fast as possible into a microphone. The second block consisted of colored bars; the participants were asked to name the colors as fast as possible. The third block consisted of color words printed in inconsistent colors; participants were asked to name the color of the ink as fast as possible, ignoring the meaning of the words. Reaction times were registered and the experimenter registered incorrect and invalid responses.

### 2.3. Statistical analysis

Invalid and incorrect responses were removed from the data. To minimize the influence of outliers, we also deleted the RTs that were longer than two standard deviations from the mean. Negative and positive affective processing bias scores were calculated using the difference in RTs between neutral and negative or neutral and positive blocks, respectively. Higher scores indicated greater negative or positive affective processing bias. General interference was calculated as the main outcome variable of the cognitive Stroop task, by subtracting the mean RT from the inconsistent word color condition from the colored-bars condition. Normality of distribution of the affective processing bias variables was examined using a Kolmogorov–Smirnov test and yielded non-significant results, indicating a normal distribution of the bias scores. Participants made very few errors (< 1%).

The differences with regard to negative and positive processing bias and general interference between remitted patients, never-depressed female siblings and controls were analyzed using one-way ANOVAs with group as the between-subjects factor. To account for potential familial clustering between patients and siblings, the analyses were repeated using Generalized Estimating Equations with a linear regression model, family identity link and robust estimators. Negative bias

**Table 1**  
Experimental stimulus words used in the emotional Stroop task.

Condition		
Negative	Neutral	Positive
Ruminate	Can	Laugh
Aversion	License number	Colorful
Pathetic	Fridge	Cheerful
Bad	Cabinet	Funny
Miserable	Shop	Joy
Lousy	Route	Happy
Melancholy	Shoe	Healthy
Sad	Duchess	Lovely
Failed	Warehouse	Pleasant
Depressing	Pudding	Optimist

was included as dependent variable and group (patient, sibling, control) as predictor. We used an alpha level of  $P=0.05$  for all statistical tests.

### 3. Results

#### 3.1. Participant characteristics

Participants originated from 42 different pedigrees. Demographic and clinical data and stroop mean reaction times are shown in Table 2. Oneway ANOVA's were conducted to analyze differences in age and educational level between remitted patients, never-depressed female siblings and controls. The three groups did not differ significantly with regard to age ( $F(2,65)=0.80$ ,  $P=0.46$ ), or education ( $F(2,65)=0.16$ ,  $P=0.86$ ).

The majority of the patients was classified with a severe (59%) and recurrent (82%) MDD. Patients reported a mean number of 4.5 (S.D.=6.4) depressive episodes. Mean age at onset was 24.1 years (S.D.=9.4 years). Current pharmacological treatment of the patients included one or more SSRIs (52%, 12/23), other antidepressants (22%, 5/23) or no treatment (26%, 6/23). One patient had lithium addition (4%, 1/23) and two used prescribed benzodiazepines (9%, 2/23). The never-depressed female siblings and the controls took no antidepressants or other psychotropic medication; except for one sibling and one control participant taking benzodiazepines. Medicated patients did not demonstrate a significant larger negative or positive bias compared to unmedicated patients ( $T=-0.94$ ,  $P=0.36$  and  $T=-1.13$ ,  $P=0.27$ , respectively).

#### 3.2. Emotional Stroop task

An analysis of variance showed that there was a significant effect of group on negative processing bias ( $F(2,65)=3.42$ ,  $P=0.04$ ). The results are presented in Fig. 1. Post-hoc comparisons using the Tukey HSD test demonstrated that the mean score of the never-depressed female siblings ( $M=11$ , S.D.=5) was significantly different from the controls' mean score ( $M=-7$ , S.D.=5,  $P=0.03$ ), indicating that never-depressed siblings showed a larger negative processing bias than never-depressed controls. The remitted patients ( $M=3$ , S.D.=5) did not differ significantly from the never-depressed female siblings, nor from the controls ( $P=0.44$  and  $P=0.34$ , respectively) and demonstrated level of negative processing bias in between the controls and the siblings. When adjusting for familial clustering using Generalized Estimating Equations, the differences in negative processing bias between the patients, siblings and controls remained significant ( $P=0.02$ ).

A second analysis of variance showed that the three participant groups did not differ significantly with regard to positive processing bias ( $F(2,65)=1.05$ ,  $P=0.36$ ).

**Table 2**  
Demographic data and Stroop mean reaction times in ms in remitted female patients, unaffected female siblings and female controls.

	Patients <i>n</i> =23	Unaffected siblings <i>n</i> =22	Controls <i>n</i> =21
<b>Age</b>			
Mean (S.D.)	45.2 (10.5)	46.1 (9.4)	42.1 (12.5)
<b>Education (1–7)<sup>a</sup></b>			
Mean (S.D.)	5.2 (0.7)	5.1 (0.8)	5.1 (0.8)
<b>Emotional Stroop (ms)</b>			
Negative condition			
Mean (S.D.)	652 (61)	663 (56)	616 (71)
Neutral condition			
Mean (S.D.)	649 (55)	652 (58)	623 (68)
Positive condition			
Mean (S.D.)	653 (55)	656 (57)	623 (66)
<b>Conventional Stroop (ms)</b>			
Word Reading			
Mean (S.D.)	527 (66)	520 (65)	491 (55)
Color Bars			
Mean (S.D.)	577 (57)	589 (73)	537 (54)
Inconsistent color words			
Mean (S.D.)	796 (124)	810 (124)	782 (148)

<sup>a</sup> Education was coded following the Verhage system (1–4 lower education, 5 high school, 6–7 higher education).


#### 3.3. Conventional Stroop task

No significant differences in general interference were found between the three participant groups ( $F(2,65)=0.44$ ,  $P=0.64$ ).

### 4. Discussion

Negative affective processing bias is proposed to be the central cognitive dysfunction found in MDD (Gotlib and Joormann, 2010; Beck, 2008). In order to extend our understanding of the cognitive factors that affect the onset of depression, we examined for the first time the presence of affective processing bias in never-depressed adult females with a family history of MDD. The results indicate that never-depressed females with a familial risk of depression demonstrated negative affective processing bias, in contrast to never-depressed controls without a family history of depression. Never-depressed females with increased familial risk showed more negative affective processing bias and had more difficulties directing their attention towards task-specific features of the negative material as compared to never-depressed controls. When we adjusted the analysis for family relatedness, the results remained significant. Analyzing within group differences yielded similar results. Our findings are consistent with findings from studies in children of mothers with MDD. At risk children generally have been found to display stronger negative biases than children with no family history of depression (Jaenicke et al., 1987; Joormann et al., 2007; Taylor and Ingram, 1999). Attentional inhibitory dysfunction of negative material after recovery from depression has been reported frequently in both behavioral (Joormann and Gotlib, 2007; Fritzsche et al., 2010; Dai and Feng, 2009) and neurophysiological responses (Dai et al., 2011; Ramel et al., 2007). This dysfunctional inhibition of negative information may be a trait characteristic that interferes with the processing of neutral information and that may increase the risk of developing MDD.

Another finding is that formerly depressed and never-depressed females with a family history of depression did not demonstrate significantly more interference in non-emotional attentional processes than never-depressed controls. These findings suggest that the affective interference effects were specific to negative information and could not be attributed to a general interference effect.



**Fig. 1.** Mean negative affective processing bias in remitted depressed female patients, never-depressed female siblings and never-depressed female controls using the emotional Stroop task (ms, 95%CI),  $F(2,65)=3.27$ ,  $P=0.04$ . A higher negative processing bias score indicates a greater bias towards negative words.

Biased attention to negative information may reflect a cognitive susceptibility factor for MDD, while a general lack of executive attentional control may become apparent during depressive episodes only as a state marker of the disease. This concurs with the current viewpoint that deficits in cognitive control in depressed patients mainly concern the processing of affective information (Gotlib and Joormann, 2010), and not neutral information.

Affective processing bias was expected to be larger in patients than in never-depressed female siblings. Contrary to these expectations, remitted patients did not show significantly greater affective processing biases than their never-depressed female siblings. The majority of the patients took antidepressant drugs at the time of testing. Recent studies report that serotonin manipulations affect affective processing bias (Merens et al., 2007) and that antidepressants may act by modifying specific neural dysfunctions correlated to negative cognitive biases (Harmer et al., 2009; Di Simplicio et al., 2011). The relatively mild affective processing bias in patients observed in this study may therefore be due to treatment effects, although medicated patient did not demonstrate less negative bias compared to unmedicated patients. Another unexpected finding was the lack of significant differences between patients, female siblings and controls with regard to positive bias. This result adds to the mixed empirical results regarding positive attentional and memory bias in patients with MDD (De Raedt and Koster, 2010). Probably, negative bias is a more substantial characteristic of MDD compared to positive bias. Indeed, negative processing bias predicted changes in anxiety and dysphoria in non-depressed students in the following semester (Osinsky et al., 2012) as well as the magnitude of the physiological response to stress up to 8 months follow up in a group of never-depressed students (Fox et al., 2010). Furthermore, negative processing bias predicted higher depressive symptom scores in soldiers with war zone stress exposure (Beavers et al., 2011).

A limitation of the current study is the cross-sectional experimental design that does not enable us to establish causal relations between affective processing bias and MDD, or to differentiate between genetic and environmental contributions to affective processing bias. Twin studies or adoption studies are better able to reveal the nature of these vulnerability factors. However, the advantage of studying the siblings instead of the children of depressed patients is that siblings are not necessarily exposed to the environmental pathogen of having a depressed parent. Family discord is more prevalent in families with a depressed parent (Nomura et al., 2002; Pilowsky et al., 2006; Hammen et al., 2004) and depressed women have been shown to have more conflictual

interactions with their children (Hammen and Brennan, 2002; Lovejoy et al., 2000). Another limitation of the study is the small sample size and subsequent limited power to detect differences between the groups, especially when adjusting for family relatedness. A strength of our study is that we included only women in order to have a homogeneous group. Different pathways to depression may exist in men and women (Kendler et al., 2002; Kendler et al., 2006) and the pathways to deficient affective information processing may differ between men and women. For example, a significant interaction effect of *BDNF* Val66Met with childhood stressful life events was found on affective memory bias in never-depressed males, while in never-depressed females *BDNF* Val66Met variant did not seem to mediate the cognitive susceptibility to depression (Van Oostrom et al., 2012). A limitation of this approach is however that it remains unclear whether our findings can be generalized to men.

In conclusion, this study is the first to demonstrate the presence of affective processing bias in never-depressed siblings with a family history of depression. Never-depressed female individuals with a family history of depression demonstrated more negative affective processing bias than never-depressed controls. Inadequate affective information processing may render unaffected family members cognitively susceptible to depression. New developments, like affective processing bias modification, are promising interventions that may also be beneficial to the first-degree relatives of depressed patients.

## Acknowledgments

We are very grateful to the patients and their family members who participated in this study. The authors would also like to thank Armand van Oosterwijck and Nienke Jansen for testing and data management.

## References

- Amico, F., Carballedo, A., Lisecka, D., Fagan, A.J., Boyle, G., Frodl, T., 2012. Functional anomalies in healthy individuals with a first degree family history of major depressive disorder. *Biology of Mood & Anxiety Disorders* 2, 1.
- Beck, A.T., 2008. The evolution of the cognitive model of depression and its neurobiological correlates. *American Journal of Psychiatry* 165, 969–977.
- Beavers, C.G., Lee, H.J., Wells, T.T., Ellis, A.J., Telch, M.J., 2011. Association of predeployment gaze bias for emotion stimuli with later symptoms of PTSD and depression in soldiers deployed in Iraq. *American Journal of Psychiatry* 168, 735–741.
- Bhagwagar, Z., Cowen, P.J., Goodwin, G.M., Harmer, C.J., 2004. Normalization of enhanced fear recognition by acute SSRI treatment in subjects with a previous history of depression. *American Journal of Psychiatry* 161, 166–168.
- Bower, G.H., 1981. Mood and memory. *The American Psychologist* 36, 129–148.
- Cannon, T.D., Keller, M.C., 2006. Endophenotypes in the genetic analyses of mental disorders. *Annual Review of Clinical Psychology* 2, 267–290.
- Chan, S.W., Goodwin, G.M., Harmer, C.J., 2007. Highly neurotic never-depressed students have negative biases in information processing. *Psychological Medicine* 37, 1281–1291.
- Dai, Q., Feng, Z., 2009. Deficient inhibition of return for emotional faces in depression. *Progress in Neuropsychopharmacology and Biological Psychiatry* 33, 921–932.
- Dai, Q., Feng, Z., Koster, E.H., 2011. Deficient distracter inhibition and enhanced facilitation for emotional stimuli in depression: an ERP study. *International Journal of Psychophysiology* 79, 249–258.
- De Raedt, R., Koster, E.H., 2010. Understanding vulnerability for depression from a cognitive neuroscience perspective: A reappraisal of attentional factors and a new conceptual framework. *Cognitive, Affective & Behavioral Neuroscience* 10, 50–70.
- Di Simplicio, M., Norbury, R., Harmer, C.J., 2011. Short-term antidepressant administration reduces negative self-referential processing in the medial prefrontal cortex in subjects at risk for depression. *Molecular Psychiatry*, 1–8.
- Endicott, J., Andreasen, N.C., Spitzer, R.L., 1975. Family History Research Diagnostic Criteria. Biometrics Research, New York State Psychiatric Institute.
- Fox, E., Cahill, S., Zougkou, K., 2010. Preconscious processing biases predict emotional reactivity to stress. *Biological Psychiatry* 67, 371–377.
- Fritzsche, A., Dahme, B., Gotlib, I.H., Joormann, J., Magnussen, H., Watz, H., Nutzinger, D.O., von Leupoldt, A., 2010. Specificity of cognitive biases in

- patients with current depression and remitted depression and in patients with asthma. *Psychological Medicine* 40, 815–826.
- Gotlib, I.H., Joormann, J., 2010. Cognition and depression: current status and future directions. *Annual Review of Clinical Psychology* 6, 285–312.
- Gottesman, I.I., Gould, T.D., 2003. The endophenotype concept in psychiatry: etymology and strategic intentions. *American Journal of Psychiatry* 160, 636–645.
- Hammen, C., Brennan, P.A., 2002. Interpersonal dysfunction in depressed women: impairments independent of depressive symptoms. *Journal of Affective Disorders* 72, 145–156.
- Hammen, C., Brennan, P.A., Shih, J.H., 2004. Family discord and stress predictors of depression and other disorders in adolescent children of depressed and nondepressed women. *Journal of the American Academy for Child & Adolescent Psychiatry* 43, 994–1002.
- Harmer, C.J., O'Sullivan, U., Favaron, E., Massey-Chase, R., Ayres, R., Reinecke, A., Goodwin, G.M., Cowen, P.J., 2009. Effect of acute antidepressant administration on negative affective bias in depressed patients. *American Journal of Psychiatry* 166, 1178–1184.
- Hayden, E.P., Dougherty, L.R., Maloney, B., Olino, T.M., Sheikh, H., Durbin, C.E., Nurnberger Jr., J.I., Lohr, D.K., Klein, D.N., 2008. Early-emerging cognitive vulnerability to depression and the serotonin transporter promoter region polymorphism. *Journal of Affective Disorders* 107, 227–230.
- Jaenicke, C., Hammen, C., Zupan, B., Hiroto, D., Gordon, D., Adrian, C., Burge, D., 1987. Cognitive vulnerability in children at risk for depression. *Journal of Abnormal Child Psychology* 15, 559–572.
- Joormann, J., Gotlib, I.H., 2007. Selective attention to emotional faces following recovery from depression. *Journal of Abnormal Psychology* 116, 80–85.
- Joormann, J., Talbot, L., Gotlib, I.H., 2007. Biased processing of emotional information in girls at risk for depression. *Journal of Abnormal Psychology* 116, 135–143.
- Kendler, K.S., Gardner, C.O., Prescott, C.A., 2002. Toward a comprehensive developmental model for major depression in women. *American Journal of Psychiatry* 159, 1133–1145.
- Kendler, K.S., Gardner, C.O., Prescott, C.A., 2006. Toward a comprehensive developmental model for major depression in men. *American Journal of Psychiatry* 163, 115–124.
- Levinson, D.F., 2005. Meta-analysis in Psychiatric Genetics. *Current Psychiatry Reports* 7, 143–151.
- Lisiecka, D.M., Carballedo, A., Fagan, A.J., Connolly, G., Meaney, J., Frodl, T., 2012. Altered inhibition of negative emotions in subjects at family risk of major depressive disorder. *Journal of Psychiatric Research* 46, 181–188.
- Lovejoy, M.C., Graczyk, P.A., O'Hare, E., Neuman, G., 2000. Maternal depression and parenting behavior: a meta-analytic review. *Clinical Psychology Review* 20, 561–592.
- Mannie, Z.N., Bristow, G.C., Harmer, C.J., Cowen, P.J., 2007. Impaired emotional categorisation in young people at increased familial risk of depression. *Neuropsychologia* 45, 2975–2980.
- Maxwell, M.E., 1992. *Manual for the FIGS (Family Interview for Genetic Studies)*. National Institute of Mental Health.
- Merens, W., Willem Van der Does, A.J., Spinhoven, P., 2007. The effects of serotonin manipulations on emotional information processing and mood. *Journal of Affective Disorders* 103, 43–62.
- Monk, C.S., Klein, R.G., Telzer, E.H., Schroth, E.A., Mannuzza, S., Moulton III, J.L., Guardino, M., Masten, C.L., Clure-Tone, E.B., Fromm, S., Blair, R.J., Pine, D.S., Ernst, M., 2008. Amygdala and nucleus accumbens activation to emotional facial expressions in children and adolescents at risk for major depression. *American Journal of Psychiatry* 165, 90–98.
- Nomura, Y., Wickramaratne, P.J., Warner, V., Mufson, L., Weissman, M.M., 2002. Family discord, parental depression, and psychopathology in offspring: ten-year follow-up. *Journal of the American Academy for Child & Adolescent Psychiatry* 41, 402–409.
- Osinsky, R., Losch, A., Hennig, J., Alexander, N., MacLeod, C., 2012. Attentional bias to negative information and 5-HTTLPR genotype interactively predict students' emotional reactivity to first university semester. *Emotion* 12, 460–469.
- Perez-Edgar, K., Bar-Haim, Y., McDermott, J.M., Gorodetsky, E., Hodgkinson, C.A., Goldman, D., Ernst, M., Pine, D.S., Fox, N.A., 2010. Variations in the serotonin-transporter gene are associated with attention bias patterns to positive and negative emotion faces. *Biological Psychology* 83, 269–271.
- Pilowsky, D.J., Wickramaratne, P., Nomura, Y., Weissman, M.M., 2006. Family discord, parental depression, and psychopathology in offspring: 20-year follow-up. *Journal of the American Academy for Child & Adolescent Psychiatry* 45, 452–460.
- Ramel, W., Goldin, P.R., Eyler, L.T., Brown, G.G., Gotlib, I.H., McQuaid, J.R., 2007. Amygdala reactivity and mood-congruent memory in individuals at risk for depressive relapse. *Biological Psychiatry* 61, 231–239.
- Sadock, B.J., Sadock, V.A., 2007. *Kaplan & Sadock's Synopsis of Psychiatry*, 10th ed. Wolters Kluwer, Philadelphia.
- Stroop, J.R., 1935. Studies of interference in serial verbal reactions. *Journal of Experimental Psychology* 18, 643–661.
- Sullivan, P.F., Neale, M.C., Kendler, K.S., 2000. Genetic epidemiology of major depression: review and meta-analysis. *American Journal of Psychiatry* 157, 1552–1562.
- Taylor, L., Ingram, R.E., 1999. Cognitive reactivity and depressotypic information processing in children of depressed mothers. *Journal of Abnormal Psychology* 108, 202–210.
- Teasdale, J.D., Dent, J., 1987. Cognitive vulnerability to depression: an investigation of two hypotheses. *British Journal of Clinical Psychology* 26 (Pt 2), 113–126.
- Thomason, M.E., Henry, M.L., Paul, H.J., Joormann, J., Pine, D.S., Ernst, M., Goldman, D., Mogg, K., Bradley, B.P., Britton, J.C., Lindstrom, K.M., Monk, C.S., Sankin, L.S., Louro, H.M., Gotlib, I.H., 2010. Neural and behavioral responses to threatening emotion faces in children as a function of the short allele of the serotonin transporter gene. *Biological Psychology* 85, 38–44.
- Van Oostrom, I., Franke, B., Rijpkema, M., Gerritsen, L., Arias-Vasquez, A., Fernandez, G., Tendolkar, I., 2012. Interaction between BDNF Val66Met and childhood stressful life events is associated to affective memory bias in men but not women. *Biological Psychology* 89, 214–219.
- Verhagen, M., van der, M.A., Franke, B., Vollebergh, W.A., de, G.R., Buitelaar, J.K., Janzing, J.G., 2008. Familiality of major depressive disorder and patterns of lifetime comorbidity. The NEMESIS and GenMood studies. A comparison of three samples. *European Archives of Psychiatry and Clinical Neuroscience* 258, 505–512.
- Voogd, H.F.J.M., 2005. *Stroop State Machine*. Radboud University Nijmegen.
- Williams, J.M., Mathews, A., MacLeod, C., 1996. The emotional Stroop task and psychopathology. *Psychological Bulletin* 120, 3–24.
- Williams, L.M., Gatt, J.M., Grieve, S.M., Dobson-Stone, C., Paul, R.H., Gordon, E., Schofield, P.R., 2010. COMT Val(108/158)Met polymorphism effects on emotional brain function and negativity bias. *Neuroimage* 53, 918–925.