

Technical University of Denmark



3D cardiac Chemical Shift Imaging of [1-13C] hyperpolarized acetate and pyruvate in pigs

Luca, Menichetti; Francesca, Frijia; Alessandra, Flori; Vincenzo, Lionetti; Matteo, Liserani; Giulio, Giovannetti; Giacomo, Bianchi; L, Romano Simone; Vincenzo, Positano; Ardenkjær-Larsen, Jan Henrik; F, Schulte Rolf; A, Recchia Fabio; Luigi, Landini; Maria, Santarelli; Massimo, Lombardi

Published in:
Journal of Cardiovascular Magnetic Resonance

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Luca, M., Francesca, F., Alessandra, F., Vincenzo, L., Matteo, L., Giulio, G., ... Massimo, L. (2013). 3D cardiac Chemical Shift Imaging of [1-13C] hyperpolarized acetate and pyruvate in pigs. *Journal of Cardiovascular Magnetic Resonance*, 15(Suppl 1), P10.

DTU Library
Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

POSTER PRESENTATION

Open Access

3D cardiac Chemical Shift Imaging of [1-13C] hyperpolarized acetate and pyruvate in pigs

Luca Menichetti^{1,2}, Francesca Frijia^{2*}, Alessandra Flori³, Vincenzo Lionetti³, Matteo Liserani⁴, Giulio Giovannetti¹, Giacomo Bianchi², Simone L Romano³, Vincenzo Positano^{2,1}, Jan Henrik Ardenkjaer-Larsen^{5,6}, Rolf F Schulte⁷, Fabio A Recchia², Luigi Landini⁸, Maria Filomena Santarelli^{1,2}, Massimo Lombardi²

From 16th Annual SCMR Scientific Sessions
San Francisco, CA, USA. 31 January - 3 February 2013

Background

¹³C Dynamic Nuclear Polarization (DNP) with rapid dissolution together with Magnetic Resonance Chemical Shift Imaging (CSI) have been used for non-invasive real-time metabolic assessment in cardiac experimental models on a clinical 3T scanner. Here, we report an in vivo comparison of hyperpolarized [1-¹³C] pyruvate and [1-¹³C] acetate perfusion and metabolism: a method based on a 3D Spiral CSI sequence is presented for obtaining spatially and spectrally-resolved information on whole heart cardiac metabolism.

Methods

In this work hyperpolarized [1-¹³C] pyruvate and [1-¹³C] acetate were injected in vivo to obtain spatially and spectrally resolved information of basal metabolism on whole heart in middle size animal models. Five healthy male farm pigs (38±2 kg) were studied in basal condition and subjected to imaging experiments performed on a 3T GE Signa HDx scanner using a ¹³C-quadrature birdcage coil (Rapid Biomedical). An HyperSense DNP polarizer (Oxford Inst.) was employed for the studies: a procedure for the hyperpolarization and dissolution of a large dose of TRIS-[1-¹³C]acetate water/glycerol mixture was set up while the preparation of a large dose of [1-¹³C] pyruvic acid was performed as recently published by this group. An anatomical region of interest covering the whole heart was first acquired with a proton reference scan and the metabolic information was then obtained using 3D IDEAL spiral CSI on the same region. Image re-slicing along cardiac short axis (SA) views and image fusion of ¹³C

metabolite maps and anatomical 1H reference images were performed by PMOD software.

Results

A graph of the γ -variate and mono-exponential fitting of hyperpolarized [1-¹³C] acetate myocardial spectroscopic signals is reported in Figure 1 while a representative map in SA orientation through the heart is shown in Figure 2: [1-¹³C] acetate is extracted inside the heart and clearly detected in the heart-chambers and myocardial wall. Representative maps of spatial distribution of [1-¹³C] bicarbonate, [1-¹³C] lactate and [1-¹³C] pyruvate in SA orientation through the heart are also produced using hyperpolarized [1-¹³C] pyruvate.

Conclusions

A comparison between acetate and pyruvate ¹³C-mapping has been realised as far as we know for the first time in pigs with this experimental approach. This ongoing study demonstrates the feasibility of whole-heart ¹³C-cardiac metabolic imaging in pigs for detecting and mapping cardiac metabolism in basal condition with hyperpolarized [1-¹³C]acetate in comparison with [1-¹³C] pyruvate.

This study is the first step towards the optimization of the [1-¹³C] acetate concentration and the acquisition sequence parameters to ensure suitable MR signals in myocardial tissue and to study its metabolic fate.

Funding

Self funding.

²Fondazione G. Monasterio CNR-Regione Toscana, Pisa, Italy
Full list of author information is available at the end of the article




Figure 1 ^{13}C dynamic spectra were acquired using a slice selective pulse-and-acquire sequence (bandwidth 5000 Hz, 2048 pts, 10° FA). A long-axis slice of 20 mm was selected during excitation. Spectra were acquired from the beginning of the injection of the hyperpolarized [^{13}C] acetate, every 2 s, for 120 s. Diagrammatic representation of the γ -variate and mono-exponential fitting of cardiac spectroscopic signal to obtain rate constants (N=4).




Figure 2 Representative maps in SA view of the heart showing the in vivo spatial distribution of hyperpolarized TRIS-[^{13}C] acetate in pigs; spectroscopic data were normalized to the maximum value of signal amplitude.

Author details

¹Institute of Clinical Physiology, National Research Council, Pisa, Italy. ²Fondazione G. Monasterio CNR-Regione Toscana, Pisa, Italy. ³Scuola Superiore Sant'Anna, Pisa, Italy. ⁴Faculty of Physics, University of Pisa, Pisa, Italy. ⁵GE Healthcare, Hillerod, Denmark. ⁶Department of Electrical Engineering, Technical University of Denmark, Kongens Lyngby, Denmark. ⁷GE Global Research, Munich, Germany. ⁸Department of Information Engineering, University of Pisa, Pisa, Italy.

doi:10.1186/1532-429X-15-S1-P10

Cite this article as: Menichetti *et al.*: 3D cardiac Chemical Shift Imaging of [^{13}C] hyperpolarized acetate and pyruvate in pigs. *Journal of Cardiovascular Magnetic Resonance* 2013 **15**(Suppl 1):P10.