

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

ENDORSE
Environmental determinants of overweight in Rotterdam schoolchildren

Klazine van der Horst - Nachtegaal

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

Colofon

ISBN: 978-90-8559-569-4

Copyright © 2009 Klazine van der Horst - Nachtegaal

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the author or the copyright-owning journals for previously published chapters.

Lay-out and Print: Optima Grafische Communicatie

Cover-illustration: Optima Grafische Communicatie

This thesis was printed with financial support of the Department of Public Health, Erasmus MC, Rotterdam.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

ENDORSE
Environmental Determinants of Overweight in Rotterdam Schoolchildren

ENDORSE
Omgevingsdeterminanten van overgewicht bij Rotterdamse scholieren

Proefschrift

ter verkrijging van de graad van doctor aan de
Erasmus Universiteit Rotterdam
op gezag van de rector magnificus

Prof.dr. H.G. Schmidt

en volgens besluit van het College voor Promoties.

De openbare verdediging zal plaatsvinden op
woensdag 7 oktober 2009 om 11.30 uur
door

Klazine-Anja van der Horst - Nachtegaal

geboren te Zwolle



1 **PROMOTIECOMMISSIE**

2

3 Promotor: Prof. dr. ir. J. Brug

4

5 Overige leden: Prof. dr. J.P. Mackenbach

6 Prof. dr. M.C.H. Donker

7 Prof. dr. W. van Mechelen

8

9 Co-promotor: Dr. A. Oenema

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

1	CONTENTS	
2		
3	Part I - Introduction to the ENDORSE project	7
4	1. General Introduction	9
5	2. The ENDORSE study: research into environmental determinants of obesity	27
6	related behaviors in Rotterdam schoolchildren	
7		
8	Part II - Environmental correlates of energy balance-related behaviors:	45
9	reviews of the literature	
10	3. A review of environmental correlates of obesity-related dietary behaviors in	47
11	youth	
12	4. Environmental correlates of physical activity in youth – A review and	79
13	update	
14		
15	Part III – Socio-demographic correlates of energy balance-related behaviors	125
16	5. Gender, ethnic and educational differences in overweight and energy balance-	127
17	related behaviors among Dutch adolescents	
18	6. Socio-demographic factors as correlates of active commuting to school in	143
19	Rotterdam, the Netherlands	
20		
21	Part IV - Individual and environmental correlates of energy balance-related	157
22	behaviors	
23	7. The school food environment: associations with adolescent soft drink and	159
24	snack consumption	
25	8. Do individual cognitions mediate the association of socio-cultural and	175
26	physical environmental factors with adolescent sports participation?	
27	9. Perceived parenting style and practices and the consumption of sugar-	193
28	sweetened beverages by adolescents	
29		
30	10. General discussion	209
31	Summary	233
32	Samenvatting	237
33	Dankwoord	241
34	Curriculum Vitae	243
35	List of publications	245
36	PhD Portfolio	247
37		
38		
39		

Part I Introduction to the ENDORSE project

1 General Introduction

1.1 INTRODUCTION

Overweight and obesity are a major threat to public health as the prevalence of overweight and obesity are rising worldwide in all age groups. Obesity in children and adolescents is of particular interest since it persists into adulthood and is associated with severe health consequences. Therefore, the prevention of overweight and obesity is important for public health. To be able to develop theory and evidence-based interventions aimed at the prevention of excess weight gain, it is essential to identify which specific energy intake and energy expenditure behaviors contribute most to excess weight gain, and which determinants are associated with engagement in such behaviors.

This thesis reports on a number of studies on the identification of individual and environmental correlates of behaviors related to the energy balance (i.e. energy intake and expenditure behaviors). These studies were part of the ENDORSE project (ENvironmental Determinants of Obesity in Rotterdam SchoolchildrEn) which was initiated to contribute to systematic, evidence based research on individual and environmental determinants of overweight and obesity. The ENDORSE project was initially a cross-sectional study and after the first data collection a follow-up data collection was funded. Data were collected at baseline (2005/2006) and two years later (2007/2008) in a cohort of adolescents aged 12-15 years. The studies in this thesis were based on the baseline data collection, as the longitudinal data was not available in time to use in this thesis. This introductory chapter describes the background, aims and theoretical framework used in the ENDORSE project and presents an overview of the individual studies that are part of this thesis.

1.2 A MODEL FOR PLANNED HEALTH PROMOTION

Overweight and obesity prevention has to target the most important risk factors and the underlying determinants. Prevention of overweight and obesity through the promotion of healthy dietary habits and a physically active lifestyle by means of health education and health promotion should therefore be carefully planned. To increase the likelihood of intervention success, careful evidence-based planning of obesity prevention interventions should be a standard procedure. The use of health promotion planning models helps to improve the quality of interventions and in these models five important steps can be distinguished (Figure 1.1) [1]. The first two steps in this model for planned health education and promotion cover the epidemiological analysis. These first two steps should (I) identify important threats to public health and (II) the risk factors including risk behaviors for these public health threats. The result is a set of priorities for preventive interventions, health change goals and specific target groups for interventions. The third step identifies important and changeable determinants of risk behaviors. In step 4 of the model, intervention strate-




Figure 1.1 A model for planned health education and promotion [1]

gies, methods and materials need to be selected and/or developed that address the most important and changeable determinants identified in the third step. In the final step the interventions should be implemented and disseminated in such a way that the target population is reached and exposed to the intervention. Each step should be based on established theory and sufficient scientific evidence.

The ENDORSE project was initiated to contribute to systematic, evidence based research on individual and environmental determinants of overweight and obesity and focused mainly on step 2 and 3 of the model for planned health education and promotion (Figure 1.1).

1.3 ANALYSIS OF HEALTH AND QUALITY OF LIFE: OBESITY

Obesity is a vast and growing public health problem as the prevalence of overweight and obesity is rising worldwide in all age groups [2, 3]. Obesity may become the most important determinant of preventable diseases within the foreseeable future [4]. Overweight and obesity often manifest early in life [5, 6] and are associated with an increased risk of serious diseases during childhood and adolescence [7, 8]. Obesity during childhood causes a clustering of cardiovascular disease risk factors such as hypertension and dyslipidaemia. Other important complications of childhood obesity are type 2 diabetes, musculoskeletal and pulmonary disorders and obesity is further associated with psychosocial problems such as a low self esteem, depression and eating disorders (Table 1.1) [9, 10]. Furthermore, obese

1 children and adolescents are likely to become obese adults, who have an increased risk for
2 various chronic diseases and premature death [9, 11].

3
4 In the Netherlands, an increasing proportion of adolescents are classified as overweight
5 and obese (Table 1.2 & 1.3). Between 2002 and 2004, in 12 to 15 year old adolescents the
6 prevalence of overweight ranged between 15.2% and 20.1% and between 2.7% and 4.7% for
7 obesity [12-14]. Youth from Moroccan and Turkish backgrounds have the highest preva-
8 lence rates of overweight [12, 15]. In Rotterdam, overweight and obesity prevalence is higher
9 compared to the general Dutch adolescent population. Especially youth in Rotterdam from
10 a Turkish background show very high prevalence rates of overweight and obesity with per-
11 centages ranging between 38% and 45% [16]. The complications that obese adolescents may
12 develop, the tracking of obesity into adulthood, and the vulnerability of obese adolescents
13 make a strong case for the prevention and treatment of overweight and obesity in youth.
14 However, treatment of overweight and obesity in adolescents is difficult as adolescents have
15 less autonomy over food and physical activity behaviors compared to adults and they are
16 more susceptible to peer pressure. The overall success of existing family and school-based
17 interventions has been disappointing [17, 18]. Only few adolescents succeed to maintain
18 their lower body weight and most of the weight loss is often regained within a few years [9].
19 Adolescents are therefore an important target group for intervention activities that prevent
20 them from gaining excess weight and becoming overweight or obese.

21
22 **Table 1.1** Complications of childhood obesity [9, 10]

23 Complications of childhood obesity	24 Examples
25 Psychosocial problems	Poor self esteem
26	Depression
27	Eating disorders
28 Pulmonary problems	Sleep apnoea
29	Asthma
30	Exercise intolerance
31 Gastrointestinal problems	Gall stones
32 Endocrine problems	Type 2 diabetes
33 Cardiovascular problems	Dyslipidaemia
34	Hypertension
35 Neurological problems	Pseudo tumor cerebri: headache, vision abnormalities
36 Renal problems	Glomerulosclerosis
37 Musculoskeletal problems	Flat feet
38	Low back pain
39	

Table 1.2 Prevalence of overweight in the Netherlands (adolescents 12-15 years old) [13, 14, 19]

Overweight (%)						
	Boys			Girls		
Age	1980	1997	2002-2004	1980	1997	2002-2004
12	3.4	7.1	16.2	6.1	9.0	17.1
13	3.6	7.1	15.3	6.0	9.1	15.2
14	3.9	7.3	15.6	6.1	9.1	16.2
15	4.2	7.7	16.8	6.2	9.4	20.1

Table 1.3 Prevalence of obesity in the Netherlands (adolescents 12-15 years old) [13, 14, 19]

Obesity (%)						
	Boys			Girls		
Age	1980	1997	2002-2004	1980	1997	2002-2004
12	0.2	0.7	2.8	0.4	1.1	3.1
13	0.2	0.7	2.8	0.4	1.0	2.7
14	0.2	0.7	3.4	0.4	1.0	2.8
15	0.2	0.7	3.9	0.4	1.1	4.7

1.4 ANALYSIS OF BEHAVIORAL RISK FACTORS FOR OBESITY

A long-term positive energy balance in which energy input through food intake exceeds energy expenditure through physical activity eventually causes obesity. Therefore, high energy intake and low physical activity can be identified as important risk behaviors for overweight and obesity. Prevention of overweight and obesity can be achieved by lowering energy intake and/or increasing energy expenditure. The increase in overweight and obesity is therefore largely related to behavioral factors that can be referred to as energy balance-related behaviors. However, it is not yet very clear which specific risk behaviors are related to overweight and obesity in children and adolescents [20]. Recent overviews have suggested a range of energy balance-related behaviors that may contribute substantially to a higher risk for unnecessary weight gain, such as high intake of energy-dense, micronutrient poor foods and a sedentary lifestyle, and behaviors that may contribute to a lower risk for weight gain such as physical activity and high fiber intake [21-23]. However, studies and reviews have also reported inconsistent results on the role of specific dietary and physical activity sub-behaviors [22]. An overview of the available evidence is given in Table 1.4. Most evidence in this overview is based on systematic reviews of observational and intervention studies.

Table 1.4 Overview of the available evidence on risk behaviors for overweight and obesity

Behaviors	Evidence	References
<i>Dietary behaviors</i>		
Snacks / fast food intake	Probable / insufficient for children	[26, 27]
Sugar-sweetened beverage intake	Convincing	[23, 28]
Breakfast consumption	Possible	[29]
Fruit / vegetable	Insufficient	[30, 31]
Fiber intake / Non-starch polysaccharide	Convincing	[23, 32-34]
Intake of dairy products	Possible	[23, 35, 36]
Portion sizes	Possible	[23, 37]
<i>Physical activity behaviors</i>		
Overall physical activity	Probable	[38, 39]
Leisure time physical activity (sports, walking, cycling)	Insufficient	[38]
Physical education	Insufficient	[40-42]
Active transport to school	Insufficient / no relationship	[43-45]
<i>Sedentary behaviors</i>		
Television viewing	Convincing (small effect)	[21, 25]
Computer use	No relationship	[25]

With respect to food intake, convincing evidence for an association with overweight and obesity exists for sugar-sweetened beverage consumption and fiber intake. The evidence for associations with overweight and obesity are less clear for other behaviors such as breakfast consumption, portion sizes and consumption of dairy products and fast food or snack consumption (Table 1.4). Snacking, fast food intake and large portion sizes have been found to be associated with energy and fat intake, but none of these factors have been found to be consistently related to obesity [20].

Furthermore, the relative importance of different aspects of physical activity is poorly understood. It is unclear whether obesity is similarly related to a reduction in physical activity behaviors and/or an increase in sedentary behaviors [22, 24]. A small but significant association has been found between television viewing and body fatness among children and adolescents [21, 25], while evidence for specific physical activity sub-behaviors such as active transport and leisure time sports, such as walking and bicycling is lacking.

In summary, the available evidence on behavioral risk factors for overweight shows that for many of the dietary and physical activity sub-behaviors the evidence is insufficient. Therefore, more and stronger studies are needed to examine the specific risk behaviors for

1 overweight and obesity in the dietary and physical activity domain, such as leisure time
2 walking and bicycling, sports and snack consumption. It has, however, been established that
3 a combination of changes in both energy balance behaviors, i.e. diet and physical activity
4 seems the most promising strategy for successful obesity prevention.

5 To curb the obesity epidemic, it is important to identify specific target groups of ado-
6 lescents who are more at risk of becoming obese by engaging in more (or specific) energy
7 balance-related behaviors. Being able to distinguish specific target groups provides the
8 opportunity to better tailor interventions to the needs and perceptions of those most at risk
9 [46]. Currently, there is insufficient insight into the occurrence of a number of overweight
10 related risk behaviors among adolescents and also whether it is possible to distinguish
11 specific subgroups that are more likely to engage in such specific risk behaviors.

14 1.5 ANALYSIS OF DETERMINANTS OF ENERGY BALANCE-RELATED 15 BEHAVIORS

17 Individual determinants

19 In determinant research the emphasis has been primarily on individual (intrapersonal)
20 cognitive determinants of behavior such as attitudes, perceived behavioral control, subjec-
21 tive norms and intentions, informed by social cognition models such as the Social Cogni-
22 tive Theory [47] and the Theory of Planned Behavior (TPB) [48]. According to the TPB,
23 if people evaluate the behavior as positive (attitude), if they think significant others want
24 them to perform the behavior (subjective norm), and if people are convinced that they can
25 successfully execute the behavior required to produce the desired outcomes (perceived be-
26 havioral control), they will be more likely to have a high intention (motivation) and a higher
27 likelihood of engaging in the behavior (Figure 1.2). The TPB has been found to be useful for
28 the identification of potential determinants of energy balance-related behaviors such as soft
29 drink consumption [49-51], snack consumption [52] total physical activity and physical ex-
30 ercise [53, 54]. On average 30% of the variance in health behaviors can be explained with the
31 TPB variables [53, 54]. The TPB is useful in examining intrapersonal determinants of obesity
32 related behaviors. However, individuals interact with people in their environment such as
33 parents and peers and the individual behavior takes place in environmental settings such
34 as the home, the neighborhood or school. Therefore, next to intrapersonal factors, other
35 factors might also be important for examining determinants of obesity related behaviors.




Figure 1.2 Theory of Planned Behavior [28]

Environmental determinants

An important development in overweight and obesity prevention research has been the recognition of the role of “the environment” in influencing health and health behavior and there has been a shift in focus from individual influences on health behaviors to environmental influences [55]. Socio-ecological models include environmental factors as an important element. The environment can be defined as everything and anything outside the person [56]. The environment is the context in which human beings act and interact. Socio-ecological models focus on how individual, intrapersonal and environmental factors interact and shape health behavior. Socio-ecological models have a complex and multidimensional nature as they describe the environment not only in terms of physical and social components, but also in terms of the objective (actual) or subjective (perceived) attributes of the environment [57].

A useful framework for the classification of environmental determinants is the Analysis Grid for Environments Linked to Obesity (ANGELO) (figure 1.3) [58]. This framework was specifically developed to conceptualize obesogenic environments, and enables the identification of potential intervention settings and strategies.

According to the ANGELO framework, environmental determinants can be classified according to two environmental levels (micro and macro) and four environmental types (physical, socio-cultural, economic, and political). Individuals interact with the environment in various micro or local environments such as schools, homes, workplaces and neighborhoods. Broader macro environments such as health and education systems, food industry, media and the government influence these micro environmental settings. Different types of the environment can be distinguished, such as the *physical* environment, which refers to

1 which foods and physical activity opportunities are available, the *economic* environment,
 2 which refers to the costs of food and activity, the *political* environment, which refers to the
 3 rules related to food and activity (e.g. laws and regulations), and the *socio-cultural* environ-
 4 ment which refers to social interactions, norms, beliefs and values in a community related
 5 to food and activity [58]. These types of environments have representations at the micro-
 6 level and on the macro-level. An example of a physical environmental factor at a micro
 7 level is the availability of physical activity equipment at home, while an urban or suburban
 8 neighborhood setting and infrastructure is an example of a physical environmental factor
 9 at the macro level.

10 As more and more studies focus on environmental determinants of health behaviors it is
 11 important to get a clear overview of the evidence these studies have provided so far, the gaps
 12 that exist in the available literature and the possibilities to improve the research in this field.

Types \ Levels	Micro-environment	Macro-environment
Physical environment		
Economic environment		
Political environment		
Socio-cultural environment		

13
14
15
16
17
18
19
20
21 **Figure 1.3** Analysis Grid for Environments Linked to Obesity [38]

22 23 Other determinants of energy balance-related behaviors

24
25 Next to cognitive and environmental determinants there are also other possible deter-
 26 minants that may influence energy balance-related behaviors such as habit strength and
 27 demographic factors. Research is needed to explore the importance of these determinants
 28 for various energy balance-related behaviors and the working mechanisms of these possible
 29 determinants.

30 Most energy balance-related behaviors occur regularly in daily life, such as the consump-
 31 tion of breakfast and walking for transportation. Therefore, such behaviors might be more
 32 or less automatic behaviors not requiring much or any cognitive efforts. If a behavior is
 33 often repeated, it might become a habit, i.e. an automatic response to a certain environ-
 34 mental cue [59]. Recent studies indicated that habit strength seems to be a useful variable to
 35 incorporate in studies on correlates of energy balance-related behaviors [60-63].

36 Demographic factors such as gender, age, educational level and ethnicity are also associ-
 37 ated with energy balance-related behaviors [21, 24, 64-66] and can be considered as more
 38 distal or upstream determinants. These variables can act also as moderators of the envi-
 39 ronment – behavior relationship as the determinants of energy balance-related behaviors

1 might vary by population sub-groups. Studies are needed that explore whether potential
2 determinants have a different impact on energy balance-related behaviors with respect to
3 gender, ethnicity, age and educational level or socio-economic status [67].
4

5 Mediating and moderating effects of individual and environmental determinants 6

7 Well-developed ecological models specify not only *that* different types of variables (indi-
8 vidual and environmental) interact, but also the working mechanisms of these variables,
9 thus *how* they interact. Kremers and colleagues proposed in their Environmental Research
10 framework for weight Gain prevention (EnRG framework) that environmental factors can
11 have a direct and an indirect influence on behavior [56, 67] (Figure 1.4).

12 The direct influence reflects an automatic, “mindless” process of the environment on
13 behaviors. For instance, dietary behaviors and the amount of foods eaten are strongly in-
14 fluenced by factors such as portion size, food visibility and the ease of obtaining foods [68].
15 The indirect influence of environments on energy balance-related behaviors is through a
16 more cognitive process in which the individual factors, such as attitudes, subjective norms
17 and perceived behavior control play a role. For example, environments that offer appealing
18 opportunities to eat unhealthy foods may result in positive attitudes regarding the con-
19 sumption of these unhealthy foods, resulting in higher intakes of these foods. In the EnRG
20 framework it is also proposed that the direct and indirect pathways can be influenced by
21 moderating factors or effect modifiers; the level of cognitive mediation or direct environ-
22 mental influence is expected to differ according to personal and behavioral attributes such
23 as habit strength, demographic factors and personality [67, 69, 70].
24

25 Only few studies have examined the relative importance of environmental determinants and
26 individual (cognitive) determinants and there is also lack of empirical evidence regarding
27 the influence of environmental determinants on energy balance-related behaviors among
28 adolescents [67, 70-72]. More research is needed on environmental determinants of en-
29 ergy balance-related behaviors in school, neighborhood and home settings with preferably
30 stronger study designs in which mediating and moderating effects can also be examined.
31

32 In the ENDORSE study, the TPB and the ANGELO framework which are both incorpo-
33 rated in the EnRG framework were used as a theoretical framework. To inform obesity
34 prevention interventions for adolescents, the ENDORSE study focused mainly on physical
35 and socio-cultural environmental factors in micro settings such as the school, home and
36 neighborhood.
37
38
39




Figure 1.4 Environmental Research framework for weight Gain prevention [47]

1.6 OUTLINE OF THE THESIS

As stated before, the ENDORSE project was initiated to contribute to systematic, evidence based research on individual and environmental correlates of overweight and obesity. The ENDORSE project focused on the analysis of risk behaviors for overweight and the analysis of determinants of these risk behaviors, step 2 and 3 of the model for planned health education and promotion (Figure 1.1). This resulted in specific recommendations for obesity prevention interventions among adolescents. The specific aims of the ENDORSE project were:

1. to identify which presumed energy balance-related behaviors are associated with overweight and obesity;
2. to examine important individual and environmental correlates of presumed energy balance-related behaviors;
3. to investigate the associations with and the interactions between these correlates and energy balance-related behaviors;
4. to formulate objectives to be targeted in interventions aimed at the prevention of overweight in adolescents aged 12-15 years.

1 The studies presented in this thesis focus on aim 2 and 3 of the ENDORSE project, and the
2 central research questions that will be addressed are:

- 3
4 I. What are important individual and environmental correlates of energy balance-related
5 behaviors?
6 II. To what extent is the association between environmental factors and energy balance-
7 related behaviors mediated by individual cognitions?
8

9 The thesis is divided in three parts. The first part of this thesis gives an introduction to the
10 ENDORSE study, with this general introduction as the first chapter. Chapter 2 presents the
11 study protocol of the ENDORSE study in which the design and methods are described, as
12 well as the results of the pilot work on the identification of risk behaviors for overweight
13 and obesity.
14

15 In the second part of this thesis environmental correlates of energy balance-related behav-
16 iors are identified by studying the existing literature. Two systematic reviews of the litera-
17 ture were conducted, one for dietary behaviors (Chapter 3) and one for physical activity
18 (Chapter 4). The review for dietary behaviors is an original review, whereas the review for
19 physical activity is an update of an existing review conducted by Sallis and colleagues [64].
20

21 In the third part of this thesis, demographic factors as correlates of energy balance-related
22 behaviors are studied. Chapter 5 presents gender, ethnic and educational differences in
23 overweight and energy balance-related behaviors. Chapter 6 describes the results of a study
24 on socio-demographic correlates of active commuting to school. Although such socio-
25 demographic factors are not easily modifiable and are therefore not easy access points for
26 intervention development, these factors can be important to identify specific target groups
27 for obesity prevention interventions.
28

29 In the fourth part of the thesis, the associations of individual and environmental corre-
30 lates with energy balance-related behaviors, and possible mediation through individual
31 correlates from the Theory of Planned Behavior are examined, with the EnRG framework
32 as the theoretical framework. In Chapters 7-9 the potentially important mediation role of
33 cognitions in the association between environmental factors and energy balance-related
34 behaviors is investigated. The study in Chapter 9 is based on another dataset to examine the
35 mediating role of cognitions in further detail with other variables. This study used data from
36 the Dutch Obesity Intervention in Teenagers Study [73].

37 In the general discussion a summary of the main findings of this thesis and recommenda-
38 tions for further research and practice are provided.
39

1 REFERENCES

- 2 1. Brug J, Oenema A, Ferreira I: Theory, evidence and Intervention Mapping to improve behavior nutri-
3 tion and physical activity interventions. *Int J Behav Nutr Phys Act* 2005, 2(1):2.
- 4 2. Lobstein T, Frelut ML: Prevalence of overweight among children in Europe. *Obes Rev* 2003, 4(4):195-
5 200.
- 6 3. Hedley AA, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM: Prevalence of overweight
7 and obesity among US children, adolescents, and adults, 1999-2002. *JAMA* 2004, 291(23):2847-2850.
- 8 4. Peeters A, Barendregt JJ, Willekens F, Mackenbach JP, Al Mamun A, Bonneux L, Nedcom tNEaD-
9 CoMRG: Obesity in adulthood and its consequences for life expectancy: a life-table analysis. *Ann
10 Intern Med* 2003, 138(1):24-32.
- 11 5. Power C, Lake JK, Cole TJ: Measurement and long-term health risks of child and adolescent fatness.
12 *Int J Obes Relat Metab Disord* 1997, 21(7):507-526.
- 13 6. Must A, Jacques PF, Dallal GE, Bajema CJ, Dietz WH: Long-term morbidity and mortality of over-
14 weight adolescents. A follow-up of the Harvard Growth Study of 1922 to 1935. *N Engl J Med* 1992,
15 327(19):1350-1355.
- 16 7. Must A, Strauss RS: Risks and consequences of childhood and adolescent obesity. *Int J Obes Relat
17 Metab Disord* 1999, 23 Suppl 2:S2-11.
- 18 8. Wabitsch M: Overweight and obesity in European children: definition and diagnostic procedures,
19 risk factors and consequences for later health outcome. *Eur J Pediatr* 2000, 159 Suppl 1:S8-13.
- 20 9. Ebbeling CB, Pawlak DB, Ludwig DS: Childhood obesity: public-health crisis, common sense cure.
21 *Lancet* 2002, 360(9331):473-482.
- 22 10. Speiser PW, Rudolf MC, Anhalt H, Camacho-Hubner C, Chiarelli F, Eliakim A, Freemark M, Gruters
23 A, HersHKovitz E, Iughetti L *et al*: Childhood obesity. *J Clin Endocrinol Metab* 2005, 90(3):1871-1887.
- 24 11. Singh AS, Mulder C, Twisk JW, van Mechelen W, Chinapaw MJ: Tracking of childhood overweight
25 into adulthood: a systematic review of the literature. *Obes Rev* 2008.
- 26 12. Schokker DF, Visscher TL, Nooyens AC, van Baak MA, Seidell JC: Prevalence of overweight and
27 obesity in the Netherlands. *Obes Rev* 2007, 8(2):101-108.
- 28 13. van den Hurk K, van Dommelen P, de Wilde JA, Verkerk PH, van Buuren S, Hirasings RA: Prevalentie
29 van overgewicht en obesitas bij jeugdigen 4-15 jaar in de periode 2002-2004. TNO Kwaliteit van
30 Leven; 2006.
- 31 14. van den Hurk K, van Dommelen P, van Buuren S, Verkerk PH, Hirasings RA: Prevalence of overweight
32 and obesity in the Netherlands in 2003 compared to 1980 and 1997. *Arch Dis Child* 2007, 92(11):992-
33 995.
- 34 15. Fredriks AM, Van Buuren S, Sing RA, Wit JM, Verloove-Vanhorick SP: Alarming prevalences of
35 overweight and obesity for children of Turkish, Moroccan and Dutch origin in The Netherlands
36 according to international standards. *Acta Paediatr* 2005, 94(4):496-498.
- 37 16. Rapportage gemeente Rotterdam 2008 [[http://www.jeugdmonitorrotterdam.nl/Rotterdam/Internet/
38 Overig/JMR/pdf/Gemeenterapport%20JMR%20maart%202008.pdf](http://www.jeugdmonitorrotterdam.nl/Rotterdam/Internet/Overig/JMR/pdf/Gemeenterapport%20JMR%20maart%202008.pdf)]
- 39 17. Shaya FT, Flores D, Gbarayor CM, Wang J: School-based obesity interventions: a literature review. *J
School health* 2008, 78(4):189-196.
18. Livingstone MB, McCaffrey TA, Rennie KL: Childhood obesity prevention studies: lessons learned
and to be learned. *Public Health Nutr* 2006, 9(8A):1121-1129.
19. Hirasings RA, Fredriks AM, van Buuren S, Verloove-Vanhorick SP, Wit JM: [Increased prevalence
of overweight and obesity in Dutch children, and the detection of overweight and obesity using
international criteria and new reference diagrams]. *Ned Tijdschr Geneesk* 2001, 145(27):1303-1308.

- 1 20. Moreno LA, Rodriguez G: Dietary risk factors for development of childhood obesity. *Curr Opin Clin Nutr Metab Care* 2007, 10(3):336-341.
- 2 21. Gorely T, Marshall SJ, Biddle SJ: Couch kids: correlates of television viewing among youth. *Int J Behav Med* 2004, 11(3):152-163.
- 3 22. Rennie KL, Johnson L, Jebb SA: Behavioural determinants of obesity. *Best Pract Res Clin Endocrinol Metab* 2005, 19(3):343-358.
- 4 23. Swinburn BA, Caterson I, Seidell JC, James WP: Diet, nutrition and the prevention of excess weight gain and obesity. *Public Health Nutr* 2004, 7(1A):123-146.
- 5 24. Van Der Horst K, Paw MJ, Twisk JW, Van Mechelen W: A brief review on correlates of physical activity and sedentariness in youth. *Med Sci Sports Exerc* 2007, 39(8):1241-1250.
- 6 25. Marshall SJ, Biddle SJ, Gorely T, Cameron N, Murdey I: Relationships between media use, body fatness and physical activity in children and youth: a meta-analysis. *Int J Obes Relat Metab Disord* 2004.
- 7 26. Rosenheck R: Fast food consumption and increased caloric intake: a systematic review of a trajectory towards weight gain and obesity risk. *Obes Rev* 2008.
- 8 27. Pereira MA, Kartashov AI, Ebbeling CB, Van Horn L, Slattery ML, Jacobs DR, Jr., Ludwig DS: Fast-food habits, weight gain, and insulin resistance (the CARDIA study): 15-year prospective analysis. *Lancet* 2005, 365(9453):36-42.
- 9 28. Malik VS, Schulze MB, Hu FB: Intake of sugar-sweetened beverages and weight gain: a systematic review. *Am J Clin Nutr* 2006, 84(2):274-288.
- 10 29. Rampersaud GC, Pereira MA, Girard BL, Adams J, Metzler JD: Breakfast habits, nutritional status, body weight, and academic performance in children and adolescents. *J Am Diet Assoc* 2005, 105(5):743-760; quiz 761-742.
- 11 30. te Velde SJ, Twisk JW, Brug J: Tracking of fruit and vegetable consumption from adolescence into adulthood and its longitudinal association with overweight. *Br J Nutr* 2007, 98(2):431-438.
- 12 31. Tohill BC, Seymour J, Serdula M, Kettel-Khan L, Rolls BJ: What epidemiologic studies tell us about the relationship between fruit and vegetable consumption and body weight. *Nutr Rev* 2004, 62(10):365-374.
- 13 32. Pereira MA, Ludwig DS: Dietary fiber and body-weight regulation. Observations and mechanisms. *Pediatr Clin North Am* 2001, 48(4):969-980.
- 14 33. Ludwig DS, Pereira MA, Kroenke CH, Hilner JE, Van Horn L, Slattery ML, Jacobs DR, Jr.: Dietary fiber, weight gain, and cardiovascular disease risk factors in young adults. *JAMA* 1999, 282(16):1539-1546.
- 15 34. Howarth NC, Saltzman E, Roberts SB: Dietary fiber and weight regulation. *Nutr Rev* 2001, 59(5):129-139.
- 16 35. Barba G, Russo P: Dairy foods, dietary calcium and obesity: a short review of the evidence. *Nutr Metab Cardiovasc Dis* 2006, 16(6):445-451.
- 17 36. Huang TT, McCrory MA: Dairy intake, obesity, and metabolic health in children and adolescents: knowledge and gaps. *Nutr Rev* 2005, 63(3):71-80.
- 18 37. Fisher JO, Kral TV: Super-size me: Portion size effects on young children's eating. *Physiol Behav* 2008, 94(1):39-47.
- 19 38. Wareham NJ, van Sluijs EM, Ekelund U: Physical activity and obesity prevention: a review of the current evidence. *Proc Nutr Soc* 2005, 64(2):229-247.
- 20 39. Connelly JB, Duaso MJ, Butler G: A systematic review of controlled trials of interventions to prevent childhood obesity and overweight: a realistic synthesis of the evidence. *Public health* 2007, 121(7):510-517.
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39

- 1 40. Kahn EB, Ramsey LT, Brownson RC, Heath GW, Howze EH, Powell KE, Stone EJ, Rajab MW, Corso
2 P: The effectiveness of interventions to increase physical activity. A systematic review. *Am J Prev Med*
3 2002, 22(4 Suppl):73-107.
- 4 41. Cleland V, Dwyer T, Blizzard L, Venn A: The provision of compulsory school physical activity: as-
5 sociations with physical activity, fitness and overweight in childhood and twenty years later. *Int J*
6 *Behav Nutr Phys Act* 2008, 5(1):14.
- 7 42. Doak CM, Visscher TL, Renders CM, Seidell JC: The prevention of overweight and obesity in chil-
8 dren and adolescents: a review of interventions and programmes. *Obes Rev* 2006, 7(1):111-136.
- 9 43. Heelan KA, Donnelly JE, Jacobsen DJ, Mayo MS, Washburn R, Greene L: Active commuting to and
10 from school and BMI in elementary school children-preliminary data. *Child Care Health Dev* 2005,
11 31(3):341-349.
- 12 44. Landsberg B, Plachta-Danielzik S, Much D, Johannsen M, Lange D, Muller MJ: Associations between
13 active commuting to school, fat mass and lifestyle factors in adolescents: the Kiel Obesity Prevention
14 Study (KOPS). *Eur J Clin Nutr* 2008, 62(6):739-47.
- 15 45. Rosenberg DE, Sallis JF, Conway TL, Cain KL, McKenzie TL: Active transportation to school over 2
16 years in relation to weight status and physical activity. *Obesity* 2006, 14(10):1771-1776.
- 17 46. Kreuter MW, Lukwago SN, Bucholtz RD, Clark EM, Sanders-Thompson V: Achieving cultural ap-
18 propriateness in health promotion programs: targeted and tailored approaches. *Health Educ Behav*
19 2003, 30(2):133-146.
- 20 47. Bandura A: *Social foundations of thought and action: a social cognitive theory.*: Englewood Cliffs, NJ:
21 Prentice-Hall; 1986.
- 22 48. Ajzen I: *Attitudes, personality, and behavior.*: Homewood, IL, US: Dorsey Press; 1988.
- 23 49. Grimm GC, Harnack L, Story M: Factors associated with soft drink consumption in school-aged
24 children. *J Am Diet Assoc* 2004, 104(8):1244-1249.
- 25 50. Kassem NO, Lee JW: Understanding soft drink consumption among male adolescents using the
26 theory of planned behavior. *J Behav Med* 2004, 27(3):273-296.
- 27 51. Kassem NO, Lee JW, Modeste NN, Johnston PK: Understanding soft drink consumption among
28 female adolescents using the Theory of Planned Behavior. *Health Educ Res* 2003, 18(3):278-291.
- 29 52. de Bruijn GJ, Kremers SP, Schaalma H, van Mechelen W, Brug J: Determinants of adolescent bicycle
30 use for transportation and snacking behavior. *Prev Med* 2005, 40(6):658-667.
- 31 53. Godin G, Kok G: The theory of planned behavior: a review of its applications to health-related
32 behaviors. *Am J Health Promot* 1996, 11(2):87-98.
- 33 54. Hagger MS, Chatzisarantis NL, Biddle SJ: The influence of autonomous and controlling motives on
34 physical activity intentions within the Theory of Planned Behaviour. *Br J Health Psychol* 2002, 7(Part
35 3):283-297.
- 36 55. Booth SL, Sallis JF, Ritenbaugh C, Hill JO, Birch LL, Frank LD, Glanz K, Himmelgreen DA, Mudd
37 M, Popkin BM *et al*: Environmental and societal factors affect food choice and physical activity:
38 rationale, influences, and leverage points. *Nutr Rev* 2001, 59(3 Pt 2):S21-39; discussion S57-65.
- 39 56. Sallis JF, Owen N: Ecological models of health behavior. In *Health behavior and health education.* 3
edition. Edited by Glanz K, Rimer BK, Lewis FM. San Fransisco: Jossey-Bass.
57. Stokols D: Establishing and maintaining healthy environments. Toward a social ecology of health
promotion. *Am Psychol* 1992, 47(1):6-22.
58. Swinburn B, Egger G, Raza F: Dissecting obesogenic environments: the development and application
of a framework for identifying and prioritizing environmental interventions for obesity. *Prev Med*
1999, 29(6 Pt 1):563-570.

- 1 59. Verplanken B, Orbell S: Reflections on past behavior: A self-report index of habit strength. *J Appl Soc Psychol* 2003, 33(6):1313-1330.
- 2 60. Brug J, de Vet E, de Nooijer J, Verplanken B: Predicting fruit consumption: cognitions, intention, and
- 3 habits. *J Nutr Educ Behav* 2006, 38(2):73-81.
- 4 61. Verplanken B, Aarts H, van Knippenberg A, Moonen A: Habit versus planned behaviour: a field
- 5 experiment. *Br J Soc Psychol* 1998, 37 (Pt 1):111-128.
- 6 62. De Bruijn G-J, Kremers SPJ, De Vet E, De Nooijer J, Van Mechelen W, Brug J: Does habit strength
- 7 moderate the intention-behaviour relationship in the Theory of Planned Behaviour? The case of fruit
- 8 consumption. *Psychol Health* 2007, 22(8):899 - 916.
- 9 63. Aarts H, verplanken B, van Knippenberg A: Predicting behavior from actions in the past: Repeated
- 10 decision making or a matter of habit? *J Appl Soc Psychol* 1998, 28:1355-1374.
- 11 64. Sallis JF, Prochaska JJ, Taylor WC: A review of correlates of physical activity of children and adoles-
- 12 cents. *Med Sci Sports Exerc* 2000, 32(5):963-975.
- 13 65. Delva J, O'Malley PM, Johnston LD: Racial/ethnic and socioeconomic status differences in over-
- 14 weight and health-related behaviors among American students: national trends 1986-2003. *J Adolesc*
- 15 *Health* 2006, 39(4):536-545.
- 16 66. te Velde SJ, Wind M, van Lenthe FJ, Klepp KI, Brug J: Differences in fruit and vegetable intake and
- 17 determinants of intakes between children of Dutch origin and non-Western ethnic minority children
- 18 in the Netherlands - a cross sectional study. *Int J Behav Nutr Phys Act* 2006, 3:31.
- 19 67. Kremers SP, De Bruijn GJ, Visscher TL, Van Mechelen W, De Vries NK, Brug J: Environmental influ-
- 20 ences on energy balance-related behaviors: A dual-process view. *Int J Behav Nutr Phys Act* 2006,
- 21 3(1):9.
- 22 68. Cohen D, Farley TA: Eating as an automatic behavior. *Preventing chronic disease* 2008, 5(1):A23.
- 23 69. Kremers SP, de Bruijn GJ, Droomers M, van Lenthe F, Brug J: Moderators of environmental interven-
- 24 tion effects on diet and activity in youth. *Am J Prev Med* 2007, 32(2):163-172.
- 25 70. de Bruijn GJ, Kremers SP, de Vries H, van Mechelen W, Brug J: Associations of social-environmental
- 26 and individual-level factors with adolescent soft drink consumption: results from the SMILE study.
- 27 *Health Educ Res* 2006.
- 28 71. de Bruijn GJ, Kremers SP, Lensvelt-Mulders G, de Vries H, van Mechelen W, Brug J: Modeling in-
- 29 dividual and physical environmental factors with adolescent physical activity. *Am J Prev Med* 2006,
- 30 30(6):507-512.
- 31 72. Brug J, van Lenthe F: *Environmental determinants and interventions for physical activity, nutrition and*
- 32 *smoking: A review*. Zoetermeer: Speed-Print; 2005.
- 33 73. Singh AS, Chin APMJ, Kremers SP, Visscher TL, Brug J, van Mechelen W: Design of the Dutch Obesity
- 34 Intervention in Teenagers (NRG-DOiT): systematic development, implementation and evaluation of
- 35 a school-based intervention aimed at the prevention of excessive weight gain in adolescents. *BMC*
- 36 *Public Health* 2006, 6:304.
- 37
- 38
- 39

2 The ENDORSE study: research into environmental determinants of obesity related behaviors in Rotterdam schoolchildren

van der Horst K, Oenema A, van de Looij-Jansen P, Brug J. The ENDORSE study: research into environmental determinants of obesity related behaviors in Rotterdam schoolchildren.

BMC Public Health 2008, 8: 142.

1 ABSTRACT

2
3 **Background:** Children and adolescents are important target groups for prevention of
4 overweight and obesity as overweight is often developed early in life and tracks into adult-
5 hood. Research into behaviors related to overweight (energy balance-related behaviors) and
6 the personal and environmental determinants of these behaviors is fundamental to inform
7 prevention interventions. In the Netherlands and in other countries systematic research
8 into environmental determinants of energy balance related behaviors in younger adoles-
9 cents is largely lacking. This protocol paper describes the design, the components and the
10 methods of the ENDORSE study (Environmental Determinants of Obesity in Rotterdam
11 SchoolchildrEn), that aims to identify important individual and environmental determi-
12 nants of behaviors related to overweight and obesity and the interactions between these
13 determinants among adolescents.

14 **Methods:** The ENDORSE study is a longitudinal study with a two-year follow-up of a
15 cohort of adolescents aged 12-15 years. Data will be collected at baseline (2005/2006) and
16 at two years follow-up (2007/2008). Outcome measures are body mass index (BMI), waist
17 circumference, time spent in physical activity and sedentary behaviors, and soft drink,
18 snack and breakfast consumption. The ENDORSE study consists of two phases, first em-
19 ploying qualitative research methods to inform the development of a theoretical framework
20 to examine important energy balance related behaviors and their determinants, and to
21 inform questionnaire development. Subsequently, the hypothetical relationships between
22 behavioral determinants, energy balance related behaviors and BMI will be tested in a
23 quantitative study combining school-based surveys and measurements of anthropometrical
24 characteristics at baseline and two-year follow-up.

25 **Discussion:** The ENDORSE project is a comprehensive longitudinal study that enables
26 investigation of specific environmental and individual determinants of overweight and
27 obesity among younger adolescents. The project will result in specific recommendations for
28 obesity prevention interventions among younger adolescents.

29
30
31
32
33
34
35
36
37
38
39

1 BACKGROUND

2
3 Adolescent overweight and obesity are important public health concerns in the Netherlands
4 as well as in other western countries, due to the increasing proportion of adolescents classi-
5 fied as overweight or obese [1]. Children and adolescents are an important target group for
6 intervention activities aimed at the prevention of overweight. Overweight and obesity often
7 manifest early in life [2, 3] and is associated with an increased the risk of serious diseases
8 during childhood and adolescence [4]. Furthermore, obese children and adolescents are
9 likely to become obese adults, who have an increased risk for various chronic diseases
10 and premature death [5]. Therefore, it is important to develop interventions that prevent
11 children and adolescents from gaining excess weight. Prevention of weight gain can best
12 be achieved by focusing on both sides of the energy balance equation; energy intake (diet)
13 and energy expenditure (physical activity). To be able to develop theory and evidence-based
14 interventions aimed at the prevention of excess weight gain, it is essential to identify which
15 specific energy intake and energy expenditure behaviors contribute most to excess weight
16 gain, and which determinants mediate or predict engagement in such behaviors.

17 An important development in overweight and obesity prevention research has been the
18 recognition of the environment as a potentially important determining factor for energy
19 balance related behaviors [6]. Currently, there is only limited scientific evidence regarding
20 the influence of environmental determinants on energy balance related behaviors among
21 adolescents [7, 8]. Few studies have examined the relative importance of environmental
22 determinants and individual (cognitive) determinants that have been the more traditional
23 focus of behavior change interventions, and there is also lack of empirical evidence regard-
24 ing the interactions between these determinants [9-11]. Research into environmental deter-
25 minants is now emerging and more rigorous and well-designed studies are needed to draw
26 stronger inferences for relationships between environmental determinants, energy balance
27 related behaviors and BMI. Such studies are also needed to identify the interactions between
28 determinants of various energy balance related behaviors and the mechanisms underlying
29 the associations between individual and environmental determinants of these behaviors
30 [6, 11]. Therefore, a comprehensive study was designed that examines key energy balance
31 related behaviors, the individual and an environmental determinants of these behaviors
32 and that contains objective measures of height, weight and waist circumference. The target
33 population of the study is adolescents aged 12-15 years. The specific aims of the study are:
34 (i) to identify important behaviors related to overweight (energy balance related behaviors),
35 (ii) to examine important individual (cognitive) and environmental determinants for the
36 energy balance related behaviors identified, (iii) to investigate the associations with and the
37 interactions between these determinants and BMI in cross-sectional and prospective analy-
38 ses with a two year follow-up, and (iv) to formulate objectives to be targeted in interventions
39 aimed at the prevention of overweight in adolescents aged 12-15 years.

1
2 The ENDORSE study (Environmental Determinants of Obesity in Rotterdam's Schoolchild-
3 drEn) is conducted within the Center for Effective Public Health In the larger Rotterdam
4 area (CEPHIR), an established collaboration between a university research center (Erasmus
5 MC) and the Municipal Health Service Organization in the Rotterdam region. The data
6 collection process takes place in close cooperation with the Municipal Health Service Rot-
7 terdam area, using an existing research infrastructure. In this article we describe the design
8 and protocol of the ENDORSE study.

9 The study comprises of two parts. The first part focused on the identification of the key
10 energy balance related behaviors and the important individual and environmental determi-
11 nants to examine. Based on this identification, measurement instruments were developed.
12 The second part consists of a combined cross-sectional and longitudinal study utilizing
13 these instruments.

14 15 16 **PILOT WORK: IDENTIFICATION OF RISK BEHAVIORS AND IMPORTANT** 17 **ENVIRONMENTAL DETERMINANTS**

18
19 This phase of the ENDORSE study involved the development of questionnaires for ado-
20 lescents and parents, interview forms for school representatives and canteen managers, an
21 audit instrument used to observe the school and the neighborhood around schools, and a
22 list of important census data on neighborhood level. To develop these instruments system-
23 atically, important behaviors and determinants were identified in the following steps.

24 25 **Identification of important energy balance related behaviors in youth**

26
27 The most important energy balance related behaviors were identified to gain insight in the
28 contribution of these behaviors to overweight and obesity in adolescents. Based on a review
29 of relevant reviews of the literature, a preliminary list of specific relevant energy balance
30 related behaviors was compiled. This list contained: watching television, computer use,
31 sports, physical education, transport to school, leisure time activities, soft drink consump-
32 tion, skipping breakfast, consumption of foods high in fat, fruit and vegetable consumption,
33 portion sizes and dining out. Subsequently national experts on energy balance related
34 behaviors were asked to review this list and suggest additional important behaviors and
35 score the behaviors on the importance and changeability of each behavior. This procedure
36 resulted in the following identification of behaviors to be examined in the present study:
37 active transport to school, leisure time activities, sports, watching television, computer use,
38 soft drink consumption, sweets/cookies/cake/chocolate bar consumption, savory snack
39 consumption and breakfast consumption.

1 Identification of environmental and individual determinants of energy balance related 2 behaviors among youth

3
4 In the ENDORSE study the environment was defined as ‘anything outside the individual’.
5 The environment can be subdivided by means of distinguishing various environmental
6 factors. A suitable framework for the classification of environmental determinants is the
7 Analysis Grid for Environments Linked to Obesity (ANGELO) [12]. This framework was
8 specifically developed to conceptualize health behavior environments, and enables the
9 identification of potential intervention settings and strategies. According to the ANGELO
10 grid, environmental determinants can be grouped in four environmental types (physical,
11 socio-cultural, economic, and political) and specific environmental levels (micro and
12 macro). To integrate important environmental types and levels in the ENDORSE study;
13 physical, socio-cultural, economic and policy determinants were examined at the micro
14 level (home, school and neighborhood level). Combinations of perceived and objectively
15 measured environmental determinants were used to investigate the interactions between
16 environmental and individual determinants.

17 Environmental determinants previously shown to be important were identified by
18 conducting two systematic reviews, one with physical activity as the outcome behavior [7]
19 and the other with specific obesity related dietary behaviors as outcome [8]. The results
20 from the reviews were categorized using the ANGELO grid. Convincing evidence of an
21 important role for physical environmental determinants was not found. However, only a
22 limited number of studies assessing physical environmental determinants of energy bal-
23 ance related behaviors were retrieved. Most consistent determinants of physical activity in
24 adolescents were support from significant others, mother’s education level, family income
25 and non-vocational school attendance and low neighborhood crime incidence [7]. The
26 most consistent determinants of obesity related dietary behaviors among adolescents were
27 parental and family influences, e.g. parental and sibling intakes, parenting style, family con-
28 nectedness and parental education [8]. The results of the reviews were used to guide the
29 design of questionnaires and observation forms. Since the evidence from the reviews itself
30 was not sufficient, potential determinants of physical activity and dietary behaviors were
31 also included in the measurement instruments.

32 The theory of planned behavior (TPB) was used for the selection of potential individual
33 determinants to be included in the study [13]. The TPB postulates that intention to perform
34 a behavior, the determinant most proximal to behavior, is determined by three conceptu-
35 ally independent constructs: attitude, subjective norms and perceived behavioral control.
36 To further explore what specific concepts, beliefs or perceptions would be important for
37 adolescents; focus group interviews with adolescents were held. A focus group interview
38 is conducted among a small group of people who, led by a moderator and following a pre-
39 determined interview scheme, discuss several topics related to a specific subject. The aim

1 of the focus groups was to gain insight in the individual and environmental determinants
2 of snacking, soft drink consumption, eating breakfast and physical activity. Three schools
3 participated and teachers were asked to select adolescents who would be able to function
4 in a group discussion (i.e. who were not too shy or too dominant). Five focus groups were
5 conducted with seven to nine adolescents aged 13-15 years old, and a total of 39 adolescents
6 participated. Two of these groups consisted of boys only; two of girls only; and one was a
7 mixed group with boys and girls. Three of the groups were composed of adolescents from
8 cultural and ethnic minorities reflecting the cultural diversity of the residents of Rotterdam.
9 Each interview was tape-recorded and lasted about 45 minutes. The focus groups were tran-
10 scribed verbatim and from these transcripts, quotes were categorized into the determinants
11 or concepts that they reflected. Many adolescents identified that seeing other people eating
12 or drinking and smelling fast food were factors that influenced their eating and drinking
13 patterns. These factors can be translated as the concept 'external cues.' Assessment of a ten-
14 dency to respond to external cues was therefore included in the adolescent questionnaire as
15 an individual determinant. Rules at home (e.g. not allowed leaving the house without eating
16 breakfast) or the lack of rules (e.g. allowed to drink as much soft drinks the adolescent
17 wants) were also mentioned by participants. Parental influences already were identified as
18 potential important determinants from the systematic reviews, and based on the results
19 of the focus group interviews, items examining parents' rules or 'parenting practices' were
20 included in the adolescent and parent questionnaires.

23 METHODS

25 Design

27 The ENDORSE study has a cross-sectional and a prospective two year follow-up compo-
28 nent. Data will be collected at baseline (2005/2006) when adolescents aged 12-15 years,
29 and two years later (2007/2008). Outcome measures are body mass index (BMI), waist
30 circumference, physical activity, sedentary behaviors, and soft drink, snack and breakfast
31 consumption. The study is an integral part of the ongoing health surveillance system of
32 the Municipal Health Service in the Rotterdam area (Youth Monitor Rotterdam), in which
33 general health, well being and related factors of youth aged 0-19 years are monitored. The
34 Medical Ethics Committee of Erasmus University Medical Center reviewed the proposal
35 and issued a "declaration of no objection" for the ENDORSE project.

1 Recruitment of schools

2
3 Schools located in the Rotterdam area that participate in the Youth Monitor Rotterdam
4 (YMR) (N=56) were invited for participation in the ENDORSE study. A letter and an
5 information sheet explaining the goals and the logistics of the study were sent to school
6 principals. The schools Principals were contacted by a researcher, upon which they could
7 express their interest in participating in the study. If necessary, additional information to
8 make a more informed decision was provided. Subsequently, a random sample of 17 school
9 locations was drawn from the pool of schools that were willing to participate, after stratifica-
10 tion of the schools according to the area in the city in which they are located. Stratification
11 was done, to ensure a range of physical and cultural environments. Rotterdam is the second
12 largest city of the Netherlands. It has approximately 600,000 inhabitants of which 46% are
13 of non-Dutch origin [14].
14

15 Recruitment of participants within schools

16
17 Five classes in each participating school were randomly selected for participation in the cross-
18 sectional study, which took place in 2005/2006. All adolescents in one class participated in
19 the study, unless they or their parents indicated that they were not willing to participate. The
20 adolescents in the first year classes were also asked to complete the questionnaires at two
21 years follow up. To have sufficient power, we assumed (conservatively) that obesity inducing
22 risk behaviors will be present among at least 40% of adolescents, and we assumed moderate
23 effect sizes of determinants on behaviors. With a significance level of .05 and 80% power, a
24 sample size of approximately 600 students would be sufficient. Since we plan to do separate
25 analyses for girls and boys we aimed at including 1200 students. First year adolescents were
26 over-sampled to 800 as these adolescents will be followed up two years later.
27

28 Procedure

29
30 The ENDORSE study follows the logistics of the YMR. The YMR routinely collects data
31 among adolescents in the first and third year of secondary school. The school levels vary
32 from lower vocational training to high school. According to the usual procedure of the
33 YMR, the ENDORSE study was announced through a letter to the parents. This letter
34 explained that the YMR was extended with an extra part, aimed to gain insight in the preva-
35 lence and causes of overweight. Parents could keep their child from participating in the
36 study by sending the attached form to the adolescent's teacher (passive consent procedure).
37 Approximately two weeks after the usual YMR questionnaire, the adolescents completed the
38 ENDORSE questionnaire confidentially during a school hour with a teacher and a research
39 assistant present. Within a month after completion of the ENDORSE questionnaire, two

1 trained research assistants measured height, weight, waist circumference and pubertal de-
2 velopment according to standardized procedures described in a measurement protocol. The
3 adolescents were asked to come in succession to a private room where they were measured
4 without shoes. After the anthropometrical measurements, the adolescents completed a Tan-
5 ner scale to assess pubertal development [15]. To guarantee confidentiality the adolescents
6 could put the Tanner scale form in an envelope before handing it over to the research as-
7 sistant. After these measurements, the adolescents received a Frisbee as a compensation
8 for their participation and were requested to give an envelope with a questionnaire to their
9 parents. The envelope contained a letter explaining the purpose of the study and the reason
10 why the parents were asked to complete the questionnaire, a pre-addressed and stamped
11 envelope and a card which they could complete to participate in a raffle to win one of five
12 I-pods. Parents were reminded twice to complete and return the questionnaires by means of
13 reminder cards delivered to parents via the adolescents. Parents were not addressed directly,
14 since the YMR procedure did not allow us to have any personal or address details.

15
16 Two observers independently conducted audits of the schools, school canteens, schoolyards,
17 and an area of 300 meter radius surrounding the schools. The observations were conducted
18 within three months from the completion of the adolescent questionnaire. A brief interview
19 with canteen managers and school representatives was part of the audit. One of the observ-
20 ers conducted the interviews with school representatives, and the other observer conducted
21 the interviews with the school canteen managers.

22 Census data (year 2005/2006) from the Center for Research and Statistics (COS), the
23 research center of the municipality of Rotterdam was collected on all neighborhoods of
24 Rotterdam.

25 In the follow-up data collection the same procedures are used. All measurements (ques-
26 tionnaire, anthropometrics, audits and interviews) are conducted within one week per
27 school. As an incentive for their participation, the adolescents received a key holder.

28 29 Measurements

30
31 The ENDORSE questionnaires were developed by using existing validated Dutch ques-
32 tionnaires where possible. If no validated questionnaires were available the ENDORSE
33 questionnaires were informed by questionnaires on related topics that were used in on-
34 going projects in the Netherlands, and questionnaires used in other countries. Relevant
35 parts of these questionnaires were adapted to tailor the specific behaviors identified. If no
36 relevant and validated questionnaires were available, new questions were developed for the
37 ENDORSE study. The ENDORSE study contained the following measurements: adolescent
38 questionnaire, parent questionnaire, interviews with school representatives and canteen
39 managers, observations of the school environment, census data collection and adolescent

body measurements. These are described in detail in the following paragraphs. All determinants measured in the ENDORSE study are listed in Table 2.1.

Table 2.1 Individual and environmental correlates measured in the ENDORSE study

	Perceived / self-reported variables			Objectively measured variables	
	Adolescent questionnaire	Parent questionnaire	Interview	Observation	Census data
Individual variables	attitude; parental norms; modeling; perceived behavior control; intention; habit; external eating behavior				
Physical environment	<p>Home*: availability of sports facilities, bicycle; soft drinks, breakfast products, snacks/ sweets, television set in bedroom.</p> <p><u>accessibility</u> of soft drinks, snacks/sweets, breakfast products, television set</p> <p>School: amount of traffic; safety for cycling; availability of sidewalks and cycle lanes; availability of a bicycle shed</p> <p>Neighborhood: amount of traffic; safety for cycling; availability of sidewalks and cycle lanes; safety and attractiveness of neighborhood; availability of playgrounds, parks, squares, sports clubs</p>	<p>Home: availability of bicycles, cars, soft drinks, breakfast products, snacks/ sweets</p> <p><u>Accessibility</u> of soft drinks, snacks/sweets, breakfast products, television set</p> <p>Neighborhood: amount of traffic; safety for cycling; availability of sidewalks and cycle lanes; safety in neighborhood; attractiveness of neighborhood</p>		<p>School : <u>Availability</u> of bicycle shed, food products in the school canteen and vending machines, PA facilities on the school playground; Shops, fast food restaurants & PA facilities in the school neighborhood. Traffic amount and safety Facilities and frequency of public transport</p>	<p>School: availability of shops, sports facilities and playgrounds for children > 12 years old; areas of sidewalks, bicycle lanes, roads, grass, plants, water; traffic accidents; criminality, crime reports</p> <p>Neighborhood: <u>availability</u> of shops, sports facilities and playgrounds for children > 12 years old; areas of sidewalks, bicycle lanes, roads, grass, plants, water; traffic accidents; criminality, crime reports</p>

table continued on next page

	Perceived / self-reported variables			Objectively measured variables	
	Adolescent questionnaire	Parent questionnaire	Interview	Observation	Census data
Economic environment	Home: income; amount of money that can be spent in 1 week	Home: having a paid job; educational level		School: pricing of school canteen products; pricing of products in shops around the school	Neighborhood: residential types; household income; educational level; % unemployment, % living on social security: % rented houses / owner-occupied properties; mean value of houses; % various ethnic groups
Political environment			School: Food & physical activity policy		

Adolescent questionnaire

Physical activity and sedentary behaviors were assessed with an adapted version of the Activity Questionnaire for Adolescents & Adults (AQuAA) [Chin A Paw MJ, Sloomaker SM, Schuit AJ, van Zuidam M, Van Mechelen W, unpublished] which is a short questionnaire to assess physical activity at school and during leisure time, active transportation to school and sedentary behaviors in leisure time. The structure of the AQuAA was obtained from the SQUASH-questionnaire [16]. The AQuAA refers to activities in the past week (7-day recall). The test-retest reproducibility was fair to moderate for this questionnaire, with intra-class correlations ranging from 0.46 to 0.59.

Dietary intake was assessed with food frequency questions referring to a general week, and a 24-hour recall question. The questionnaire included TPB items for all behaviors. All the questions on TPB variables were measured on a five-point bipolar scale. Attitude was assessed with two items by asking if the adolescent considered the behavior as good or bad, and as pleasant or unpleasant ('e.g. Regular physical activity is very good (+1) – very bad (-1)'). Subjective norm was assessed with one item, for example 'my parents consider eating breakfast as very good (+2) – very bad (-2)'. Modeling was assessed with two items by asking if the parents and friends perform the behavior ('My friends eat snacks...a lot (+2) – very little (-2)'). Perceived behavioral control was assessed with two items by asking how easy or

1 difficult the behavior is to perform (How easy/difficult is it for you to eat breakfast? Very easy
2 (+2) – very difficult (-2)), and by asking if the decision to perform a behavior is completely
3 under the control of the adolescent (Do you decide by yourself if you eat breakfast? Yes, that
4 is completely my own decision (+2) – no, that is not fully my own decision (-2)). Intention
5 to perform the behavior was assessed with one item asking how certain the adolescent is to
6 perform the behavior in the coming six months (Do you intend to eat breakfast the next six
7 months? Yes, certainly do (+2) – no certainly do not (-2)).

8 Habit strength of dietary and physical activity behaviors was measured by means of the
9 Self Report Habit Index [17]. This questionnaire assesses three features of habitual behavior:
10 the extent to which a behavior is automatic, the repeated character of the behavior and
11 the sense of identity the behavior reflects. Three items assessed these features, namely:
12 the behavior 'x' is something.... 'I do frequently', 'is something I do automatically' and 'is
13 something that's typically 'me". These items were measured on a five-point scale, ranging
14 from 'I completely agree' (+2) to 'I completely disagree' (-2).

15 External cues that can influence eating and drinking patterns were questioned with nine
16 items on a four point Likert scale (always (+2) – never (-2)), for example 'I get hungry
17 when I see snacks or candy' or 'When I walk past a fast-food restaurant, I feel like buying
18 something. These questions were based on the external eating behavior questions from the
19 Dutch Eating Behavior Questionnaire [18] and adapted to address the topics adolescents
20 mentioned in the focus group interviews.

21 In the adolescent questionnaire the following perceived environmental determinants were
22 assessed: availability and accessibility of facilities for physical activity and food, school fac-
23 tors, neighborhood factors, parenting factors and economic factors. Demographic factors
24 (gender, age, ethnicity) were available for each adolescent through the YMR questionnaire.

25 The adolescent questionnaire was pre-tested among ten adolescents by means of cognitive
26 interviewing. Subsequently, the questionnaire was completed twice by 89 schoolchildren
27 (aged 13-14) ten-days apart to assess the test-re-test reliability and other psychometrics of
28 the questionnaire. Items with low reliability were adjusted or deleted from the questionnaire.
29

30 *Parent questionnaire*

31 Parental behavior, family and household environmental determinants were assessed in the
32 parent questionnaire. Parental physical activity and sedentary behaviors were assessed with
33 the adapted version of the AQuAA [Chin A Paw M], Sloomaker SM, Schuit AJ, van Zuidam
34 M, Van Mechelen W, unpublished], referring to activities in the past week (7-day recall).
35 Dietary behaviors were assessed with food frequency questions referring to a general week.
36 Neighborhood factors as perceived by the parents, such as safety in neighborhood and
37 attractiveness of neighborhood, parenting practices, parental allowance, availability and
38 accessibility of soft drinks, breakfast products, snacks/sweets, television set and parental
39

1 self-reported body weight and height, and demographics (gender, educational level, having
2 a paid job) were assessed. One parent completed this questionnaire.

3 4 *Interview questionnaires*

5 To assess school food and physical activity factors, pre-structured interview forms were
6 developed for interviews with the school canteen managers and with a school representa-
7 tive. The interview form for canteen managers contained questions on the availability of
8 food products, opening hours of the canteen, pricing policy, who will receive profits from
9 the foods sold, and canteen policies, for example if there are agreements on the assortment
10 with the catering organization. The form also included information about the soft drink and
11 snack vending machines, e.g. how often the vending machines are filled. The interviews with
12 a school representative were aimed at gaining insight in school policies regarding diet and
13 physical activity. As a basis for the interview form, the 7-item school-wide food practices
14 scale was used [19]. This scale assesses food practices allowed at school with the following
15 items: 'Are students allowed to have food in the classroom?', 'Are students allowed to have
16 beverages in the classroom?', 'Are students allowed to have snacks in the hallways?', 'Are
17 students allowed to have beverages in the hallways?', 'Are food or food coupons used as
18 reward or incentive for students?', 'Do you have classroom fundraising that includes food
19 sales?', and 'Do you have school wide fundraising that includes food sales?'. The question-
20 naire furthermore contained questions about whether or not the school has a formal food or
21 physical activity policy and if yes, to indicate what this policy is. Questions on what health
22 education programs they use in schools and possibilities and promotion activities for the
23 adolescents to be active before, during and after school time, were also included.

24 25 *Audit instrument for area observations*

26 An audit instrument was developed to assess the availability and accessibility of foods and
27 physical activity facilities in the schools, in the schoolyards and in the neighborhood around
28 schools. The audit instrument consists of a pre-structured form with five parts: school infor-
29 mation, school building, nutrition, physical activity and school environment. As much as
30 possible, the instrument had a 'tick box' answering format and included observation of 'ob-
31 jective' characteristics. When more subjective characteristics such as 'state of maintenance
32 of the school yard', or 'traffic situation around the school' were reported, photographs were
33 taken from pre-defined angles. The audit instrument included also a description of the item
34 to be observed. The neighborhoods around schools that were observed were defined as a
35 radius of 300 meters from the school. This definition was based on the basic assumption that
36 the facilities in the neighborhood around schools should be accessible in a general school
37 lunch break of approximately 30 minutes, and that adolescents use facilities that are close by
38 the school. The audit instrument included maps of the 300-meter radius around the schools,
39 on which the route walked to observe the area could be drawn and the location of green

1 spaces could be indicated. The first part (A) of the audit instrument involved some general
2 school information e.g. the address and number of pupils. Part B involved items on the
3 school building e.g. number of floors, entrance for schoolchildren, availability of an elevator
4 and location and visibility of stairs. Part C involved observation of the school canteens e.g.
5 counting the number of soft-drink and snack vending machines in the canteen, checking
6 the items that are available in these machines, how well they were filled and advertising
7 in the canteen. Part D involved observation of the school physical activity facilities e.g.
8 the bicycle shed, facilities for activity and aesthetics of the schoolyards. Part E involved
9 observation of the neighborhood around schools e.g. the facilities for physical activity (e.g.
10 parks, fields, playing and sports fields) that were visible from the schoolyards and that were
11 present in the neighborhood (300 meters). The component of the neighborhood observa-
12 tion related to dietary intake involved observation of the food retail outlets (baker's shops,
13 snack bars, fast-food chains, supermarkets, kiosks, gas stations, tobacco shops, chemist's
14 shops) that were visible from the schoolyards and that were present in the neighborhood
15 around schools.

16 The audit instrument first was reviewed by experts on accuracy and completeness of the
17 instrument for its intended purpose. Secondly, the instrument was pilot tested by conduct-
18 ing the observations at two schools and in the corresponding neighborhoods with two ob-
19 servers. Important aspects of the pilot test were the completeness of the forms and feasibility
20 and suitability of using the definition of 300-meter radius for school environment. After the
21 pilot tests at the two schools, the audit forms were adapted if needed. The adapted forms
22 were tested at a third school, by three observers.

23 24 *Census data*

25 Census data was utilized to gather additional environmental data regarding the neighbor-
26 hoods around schools and the neighborhoods in which the children live. The data included
27 area-level household income, educational level, residential types, percentage of residents
28 aged 10-19, percentage unemployment, percentage living on social security, percentage of
29 rented houses and owner-occupied properties, mean value of houses, percentages of various
30 ethnic groups, number of stores, fast food restaurants and the amount and type of green
31 spaces, water, bicycle tracks and foot paths. The census data could be linked to the home
32 environment of the adolescents with information on the ZIP code, which was asked in the
33 adolescent and parent questionnaires. Neighborhoods were defined based on a formal clas-
34 sification from Statistics Netherlands.

35 36 *Body measurements*

37 Body height was measured without shoes with a Seca 225 mobile height rod with an ac-
38 curacy of 0.1 cm. A calibrated electronic digital floor scale (SECA 888 class III) was used to
39 determine body weight of the participant in street clothes, without shoes, with an accuracy

1 of 0.1 kg. Waist circumference was measured using a spring loaded measuring tape (SECA
2 200) to the nearest 0.1 cm. The waist circumference was measured twice. In case of a differ-
3 ence of more than 1.0 cm between these two measurements, the waist circumference was
4 measured twice again. Adolescents self-reported on their stage of pubertal development
5 using drawings of Tanner stages [15] in the baseline data collection only.
6

7 8 **DISCUSSION**

9
10 The ENDORSE study is a comprehensive, longitudinal study in which both individual
11 and environmental determinants of selected obesity related behaviors in adolescents are
12 examined. The study has several strengths. It examines both sides of the energy balance
13 equation. It was designed to examine the influence of environmental factors on obesity
14 related behaviors and BMI and objective measures for mapping the environment were in-
15 cluded. Moreover, the study includes environmental factors in various settings, including
16 the home, school and neighborhood. The study involves assessments of both individual and
17 environmental determinants, as opposed to many previous studies, that focused on one or
18 the other. The study has a longitudinal design, allowing analyses of prediction rather than
19 cross-sectional associations only. To date, there are very few studies that examine environ-
20 mental factors of energy balance related behaviors longitudinally. There is an urgent need
21 for such studies, in order to be able to draw stronger inferences for relationships between
22 environmental factors and BMI. However, there were also some limitations in the study
23 protocol. For instance previously validated instruments were not available for all neces-
24 sary measures. Another limitation is that assessments of adolescents' and parental physical
25 activity and dietary behaviors are self-reported. The definition of environment and neigh-
26 borhood is also somewhat arbitrary. The scale of environment to be studied needs further
27 conceptual development [20]. A clear definition of 'the neighborhood' is needed in terms
28 of measurement of respondent perceptions and objective measures of the environment.
29 However there is to date little evidence or consensus as to what constitutes a neighborhood.
30 There is poor agreement about which boundary or scale to use, and how this might impact
31 on the association between predictor and outcome variables is unknown. Moreover the
32 boundary to be used might differ for different target groups and different settings (school
33 or home environments) [20].

34 The ENDORSE study contains rich data examining individual and environmental de-
35 terminants of energy balance related behaviors among adolescents. With this information
36 the influence of risk behaviors for overweight and relationships with socio-economic status
37 and ethnicity can be investigated. Individual and environmental determinants of obesity-
38 related behaviors among adolescents can be examined as well as the interactions between
39 individual and environmental determinants of obesity inducing behaviors. Therefore,

1 data will be analyzed by means of multi-level regression analyses and structural equation
2 modeling. Eventually, the ENDORSE study will provide objectives and entry points for
3 prevention of overweight interventions in younger adolescents. In 2008 the questionnaires
4 for adolescents and parents, the school policy interview forms and the audit instrument will
5 be made available on the Internet [21].
6

7 8 **ACKNOWLEDGEMENTS**

9
10 This study was financially supported by a grant from ZonMw, The Netherlands Organiza-
11 tion for Health Research and Development (grant ID no 2100.0103).
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

1 REFERENCES

- 2 1. Lobstein T, Frelut ML: Prevalence of overweight among children in Europe. *Obes Rev* 2003, 4(4):195-
3 200.
- 4 2. Power C, Lake JK, Cole TJ: Measurement and long-term health risks of child and adolescent fatness.
5 *Int J Obes Relat Metab Disord* 1997, 21(7):507-526.
- 6 3. Must A, Jacques PF, Dallal GE, Bajema CJ, Dietz WH: Long-term morbidity and mortality of over-
7 weight adolescents. A follow-up of the Harvard Growth Study of 1922 to 1935. *N Engl J Med* 1992,
8 327(19):1350-1355.
- 9 4. Must A, Strauss RS: Risks and consequences of childhood and adolescent obesity. *Int J Obes Relat*
10 *Metab Disord* 1999, 23 Suppl 2:S2-11.
- 11 5. Ebbeling CB, Pawlak DB, Ludwig DS: Childhood obesity: public-health crisis, common sense cure.
12 *Lancet* 2002, 360(9331):473-482.
- 13 6. Booth SL, Sallis JF, Ritenbaugh C, Hill JO, Birch LL, Frank LD, Glanz K, Himmelgreen DA, Mudd
14 M, Popkin BM *et al*: Environmental and societal factors affect food choice and physical activity:
15 rationale, influences, and leverage points. *Nutr Rev* 2001, 59(3 Pt 2):S21-39; discussion S57-65.
- 16 7. Ferreira I, van der Horst K, Wendel-Vos W, Kremers S, van Lenthe F, Brug J: Environmental cor-
17 relates of physical activity in youth - A review and update. *Obesity Reviews* 2007, 8(2):129-154.
- 18 8. van der Horst K, Oenema A, Ferreira I, Wendel-Vos W, Giskes K, van Lenthe F, Brug J: A systematic
19 review of environmental correlates of obesity-related dietary behaviors in youth. *Health Educ Res*
20 2007, 22:203-226.
- 21 9. de Bruijn GJ, Kremers SP, de Vries H, van Mechelen W, Brug J: Associations of social-environmental
22 and individual-level factors with adolescent soft drink consumption: results from the SMILE study.
23 *Health Educ Res* 2006.
- 24 10. de Bruijn GJ, Kremers SP, Lensvelt-Mulders G, de Vries H, van Mechelen W, Brug J: Modeling in-
25 dividual and physical environmental factors with adolescent physical activity. *Am J Prev Med* 2006,
26 30(6):507-512.
- 27 11. Kremers SP, De Bruijn GJ, Visscher TL, Van Mechelen W, De Vries NK, Brug J: Environmental influ-
28 ences on energy balance-related behaviors: A dual-process view. *Int J Behav Nutr Phys Act* 2006,
29 3(1):9.
- 30 12. Swinburn B, Egger G, Raza F: Dissecting obesogenic environments: the development and application
31 of a framework for identifying and prioritizing environmental interventions for obesity. *Prev Med*
32 1999, 29(6 Pt 1):563-570.
- 33 13. Ajzen I: *Attitudes, personality, and behavior*: Homewood, IL, US: Dorsey Press; 1988.
- 34 14. The Centre for Research and Statistics [[http://cos.rotterdam.nl/Rotterdam/Openbaar/Diensten/
35 COS/Publicaties/PDF/KC2006UK.pdf](http://cos.rotterdam.nl/Rotterdam/Openbaar/Diensten/COS/Publicaties/PDF/KC2006UK.pdf)]
- 36 15. Coleman L, Coleman J: The measurement of puberty: a review. *J Adolesc* 2002, 25(5):535-550.
- 37 16. Wendel-Vos GC, Schuit AJ, Saris WH, Kromhout D: Reproducibility and relative validity of the short
38 questionnaire to assess health-enhancing physical activity. *J Clin Epidemiol* 2003, 56(12):1163-1169.
- 39 17. Verplanken B, Orbell S: Reflections on past behavior: A self-report index of habit strength. *Journal of
Applied Social Psychology* 2003, 33(6):1313-1330.
18. Van Strien T, Frijters JER, Bergers GPA, Defares PB: The Dutch Eating Behaviour Questionnaire
(DEBQ) for assessment of restrained, emotional and external eating behaviors among Dutch and
American college students. *International Journal of Eating Disorders* 1986, 5:295-315.
19. Kubik MY, Lytle LA, Story M: Schoolwide food practices are associated with body mass index in
middle school students. *Arch Pediatr Adolesc Med* 2005, 159(12):1111-1114.

- 1 20. Giles-Corti B, Timperio A, Bull F, Pikora T: Understanding physical activity environmental cor-
2 relates: increased specificity for ecological models. *Exerc Sport Sci Rev* 2005, 33(4):175-181.
- 3 21. Department of Public Health, Erasmus MC [<http://survey.erasmusmc.nl/intern/actreport/phpwcms/>
4 [index.php?index](http://survey.erasmusmc.nl/intern/actreport/phpwcms/index.php?index)]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

**Part II Environmental
correlates of energy balance-
related behaviors: reviews of
the literature**

3 A systematic review of environmental correlates of obesity-related dietary behaviors in youth

van der Horst K, Oenema A, Ferreira I, Wendel-Vos W, Giskes K, van Lenthe F, Brug J. A systematic review of environmental correlates of obesity-related dietary behaviors in youth.

Health Education Research 2007, 22(2): 203-226.

1 ABSTRACT

2
3 **Background:** There is increasing interest in the role the environment plays in shaping the
4 dietary behavior of youth, particularly in the context of obesity prevention. An overview
5 of environmental factors associated with obesity-related dietary behaviors among youth is
6 needed to inform the development of interventions.

7 **Methods:** A systematic review of observational studies on environmental correlates of
8 energy, fat, fruit/vegetable, snack/fast food, and soft drink intakes in children (4-12) and
9 adolescents (13-18) was conducted. The results were summarized using the Analysis Grid for
10 Environments Linked to Obesity (ANGELO).

11 **Results:** The 58 papers reviewed mostly focused on socio-cultural and economical environ-
12 mental factors at the household level. The most consistent associations were found between
13 parental intake and children's fat, fruit/vegetable intakes, parent and sibling intake with
14 adolescent's energy and fat intakes, and parental education with adolescent's fruit/vegetable
15 intake. A less consistent but positive association was found for availability and accessibility
16 on children's fruit/vegetable intake.

17 **Conclusion:** Environmental factors are predominantly studied at the household level and
18 focus on socio-cultural and economic aspects. Most consistent associations were found for
19 parental influences (parental intake and education). More studies examining environmental
20 factors using longitudinal study designs and validated measures are needed for solid evi-
21 dence to inform interventions.

22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

1 INTRODUCTION

2
3 The promotion of healthful eating in children and adolescents has become an increasingly
4 important public health and research priority as the prevalence of overweight and obesity
5 among children and adolescents continues to rise [1, 2]. Preventing the onset of obesity
6 inducing dietary behaviors or modifying these behaviors at an early age is likely to contrib-
7 ute to the prevention of overweight and obesity. A detailed understanding of factors that
8 determine these behaviors is essential, to be able to effectively prevent or modify obesity
9 inducing eating patterns. The research of determinants of dietary intake in children and
10 adolescents has predominantly focused on individual level determinants of these behaviors,
11 such as attitudes, taste preferences, social influences and perceived behavioral control.
12 However, more recently a shift in attention to environmental determinants of behavior has
13 occurred as it has been acknowledged that a major driving force for the increasing obesity
14 prevalence may be the environment that encourages eating and discourages physical activ-
15 ity [3, 4]. These environmental factors are highlighted in so-called ecological models, and
16 are conceptualized as being interrelated with factors at the individual level [5]. As stated
17 by Rothschild [6], the likelihood that an individual will engage in a healthy behavior is
18 largest when someone is motivated to act healthily, has the abilities to engage in the healthy
19 behavior, and the environment offers the right opportunities to engage in the healthy behav-
20 ior. Motivation and abilities can be regarded as individual determinants of health behavior,
21 whereas opportunities depend on environmental factors.

22 Child and adolescent dietary behavior is likely to be strongly influenced by environmental
23 factors, since children may have less autonomy in food choice. From the age of about three
24 years, children's eating behavior is influenced by their responsiveness to environmental
25 cues, and a variety of family and social factors start to influence children's eating behaviors
26 [7]. The role of parents is considered to be of particular importance, since parents directly
27 determine the child's physical and social environment, and indirectly influence behavior
28 and habits through socialization processes and modeling [8]. When children grow older and
29 move into adolescence they become more autonomous, and lifestyle, developmental, social
30 and environmental changes take place. During this transition to adolescence, dietary intake
31 patterns change and decline in quality compared to childhood. Intakes of fruit, vegetables,
32 milk and fruit juice decrease, whereas intake of soft drink increases during this time [9].

33
34 The expected importance of the environment for obesity related behaviors in children
35 and adolescents is well documented in position papers and narrative reviews [3, 7-11]. The
36 number of studies examining the influence of environmental factors on behavior is expand-
37 ing, but there is no systematic overview of which environmental factors have been studied
38 extensively, and what aspects of the environment are more influential than others. Such an
39

1 overview is needed to identify a research agenda for further investigation and to inform
2 interventions that take environmental factors into account.

3
4 We conducted a systematic review of environmental factors that may potentially influence
5 obesity related dietary behaviors of children and adolescents. We focused our review on
6 energy, fat (total and percent energy), fruit/vegetable, snack/fast food, and soft drink intake.
7 These behaviors have been identified as factors most strongly associated with obesity in
8 adults [4], and are considered to be important obesity inducing behaviors in children and
9 adolescents as well [12, 13]. The environment was defined as ‘anything outside the individual’.
10 Many classifications have been proposed to order the complexity of potential environmental
11 factors. We chose to use the ANGELO framework (Analysis Grid for Environments Linked
12 to Obesity) [14] as a tool to classify the various environmental determinants. The ANGELO
13 framework dissects the environment by two dimensions: the size (micro and macro) and
14 the type of environment. Micro-environments are environmental settings where groups of
15 people meet and gather (e.g. homes, schools, restaurants, neighborhoods). Macro-envi-
16 ronments include the broader infrastructure that may support or hinder health behaviors
17 (e.g. town planning, transport infrastructure, the health system, the media). The ‘types’ of
18 environments distinguished in the ANGELO framework are the physical, socio-cultural,
19 economic, and political environment. The physical environment refers to the availability of
20 opportunities for healthy and unhealthy choices, for instance the availability and accessibil-
21 ity of healthy and unhealthy foods. The socio-cultural environment refers to the social and
22 cultural subjective and descriptive norms and other social influences such as parental influ-
23 ences and peer pressure. The economic environment refers to the costs related to healthy
24 and unhealthy behaviors for instance costs of fruit and vegetables and household income.
25 The political environment refers to the rules and regulations that may influence food choice
26 or availability, for example bans on snack vending machines in schools.

27
28 The review aimed to address the following specific research questions:

29 Which environmental correlates have been studied in relation to child and adolescent en-
30 ergy, fat (total and energy percent), fruit, vegetable, snack, fast food, and soft drink intake?
31 Which environmental factors are consistently associated with these obesity-related dietary
32 behaviors?

33
34
35
36
37
38
39

1 METHODS

3 Data sources and search strategy

5 Studies eligible for inclusion in the review were located from the Medline (PubMed),
6 PsychInfo, Web of Science, and Human Nutrition databases, from January 1980 to December
7 2004. Our search strategy involved using a combination of the broad indexing terms of each
8 database and searching for terms in article titles and abstracts. We used the combination of
9 dietary intake keywords with environmental factor keywords to locate suitable articles. For
10 dietary intake the following keywords were used: energy intake, caloric intake, fat intake,
11 fat consumption, soft drink, soft drink consumption, soft drink intake, sweetened beverage,
12 fruit, fruit consumption, fruit intake, vegetable, vegetable consumption, vegetable intake,
13 eating, diet, nutrition, food habits, food preferences. For environmental factors the follow-
14 ing keywords were used: physical environment, social environment, cultural environment,
15 socio-cultural environment, socio-economic environment, social influences, neighborhood,
16 political environment, built environment, urban environment, rural environment, local
17 environment, school environment, home environment, availability, accessibility, residence
18 characteristics, environment design, parental influence, parenting. Key terms were matched
19 to database specific indexing terms. The sensitivity of the search strategy was verified by
20 checking whether key articles from our personal databases that should be selected through
21 the search strategy, were actually retrieved. In addition to database searches, reference lists
22 of review studies and of articles included in the review were screened for titles that included
23 key terms.

25 Inclusion / exclusion criteria

27 A study had to meet the following criteria to be eligible for inclusion: healthy young people
28 in the age range of 3-18 years (or mean age within this range) as subjects of study; a measure
29 of energy and/or fat intake (total or percent energy), fruit, vegetable, snack, fast food or soft
30 drink consumptions as the dependent variable(s); an outcome measure that was assessed for
31 at least one complete day (for example, studies assessing fruit intakes at just one meal were
32 not eligible). The study samples had to be drawn from countries with established market
33 economies as defined by the World Bank, and the paper had to be published in international
34 peer-reviewed journals in English. Intervention studies and studies that included only over-
35 weight/obese children were excluded.

39

1 Identification of relevant studies

2
3 Potentially relevant papers were selected by screening the titles (first step), abstracts
4 (second step) and the entire article (third step) retrieved through the database searches.
5 Two researchers (KVDH, IF) independently conducted this screening. Disagreement about
6 eligibility between the reviewers was solved through discussion with a third co-author (JB).

8 Data extraction

9
10 Two authors (KVDH and AO) extracted the data from the included studies. Each study's
11 findings and methodological details, such as study design, sample size, dietary outcome(s),
12 environmental determinant(s) assessed, assessment methodology (child and/or parent-
13 report, objectively measured), and statistical analysis methods were listed in tables.

15 Summarizing study findings

16
17 Associations between environmental factors and dietary outcomes were coded as '+' for a
18 positive association, '-' for an inverse association and 'o' for no association. Associations
19 were regarded significant when the p-value reported in the study was smaller than .05. In
20 studies that reported results from univariate and multivariate analysis, only the multivariate
21 results were included. To reduce the number of specific environmental correlates studied,
22 conceptually similar environmental factors were combined (e.g. intakes from father and
23 mother to parental intake). An independent sample was used as the unit of analysis and was
24 defined as the smallest independent sub-sample for which relevant data were reported (e.g.
25 boys/girls) [15].

27 Categorization of variables

28
29 Study findings were tabulated by categorizing the distinct dietary outcomes in a grid dis-
30 secting different environmental settings, i.e. home/household, educational institutions,
31 neighborhoods, city/municipality, and the various types of environmental factors: physical,
32 socio-cultural, economic and political, following the ANGELO framework [14] (Tables 3.2
33 and 3.3).

34
35
36
37
38
39

1 RESULTS

2 3 Search and selection of studies

4
5 The databases search located 6616 titles (Pubmed 1975; PsychInfo 317; Web of Science 2932;
6 Human Nutrition 1392), resulting in 6121 unique titles of potentially relevant articles. Refer-
7 ence sections of earlier reviews and primary studies added 44 titles. Screening the titles and
8 abstracts resulted in a selection of 81 articles, for full text review. Twenty-three of these articles
9 did not meet the inclusion criteria, resulting in a final inclusion of 58 articles with 77 samples.

10 11 Characteristics of included studies

12
13 Most of the studies were cross-sectional (n=55) (Table 3.1). Twenty-nine studies (37 samples)
14 had children as the study population [16-44], and 27 (40 samples) included adolescents
15 [45-71]. One study included a child and adolescent sample [72]. In one paper the age of
16 the population was unclear, and this study was reviewed under an adolescent sample [73].
17 Environmental determinants of fruit/vegetable intake were examined in 34 studies, deter-
18 minants of fat intake in 23 studies, determinants of fast food/snack intake in 21 studies,
19 determinants of energy intake in 17 studies and determinants of soft drink intake in 10
20 studies. Only five studies reported the validity, and seven studies reported the reliability of
21 the dietary intake measurements used.

22 23 Potential environmental correlates of children's dietary behaviors

24
25 The findings from the studies are summarized in Tables 3.2 and 3.3. Table 3.4 provides
26 a summary of the number of studies and the consistent associations in each cell of the
27 ANGELO framework. In the following sections a summarized description of the results is
28 provided for the various behaviors. The factors examined on each environmental level and
29 the environmental factors that showed consistent associations with dietary behaviors in at
30 least two replicated studies are described.

31 32 *Environmental correlates of energy intake*

33 At the household environmental level, physical factors (one study/sample), socio-cultural
34 factors (five studies, six samples), and economic factors (five studies/samples) were ex-
35 amined in relation to energy intake (Table 3.2). One study examined factors in the school
36 environment, no studies examined factors in the neighborhood environment, and two stud-
37 ies examined factors at the city/municipality level. At the household socio-cultural level, an
38 inverse association with energy intake was found for encouragement, offering assistance and
39 giving prompts to increase food intake during meals in two out of three samples [35, 36, 40].

Table 3.1 Characteristics of studies included in the review: sample size; sex; study design; assessment of dietary intake; data analysis; country

	Children (3-12 years)				Adolescents (13-18 years)			
	References		Samples		References		Samples	
			Number	%			Number	%
Sample size								
<100	[25M/I, 29, 35, 37, 40, 43M/F]	37	100		[61M]	40	100	
100-199	[19, 25F, 26F, 27F, 32, 33F]	8	21.6		[45M/F, 60]	1	2.5	
200-299	[28]	6	16.2		[46]	3	7.5	
300-499	[16, 24, 36, 38, 44]	1	2.7		[56, 68M/F, 71M/F]	1	2.5	
500-999	[17, 18]	5	13.5		[47M/F, 48, 49I, 50, 51M/F, 54M/F, 62-64, 73M/F]	5	12.5	
1,000-2,999	[21I/II, 23, 30, 34M/F, 39F, 41, 42]	2	5.4			14	35.0	
3,000-4,999	[22M/F, 72M/F]	9	24.3			3	7.5	
≥5000	[20, 31]	4	10.8		[49II, 59, 65-67, 70]	6	15.0	
		2	5.4		[52M/F, 53, 57M/F, 58, 69]	7	17.5	
Sex								
Girls only	[26, 27, 33, 39]	4	10.8			-	-	
Boys only		-	-		[61]	1	2.5	
Boys and girls combined	[16-20, 21I/II, 23, 24, 25I, 28-32, 35-38, 40-42, 44]	23	62.2		[46, 48, 49I/II, 50, 53, 55, 56, 58-60, 62-67, 69, 70]	19	47.5	
Boys and girls, separately	[22M/F, 25I/M/F, 34, 43M/F, 72M/F]	10	27.0		[45M/F, 47, 51, 52, 54, 57, 68, 71, 72M/F, 73]	20	50.0	
Study design								
Cross-sectional	[16-20, 21I/II, 22M/F, 23, 24, 25M/F/I, 26F, 27F, 28-32, 33F, 34M/F, 35, 37, 38, 39F, 40-42, 43M/F, 44, 72M/F]	36	97.3		[46, 47M/F, 48, 49I/II, 50, 51M/F, 52M/F, 53, 54M/F, 55, 56, 57M/F, 58-60, 61M, 63-67, 68M/F, 69, 70, 71M/F, 72M/F, 73M/F]	37	92.5	
Longitudinal (length of study)	[36] (2.5 years)	1	2.7		[62] (6 years)	1	2.5	
Case-control		-	-		[45M/F]	2	5.0	
Dietary outcome								
Energy intake	[23, 33F, 35, 36, 38, 39F, 40, 41, 43M/F]	10	27.0		[45M/F, 46, 50, 54M/F, 56, 67, 70, 73M/F]	11	27.5	
Energy from fat (%)	[23, 27F, 36, 38, 39F, 41, 44]	7	18.9		[47M/F, 50, 51M/F, 54M/F, 56, 60, 64, 67, 70, 71M/F, 73M/F]	16	40.0	
Total fat intake (g)	[22M/Fa, 43M/F]	4	10.8		[45M/F, 47M/F, 57M/F, 68M/F, 71M/F, 73M/F]	12	30.0	
Fruit intake	[16, 18, 24, 28, 32, 37, 72M/F]	8	21.6		[48, 50, 53b, 57M/F, 58, 60, 61M, 63, 64, 67, 69, 70, 72M/F]	15	37.5	

	Children (3-12 years)			Adolescents (13-18 years)		
	References	Samples		References	Samples	
		Number	%		Number	%
Vegetable intake	[16, 18, 24, 28, 32, 37, 42, 72M/F]	9	24.3	[50, 53b, 57M/F; 58, 60, 61M, 64, 67, 69, 70, 72M/F]	13	32.5
Juice intake	[28, 32]	2	5.4	[48, 61M]	2	5.0
Composite measure FV intake	[21M/F, 22M/F, 26F, 30, 31, 34M/F, 42]	10	27.0	[52M/F, 62, 64-67]	7	17.5
Composite measure of FJV intake	[25I/M/F, 28, 29c, 32]	6	16.2	[61M]	1	2.5
Fast food consumption	[20, 24]	2	5.4	[57M/F, 59d]	3	7.5
Snack food intake	[19, 31, 42, 72M/F]	5	13.5	[49I/IIe, 52M/Fe, 60, 67, 72M/F]	8	20.0
Pizza & snack	[29f]	1	2.7	[56]	1	2.5
Soft drink consumption	[16, 17, 24, 29f, 72M/F]	6	16.2	[48, 55, 57M/F, 60, 67, 72M/F]	8	20.0
Assessment of dietary outcome						
Self-report	[17, 19, 21I, 22M/F, 25M/F, 28-32, 34M/F, 36, 38, 39F, 42]	18	48.6	[46, 47M/F, 48, 49I/II, 52M/F, 53, 54M/F, 55, 56, 57M/F, 58-60, 61M, 62-67, 68M/F, 69, 70, 71M/F, 72M/F, 73M/F]	35	87.5
Parent-report	[16, 18, 21II, 25I, 26F, 33F, 35, 40, 41, 44, 72M/F]	12	32.4		-	-
Self- or parent-report	[20, 23]	2	5.4	[45M/F, 50]	3	7.5
Parent- and self-report (together)	[24, 27F, 37, 43M/F]	5	13.5	[51M/F]	2	5.0
Measurement instrument dietary outcome						
24-hour recall	[20, 23, 26F, 27F, 29, 30, 33F, 34M/F, 38, 42]	11	29.7	[46, 49I, 61M, 64]	4	10.0
48-hour recall		-	-	[50]	1	2.5
Food frequency questionnaire	[16, 18, 19, 21M/F, 22M/F, 24, 31]	9	24.3	[51M/F, 53, 56, 57M/F, 58, 60, 63, 65-67, 68M/F, 70, 71M/F]	17	42.5
2-day food record	[28, 32]	2	5.4		-	-
3-day food record	[37, 39F, 43M/F]	4	10.8	[54M/F, 73M/F]	4	10.0
7-day food record	[25I/M/F, 40]	4	10.8	[45M/F]	2	5.0
Questionnaires	[17, 35, 36, 72M/F]	5	13.5	[48, 49II, 52M/F, 55, 59, 62, 69, 72M/F]	10	25.0
24-hour recall & 2-day food record	[41, 44]	2	5.4	[47M/F]	2	5.0

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

	Children (3-12 years)		Adolescents (13-18 years)	
	References	Samples Number %	References	Samples Number %
Reliability of dietary intake measurement				
Unknown / not reported	[16-20, 22M/F, 23, 24, 25I/M/F, 26F, 30, 31, 33F, 35-38, 39F, 40-42, 43M/F, 44, 72M/F]	29 78.4	[45M/F, 46, 47M/F, 48, 49I/II, 50, 51M/F, 52M/F, 53, 54M/F, 57M/F, 58, 60, 63-65, 69, 72M/F]	26 65.0
Reported elsewhere	[27F, 28, 29, 32, 34M/F]	6 16.2	[56, 59, 67, 71M/F, 73M/F]	7 17.5
< 0.7		- -	[61M, 66, 68M/F, 70]	5 12.5
≥ 0.7	[21M/F]	2 5.4	[55, 62]	2 5.0
Validity of dietary intake measurement				
Unknown / not reported	[17-20, 23, 24, 26F, 30, 33F, 35-38, 39F, 40-42, 43M/F, 44, 72M/F]	22 59.5	[45M/F, 46, 47M/F, 48, 49I/II, 50, 52M/F, 53, 54M/F, 55, 57M/F, 58, 61M, 62, 69, 72M/F, 73M/F]	25 62.5
Reported elsewhere	[22M/F, 25I/M/F, 27F, 28, 29, 32, 34M/F]	11 29.7	[51M/F, 56, 59, 60, 63-65, 67, 68M/F, 71M/F]	13 32.5
< 0.6	[16fruit, veg., 21M/F, 31]	4 10.8	[66, 70]	2 5.0
≥ 0.6	[16soft drinks]	1 2.7		-
Data analysis				
Univariate	[19, 24, 28, 32, 35, 38, 39F, 72M/F]	9 24.3	[45M/F, 48, 50, 51M/F, 54M/F, 56, 57M/F, 68M/F, 71M/F, 72M/F, 73M/F]	19 47.5
Multiple	[20, 23, 31, 36, 40, 41]	6 16.2	[47M/F, 49I/II, 52M/F, 55, 58-60, 63-65, 67, 69, 70]	16 40.0
Univariate & multiple	[16-18, 21M/F, 22M/F, 27F, 29, 37, 42, 43M/F, 44]	14 37.8	[46, 53, 61M, 62]	4 10.0
Model testing (Structural equation modeling)	[25I/M/F, 26F, 30, 33F, 34M/F]	8 21.6	[66]	1 2.5
Country				
North America	[17, 20, 23, 25M/F/I, 26F, 27F, 28-32, 33F, 34M/F, 35, 36, 38, 39F, 41, 42, 43M/F, 44]	25 67.6	[46, 47M/F, 49I/II, 52M/F, 53, 58, 61M, 64-67, 68M/F, 69, 70, 73M/F]	20 50.0
Europe	[16, 18, 19, 21I/II, 22M/F, 24, 37, 40]	10 27.0	[45M/F, 48, 50, 51M/F, 54M/F, 56, 57M/F, 60, 62, 63, 71M/F]	16 40.0
Oceania			[55, 59]	2 5.0

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

	Children (3-12 years)		Adolescents (13-18 years)	
	References	Samples Number %	References	Samples Number %
Asia	[72M/F]	2 5.4	[72M/F]	2 5.0

F = girls only; M = boys only; M/F = boys and girls analyzed separately; I/II = two independent samples based on different age groups

^a Frequency of high fat food consumption

^b Inadequate fruit / vegetable consumption (less than once a day)

^c Percent of total daily energy intake contributed to fruit, vegetable and juice intake

^d Frequency of take-away food consumption

^e Low nutrient dense / high fat snacks consumption

^f Percent of total daily energy intake contributed by that food group & consumption frequency per day

^g Unclear whether dietary intake was measured with a food-frequency questionnaire or with another questionnaire

Table 3.2 Summary of micro and macro environmental correlates of dietary intakes among children (3 to 12 year olds)

Correlate	Related to dietary behaviors		Assoc (+ or -)†	References	Unrelated to dietary behaviors	# Samples	Summary (n)	
	References	References					+	-
Physical								
Energy								
Accessibility to food	[36]		-			1	0	1
Minutes foods present at home		[36]				1	0	0
Fat (total fat, en% fat)								
Accessibility to food		[36]				1	0	0
Minutes foods present at home		[36]				1	0	0
Fruit, Juice, Vegetables								
Availability	[25F, 28, 30, 34F]		+	[25M/I, 34M]		7	4	0
Accessibility	[21I/II, 25F/I]		+	[25M, 28]		6	4	0
Home FIV barriers	[28]		-			1	0	1
Television on during meals	[29]		-			1	0	0
Snacks, Fast food								
Television on during meals	[29]		+			1	1	0
Soft drink								
Television on during meals	[29]		+			1	1	0
Availability	[17]		+			1	1	0
Socio-Cultural								
Energy								
Parental intake		[43M/F]				2	0	0
Parenting practices								
Control/restriction/discouragement	[33F]		-	[35, 36]		3	1	0
Encouragement/assistance/prompts to increase food intake	[35, 40]		-	[36]		3	0	2
Food as reward		[36]				1	0	0
Parents negative statements about foods	[40]		-			1	0	1
Family support		[36]				1	0	0
Minutes spent eating at home		[36]				1	0	0
No. meals eaten out		[36]				1	0	0
Marital status parents		[36]				1	0	0
Food presentations/ food offers	[35]		+			1	1	0

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

Correlate	Related to dietary behaviors	Assoc (+ or -)†	Unrelated to dietary behaviors		# Samples	Summary (n)	
	References		References	+		-	0
Fat (total fat, en% fat)							
Parental intake	[27F, 43M/Fmother]	+	[43M/Ffather]		3	3*	0 2*
Parenting practices	[36**]	-			2	2*	1* 0
Control/restriction	[27F, 36\$] [36increase]	+					
Prompts to increase/decrease food intake		-	[36decrease]		1	0	1* 1*
Food as reward		-	[36]		1	0	0 1
Pressure to eat	[27F]	+			1	1	0 -
Mothers monitoring			[27F]		1	0	0 1
Family support			[36]		1	0	0 1
Minutes spent eating at home			[36]		1	0	0 1
No. meals eaten out			[36]		1	0	0 1
Marital status parents			[36]		1	0	0 1
Single parent family	[38]	+			1	1	0 0
Fruit, Juice, Vegetables							
Modeling (parents, important others)	[21I, 28]	+			2	2	0 0
Mothers intake of soft drinks and sweets			[16]		1	0	0 1
Avoidance of negative modeling			[16]		1	0	0 1
Parental intake	[16, 18, 21II, 26F, 37fruit]	+	[37veg]		5	5*	0 1*
Parental intake if FV are highly available	[30]	+			1	1	0 0
Parent FV intake if FV are low available			[30]		1	0	0 1
Parenting style	[32negative]	-	[32authoritative]		1	0	1* 1*
Parenting practices			[18]		1	0	0 1
Food as reward			[16]		1	0	0 0
Encouragement / verbal praise	[16veg]	+	[16fruit, 32]		2	1*	0 2*
Discouragement to eat sweets, soft drinks			[16]		1	0	0 1
Control / restriction	[28, 32]				2	0	0 2
Permissiveness			[16]		1	0	0 1
Negotiation			[16]		1	0	0 1
Pressure to eat	[26F]	-	[16]		2	0	1 1
Catering on children's demands			[16]		1	0	0 1

Correlate	Related to dietary behaviors		Assoc (+ or -)†	Unrelated to dietary behaviors		# Samples	Summary (n)	
	References	behaviors		References	behaviors		+	-
Parental preparation practices				[32]		1	0	0
Age of introduction of FIV (late)	[18fruit]		-	[18Veg]		1	0	1*
Breastfeeding	[18fruit]		+	[18Veg]		1	1*	0
Parent preparation of quick & easy food				[29]		1	0	0
Two parent family	[42]			[29]		1	0	0
One parent family	[31]		-	[72Ffruit] [72Mfruit/veg]		1	0	1
Family dinner	[72Fveg]		+			3	1*	1*
Family breakfast	[72Ffruit] [72Mfruit/veg]		+	[72Fveg]		2	3*	0
Snacks, Fast food								
Parental intake	[19]		+			1	1	0
Parenting practices (general)	[19]		+			1	1	0
Control / reward				[19]		1	0	0
Parent preparation of quick & easy food				[29]		1	0	0
Two parent family				[29]		1	0	0
One parent family				[42]		1	0	0
Family dinner			-	[31 snack foods, 72M/F]		3	0	1*
Family breakfast	[31 fried snack foods]		-	[72M]		2	0	1
Soft drink								
<i>Parental intake</i>								
Refraining from negative modeling	[16, 17]		+			2	2	0
Parenting practices				[16]		1	0	0
Food as reward				[16]		1	0	0
Discouragement to drink soft drinks				[16]		1	0	0
Encouragement to eat fruit, vegetables				[16]		1	0	0
Control / pressure				[16]		1	0	0
Verbal praise				[16]		1	0	0
Permissiveness	[16]		+			1	1	0
Negotiation				[16]		1	0	0
Catering on children's demands				[16]		1	0	0
Parent preparation of quick & easy food				[29]		1	0	0

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

Correlate	Related to dietary behaviors	Assoc (+ or -)†	Unrelated to dietary behaviors		Summary (n)
	References		# Samples	+	
Two parent family	[31]		[29]	1	0 0 1
Family dinner	[72F]	-	[72M]	3	1 1 1
Family breakfast	[72M/F]	+		2	0 0 2
Economic					
Energy					
Income sufficiency			[38]	1	0 0 1
Household income (high)	[23]	-	[39E, 41]	3	0 0 1
Parents educational level (high)	[23]	+	[39F]	2	1 0 1
# persons/household			[36, 38, 41]	3	0 0 3
Occupation			[23]	1	0 0 1
Parents live less than 10y in present house	[23]	+		1	1 0 0
Fat (total fat, en% fat)					
Income sufficiency			[38]	1	0 0 1
Household income (high)	[39F]	-	[27E, 41]	3	0 1 2
Parents educational level (high)	[23, 39F]	-		2	0 0 2
Maternal employment			[44]	1	0 0 1
# persons/household	[38]	-	[36, 41]	3	0 1 2
Parents live less than 10y in present house	[23]	-		1	0 1 0
Fruit, Juice, Vegetables					
Household income (high)			[29]	1	0 0 1
Deprivation index (high)	[37fruit]	-	[18, 37veg]	2	0 0 1*
Parents educational level (high)	[18veg, 37fruit]	+	[16, 18fruit, 29, 37veg]	4	2* 0 4*
Number of hours/week worked by mother			[29, 42]	2	0 0 2
SES / occupational class	[24veg]	+	[24fruit, 42]	1	1* 0 2*
Snacks, Fast food					
Household income (high)	[20]	+	[23, 29]	3	1 0 2
Parents educational level			[29]	1	0 0 1
Number of hours/week worked by mother	[42white children]	-	[29]	2	1* 1* 1*
SES / occupational class	[42black children]	+			
	[42]	+	[23, 24]	3	1 0 2

Correlate	Related to dietary behaviors		Assoc (+ or -)†	Unrelated to dietary behaviors		Summary (n)		
	References	# Samples		References	# Samples			
Household income				[29]	1	0	0	1
Parents educational level				[16]	1	0	0	1
Occupational class				[24]	1	0	0	1
Number of hours/week worked by mother				[29]	1	0	0	1
Physical								
Energy								
Minutes food present at school				[36]	1	0	0	1
Fat (total fat, en% fat)								
Minutes food present at school				[36]	1	0	0	1
Soft drink								
Availability	[17]		+		1	1	0	0
Socio-Cultural								
Energy								
Prompts to increase food at school lunch				[36]	1	0	0	1
Fat (total fat, en% fat)								
Prompts to increase food at school lunch	[36]		-		1	0	1	0
Fruit, Juice, Vegetables								
Modeling peers FJV				[28]	1	0	0	1
Soft drink								
Friends intake	[17]		+		1	1	0	0
Economic								
Fruit, Juice, Vegetables								
Area deprivation index (most deprived)	[22F]		-	[22M]				
Snack, Fast food								
Area deprivation index (most deprived)	[22M/F]		+		2	2	0	0
Physical								
Energy								
Non-metropolitan residence	[41]		+	[23]	2	1	0	1
Fat (total fat, en% fat)								
Non-metropolitan residence	[41]		+	[23]	2	1	0	1
Snack, Fast food								
Region (southern USA vs. other)	[20]		+		1	1	0	0
Urbanization				[20]	1	0	0	1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

† Associations between environmental factors and dietary outcomes were coded '+' for a positive association, '-' for an inverse association

* If in one study, a determinant is examined in relation to two outcomes (e.g. fruit intake and vegetable intake), and the results differ for the two outcomes (e.g. a positive association was found for fruit intake, and no association was found for vegetable intake), the study is counted once in the column '# of samples, and twice in the summary column.

** Parental control over child's fat intake

\$ Parental control over child's food intake

Italic printed text, indicate factors with consistent associations.

1 *Environmental correlates of fat intake*

2 At the household environmental level, physical factors (one study/sample), socio-cultural
3 factors (four studies, five samples), and economic factors (seven studies/samples) were
4 examined in relation to fat intake. One study examined factors in the school environment,
5 no studies examined factors in the neighborhood environment, and two studies examined
6 factors at the city/municipality level. At the household socio-cultural level a positive as-
7 sociation was found for parental fat intake (three out of three samples) [27, 43] , and parental
8 control over food intake (two out of two samples) [27, 36]. Parental education was inversely
9 associated with fat intake in two out of two samples [23, 39].

10

11 *Environmental correlates of fruit and vegetable intake*

12 At the household environmental level, physical factors (six studies, eleven samples), socio-
13 cultural factors (twelve studies, 14 samples), and economic factors (six studies/samples)
14 were examined in relation to fruit/vegetable intake. One study examined factors in the
15 school environment, one study (two samples) examined factors in the neighborhood envi-
16 ronment, and no studies examined factors at the city/municipality level. At the household
17 physical level, availability of fruit/vegetables was associated with higher fruit/vegetable
18 intake in four out of seven samples [25, 28, 30, 34]. Accessibility of fruit/vegetables was
19 positively associated with fruit/vegetable intake in four out of six samples [21, 25, 28]. At the
20 household socio-cultural level positive associations were found for modeling (two out of
21 two samples/studies) [21, 28] and parental intake of fruit/vegetables (six out of six samples)
22 [16, 18, 21, 26, 30, 37].

23

24 *Environmental correlates of snack/fast food intake*

25 At the household environmental level, physical factors (one study/sample), socio-cultural
26 factors (five studies, six samples), and economic factors (five studies/samples) were ex-
27 amined in relation to snack/fast food intake. No studies examined factors in the school
28 environment, one study (two samples) examined factors in the neighborhood, and one
29 study examined factors at the city/municipality level. None of the factors examined showed
30 consistent associations with snack/fast food intake.

31

32 *Environmental correlates of soft drink intake*

33 At the household environmental level, physical factors (two studies/samples), household
34 socio-cultural factors (five studies, six samples), and household economic factors (three
35 studies/samples) were examined in relation to soft drink intake. One study examined fac-
36 tors in the school environment, no studies examined factors in the neighborhood or city/
37 municipality environment. At the household socio-cultural level, parental soft drink intake
38 was positively associated with children's soft drink intake in two out of two samples [16, 17].

39

1 Potential environmental correlates of adolescent's dietary behaviors

3 *Environmental correlates of energy intake*

4 At the household environmental level, socio-cultural factors (six studies, nine samples) and
5 economic factors (two studies/samples) were examined as potential correlates of energy
6 intake (Table 3.3). No studies examined factors in the school, neighborhood or city/mu-
7 nicipality environment. At the household socio-cultural level parental energy intake was
8 positively associated with adolescent's energy intake (six out of six samples) [46, 54, 56, 73].
9 A positive association was also found for sibling intake (four out of four samples) [54, 73].

11 *Environmental correlates of fat intake*

12 At the household environmental level, nine studies (15 samples) examined socio-cultural
13 factors and five studies (eight samples) examined economic factors as potential correlates
14 of fat intake. One study examined factors in the school environment, no studies examined
15 factors in the neighborhood. One study (two samples) examined factors at the city/mu-
16 nicipality level. At the household socio-cultural level parental fat intake was found to be
17 positively associated with adolescent's fat intake (eight out of nine samples) [51, 54, 56, 68,
18 73]. A positive association was also found for sibling intake (four out of four samples) [54,
19 73].

21 *Environmental correlates of fruit and vegetable intake*

22 At the household environmental level, physical factors (two studies/samples), socio-cultural
23 factors (ten studies, eleven samples) and economic factors (eight studies, ten samples) were
24 examined as potential correlates of fruit/vegetable intake. One study examined factors in
25 the school environment, one study examined factors in the neighborhood environment, and
26 one study (two samples) examined factors at the city/municipality level. At the household
27 socio-cultural level an authoritative parenting style was positively associated with fruit/
28 vegetable intake (two out of two samples) [63, 65]. Family connectedness was positively
29 associated with adolescent fruit/vegetable intake (two out of two samples) [53, 58]. At the
30 household economic level parent educational level was found to be positively associated
31 with fruit/vegetable intake (six out of six samples) [50, 52, 65, 69, 70].

33 *Environmental correlates of snacks/fast food intake*

34 At the household environmental level, socio-cultural factors (four studies, six samples) and
35 economic factors (three studies, five samples) were studied in relation to snack and fast food
36 intake. One study examined factors in the school environment, no studies examined factors
37 in the neighborhood environment, and one study (two samples) examined factors at the
38 city/municipality level. None of the factors examined showed consistent associations with
39 snack/fast food intake.

Table 3.3 Summary of micro and macro environmental correlates of dietary intakes among adolescents (>12 to 18 year olds)

Correlate	Related to dietary behaviors References	Assoc (+ or -)†	Unrelated to dietary behaviors References	# Samples	Summary (n)
					+ -
Physical					0
Fruit, Juice, Vegetables					
Availability	[66]	+	[61M]	2	1 0 1
Socio-Cultural					
Energy					
Parental intake	[46, 54M/F, 56, 73M/F]	+		6	6 0 0
Siblings intake	[54M/F, 73M/F]	+		4	4 0 0
Single mother family	[45M]	+	[45F]	2	1 0 1
Frequency of family meals	[67]	+		1	1 0 0
Fat (total fat, en% fat)					
Parental intake	[51M, 54M/F, 56, 68F, 73M/F] [51Fmother]	+	[51F]father, 68M]	9	8* 0 2*
Siblings intake	[54M/F, 73M/F]	+		4	4 0 0
Friends intake	[56]			1	0 0 1
Communication strategies	[60]			1	0 0 1
Family food routines	[60]			1	0 0 1
Food rules	[60]			1	0 0 1
Frequency of family meals	[67]			1	0 0 1
Shopping:					
Healthy food is asked for	[60]	-		1	0 1 0
Food asked for is bought	[60]			1	0 0 1
Shopping alone / family	[60]			1	0 0 1
Single mother family	[45M/F]			2	0 0 2
Head of household status (male vs. other)	[47M] [47F]	+		2	1 1 0
Fruit, Juice, Vegetables					
# of evening meals eaten with parent present	[69]	+		1	1 0 0
Frequency of family meals	[67]	+		1	1 0 0
Breakfast with family	[72M/F]			2	- 0 2
Dinner with family	[72Ffruit]	+	[72M/Fveg] [72Mfruit]	2	1* 0 3*

Home/household

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

Correlate	Related to dietary behaviors		Unrelated to dietary behaviors	# Samples	Summary (n)	
	References	Assoc (+ or -)†			References	+
Shopping:						
Healthy food is asked for	[60fruit]	+	[60veg]	1	1*	0 1*
Food asked for is bought	[60fruit/veg]	-		1	0	2* 0
Shopping alone/ family			[60fruit/veg]	1	0 0	2* 2*
Negative communication strategies	[60veg]	-	[60fruit]	1	0	1* 1*
Family food routines			[60]	1	0 0	0 1
Food rules			[60]	1	0 0	0 1
Parent present during leave/return from school			[69]	1	0 0	0 1
Parental control on food choice			[69]	1	0 0	0 1
Parenting style (authoritative, indulgent vs. authoritative, neglective)	[63, 65authoritative]	+		2	2	0 0
Residence other than with family:						
J, F in summer	[48]	+		1	1	0 0
F in winter			[48]	1	0 0	0 1
Family connectedness (high vs. mod/low)	[53, 58]	+		2	2	0 0
Positive relation with parents	[62]	+		1	1	0 0
Positive relation with peers						
F&V intake at 15 years	[62]	+		1	1	0 0
F&V intake at 21 years			[62]	1	0	0 1
Snacks, Fast food						
Parental and friends' intake of foods	[56]	+		1	1	0 0
Frequency of family meals, breakfast/dinner with family			[67, 72M/F]	3	0 0	0 3
Breakfast, lunch, dinner at home vs. school			[49I/II]	2	0	0 2
Breakfast at other site than home or school	[49I]	+		1	1	0 0
Lunch at other site than home or school	[49I/II]	+		2	2	0 0
Breakfast/dinner at other site than home or school			[49II]	1	0	0 1
Dinner at other site than home			[49I]	1	0	0 1
Soft drink						
Parental intake	[55]	+		1	1	0 0

Correlate	Related to dietary behaviors		Assoc (+ or -)†	Unrelated to dietary behaviors		Summary (n)
	References			References		
Friends intake	[55]		+			0
Frequency of family meals	[67]		-			0
Breakfast with family	[72M]		-	[72F]		1
Dinner with family	[48]		-	[72M/F]		2
Residence other than with family			-			0
Shopping:						
Healthy food is asked for				[60]		1
Food asked for is bought				[60]		1
Shopping alone / family				[60]		1
Communication strategies				[60]		1
Family food routines				[60]		1
Food rules	[60]		-			0
Economic						
Energy						
Family income				[70]		1
Fathers occupation				[50]		1
Parental education				[50, 70]		2
Fat (total fat, en% fat)						
Family income / household income				[47M/F, 70]		3
Fathers occupation	[50]		-			0
Parental education	[70]		-	[50, 71M/F]		3
Socio-economic index				[57M/F]		2
Household size				[47M/F]		2
Fruit, Juice, Vegetables						
Family income				[52M/F, 61M, 70]		4
Fathers occupation				[50]		1
Parental education				[50veg]		6
Socio-economic index / SES				[57Mfruit/Fveg]		3
Snacks, Fast food						
Household income				[52M/F]		2
Education of responsible parent	[52F]		-	[52M]		1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

Correlate	Related to dietary behaviors		Assoc (+ or -)†	Unrelated to dietary behaviors		Summary (n)
	References			References	# Samples	
Socio-economic index				[57M/F]	2	0 0 0 2
Pocket money > \$20/week	[59]		+		1	1 0 0 0
Physical						
Fat (total fat, en% fat)						
A la carte program	[64]		+		1	1 0 0 0
Snack vending				[64]	1	0 0 0 1
Fruit, Juice, Vegetables						
A la carte program	[64fruit] [64fruit&veg]		-	[64veg]	1	0 0 2* 1*
Snack vending	[64fruit]		-	[64veg] [64fruit&veg]	1	0 0 1* 2*
Beverage vending				[64veg] [64fruit] [64fruit&veg]	1	0 0 0 3*
Socio-Cultural						
Snacks, Fast food						
School social environment				[59]	1	0 0 0 1
Teacher support						
Highly supportive – fairly unsupportive				[59]	1	0 0 0 1
Highly unsupportive	[59]		+		1	1 0 0 0
Physical						
Fruit, Juice, Vegetables						
Availability in restaurant menu				[61Mfruit]	1	3* 0 0 1*
Availability in grocery stores				[61M]	1	0 0 0 1
Physical						
Fat (total fat, en% fat)						
Geographic region (southern USA vs. other)	[47F]		-	[47M]	2	0 1 1 1
Degree of urbanization				[47M/F]	2	0 0 0 2
Fruit, Juice, Vegetables						
City vs. county	[57M/F]		+		2	2 0 0 0
Snacks, Fast food						
Geographic region (west, Midwest, south)	[49II]		-	[49I]	2	0 1 1 1
Soft drink						
City vs. County				[57M/F]	2	0 0 0 2

† Associations between environmental factors and dietary outcomes were coded '+', for a positive association, '-' for an inverse association
 * If in one study, a determinant is examined in relation to two outcomes (e.g. fruit intake and vegetable intake), and the results differ for the two outcomes (e.g. a positive association was found for fruit intake, and no association was found for vegetable intake), the study is counted once in the column '# of samples', and twice in the summary column.

Italic printed text, indicate factors with consistent associations.

Table 3.4 ANGELO framework with the number of studies and the associations found in each cell.

	Physical		Socio-cultural		Economic		Political	
	# studies	Assoc.	# studies	Assoc.	# studies	Assoc.	# studies	Assoc.
Home/ household								
Energy	1		11	Encouragement/ assistance (-;c) Parental intake (+;a)** Sibling intake (+;a)	7			
Fat	1		13	Control over intake (+;c) Parental intake (+;c) Parental intake (+;a) Sibling intake (+;a)	12	Parental education (+;c)		
Fruit/ vegetables	8	Availability (+;c)* Accessibility (+;c)	16	Modeling parents (+;c) Parental intake (+;c) Parenting style (+;a) Family connectedness (+;a)	12	Parental education (+;a)		
Snack/fast food	1		9		8			
Soft drink	2		10	Parental intake (+;c)	3			
Educational institutions								
Energy	1		1					
Fat	2		1					
Fruit/ vegetables	1		1					
Snack/fast food			1					
Soft drink	1		1					
Neighborhood								
Energy								
Fat								
Fruit/ vegetables	1				1			
Snack/fast food					1			
Soft drink								
City/ municipality								
Energy	2							
Fat	3							
Fruit/ vegetables	1							
Snack/fast food	2							
Soft drink	1							

c Positive association found for children

a Positive association found for adolescents

1 *Environmental correlates of soft-drink intake*

2 At the household environment level, socio-cultural factors (five studies, six samples) were
3 examined as potential environmental correlates of soft drink intake. No studies examined
4 factors in the school and neighborhood environment. One study (two samples) examined
5 factors at the city/municipality level. None of the factors examined showed consistent as-
6 sociations with soft drink intake.

7 8 **Summary in ANGELO framework**

9
10 Socio-cultural factors on the household level are the most studied environmental factors
11 for all dietary behaviors, followed by economic factors on the household level (Table 3.4).
12 Factors studied on the school environmental level (physical and socio-cultural) were mostly
13 single studies. At the city/municipality level only physical factors were studied.

14 15 16 **DISCUSSION**

17
18 The present systematic review of the literature on environmental correlates of energy, fat,
19 fruit, vegetable, snack/fast food, and soft drink intake in children and adolescents showed
20 that household socio-cultural factors (e.g. parental and sibling intake, parenting practices)
21 and household economic factors (e.g. household income, parent educational level) were
22 studied most extensively as potential environmental determinants. Few studies examined
23 the influence of physical environmental factors, few looked at environmental factors in
24 schools, neighborhoods and city/municipality, and none looked at political factors. This
25 review showed consistent evidence (findings replicated in multiple studies), for the relation-
26 ship between parental intake and children's fat, fruit and vegetable intake, for parent and
27 sibling intakes with adolescent's energy and fat intake, and for parent educational level with
28 adolescent's fruit and vegetable intake. A positive association was found for the relationship
29 between availability and accessibility with children's fruit and vegetable intake, even though
30 the samples that found a positive association only slightly outnumbered the samples that
31 found no association. Further positive associations were found for controlling/restrictive
32 practices (fat), parent educational level (fat), modeling (fruit/vegetable), parental intake
33 (soft drink) parenting style (fruit/vegetable), family connectedness (fruit/vegetable) and
34 encouragement to increase food intake (fruit/vegetable). A negative association was found
35 for encouragement/assistance/prompts (energy). These factors were examined in only two
36 studies, which limits the possibility to draw firm conclusions regarding consistency of as-
37 sociations. The direction of the association for encouragement/assistance/prompts seems
38 unexpected. However, since these studies were cross-sectional, it could also be that a low
39 child food intake provokes parental encouragement, assistance and prompts to increase in-

1 takes. All other associations studied between dietary behaviors and potential environmental
2 factors were inconsistent, appeared non-existent, or were not replicated.

3
4 A major limitation of the currently available published research papers is that many po-
5 tential environmental determinants have been examined for a variety of dietary behaviors,
6 but that only few studies have been conducted on the same specific environmental factor
7 – dietary behavior combination. Replication of studies on such combinations is necessary,
8 to generate more compelling evidence for associations between environmental factors
9 and dietary intake. With regard to the strength of the study designs, most of the included
10 studies were cross-sectional, making conclusions about direction and possible causality of
11 associations impossible. Furthermore, most studies relied on self-reported data, of which
12 the validity and reliability of the instruments used was hard to judge, since this information
13 was not reported in the majority of the studies. We retrieved few studies that used objec-
14 tive observation instruments to assess factors in the physical environment or to measure
15 the behavioral outcome. The behavioral outcome measures in the studies included, may be
16 somewhat biased because the studies mostly relied on self-reports.

17
18 There are some issues and limitations that have to be taken into account in interpreting
19 the results of the review. In order to summarize the findings of the studies we collapsed
20 conceptually similar environmental determinants into one category, although potential
21 determinants in the same category were often dissimilar or measured in different ways.
22 Our search strategy only included studies that were published in English in peer-reviewed
23 journals and referenced in electronic databases; therefore our findings may be influenced by
24 a publication bias. However, the high number of non-associations reported in the included
25 studies may indicate that a bias towards publication of significant results only, was not very
26 strong. The studies included were heterogeneous in the conceptualization, measurement
27 of the environmental determinant and/or dietary intakes, samples and analyses used, and
28 therefore it was not possible to assess the overall strength of associations. Finally, we in-
29 cluded multiple environmental factors examined in one study in the review, and it must be
30 kept in mind that these associations are not independent.

31
32 Previously published reviews on the associations of environmental factors regarding eat-
33 ing behaviors in children and adolescents were narrative as opposed to systematic reviews
34 [7-9, 11, 74, 75]. The main conclusions from those reviews were that the role of parents is
35 particularly important, that parents should create supportive food environments for their
36 children [7-9, 74], and that school food environments may have a large impact on food
37 choices [9, 11, 74]. In the present review we indeed found that parental intake and to a lesser
38 extent availability and accessibility were associated with intakes in adolescents and children.
39 Furthermore, some evidence (examined in two studies) was found for a positive association

1 between an authoritative parenting style and adolescent's fruit and vegetable intake, and for
2 specific parenting practices and children's energy and fat intakes. Only very few studies on
3 peer influences were retrieved in this systematic review. The importance of the school food
4 environment can also not be substantiated with the evidence from the studies included in
5 the present review.

6
7 We identified several gaps in the currently available evidence of relationships between en-
8 vironmental factors and child and adolescent dietary intakes. First, we were able to identify
9 very few studies examining associations between micro environmental factors in school and
10 neighborhood settings, and on macro environmental factors in city/municipality settings:
11 the broader, more anonymous infrastructure that may support or hinder health behaviors.
12 It must, however, be noted that there are studies available that examine the effects of adver-
13 tising and marketing on eating behaviors of youngsters [10, 76], but since these studies are
14 mostly intervention studies, these were not included in the review. Secondly, the studies
15 mainly depended on perceived and self-reported environmental information, as opposed
16 to more objective observations of environments. Objectively assessing characteristics of the
17 environment (observations or Geographic Information System), is a topic of recent interest
18 [77, 78]. Furthermore, we retrieved only a limited number of studies assessing environ-
19 mental determinants of snack and soft drink intakes, while these two behaviors may be of
20 specific importance in obesity prevention [4, 12]. An important reason for some of the gaps
21 may be that attention to the role of the physical environment is of recent interest, and many
22 studies that examine possible influences of the physical environment may be underway.

23
24 The current evidence of associations between environmental determinants and dietary
25 intakes among children and adolescents suggests that parental intakes, sibling intakes and
26 educational level of parents are environmental determinants most consistently associated
27 with intakes. A less consistent repeated but positive association was found for availability
28 and accessibility on child fruit and vegetable intake. The finding that parental behavior
29 is associated with child and adolescent intakes implies that interventions should take the
30 behavior of parents into account, or desensitize adolescents for the (unfavorable) behavior
31 of their parents. Parents should be more strongly encouraged to give the right example,
32 especially where fat and energy intakes are concerned. Fruit and vegetable promotion
33 should focus especially on adolescents from parents with lower levels of education. To get
34 a broad understanding of the influence of environmental factors associated with obesity
35 inducing behaviors in children and adolescents at the various levels distinguished in the
36 ANGELO framework, studies are needed that target the environmental levels and factors
37 that have found to be (nearly) empty in the ANGELO grid (Table 3.4), such as physical,
38 socio-cultural, economic and political factors in the school (e.g. school food policy and
39 food prices), neighborhood (e.g. availability and accessibility of foods in shops) and city/

1 municipality environment (e.g. food policy, food prices, marketing). Furthermore factors
2 such as availability and accessibility at home, school and neighborhood should be studied in
3 relation to energy, fat, soft drink, snacks and fast food intake. For all environmental factors,
4 including the factors that have already been studied, there is a need for longitudinal studies
5 in which valid or objective measurement instruments are used.

8 ACKNOWLEDGEMENTS

9
10 The authors wish to thank Carlijn Kamphuis and Gert Jan de Bruijn for their assistance in
11 the review process.

12 This study was financially supported by a grant from ZonMw The Netherlands Organiza-
13 tion for Health Research and Development. Dr. Katrina Giskes is supported by an Australian
14 National Health and Medical Research Council Sidney Sax Fellowship (grant ID number:
15 290540).

1 REFERENCES

- 2 1. Lobstein T, Frelut ML: Prevalence of overweight among children in Europe. *Obes Rev* 2003, 4(4):195-
- 3 200.
- 4 2. Hedley AA, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM: Prevalence of overweight
- 5 and obesity among US children, adolescents, and adults, 1999-2002. *JAMA* 2004, 291(23):2847-2850.
- 6 3. French SA, Story M, Jeffery RW: Environmental influences on eating and physical activity. *Annu Rev*
- 7 *Public Health* 2001, 22:309-335.
- 8 4. Swinburn BA, Caterson I, Seidell JC, James WP: Diet, nutrition and the prevention of excess weight
- 9 gain and obesity. *Public Health Nutr* 2004, 7(1A):123-146.
- 10 5. Sallis JF, Owen N: Ecological models of health behavior. In *Health behavior and health education*
- 11 *Theory, research, and practice*. 3 edition. Edited by Glanz K, Reimer BK, Lewis FM. San Francisco: CA:
- 12 Jossey-Bass; 2002:462-484.
- 13 6. Rothschild ML: Carrots, sticks, and promises: a conceptual framework for the management of public
- 14 health and social issue behaviors. *Journal of Marketing* 1999, 63:24-37.
- 15 7. Patrick H, Nicklas TA: A review of family and social determinants of children's eating patterns and
- 16 diet quality. *J Am Coll Nutr* 2005, 24(2):83-92.
- 17 8. Ritchie LD, Welk G, Styne D, Gerstein DE, Crawford PB: Family environment and pediatric over-
- 18 weight: what is a parent to do? *J Am Diet Assoc* 2005, 105(5 Suppl 1):S70-79.
- 19 9. Story M, Neumark-Sztainer D, French S: Individual and environmental influences on adolescent
- 20 eating behaviors. *J Am Diet Assoc* 2002, 102(3 Suppl):S40-51.
- 21 10. Story M, French S: Food Advertising and Marketing Directed at Children and Adolescents in the US.
- 22 *Int J Behav Nutr Phys Act* 2004, 1(1):3.
- 23 11. Crockett SJ, Sims LS: Environmental-Influences on Childrens Eating. *Journal of Nutrition Education*
- 24 1995, 27(5):235-249.
- 25 12. Rennie KL, Johnson L, Jebb SA: Behavioural determinants of obesity. *Best Pract Res Clin Endocrinol*
- 26 *Metab* 2005, 19(3):343-358.
- 27 13. Pereira MA, Ludwig DS: Dietary fiber and body-weight regulation. Observations and mechanisms.
- 28 *Pediatr Clin North Am* 2001, 48(4):969-980.
- 29 14. Swinburn B, Egger G, Raza F: Dissecting obesogenic environments: the development and application
- 30 of a framework for identifying and prioritizing environmental interventions for obesity. *Prev Med*
- 31 1999, 29(6 Pt 1):563-570.
- 32 15. Cooper H: *Synthesizing research: a guide for literature reviews*. 3rd ed edition. London: Sage; 1998.
- 33 16. Vereecken CA, Keukelier E, Maes L: Influence of mother's educational level on food parenting prac-
- 34 tices and food habits of young children. *Appetite* 2004, 43(1):93-103.
- 35 17. Grimm GC, Harnack L, Story M: Factors associated with soft drink consumption in school-aged
- 36 children. *J Am Diet Assoc* 2004, 104(8):1244-1249.
- 37 18. Cooke LJ, Wardle J, Gibson EL, Sapochnik M, Sheiham A, Lawson M: Demographic, familial and
- 38 trait predictors of fruit and vegetable consumption by pre-school children. *Public Health Nutr* 2004,
- 39 7(2):295-302.
19. Brown R, Ogden J: Children's eating attitudes and behaviour: a study of the modelling and control
- theories of parental influence. *Health Educ Res* 2004, 19(3):261-271.
20. Bowman SA, Gortmaker S, Ebbeling CB, Pereira MA, Ludwig DS: Effects of fast-food consumption
- on energy intake and diet quality among children in a national household survey. *Pediatrics* 2004,
- 113:112-118.

21. Bere E, Klepp KI: Correlates of fruit and vegetable intake among Norwegian schoolchildren: parental and self-reports. *Public Health Nutr* 2004, 7(8):991-998.
22. Wardle J, Jarvis MJ, Steggle N, Sutton S, Williamson S, Farrimond H, Cartwright M, Simon AE: Socioeconomic disparities in cancer-risk behaviors in adolescence: baseline results from the Health and Behaviour in Teenagers Study (HABITS). *Prev Med* 2003, 36(6):721-730.
23. Mazur RE, Marquis GS, Jensen HH: Diet and food insufficiency among Hispanic youths: acculturation and socioeconomic factors in the third National Health and Nutrition Examination Survey. *Am J Clin Nutr* 2003, 78(6):1120-1127.
24. Haapalahti M, Mykkanen H, Tikkanen S, Kokkonen J: Meal patterns and food use in 10-to 11-year-old Finnish children. *Public Health Nutr* 2003, 6(4):365-370.
25. Cullen KW, Baranowski T, Owens E, Marsh T, Rittenberry L, de Moor C: Availability, accessibility, and preferences for fruit, 100% fruit juice, and vegetables influence children's dietary behavior. *Health Educ Behav* 2003, 30(5):615-626.
26. Fisher JO, Mitchell DC, Smiciklas-Wright H, Birch LL: Parental influences on young girls' fruit and vegetable, micronutrient, and fat intakes. *J Am Diet Assoc* 2002, 102(1):58-64.
27. Lee Y, Mitchell DC, Smiciklas-Wright H, Birch LL: Diet quality, nutrient intake, weight status, and feeding environments of girls meeting or exceeding recommendations for total dietary fat of the American Academy of Pediatrics. *Pediatrics* 2001, 107(6):E95.
28. Cullen KW, Baranowski T, Rittenberry L, Cosart C, Hebert D, de Moor C: Child-reported family and peer influences on fruit, juice and vegetable consumption: reliability and validity of measures. *Health Educ Res* 2001, 16(2):187-200.
29. Coon KA, Goldberg J, Rogers BL, Tucker KL: Relationships between use of television during meals and children's food consumption patterns. *Pediatrics* 2001, 107(1):E7.
30. Kratt P, Reynolds K, Shewchuk R: The role of availability as a moderator of family fruit and vegetable consumption. *Health Educ Behav* 2000, 27(4):471-482.
31. Gillman MW, Rifas-Shiman SL, Frazier AL, Rockett HR, Camargo CA, Jr., Field AE, Berkey CS, Colditz GA: Family dinner and diet quality among older children and adolescents. *Arch Fam Med* 2000, 9(3):235-240.
32. Cullen KW, Baranowski T, Rittenberry L, Cosart C, Owens E, Hebert D, de Moor C: Socioenvironmental influences on children's fruit, juice and vegetable consumption as reported by parents: reliability and validity of measures. *Public Health Nutr* 2000, 3(3):345-356.
33. Birch LL, Fisher JO: Mothers' child-feeding practices influence daughters' eating and weight. *Am J Clin Nutr* 2000, 71(5):1054-1061.
34. Reynolds KD, Hinton AW, Shewchuk RM, Hickey CA: Social cognitive model of fruit and vegetable consumption in elementary school children. *J Nutr Educ* 1999, 31(1):23-30.
35. Drucker RR, Hammer LD, Agras WS, Bryson S: Can mothers influence their child's eating behavior? *J Dev Behav Pediatr* 1999, 20(2):88-92.
36. Zive MM, Frank-Spohrer GC, Sallis JF, McKenzie TL, Elder JP, Berry CC, Broyles SL, Nader PR: Determinants of dietary intake in a sample of white and Mexican-American children. *J Am Diet Assoc* 1998, 98(11):1282-1289.
37. Gibson EL, Wardle J, Watts CJ: Fruit and vegetable consumption, nutritional knowledge and beliefs in mothers and children. *Appetite* 1998, 31(2):205-228.
38. Johnson-Down L, O'Loughlin J, Koski KG, Gray-Donald K: High prevalence of obesity in low income and multiethnic schoolchildren: a diet and physical activity assessment. *J Nutr* 1997, 127(12):2310-2315.

- 1 39. Crawford PB, Obarzanek E, Schreiber GB, Barrier P, Goldman S, Frederick MM, Sabry ZI: The effects
2 of race, household income, and parental education on nutrient intakes of 9- and 10-year-old girls.
3 NHLBI Growth and Health Study. *Ann Epidemiol* 1995, 5(5):360-368.
- 4 40. Koivisto UK, Fellenius J, Sjoden PO: Relations between parental mealtime practices and children's
5 food intake. *Appetite* 1994, 22(3):245-257.
- 6 41. Johnson RK, Guthrie H, Smiciklas-Wright H, Wang MQ: Characterizing nutrient intakes of children
7 by sociodemographic factors. *Public Health Rep* 1994, 109(3):414-420.
- 8 42. Wolfe WS, Campbell CC: Food pattern, diet quality, and related characteristics of schoolchildren in
9 New York State. *J Am Diet Assoc* 1993, 93(11):1280-1284.
- 10 43. Oliveria SA, Ellison RC, Moore LL, Gillman MW, Garrahe EJ, Singer MR: Parent-child relationships
11 in nutrient intake: the Framingham Children's Study. *Am J Clin Nutr* 1992, 56(3):593-598.
- 12 44. Johnson RK, Smiciklas-Wright H, Crouter AC, Willits FK: Maternal employment and the quality of
13 young children's diets: empirical evidence based on the 1987-1988 Nationwide Food Consumption
14 Survey. *Pediatrics* 1992, 90(2 Pt 1):245-249.
- 15 45. Darke SJ, Disselduff MM, Try GP: A nutrition survey of children from one-parent families in New-
16 castle upon Tyne in 1970. *Br J Nutr* 1980, 44(3):237-241.
- 17 46. Laskarzewski P, Morrison JA, Khoury P, Kelly K, Glatfelter L, Larsen R, Glueck CJ: Parent-child
18 nutrient intake interrelationships in school children ages 6 to 19: the Princeton School District Study.
19 *Am J Clin Nutr* 1980, 33(11):2350-2355.
- 20 47. Johnson RK, Johnson DG, Wang MQ, Smiciklas-Wright H, Guthrie HA: Characterizing nutrient
21 intakes of adolescents by sociodemographic factors. *J Adolesc Health* 1994, 15(2):149-154.
- 22 48. Sweeting H, Anderson A, West P: Socio-demographic correlates of dietary habits in mid to late
23 adolescence. *Eur J Clin Nutr* 1994, 48(10):736-748.
- 24 49. Dausch JG, Story M, Dresser C, Gilbert GG, Portnoy B, Kahle LL: Correlates of high-fat/low-
25 nutrient-dense snack consumption among adolescents: results from two national health surveys. *Am*
26 *J Health Promot* 1995, 10(2):85-88.
- 27 50. Laitinen S, Rasanen L, Viikari J, Akerblom HK: Diet of Finnish children in relation to the family's
28 socio-economic status. *Scand J Soc Med* 1995, 23(2):88-94.
- 29 51. De Bourdeaudhuij I: Resemblance in health behaviours between family members. *Arch Public Health*
30 1996, 54:251-273.
- 31 52. Lowry R, Kann L, Collins JL, Kolbe LJ: The effect of socioeconomic status on chronic disease risk
32 behaviors among US adolescents. *JAMA* 1996, 276(10):792-797.
- 33 53. Neumark-Sztainer D, Story M, Resnick MD, Blum RW: Correlates of inadequate fruit and vegetable
34 consumption among adolescents. *Prev Med* 1996, 25(5):497-505.
- 35 54. Vauthier JM, Lluch A, Lecomte E, Artur Y, Herbeth B: Family resemblance in energy and macronutri-
36 ent intakes: the Stanislas Family Study. *Int J Epidemiol* 1996, 25(5):1030-1037.
- 37 55. Woodward DR, Boon JA, Cumming FJ, Ball PJ, Williams HM, Hornsby H: Adolescents' reported
38 usage of selected foods in relation to their perceptions and social norms for those foods. *Appetite*
39 1996, 27(2):109-117.
56. Feunekes GI, de Graaf C, Meyboom S, van Staveren WA: Food choice and fat intake of adolescents
and adults: associations of intakes within social networks. *Prev Med* 1998, 27(5 Pt 1):645-656.
57. Høglund D, Samuelson G, Mark A: Food habits in Swedish adolescents in relation to socioeconomic
conditions. *Eur J Clin Nutr* 1998, 52(11):784-789.
58. Story M, Neumark Sztainer D, Resnick MD, Blum RW: Psychosocial factors and health behaviors
associated with inadequate fruit and vegetable intake among American-Indian and Alaska-Native
adolescents. *J Nutr Educ* 1998, 30:100-106.

59. McLellan L, Rissel C, Donnelly N, Bauman A: Health behaviour and the school environment in New South Wales, Australia. *Soc Sci Med* 1999, 49(5):611-619.
60. De Bourdeaudhuij I, Van Oost P: Personal and family determinants of dietary behaviour in adolescents and their parents. *Psychology and Health* 2000, 15:751-770.
61. Edmonds J, Baranowski T, Baranowski J, Cullen KW, Myres D: Ecological and socioeconomic correlates of fruit, juice, and vegetable consumption among African-American boys. *Prev Med* 2001, 32(6):476-481.
62. Lien N, Jacobs DR, Jr., Klepp KI: Exploring predictors of eating behaviour among adolescents by gender and socio-economic status. *Public Health Nutr* 2002, 5(5):671-681.
63. Kremers SP, Brug J, de Vries H, Engels RC: Parenting style and adolescent fruit consumption. *Appetite* 2003, 41(1):43-50.
64. Kubik MY, Lytle LA, Hannan PJ, Perry CL, Story M: The Association of the School Food Environment With Dietary Behaviors of Young Adolescents. *Am J Public Health* 2003, 93(7):1168-1173.
65. Lytle LA, Varnell S, Murray DM, Story M, Perry C, Birnbaum AS, Kubik MY: Predicting adolescents' intake of fruits and vegetables. *J Nutr Educ Behav* 2003, 35(4):170-178.
66. Neumark Sztainer D, Wall M, Perry C, Story M: Correlates of fruit and vegetable intake among adolescents: Findings from Project EAT. *Prev Med* 2003, 37(3):198-208.
67. Neumark-Sztainer D, Hannan PJ, Story M, Croll J, Perry C: Family meal patterns: associations with sociodemographic characteristics and improved dietary intake among adolescents. *J Am Diet Assoc* 2003, 103(3):317-322.
68. Stanton CA, Fries EA, Danish SJ: Racial and gender differences in the diets of rural youth and their mothers. *Am J Health Behav* 2003, 27(4):336-347.
69. Videon TM, Manning CK: Influences on adolescent eating patterns: The importance of family meals. *J Adolesc Health* 2003, 32(5):365-373.
70. Xie B, Gilliland FD, Li YF, Rockett HRH: Effects of ethnicity, family income, and education on dietary intake among adolescents. *Prev Med* 2003, 36(1):30-40.
71. Tur JA, Puig MS, Benito E, Pons A: Associations between sociodemographic and lifestyle factors and dietary quality among adolescents in Palma de Mallorca. *Nutrition* 2004, 20(6):502-508.
72. Kusano-Tsunoh A, Nakatsuka H, Satoh H, Shimizu H, Sato S, Ito I, Fukao A, Hisamichi S: Effects of family-togetherness on the food selection by primary and junior high school students: family-togetherness means better food. *Tohoku J Exp Med* 2001, 194(2):121-127.
73. Perusse L, Tremblay A, Leblanc C, Cloninger CR, Reich T, Rice J, Bouchard C: Familial resemblance in energy intake: contribution of genetic and environmental factors. *Am J Clin Nutr* 1988, 47(4):629-635.
74. Jenkins S, Horner SD: Barriers that influence eating behaviors in adolescents. *J Pediatr Nurs* 2005, 20(4):258-267.
75. Wardle J: Parental influences on children's diets. *Proc Nutr Soc* 1995, 54(3):747-758.
76. Young B, Hetherington M: The literature on advertising and children's food choice. *Nutrition & Food Science* 1996, 96(5):15-18.
77. Richter KP, Harris KJ, Paine-Andrews A, Fawcett SB, Schmitd TL, Lankenau BH, Johnston J: Measuring the health environment for physical activity and nutrition among youth: a review of the literature and applications for community initiatives. *Prev Med* 2000, 31:S98-S111.
78. Glanz K, Sallis JF, Saelens BE, Frank LD: Healthy nutrition environments: concepts and measures. *Am J Health Promot* 2005, 19(5):330-333, ii.

4 Environmental correlates of physical activity in youth – a review and update

Ferreira I, van der Horst K, Wendel-Vos W, Kremers S, van Lenthe FJ, Brug J. Environmental correlates of physical activity in youth – a review and update.

Obesity Reviews 2007, 8(2): 129-154.

1 **ABSTRACT**

2
3 **Background:** Obesogenic environments are thought to underlie the increased obesity
4 prevalence observed in youth during the past decades. Understanding the environmental
5 factors that are associated with physical activity (PA) in youth is needed to better inform
6 the development of effective intervention strategies attempting to halt the obesity epidemic.

7 **Methods:** We conducted a systematic semi-quantitative review of 150 studies on environ-
8 mental correlates of youth PA published in the past 25 years. The ANalysis Grid for Environ-
9 ments Linked to Obesity (ANGELO) framework was used to classify the environmental
10 correlates studied.

11 **Results:** Most studies retrieved used cross-sectional designs and subjective measures of
12 environmental factors and PA. Especially variables of the home and school environments
13 were associated with PA in youth. Most consistent positive correlates of PA were father's PA,
14 school PA-related policies (in children), and support from significant others, mother's edu-
15 cation level, family income, and non-vocational school attendance (in adolescents). Time
16 spent outdoors (in children) and low crime incidence (in adolescents) were characteristics
17 of the neighborhood environment associated with higher PA. Convincing evidence of an
18 important role for many other environmental factors was however not found.

19 **Discussion:** Further research should aim at longitudinal study designs and use more objec-
20 tive measures of PA and its potential (environmental) determinants.

21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

1 INTRODUCTION

2
3 Physical activity (PA) is a health enhancing behavior: when practiced regularly, PA reduces
4 the risk for a range of chronic disease [1-4]. Also among the young current and future health
5 benefits can be obtained through engaging in physically active lifestyles [5]: it helps building
6 strong bones, healthy joints, a strong heart, a good mental health and prevents today's major
7 public health concern – obesity[6-9]. Despite these health benefits, many young people are
8 not engaging in recommended levels of PA [10-12]. In addition, longitudinal studies have
9 shown that a steep decrease in PA levels occurs during adolescence [13-15] and that PA levels
10 established in youth tend to track into adulthood[16-20]. PA promotion in youth is thought
11 to facilitate a carryover of healthful habits into adulthood and a lifelong protection from
12 other risk factors, and is therefore a priority in current public health policies [4, 21].

13 Given the short time frame in which the obesity prevalence has increased to epidemic
14 scales many scientists postulate that this is more likely due to changes in environments
15 than in biology [22-26]. In this vein, recent studies have indeed demonstrated associations
16 between childhood obesity and environmental features, namely at the home and neighbor-
17 hood [27-32]. Consequently, it is important to understand, measure and alter environments
18 that promote or hinder obesity-inducing *behaviors*, such as low physical activity [7, 33-38].
19 Environmental influences can be especially relevant to children and adolescents since they
20 have less autonomy in their behavioral choices [39]. Specific recommendations for research
21 on the determinants of PA in youth have emphasized the need to examine environmental
22 influences on youth PA at different levels (e.g. home, neighborhood, school)[40-42] to bet-
23 ter inform the development of interventions attempting to improve PA levels [43, 44].

24 Now that more and more studies focus on potential environmental influences on
25 children's and adolescents' PA behavior, it is important to get a detailed overview of the
26 evidence these studies have provided so far, to define a research agenda in this area. In
27 the year 2000, a comprehensive review of personal and environmental correlates of PA in
28 children and adolescents[45] identified several variables which were consistently associated
29 with children/adolescent's PA levels, including social and physical environmental factors
30 such as direct help and support from parents and significant others, access to programs/
31 facilities, opportunities to be active and time spent outdoors. We now update the review
32 of evidence provided by Sallis *et al.* focusing specifically on, and characterizing into more
33 detail, the *environmental* correlates of PA in children and adolescents.

34
35
36
37
38
39

1 METHODS

3 Search strategies and procedures

5 Relevant studies were located from 2 main sources. Firstly, the computerized literature
6 databases MedLine (PubMed), PsychInfo, Web of Science, EMBASE and SportDiscuss were
7 searched. The following keyword combinations were used: physical activity, physical active
8 lifestyle, vigorous activity, leisure activities, recreation, exercise, sport(s), motor activity,
9 physical education, walking, running, (bi)cycling, commuting, determinants, correlates,
10 influences, associations, environment, physical environment, built environment, psychoso-
11 cial determinants, social environment, social norms, socio-economic status, socio-cultural
12 environment, parents, peers, neighborhood, school, facilities, recreation, equipment, safety.
13 These searches were restricted to studies performed in humans aged up to 18 years, and
14 published between January 1980 and December 2004. After excluding duplicate studies,
15 over 5,000 articles were hereby identified. Two independent reviewers (IF, KvdH) screened
16 and selected the articles retrieved whenever it could be ascertained first, from the title (304
17 articles), second from the abstract (88 articles), and finally from the full text (84 articles),
18 that the selection criteria (see below) were met. These stepwise analyses were performed
19 separately by each reviewer, and at each step an article was kept whenever selected by at
20 least one of the reviewers.

21 Secondly, manual searches using the reference of the previous systematic review from
22 Sallis *et al.*, [45] primary studies located from the previous source and our personal databases
23 were performed and cross-checked with the articles found through the previous source.
24 This led to the inclusion of 66 additional articles. Together, these search strategies resulted
25 in a total of 150 articles, which are reviewed herein.

27 Inclusion/exclusion criteria

29 *Types of studies*

30 The present review was concerned with PA levels occurring ‘naturally’ in populations of
31 children and adolescents. Therefore, only observational studies (either cross-sectional or
32 longitudinal) were included, whereas studies investigating samples of PA-related interven-
33 tions or with a quasi-experimental design were excluded (with exception of studies report-
34 ing on baseline data from intervention studies). Qualitative studies or studies that were
35 solely descriptive in nature (i.e., reporting only frequency data), abstracts, case reports,
36 expert opinions, dissertations and unpublished data were also excluded.

1 *Participants and country*

2 Subjects (or the majority of the participants) had to be in the age range of 3-18 years old;
3 similarly to the review by Sallis et al., we have divided studies among children (i.e., 3-12 years
4 old) and adolescents (>12-18 years old). Studies on children and adolescents with chronic
5 diseases (that may affect PA levels) or children participating in top-level competitive sports
6 were not included. Only studies from samples drawn in countries with established market
7 economies (as defined by the World Bank) and published in English as papers in interna-
8 tional peer-reviewed journals were included.

9

10 *The dependent variable(s) - PA*

11 The dependent variable was any measure of (overall) PA of various types (i.e., play, games,
12 sports, work, transportation, recreation, physical education, or planned exercise) performed
13 in the context of home/family, school and community, and expressed in terms of duration
14 (e.g. in minutes), or frequency (e.g. times per week), or intensity (e.g. vigorous) or a combi-
15 nation of these, i.e. in terms of volume (e.g. METs or Kcal) [46]. When studies had multiple
16 dependent measures of PA; the correlates of mutually exclusive outcomes (e.g. habitual
17 levels of moderate- and vigorous-intensity PA) were investigated and reported separately.

18 Studies in which the dependent variable was aerobic fitness, intention, self-efficacy, or
19 other intermediate (non-behavioral) measures were not included; physical inactivity/
20 sedentary behavior was not considered as outcome because PA and inactivity are distinct
21 behaviors, often unrelated and with distinct determinants [47-51]. In addition, although
22 we acknowledge physical inactivity as an important health-impairing behavior, a recent
23 systematic review of its determinants among youth has been published recently [52].

24

25 *The predictor variable(s) - environmental characteristics*

26 Environmental variables were broadly defined as 'anything outside the individual that can
27 affect its PA behavior'. To structure our review we were in need of a conceptual framework
28 to categorize the various environmental factors studied. Different classifications of possible
29 environmental determinants of health behaviors have been proposed [36, 38, 50, 53, 54], all
30 of them showing great overlap and similarities. In the present review we have adopted The
31 ANalyses Grid for Environments Linked to Obesity (ANGELO) conceptual framework [55]
32 to classify potential environmental determinants of PA in children and adolescents. This
33 framework was specifically developed to conceptualize 'obesogenic' environments (i.e. those
34 that promote excessive energy intake and low PA), enabling the identification of specific
35 areas and settings to be targeted by intervention programs. Specifically, environmental vari-
36 ables can be distinguished within two 'sizes' (micro and macro) and four types (physical,
37 socio-cultural, economic and political) of environment. Micro-environments are defined
38 as environmental settings where groups of people meet and gather. Such settings are often
39 geographically distinct and allow direct mutual influences between individuals and the

1 environment. Examples of micro-environmental settings are homes, schools, and neighbor-
2 hoods. Macro-environments, on the other hand, include the broader, more anonymous in-
3 frastructure that may support or hinder health behaviors. Examples of macro-environments
4 are the town planning, the transport infrastructure, the media and the health-care system.
5 All studies reviewed herein were required to examine at least one environmental variable
6 (the independent variable), and this variable needed to be tested for its association with a
7 measure of PA (the dependent variable), obtained at the individual level.

9 Data analyses

11 Due to the great variety of variables and methods drawn from diverse samples, a meta-
12 analytical review was not possible to conduct. We have therefore adopted the same semi-
13 quantitative approach outlined by Sallis *et al.*, [45] recently also used by Gorely *et al.* [52], in
14 a review of the correlates television viewing among youth. An independent sample, i.e. the
15 smallest independent sub-sample (based on age and gender) for which relevant data was
16 reported (e.g. studies reporting findings for boys (M) and girls (F) separately, provide 2
17 independent samples) was used as the unit of analyses [56].

19 *Study characteristics*

20 The relevant characteristics from all the selected publications listed in the Bibliography sec-
21 tion were retrieved and registered in detailed tables (which are available upon request from
22 the corresponding author), according to current review guidelines [57, 58]. This extensive
23 information was then summarized in one background table (Table 4.1).

25 *Categorization of variables*

26 Correlates of PA investigated in the studies reviewed were categorized in the ANGELO grid,
27 i.e. were grouped in 4 environment types (physical, socio-cultural, economic, and political)
28 for each environmental setting (Micro and Macro) with a further distinction in specific
29 levels (home, educational institution, neighborhood, city/municipality, region). These data
30 was then summarized in two tables providing an overview of the potential determinants of
31 PA of children and adolescents separately (Tables 4.2 and 4.3, respectively).

33 *Coding and summarizing associations with PA*

34 A variety of statistical techniques (e.g. correlations, t-tests, linear or logistic regression
35 analyses, ANOVA and structured equation models) were used to evaluate the associa-
36 tions. Most studies not only reported univariate but also multivariate analyses (e.g., with
37 adjustment for demographic and/or other potential correlates investigated); whenever
38 possible findings reported here were those from the fully adjusted models. As with regard
39 to prospective studies, the associations found within the shortest follow-up period were

1 the ones considered, and the cross-sectional findings embedded within these studies were
2 disregarded.

3 Studies reporting positive (coded as '+') or inverse (coded as '-') association(s) between
4 the independent variable and PA were registered under the column '*related to PA*'; non-sig-
5 nificant associations were coded under the column '*unrelated to PA*' (coded as 'o'). Findings
6 for each independent variable were summarized by adding the number of associations in a
7 given direction (+, -, o); a final summary association code for each correlate examined was
8 derived as follows: $\geq 60\%$ of the associations in any direction was considered evidence for
9 a positive (summary code '+'), negative (summary code '-') or non-association (summary
10 code 'o'); a mixed pattern of associations $< 60\%$ (but above 50%) was considered evidence
11 for probable but inconsistent association (summary code '+?' or '-?' or 'o?'); a variable that
12 has been frequently studied (i.e., in ≥ 10 independent samples) but with considerable lack
13 of consistence was attributed a summary code of two question marks ('?'); where findings
14 were consistent, the codes '++', '--' or 'oo' were attributed. Final summary codes were only
15 computed for variables that have been studied in at least 3 independent samples; otherwise
16 a 'non-applicable' (N/A) summary code was attributed

19 RESULTS

21 General characteristics of the studies reviewed (Table 4.1)

22
23 We have identified a total of 150 publications that presented an empirical association be-
24 tween PA and at least one environmental correlate. The vast majority of studies (71.3%)
25 were published in the last decade (Fig.4.1) and a steep, almost 3-fold increase in adolescent
26 studies was noticed in the last 5 years. The overall studies reported data on 225 independent
27 samples. Sixty-six studies (91 independent samples) of children were reviewed, represent-
28 ing 40.4% of the total independent samples; only 16 (17.6%) of those independent samples
29 included more than 1,000 subjects. Eighty-four studies of adolescents (134 independent
30 samples; 59.6%) were reviewed (4 of which provided also data on children); about one third
31 included more than 1,000 subjects. In both children and adolescents, the vast majority of
32 the studies used a cross-sectional design reported results for boys and girls separately, relied
33 on child and/or parental self-reports as method of PA data collection (about half of which
34 with acceptable reliability/validity), and were mostly conducted in North America. Studies
35 that used objective methods of PA assessment were in the great majority restricted to stud-
36 ies among children; direct observation and doubly labeled water assessment were never
37 used in studies of adolescents.

38
39




Figure 4.1 Distribution of the 150 publications by year of publication (1980-2004).

Potential environmental determinants of children's PA (Table 4.2)

Potential determinants at the home level

We have identified a total of 17 independent samples investigating the associations between variables of the home *physical environment*, namely the amount of cars in the family and the availability and access of exercise equipment (e.g. PA promoting toys), and PA levels of children. Both variables were unrelated to children's PA. *Socio-cultural environmental* correlates of children's PA at the home/family level were the most frequently investigated. Family structure variables such as single-parent family, household size or number of children in the family, dog ownership and level of acculturation to the country of residence, were unrelated to children's PA. Modeling of PA from parents, siblings and friends were extensively examined (96 independent samples in total). Studies that have examined the relationship between children's PA levels and those of their parents, not disentangling those of the father from those of the mother, as well as those from other significant others (e.g. parents, siblings or friends), found no relevant associations. However, in studies where father's and mother's PA levels were disentangled from each other, father's PA levels emerged as a probable positive correlate (in 52% of the cases), whereas mother's PA levels were mostly unrelated to children's PA. Studies investigating potential familial influences other than modeling, namely support, encouragement and PA-related social norms of parents, friends and significant others, have also been numerous (a total of 99 independent samples). These variables were generally unrelated to children's PA. The *economic environment* of children's

Table 4.1 Child and adolescents studies categorized by sample size, sex, study design, physical activity measurement issues, and country

SAMPLE SIZE	CHILDREN (3-12 years)		ADOLESCENTS (>12-18 years)		N Samples	%
	Bibliography no.	N Samples	Bibliography no.	N Samples		
<100	5F, 15M/F, 25, 32, 39, 66, 71, 82, 89M/F, 111, 113MI, V/F, II, V, 129, 134M	91	20I, 23M/F, 24M/F, 26M/F, 30F, 50M, 68, 69, 90M/F, 106F, 113MIV, VI/F, VI, 123, 126M, 133M, 138I, II	134	100	16.4
100-199	10*, 21, 22F, 31*, 42, 46, 51, 58, 75M, 76, 103M/F, 108M/F, 113MII, 124M/F, 131MI/FI, 134F, 150	20	14, 44M/F, 45M/F, 48II, 49, 50F, 101, 102F, 110II, 113MIII/FIII, IV, 126F, 128, 133F	17	12.7	
200-299	13F, 28, 37, 38, 41, 63M/F, 70, 75F, 96, 109, 110I, 112MI/FI, 118	15	12, 16M/F, 20II, 93F, 102M, 112MIII, III/FII, III, 125, 148MIII	12	8.9	
300-499	27M/F, 56F, 81, 97, 107M/F, 115, 131MII/FII, 132, 148MII/FI	13	8M/F, 34M/F, 40F, 43, 67M, 79M/F, 105, 116F, 135, 147, 148MII/FII, III	16	11.9	
500-999	56M, 64, 84M/F, 100, 104, 119, 144	8	4M/F, 7M/F, 17F, 18F, 29, 33, 36, 40M, 47, 48I, 61M/F, 62, 67F, 87F, 99, 145, 149M/F	21	15.7	
1,000-2,999	11M/F, 57, 73, 85, 86M/F, 88, 95M/F, 120, 122, 137, 143M/F	15	1M/F, 3M/F, 6, 35F, 54, 65M/F, 80M/F, 87M, 91F, 94I, II, 114F, 117M/F, 136, 139, 141, 142M/F, 146M/F	25	18.7	
3,000-4,999	19	1	2M/F, 72, 78, 83, 121, 140	7	5.2	
≥5,000			9, 52, 53, 55, 59M/F, 60, 74, 77, 92, 98, 127M/F, 130	14	10.5	
SEX						
Girls only	5F, 13F, 22	3	17, 18, 30, 35, 91, 93, 106, 114, 116	9	6.7	
Boys and girls combined	10/31, 19, 21, 25, 28, 32, 37, 38, 39, 41, 42, 46, 51, 57, 58, 64, 66, 70, 71, 73, 76, 81, 82, 85, 88, 96, 97, 100, 104, 109, 110I, 111, 115, 118, 119, 120, 122, 129, 132, 137, 144, 150	42	6, 9, 12, 14, 20II, 29, 33, 36, 43, 47, 48I/II, 49, 52, 53, 54, 55, 60, 62, 68, 69, 72, 74, 77, 78, 83, 92, 94I, II, 98, 99, 101, 105, 110II, 121, 123, 125, 128, 130, 135, 136, 138, 139, 140, 141, 145, 147	49	36.6	

	ADOLESCENTS (>12-18 years)			
	CHILDREN (3-12 years)	N Samples	%	Bibliography no.
	Bibliography no.			
Boys and girls, separately	11, 15, 27, 56, 63, 75, 84, 86, 89, 95, 103, 107, 108, 112I, 113I,II,V, 124, 131II, 134, 143, 148I	46	50.5	1, 2, 3, 4, 7, 8, 16, 23, 24, 26, 34, 40, 44, 45, 50, 59, 61, 65, 67, 79, 80, 87, 90, 102, 112II,III, 113III,IV,VI, 117, 126, 127, 133, 142, 146, 148II,III, 149
STUDY DESIGN				
Cross-sectional	5F, 10/3I, 13F, 15M/F, 19, 21, 22F, 25, 27M/F, 28, 32, 37, 38, 39, 41, 46, 51, 56M/F, 57, 58, 63M/F, 64, 66, 70, 71, 73, 75M/F, 76, 81, 84M/F, 85, 86M/F, 88, 89M/F, 93M/F, 96, 97, 100, 103M/F, 104, 108M/F, 109, 110I, 111, 112MI/FI, 113MII,IV/FII,IV, 115, 119, 120, 122, 124M/F, 129, 131MI,II/FI,II, 132, 134M/F, 143M/F, 144, 150	81	89.0	1M/F, 3M/F, 4M/F, 6, 7M/F, 8M/F, 9, 12, 14, 16M/F, 17F, 18F, 23M/F, 24M/F, 29, 30F, 33, 34M/F, 35F, 36, 43, 44M/F, 47, 48I/II, 49, 50M/F, 52, 53, 54, 55, 59M/F, 60, 61M/F, 62, 65M/F, 67M/F, 68, 69, 72, 74, 77, 78, 79M/F, 80M/F, 83, 87M/F, 90M/F, 91F, 92, 94I,II, 98, 99, 101, 105, 106F, 110II, 112MII,III/FII,III, 113MIII,IV,VI/FIII,IV,VI, 114F, 116F, 117M/F, 121, 123, 125, 126M/F, 127M/F, 128, 130, 135, 136, 138I,II, 139, 140, 141, 142M/F, 145, 146M/F, 147, 149M/F
Longitudinal (length of study)	11M/F (1 year), 42(1 year), 82(8 weeks), 107M/F (2 years), 118(1 year), 137(1 year), 148MI/FI (3 years)	10	11.0	2M/F (2.5 years), 20I (1 week), II (9 months), 26M/F (3 years), 40M/F (1 year), 45M/F (3 years), 93F (8 months), 102M/F (4 months), 133M/F (1 year), 148MII,III/FII,III (3 years)
ASSESSMENT OF PHYSICAL ACTIVITY				
Collection method				
Self-report	11M/F, 25, 27M/F, 38, 41, 42, 46, 56M/F, 57, 58, 64, 75M/F, 84M/F, 85, 86M/F, 95M/F, 97, 103M/F, 104, 110I, 115, 118, 119, 120, 124M/F, 132, 137, 143M/F, 144, 148MI/FI	41	45.1	1M/F, 2M/F, 3M/F, 4M/F, 6, 7M/F, 8M/F, 9, 14, 16M/F, 17F, 18F, 20I,II, 26M/F, 30F, 33, 34M/F, 35F, 36, 40M/F, 43, 44M/F, 45M/F, 47, 48I/II, 49, 50M/F, 52, 53, 54, 55, 59M/F, 60, 61M/F, 62, 65M/F, 67M/F, 68, 69, 72, 74, 77, 78, 79M/F, 80M/F, 83, 87M/F, 91F, 92, 93F, 94I,II, 98, 99, 102M/F, 105, 106F, 110II, 114F, 116F, 117M/F, 121, 125, 126M/F, 127M/F, 128, 130, 133M/F, 135, 136, 139, 140, 141, 142M/F, 145, 146M/F, 147, 148MII,III/FII,III, 149M/F

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

1 home/family in relation to their PA levels was studied in 102 independent samples. Dif-
 2 ferent estimates of family/parental SES were generally unrelated to children's PA. Finally,
 3 and within the household's '*political*' environment, parenting styles were also unrelated to
 4 children' PA.

5
 6 *Potential determinants at the school level*

7 Aspects of the school environment were studied seldom (most of them only once or twice,
 8 which has not enabled us to calculate a summary association). Only one aspect of the school
 9 *political environment* – PA policies (i.e., time allowed from free play, time spent outdoors,
 10 and number of field trips) – was investigated in three or more independent samples, with
 11 60% of the cases showing a positive association with children's PA levels.

12
 13 *Potential determinants at the neighborhood level*

14 A total of 90 independent samples have examined associations between environmental
 15 characteristics at the neighborhood levels and PA levels of young children. We have iden-
 16 tified a total of 8 potential correlates of PA at the neighborhood *physical environment*, 5
 17 of which studied more than 3 times. Among these, time spent outdoors was consistently
 18 associated with higher PA levels of children, whereas availability and accessibility of PA
 19 programs or facilities, neighborhood safety and neighborhood hazards (e.g., many roads,
 20 no lights crossings, heavy traffic, physical disorder and pollution – estimated as perceived
 21 by parents in almost all studies) were consistently unrelated to children's PA. Aspects of the
 22 *social and economic environments* were unrelated to children's PA.

23
 24 *Potential determinants at the city/municipality and region/country level*

25 Only few studies have investigated differences in PA levels between children living in urban
 26 vs. suburban (only examined twice) and coastal vs. mountainous locations (only examined
 27 once). Whether residence in urban vs. rural regions is associated with children's PA levels
 28 was undetermined by the available studies. Seasonal 'effects' on children's PA were also
 29 undetermined by the available literature.

30
 31 **Potential environmental determinants of adolescents' PA (Table 4.3)**

32
 33 *Potential determinants at the home level*

34 We have identified a total of 20 independent samples investigating the associations between
 35 variables of the home *physical environment*, namely the availability and accessibility of
 36 exercise equipment, and PA levels of adolescents; these variables were mostly unrelated
 37 to adolescents' PA. *Socio-cultural environmental* correlates of adolescents' PA at the home/
 38 family level were the most frequently investigated. Family structure variables such as single-
 39 parent family and household size or number of children in the family were unrelated to

Table 4.2 Summary of correlates of physical activity among children (3 to 12 year olds)

Correlate	Related to PA		Assoc. (+ or -)	Biblio. no.	Unrelated to PA	# Samples	Summary (n)			
	Biblio. no.						+	-	0	Assoc
MICRO ENVIRONMENT										
Physical										
# cars in household	19, 131FI		-	131MI,II,FI		5	-	2	3	0
Access/availability of exercise equipment	30F, 124F, 134F		+	30F, 81, 97, 109, 124M, 132, 134E, 134M/F,		12	3	-	9	00
Socio-Cultural										
Single-parent family	103F, 108M, 113MI		+	95M/F, 95M/F, 103M, 107M/F, 108F, 108M/F, 112MI/FI, 113MI,II,V/FI,II,V		20	3	-	17	00
# household residents/children	-			38, 95M/F, 95M/F, 113MI,II,V/FI,II,V		11	-	-	11	00
Acculturation (language spoken at home; lifetime in the county; index)	11M, 95M/F, 19, 137		+	11F, 11M/F, 13F, 95M/F, 137		12	3	2	7	0
Dog ownership			-	131MI,II/FI,II		4	-	-	4	0
Parents' PA	32, 63M, 89M/F, 100, 107M, 111, 112MI, 144, 150, 124F		+	11M/F, 25, 32, 63F, 107F, 108M/F, 108M/F, 112FI, 113MI,II,V/FI,II,V, 124M		29	10	1	18	00
Father's PA	22F, 38, 39, 39, 46, 89M/F, 95M, 119M/F, 134M, 148MI/FI, 148MI/FI		+	15M/F, 15M/F, 84M/F, 95M/F, 95F, 97, 110I, 134M/F, 134F		29	15	-	14	+?
Mother's PA	15F, 38, 39, 39, 95F, 110I, 124F, 134M, 148FI, 148FI		+	15M/F, 15M, 22F, 84M/F, 89M/F, 95M, 95M/F, 97, 109, 119M/F, 124M, 134M/F, 134E, 148MI, 148MI		31	10	-	21	00
Siblings' PA	110I		+			1	1	-	-	N/A
Friend's PA	46		+	97, 134M/F, 134M/F		6	1	-	5	0
PA from significant others (parents, siblings, friends)				4I		1	-	-	1	N/A
Encouragement from parents	71, 82, 95F, 95F, 107M, 144		+	11M/F, 63M/F, 70, 95M, 95M, 95M/F, 95M/F, 107F, 108M/F, 108M/F		22	6	-	16	00

Home/household

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

Correlate	Related to PA		Assoc. (+ or -)	Unrelated to PA		# Samples	Summary (n)			
	Biblio. no.			Biblio. no.			+	-	Assoc	
Support (logistic) from parents (transports child to play, plays with child, pays fees)	5F, 22F, 107M, 107M, 108M/F, 108M, 144, 144		+	5F, 22F, 63M/F, 63M/F, 70, 107F, 107F, 107F, 108F, 108M/F, 108M/F, 112MI/FI		28	10	1	17	00
Support/encouragement from significant others (family, peers, teachers)	112MI/FI, 113MI/FI, 115, 115, 120, 124M		+	41, 97, 97, 109, 113MI, II, V, FI, II, V, 113MII, V/FI, 124F		24	9	-	15	0?
Social norms (value/enjoyment of PA of significant others - parents, siblings, peers)	25, 75M/F, 75M/F, 134M, 134M, 150		+	25, 25, 41, 84M/F, 97, 100, 112MI/FI, 112MI/FI, 124M/F, 129, 129, 134F, 134F		25	8	-	17	00
Economic										
Parental SES	27F, 27M/F, 32, 72, 88, 95F, 122		+	27M, 27M/F, 32, 71, 72, 95M, 95M/F, 103, 109, 119, 137		22	8	1	13	0?
Parental occupational status	148FI 19		+	123, 148MI/FI, 148MI		6	1	1	4	0
Father occupational status			-			6	-	-	6	0
Mother occupational status	11F, 56M/F		+	11M/F, 95M/F, 95M/F		7	2	-	5	0
Parental education	63M, 112MI 108F		+	37, 37, 37, 63F, 96, 103, 107M/F, 108M, 38, 46, 108M/F, 112FI, 113MI, II, V, FI, II, V, 137		24	2	1	21	00
Father's educational level	56M/F, 95F, 148MI, 148MI 57		+	95M, 95M/F, 148FI, 148FI		11	5	1	5	??
Mother's education level	148MI		+	19, 46, 95M/F, 19, 46, 95M/F, 131MI, II, FI, II, 148FI		12	1	-	11	00
# hours parents work			+	95M/F, 108M/F, 108M/F, 150		7	-	-	7	0
House owned				19		1	-	-	1	N/A
Political										
Parenting styles (PA rules, control)	109, 109		-	43, 109, 112MI/FI		6	2	-	4	0

Correlate	Related to PA		Assoc. (+ or -)	Unrelated to PA		# Samples		Summary (n)		
	Biblio. no.			Biblio. no.		+	-	0	Assoc	
Physical										
Distance (from home)	150		-			1	-	1	-	N/A
Availability of PA equipment				81		1	-	-	1	N/A
Socio-Cultural										
Teacher's PA				100		1	-	-	1	N/A
Teacher's attitudes toward PA				100		1	-	-	1	N/A
Teachers specific education level	28, 100		+			2	2	-	-	N/A
Economic										
School type attended (public vs. private; nursery vs. day care)	19, 100		+			2	2	-	-	N/A
Political										
Support from community PA organizations				28		1	-	-	1	N/A
PA related policies (e.g. time allowed for free play/spent outside, # field trips)	28, 81, 96		+	28, 28		5	3	-	2	+
Class size	28		+			1	1	-	-	N/A
School quality				28		1	-	-	1	N/A
Physical										
Distance to destinations	58		-			1	-	1	-	N/A
Access/availability to PA facilities/ programs	41, 109, 131FII		+	5F, 5F, 30F, 131MI,II/F,II, 131MI,II/ FI, 113MI,II,IV/FI,II,V		20	3	-	17	00
Available shelters/foot path conditions				150, 150		2	-	-	2	N/A
Time spent outdoors	10, 70, 81, 109, 109		+			5	5	-	-	+

Educational Institutions (Schools,...)

Neighbourhood

Correlate	Related to PA		Unrelated to PA	# Samples		Summary (n)	
	Biblio. no.	Assoc. (+ or -)		Biblio. no.	+	-	0
Neighbourhood hazards (e.g. many roads/no lights crossings; heavy traffic; physical disorder; pollution)	123 58, 131MII, 131MII, 131MI	+	58, 88, 113MI,II,V,FI,II,V, 150, 131MI/ FI,II, 131MI/FI,II 131MII/FI,II, 150	24	1	4	19 00
Neighbourhood physical disorder			88	1	-	-	1 N/A
Limited public transport	131FI,II	-	131MI,II	4	2	-	2 ?
Social							
Neighbourhood social disorder	88	-		1	1	-	- N/A
Involvement in community PA organizations	132, 134M, 134M	+	11M/F, 132, 134F, 134F	8	3	-	5 0
Length of residence in community	30F	+	30F	2	1	-	1 N/A
Safety	88	-	5F, 5F, 107M/F , 113MI,II,V/FI,II,V, 131MI,II/FI,II, 150	16	-	1	15 00
Economic							
Neighbourhood SES/education level	143F 64, 64	+	30F, 30F, 88, 143M,	6	1	2	3 0
MACRO ENVIRONMENT							
Physical							
Urban vs. suburban	21	-	66	2	1	1	- N/A
Urban vs. rural	27M/F, 56M/F, 66, 72, 72, 118 27M/F, 27M/F	+	46, 57, 85, 86M/F	17	8	4	5 ??
Coastal vs. mountains	46	-		1	1	-	- N/A
Season (spring, summer)	42, 51, 100, 118 10&31, 118	+	21, 37, 37, 37	10	4	2	4 ??

Biblio. no., reference number under the *Bibliography* section; Assoc., association; +, positive; -, negative; 0, no relation; ?, indeterminate; N/A, summary code not applicable because the number of independent samples investigating the relationship is below 3; PA, physical activity; M, boys only; F, girls only; SES, social-economic status; studies with prospective study designs are highlighted in bold.

1 adolescents' PA as were indicators of acculturation. Modeling of PA from parents, siblings
2 and friends were extensively examined (in 149 independent samples). Overall, all these
3 studies found no relevant associations. However, this lack of associations was somewhat
4 undetermined with regard to father's PA levels and those from significant others, since
5 they were observed in less than 60% of the cases. Studies investigating potential familial
6 influences other than modeling were also numerous (a total of 127 independent samples)
7 but mostly unrelated to adolescents' PA. However, a trend toward a positive association was
8 found with regard to general support from significant others. The relationship between the
9 *economic environment* of adolescents' home/family and their PA levels was examined in 100
10 independent samples. Studies in which parental SES was defined as a composite of parent's
11 education and income levels/occupational status were generally unrelated to children's PA.
12 However, studies in which the specific association between parent's education levels was
13 analyzed separately from parent's occupational status or income level revealed that higher
14 mother's education levels and family (per capita) income were positively associated with
15 PA; occupational status of the household's head emerged as an undetermined correlate of
16 PA. With regard to the *political environment*, parenting styles were unrelated to adolescents'
17 PA.

18 19 *Potential determinants at the school level*

20 Similarly to what we have described in children, aspects of the school physical, socio-
21 cultural, economic or political environment were studied relatively seldom in adolescents.
22 Regarding the *socio-cultural environment*, role modeling and support from teachers were
23 generally unrelated to adolescents' PA, whereas the existence of problems with (or teasing
24 from) classmates was undetermined. Finally, the type of school attended, namely high- vs.
25 vocational school, was positively, whereas the provision of instruction on PA or sport-
26 related health benefits and special Physical Education programs and/or school sports, were
27 unrelated to adolescents' PA.

28 29 *Potential determinants at the neighborhood level*

30 A total of 92 independent samples have examined associations between environmental
31 characteristics at the neighborhood level and PA levels of adolescents. Although we have
32 identified a wide range of potential correlates at the physical, socio-cultural and economical
33 level, only few were examined in more than 3 independent samples. Among these, and
34 within the *physical environment*, the availability and/or accessibility of PA equipment or
35 facilities, was unrelated to PA. Within the *socio-cultural environment*, crime incidence
36 (measured objectively) was inversely associated with adolescents' PA in 2 out of the 3 studies
37 available, a finding that was at odds with the lack of association between adolescents' PA
38 and neighborhood safety estimates perceived by them.

1 *Potential determinants at the city/municipality and region/country level*

2 Only few studies have investigated differences in PA levels between adolescents' residence
3 location. Residence in urban vs. rural regions was not associated with adolescents' PA levels;
4 seasonal 'effects' on adolescents' PA were undetermined; and exposure to or interest in
5 sports media was not associated with adolescents' PA.

6
7
8 **DISCUSSION**

9
10 Overall, the current review of the literature on environmental correlates of PA in children
11 and adolescents provided us with a broader and more detailed overview of the specific
12 research performed through the course of the past 25-years. In the past 5 years in particular
13 an increased attention to this field was observed, that may reflect a paradigm shift from
14 intra-personal to ecological conceptual models in the study of health-related behaviors such
15 as PA.

16
17 **Updating the previous review: current vs. previous findings**

18
19 We have updated the review of Sallis *et al.* by merging 51 of its original studies (those report-
20 ing on environmental potential determinants of PA, as defined in the present study) with
21 99 additional publications; twenty-three of the 99 additional studies had not been included
22 in the previous review although they were published within the same period covered by it
23 (1970-1998); interestingly half of those studies (12 out of 23) were performed in Europe, a
24 region that may have thus been under-represented in that review. With regard to the main
25 findings, a comparative summary between the two reviews is presented on Table 4.4; in
26 children, time spent outdoors remained a main correlate of children's PA, although this was
27 due to the fact that no additional studies in this regard were included in the present review.
28 The correlates of children and adolescents' PA that have emerged in the present review differ
29 considerably from those in the previous review. Overall, we can argue that the additional
30 publications of which 76 were published in the last 5 years), have thus contributed signifi-
31 cantly to a better understanding of factors associated with the PA behaviors of children and
32 adolescents, and have led to the identification and addition of new potential determinants
33 to the body of knowledge in the field. However, the fact that the associations coded and
34 summarized in our review were those derived, whenever possible, from multivariate rather
35 than from univariate analyses may also have contributed to the differences between the two
36 reviews. The previous review, which drew exclusively from univariate models may have
37 thus been somewhat inflated (since significant correlates are generally more abundant in
38 univariate analyses).

Table 4.3 Summary of correlates of physical activity among adolescents (13 to 18 year olds)

Correlate	Bibliography no.		Assoc + or -	Unrelated to PA	N Samples		Summary (%)	
	Related to PA	Bibliography no.			+	-	0	Assoc.
MICRO ENVIRONMENT								
Physical								
Access/availability of PA equipment	18F, 18F, 33 24F		+	18F, 23M/F, 23M/F, 24M, 24M/F, 26M/F, 93F, 93F 133M/F, 133M/F	20	3	1	16 00
Socio-Cultural								
Single-parent family	29, 76, 113MIV 76, 130		+	45M/F, 61M/F, 67M/F, 76, 112MII,III/ FII,III, 113MIII,VI/FIII,IV,VI, 128, 142M/F	24	3	2	19 00
# household residents/children			-	61M/F, 113MIII,IV,VI/FIII,IV,VI, 142M/F, 149M/F	12	-	-	12 00
Acculturation (adolescent/ parent born abroad; generation of residence in country)	45F, 52		+	45M, 52, 53, 116F	6	2	-	4 0
Parents' PA	33, 54, 98, 99, 142M/F		+	17F, 26M/F, 68, 79M/F, 79M/F, 90M/F, 90M/F, 112MII,III/FII,III, 113MIII,IV,VI/FIII,IV,VI, 135, 149M/F	31	6	-	25 00
Father's PA	23M, 24F, 48I, 49, 98, 105, 110II, 140, 140, 141, 142M/F, 148MII,III		+	3M/F, 23F, 23M/F, 24M, 24M/F, 48I, 48II, 49, 133M/F, 133M/F, 148FII,III	31	14	-	17 0?
Mother's PA	3F, 23F, 48I, 49, 98, 106F, 110II, 133F, 142M/F, 148FII,III		+	3M, 23M, 23M/F, 24M/F, 24M/F, 26M/F, 48II, 48II, 49, 105, 133M, 133M/F, 140, 141, 148MII,III	33	12	-	21 00
Sibling's activity	3M/F, 98, 99, 110II, 141			23M/F, 23M/F, 24M/F, 24M/F, 110II, 140, 140, 141	18	6	-	12 00
Friend's PA	24M, 33, 116F, 140, 140		+	17F, 23M/F, 23M/F, 24F, 24M/F, 133M/F, 133M/F, 141, 149M/F	20	5	-	15 00

Home/household

Correlate	Related to PA		Assoc + or -	Unrelated to PA		N Samples		Summary (%)	
	Bibliography no.			Bibliography no.		+	-	0	Assoc.
PA from significant others (parents, friends, other adults)	8M/F, 9, 18F, 24M, 102E, 126M		+	14, 18E, 18F, 24M/F, 24F, 102M, 126F, 141		16	7	-	9 0?
Support/encouragement from parents	8M/F, 18F, 18E, 29, 44M, 61M/F, 68, 79M/F, 79M/F, 90F, 112MII, III/ FII, III, 112MII, III, 113FIII, 114E, 114F, 135, 139, 149F		+	17E, 44M, 101, 113MIV/FIII, IV, VI, 139, 149F		52	26	-	26 ??
Support/encouragement from friends	44F, 83, 101, 113MIII, VI, 149M		+	17E, 44M, 101, 113MIV/FIII, IV, VI, 139, 149F		15	6	-	9 00
Support/encouragement from significant others	8M/F, 12, 14, 18E, 18F, 24F, 24F, 44M/F, 93F, 114E, 114F		+	18E, 24M, 24M, 60, 93F, 133M/F, 133M/F, 139		18	12	-	10 +?
Social norms (value/enjoyment of PA of significant others - parents, siblings, peers)	9, 26F, 47, 47, 48I/II, 80M/F, 80M/F, 87M, 91E, 91E, 112FIII, 123, 127M/F		+	8M/F, 16M/F, 16M/F, 17F, 26M, 47, 68, 68, 69, 79M/F, 87F, 112MII, III/FII, 112MII, III/FII, III, 114F, 114F, 123		42	17	-	25 0?
Economic									
Parental SES	9, 12, 18E, 18E, 121, 145, 147		+	4M/F, 7M/F, 7M/F, 17E, 18E, 48II, 76, 76, 76, 128		19	7	-	12 00
Occupational status of household head	45F, 65M/F, 73, 140		+	45M, 67M/F, 140, 141		10	5	-	5 ??
Father's occupational status	54, 94I, 136		+	2M/F, 94II, 148MII, III/FII, III, 148MII/ FII		12	3	-	9 00
Mother's occupational status	2M, 94I		+	2F, 94II, 136		5	2	-	3 0
Parents' educational level	74, 77, 112MIII, 117E, 142M/F		+	61M/F 112MII/FI, II, III, 113MIII, IV, VI/ FIII, IV, VI, 117M		19	6	-	13 00
Father's educational level	136		+	48II, 148MII, III/FII, III, 148MII/FII		8	1	-	7 0

Correlate	Related to PA		Assoc + or -	Unrelated to PA		N Samples		Summary (%)	
	Bibliography no.	Bibliography no.		Bibliography no.	Bibliography no.	+	-	0	Assoc.
Mother's educational level	53, 92, 136	48II, 116F	+	5	3	-	2	+	
Family (per capita) income	29, 53, 74, 77, 142M/F	50M/F, 60, 73	+	10	6	-	4	++	
# parents working full time	117M	117F	+	2	1	-	1	N/A	
Adolescent's paid work/pocket money	34M/F, 83, 141	53, 116F, 125, 149M/F	+	9	4	-	5	0?	
Political									
Parenting styles (authoritative; PA rules)	87M, 117M, 90M	87F, 90F, 117F, 90M/F, 112MII,III/FII,III	-	12	3	-	9	00	
Physical									
School facilities/resources	33		+	1	1	-	-	N/A	
Socio-Cultural									
Main teacher's/coach PA	90M	90M/F, 90F, 140, 140, 141, 149M/F	+	9	1	-	8	0	
Support from teacher/coach	45M, 83, 149F	45F, 149M/F, 149M	+	7	3	-	4	0?	
Classmates problems/teasing	45F, 45F	45M, 45M	-	4	2	-	2	?	
School support		44M/F, 83		3	-	-	3	0	
Relationship with PE teacher		33		1	-	-	1	N/A	
Economic									
Public vs. private school	34M/F		-	2	-	2	-	N/A	
Political									
School type (high school vs. vocational/alternative)	1M/F, 2M/F, 7M/F, 55, 55, 55, 55 40M	7M/F, 40F, 55	+	15	10	1	4	++	
School provide (special) PE program/sport teams	53	36, 133M/F, 133M/F	+	6	1	-	5	0	

Education Institutions (childcare, schools)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

Correlate	Related to PA		Assoc + or -	Unrelated to PA		N Samples	Summary (%)	
	Bibliography no.	Bibliography no.		Bibliography no.	Bibliography no.		+ -	0 Assoc.
Instruction on sport/health benefits	140		+	140, 141, 141		4	1 -	3 0
Physical								
Distance to PA facilities	50M		-	50F		2	1 -	1 N/A
Access/availability to PA equipment/facilities/programs	33, 44M/F, 61M/F, 61M/F, 61F, 61M/F, 113FVI 29, 29		+	17F, 23M/F, 23M/F, 24M/F, 24M/F, 61M/F, 90M/F, 112MII,III/ FII,III, 113MIII,IV,VI/FIII,IV,VI, 113MIII,IV,VI/FIII,IV, 125, 149M/F, 61M		45	11 2	32 00
Level of urbanization				74		1	- -	1 0
Dogs unattended				90M/F, 90M/F		4	- -	4 0
Socio-cultural								
% married couples				67M/F		2	- -	2 N/A
% youth				67M/F		2	- -	2 N/A
Neighbourhood exercisers				90M/F, 90M/F, 149M/F		6	- -	6 0
Social disorganization				74		1	- -	1 N/A
Ethnic minority concentration				74		1	- -	1 N/A
Crime incidence	50F, 53		-	50M		3	- 2	1 -
Safety	50F		-	50M, 90M/F, 90M/F, 113MIII,IV,VI/ FIII,IV,VI, 149M/F		14	- 1	13 00
Economical								
SES				74		1	- -	1 N/A
% upper occupational status				67M/F		2	- -	2 N/A
% owner occupied housing	67F		+	67M		2	1 -	1 N/A

Neighbourhood

Correlate	Related to PA		Assoc + or -	Unrelated to PA		N Samples		Summary (%)		
	Bibliography no.	Bibliography no.		Bibliography no.	Bibliography no.	+	-	0	Assoc.	
% dwellings provided by employer		67M/F				2	-	-	2	N/A
% unemployment among residents	67F		-			2	-	1	1	N/A
Length of unemployment		67M/F				2	-	-	2	N/A
MACRO ENVIRONMENT										
Physical										
Urban vs. suburban	67F		-		67M	2	1	-	1	N/A
Town size	73		+			1	1	-	-	N/A
Urban vs. rural	140		+		35F, 53, 140, 141	5	1	-	4	0
Season	20II, 138I				53, 138II	4	2	-	2	?
Unsuitable weather	20I				125	2	1	-	1	N/A
Socio-cultural										
Exposure to/interest in sports media	62, 62		+		17F, 26M/F	5	2	-	3	0
Wanting to look like media figures	127M/F		+			2	2	-	-	N/A

Biblio. no., reference number under the *Bibliography* section; Assoc., association; +, positive; -, negative; 0, no relation; ?, indeterminate; N/A, summary code not applicable because the number of independent samples investigating the relationship is below 3; PA, physical activity; M, boys only; F, girls only; SES, social-economic status; PE, physical education; studies with prospective study designs are highlighted in bold.

1 Home/family correlates of children's and adolescents PA levels

2
3 Characteristics of the home environment, particularly those related to parental influences,
4 were by far the most explored in the literature, in both children and adolescents.

5 6 *Parents as roles models*

7 Research findings regarding the relationship between PA levels of parents and those of their
8 children have been mixed. Most of the studies have in fact failed to find any association.
9 Nevertheless, fathers appear to be more important role models as compared to mothers,
10 especially in childhood; fathers' PA may be related to their child's PA regardless of their
11 gender, whereas mothers' PA appears to be more often associated with girls' rather than
12 boys PA; however, parents' PA has been generally unrelated to children's future PA levels (as
13 could be ascertained by the few prospective studies examining this issue).

14 In samples of children, parental PA levels were almost always assessed by the parents
15 themselves (self-reports) whereas in the adolescent samples they were assessed by both
16 adolescents' reports ('perceived' parental PA levels) and parents' self-reports. It is thus pos-
17 sible that differences in the agent reporting on parental PA levels (parent vs. offspring) may
18 explain some of the lack of associations found. Indeed, there is some evidence that a low
19 agreement exist between parents and children reports with regard to the levels of parental
20 PA [59] and we have noticed that associations between PA of mothers or fathers and those of
21 their offspring (adolescents) tended to be more often positive when the mothers or fathers
22 reported their own level of PA (Table 4.5).

23 24 *Parental support, encouragement and beliefs*

25 It has been hypothesized that the support and encouragement parents provide, rather than
26 their own PA behavior, may influence the PA behavior of their offspring. In the present
27 review, these potential influences could not be clearly found, particularly among children;
28 however, as many studies have shown parental support to be positively or not to be associ-
29 ated with PA levels of adolescents. Taken together, these findings lend some support to the
30 view that parents may need to be more than just active role models if their child is to lead a
31 physical active lifestyle [32, 60]. This is supported by several (school-based) risk-reduction
32 programs that have included and evaluated (generally positively) parental involvement as a
33 means to enhance program effectiveness (e.g. The San Diego Family Heart Project [61]; the
34 Children and Adolescent Trial for Cardiovascular Health (CATCH I and II) [62, 63]; The
35 Minnesota Home Team [64, 65]).

36 37 *Parental Socioeconomic status (SES)*

38 Parental/family SES is associated with a wide array of health, cognitive and socio-emotional
39 outcomes in children, throughout their development from (even before) birth to adulthood

Table 4.4 Comparative summary of the main environmental correlates of physical activity in children and adolescents: earlier vs. current review

Children		Adolescents	
Previous Review	Current review	Previous Review	Current review
Program /facility access (+)	Father's PA (+?)	Support from significant others (++)	Support from significant others (+?)
Time spent outdoors (+)	School PA-related policies (+)	Parent support (++)	Mother's education level (+)
	Time spent outdoors (+)	Sibling PA (++)	Family income (++)
		Direct help from parents (+)	Non-vocational school (++)
		Opportunities to exercise (+)	Neighbourhood crime incidence (-)

PA, physical activity; +, positive association; -, inverse association; ?, indeterminate.

[66, 67]. In the studies reviewed herein, several measures of SES have been used, most including some quantification of family income, parental education and occupational status (or a combination of these). Mother's education level and family income emerged as independent correlates of adolescents' (but not children's) PA levels. These findings not only emphasize the need to disentangle such aspects as education, occupational status and income levels, but suggest also that on reaching adolescence and young adulthood, those who have lower income may be more restricted in their PA choices and opportunities. In younger children, PA is mostly of informal nature, and may therefore not involve much extra financial cost. Possibly, with increasing age participation in physical activities becomes more elaborate and financial costly (e.g. sport clubs fees), which may reduce the likelihood of PA in adolescents from lower income families [68]. This needs further investigation.

School influences on children and adolescents' PA

Schools offer many opportunities for young people to engage in physical activities, such as Physical Education classes, recess periods, extracurricular sports or PA programs, leisure time free use of its playing fields and playgrounds. Schools have also the personnel who, with sufficient training and commitment can define and deliver PA programs and policies that support the adoption of healthy lifestyles. The literature showing that well-designed and well implemented school-based programs can improve PA of young people is paramount [69-71] and guidelines for school programs to promote lifelong PA actually exist [72-74]. Despite this, little research has investigated specific features of the school environment that impact on youth PA. Indeed, although most studies reviewed herein, have recruited their target populations from school settings, aspects of the school physical, socio-cultural, economic or political environment, remained however relatively unexplored. Most of the

Table 4.5 Analyses of the review findings regarding the association between physical activity (PA) levels of parents and their offspring (adolescents) according to the agent reporting on parental PA levels

	Association		Chi-squared (P value)
	+	0	
(a) studies examining parental associations (total of 31 independent samples)			
Assessment of parents' PA			
Parent self-report	4 (98, 99, 142M/F)	16 (68, 79 M/F, 79M/F, 112 MII,III/FII,III, 113MIII,IV,VI/FII,III)	0.02 (0.90)
Perceived by the child	2 (33, 54)	9 (17F, 26M/F, 90M/F, 90M/F, 149M/F)	
(b) studies examining paternal associations (total of 31 independent samples)			
Assessment of father's PA			
Father's self-report	7 (98, 105, 110II, 142M/F, 148 MII,III)	4 (3M/F, 148 FII,III)	2.35 (0.13)
Perceived by the child	7 (23M, 24F, 48I, 49, 140, 140, 141)	13 (23F, 23M/F, 24M, 24M/F, 48II, 48II, 49, 133M/F, 133M/F)	
(c) studies examining maternal associations (total of 33 independent samples)			
Assessment of mothers' PA			
Mother's self-report	7 (3F, 98, 110II, 142M/F, 148FII,III)	6 (3M, 26M/F, 105, 148MII,III)	2.83 (0.09)
Perceived by the child	5 (23F, 48I, 49, 106F, 133F)	15 (23M, 23M/F, 24M/F, 24M/F, 48II, 48II, 49, 133M, 133M/F, 140, 141)	

Data are number of independent samples (bibliography #).

characteristics of the school environment identified were almost never tested in more than 10 and often in less than 3 independent samples. Despite this, the present review has identified 'school policies related to PA' to be positively associated with children's PA and 'school type' (i.e. attending high- rather than vocational-schools) to be a positive correlate of adolescent's PA.

Additionally, we have identified an interesting set of studies that have investigated PA levels of classes/groups of youngsters in the context of PE lessons or recess time. One study

1 found that: classes of children taught by PE specialists (as compared to generalists) received
2 longer as well as more very active lessons, leading to higher energy expenditure rates;
3 outdoor lessons generated more time spent in vigorous activities and higher total energy
4 expenditures than indoor classes [75]. In another study, school size, length of recess and
5 the availability of balls in the playground were identified as additional correlates of higher
6 engagement in physical activities by children [76]. In adolescents, teacher's specialty and
7 gender were not associated with classes PA levels, neither was the location where the lesson
8 were taught; the only significant correlates were class size and lesson specific context (fitness
9 activities; free play, game play and skill drills; management time, and knowledge) (inversely
10 associated with class PA) [77]. Another study found that, despite the availability of the PA
11 facilities, they were used by very few students during their leisure time at school (i.e., before
12 and after school classes, and lunch break) [78]. These findings were then further explored
13 and followed by the observation that not only the availability of PA facilities, but its size
14 and state of conservation, and particularly the existence of supervision/organized activities,
15 were decisive of adolescents' engagement in physical activities during their leisure time at
16 school [79]. These findings and those of the present review, together with the observation
17 that many schools are not providing enough time for physical activities [80, 81], emphasize
18 the important role school's environments may play in children and adolescents PA levels
19 [74, 82]. Further, school-based PA may represent an important equalizing factor for op-
20 portunities for PA in children and adolescents of different SES backgrounds [83].

22 Neighborhood influences on children and adolescents' PA

24 Recently, the importance of neighborhood physical and socio-cultural characteristics in
25 shaping PA of individuals has been increasingly investigated, but relatively few studies in
26 the current review had already addressed these possible associations. Among these studies,
27 features of the physical environment (also commonly referred in the literature as the 'built
28 environment'), in particular the availability and accessibility to PA equipment, facilities or
29 programs were investigated more often, but were generally unrelated to youth PA. The pres-
30 ent review identified time spent outdoors to be positively associated with children's activity
31 levels; in adolescents, crime incidence, as measured through objective police reports, was
32 inversely associated with adolescents PA levels, a finding that apparently contrasted with
33 the lack of association between perceived neighborhood safety levels and adolescents PA.
34 This contradiction suggests that the differential associations with youth PA may depend on
35 the method assessment (perceived vs. objective) of environmental characteristics. Which
36 features are more important remains unknown, an issue that therefore deserves further
37 investigation within the same population (see methodological considerations below).

38 The importance of understanding neighborhood effects on health-related behaviors rely
39 on their potential to influence large populations [84, 85]. Although researchers are start-

1 ing to address the potential effects of communities and neighborhoods in individuals' PA
2 behavior, few empirical studies have determined, using appropriate multilevel statistical
3 techniques, whether relations between the environment and PA actually exist at the neigh-
4 borhood rather than the individual level [86, 87].

5 6 Methodological considerations

7 8 *Measurement of PA and environmental characteristics*

9 The selection of the an appropriate instrument depends on the specific research question(s)
10 to be addressed and on an 'accuracy-practicality' trade off' [88-90]. The majority of the
11 research on the potential determinants of PA reviewed herein relied on (parental or child/
12 adolescent) self-reports, which included diaries and recall instruments; these methods may
13 pose serious limitations since they provide less accurate estimates of PA levels than those
14 obtained by more objective methods such as direct observation, motion sensors, heart rate
15 monitors, and doubly-labeled water [91]. In addition, because the degree of the relation-
16 ship between objectively and self-report measures of PA is only moderate, notably among
17 self-report methods with 'acceptable' validity [92], there may be a substantial amount of
18 variance not shared by the two methods; in other words, different instruments (objective
19 vs. self-report) may have measured different aspects of the PA behavior, and therefore those
20 measures are not interchangeable. As such, the correlates of PA may also differ as a function
21 of the method used to measure the behavior, thereby impairing the generalization of the
22 findings obtained with the use of one or the other method [93]. In the present review we
23 were able to identify seven publications (10 independent samples – 3 in children and 7 in
24 adolescents; all with a cross-sectional design) which enable a more close examination of this
25 issue, by providing self-report and objective data in the same samples (Table 4.6). In these
26 studies, the magnitude of the associations between the two measures of PA was at the most
27 moderate. Furthermore, clear discrepancies between correlates of objectively measured and
28 self-reported PA levels were found. Several factors may explain these discrepancies: the
29 proposed correlates investigated in each study may have more explanatory power for self-
30 reported measures (e.g. # of vigorous activities) than for total PA levels (mostly computed
31 by the objective measures); in addition, accelerometers, the most frequently used objective
32 measure, are unable to access common activities such as bicycling riding and swimming that
33 could have been (self-) reported, but pick-up incidental physical activities throughout the
34 day, which in turn could have been forgotten on self-reports that usually refer specifically
35 to intentional physical activities; finally, there may be a shared method variance between
36 self-reported PA and self-reported potential determinants, which then leads to an inflated
37 association between the two.

38 Furthermore, self-reports of environmental factors represent *perceived* rather than '*real*'
39 features of the physical, socio-cultural, economic and political environments. Little is know

1 about the accuracy of such perceived features [94]. In adults, some studies have shown
2 objective environmental measures to be associated with PA whereas the same features
3 measured through self-reports were not [95, 96].

4 5 *Limitations of study design and data analyses methodologies*

6 The studies incorporated in the present review had mostly a cross-sectional design and
7 therefore their findings were limited in that only association could be established and not
8 prediction or causation. Nevertheless, all those cross-sectional studies have interpreted the
9 results as if ‘causality’ existed and to be uni-directional (e.g., parents may influence their
10 children). It is of course possible that reverse or reciprocal influences are operative as well
11 (e.g., children influence their parents), an issue that needs to be further explored.

12 In an attempt to disentangle the information provided by prospective from cross-sectional
13 studies we have highlighted those studies in tables 4.2 and 4.3. However, their low number
14 does not enable solid conclusion with regard to the potential environmental predictors of
15 PA change.

16 The main question of how such environmental features influence youth PA remained
17 further largely unanswered due to the data analytical methods used. Conceptually, envi-
18 ronmental influences can play a direct role in shaping PA behavior or can be mediated by
19 cognitive processes [97-99]. In order to understand these mechanistic processes better data
20 analytical methods (and study designs) are needed (for details see Bauman et al. [100]).
21 The majority of the findings reviewed herein were those that resulted from adjusted models
22 (most often, for potential confounders such as age, sex, and ethnicity, but in many studies
23 for potential mediators such as self-efficacy and attitudes), and thus concern the indepen-
24 dent contribution of environmental characteristics in the explanation of PA behavior.

25 Further, although most of the data included in the present review have an intrinsic mul-
26 tilevel structure, they were most frequently analyzed as obtained in simple random samples
27 of a single population. As such, the potential inter-dependence within clusters (e.g. schools
28 and/or neighborhoods) has been ignored, which can have led to inflated estimation of the
29 associations, and multilevel or hierarchical analytic approaches are thus needed.

30 31 *Limitations of the present review*

32
33 We acknowledge several limitations of our current review. First, the search terms used to re-
34 trieve studies from existing databases may have not been sensitive enough. This is sustained
35 by the fact that almost half of the studies included in this review were found through the
36 literature sections of articles primarily retrieved in those databases. This may have been due
37 to the fact that some articles included are simply not registered within those databases, and/
38 or in many articles retrieved, environmental correlates of children/adolescents’ PA were not
39 the primarily research goal but were embedded within a broader (i.e. health-enhancing be-

1 haviors in general) or related research question. Nevertheless, better search terms may still
2 need to be defined. However, the vast amount of studies included suggests we have covered
3 the existing literature in a quite satisfactory way. Second, the use of only English published
4 data may have discarded some studies that could have added relevant information into the
5 field. Third, the main outcome was any form of PA. In most studies this was measured
6 across several settings (e.g. the total amount of moderate-to-vigorous PA, performed at
7 school and during leisure time – either at home or in the neighborhood, or in sport clubs,
8 accumulated throughout the day or the past week), not enabling us to determine the specific
9 environmental correlates of specific physical activities. Fourth, the conceptual framework
10 we have used may have led to disputable categorizations of the correlates of PA investigated.

11 12 **Conclusions, implications and recommendations**

13
14 Clearly, many factors influence the complex behavior of youth PA. We have identified
15 father's PA habits, school PA-related policies and time spent outdoors as potential deter-
16 minants of PA in children; in adolescents, such potential determinants were support from
17 significant others, mother's education levels, family income, attendance of a non-vocational
18 school and low neighborhood crime incidence have emerged as potential determinants of
19 adolescents' PA. These variables need to be target by multi-level interventions aiming at the
20 increase of youth PA. The other variables, however, should not be discarded without further
21 investigation, namely those whose associations with PA were undetermined or not possible
22 to infer from the limited number of existing studies (particularly those at the neighborhood
23 and school settings as well as at the macro-environment level).

24 Future studies that use prospective or intervention designs enabling the analyses of
25 whether the environment-PA behaviors of children and adolescents associations are casual
26 and which (if any) cognitive processes may mediate or contextual variables may moderate
27 such associations, are in great need. In addition, it is important to conduct future research
28 with clear, possibly standardized definitions and objective methods of environmental at-
29 tributes and PA behavior assessment, within the strongest study design possible.

30
31
32
33
34
35
36
37
38
39

Table 4.6 Determinants of objective vs. self-report measured physical activity - summary of findings

Bibliography no.	Method of Physical Activity (PA) Assessment			Environmental correlates of PA*	
	Objective	Self-report	Correlation between PA assessed by the 2 methods	Objective	Self-report
23	Heart-rate monitoring (1 week – time spent in moderate-to-vigorous PA, >140 beats min ⁻¹)	Recall of PA and sport participations (1 week; hours)	'not associated' (estimate size not reported)	Father's PA (M)	Mother's PA (F)
24	Heart-rate monitoring (1 week – time spent in moderate-to-vigorous PA, >140 beats min ⁻¹)	Recall of PA and sport participations (1 week; hours)	'not associated' (estimate size not reported)	-	Father's PA (F) Friends' PA (M) Parental encouragement (F) Parental support (F) Home equipment (F)
32	Accelerometer (2 week days + 1 weekend day); METs	Frequency, duration and types of PA (2 week days + 1 weekend day; METs)	r=0.46	Parental PA Parental SES	-
39	Accelerometer (2 weekdays + 1 weekend day; counts d ⁻¹)	Frequency, duration and types of PA (2 week days + 1 weekend day; METs)	r=0.39 (Light PA) r=0.35 (moderate-to-high intensity PA)	Father's PA Mother's PA	Father's PA Mother's PA
90	Accelerometer (up to 8 d; counts h ⁻¹)	PA Record of hard and very hard intensity PA (7 d; h week ⁻¹)	'Not associated' (estimate not reported)	Teacher's PA (M) PA rules (M)	Parent transports child to PA location (F)
101	Accelerometer (5-day period; min d ⁻¹)	Participation in PA for ≥60 min (PACE+) (past week; d week ⁻¹)	r=0.46	-	Parent support Peer support
110	Accelerometer (1 week day + 2 weekend days); score	Recall checklist of PA performed for at least 15 min (1 week day + 1 weekend day; score)	? (Not reported)	Parental education (F) Single-parent status (M)	Parent transports child to PA location (F) Parent plays with child (M)

* Only the environmental variables that were correlated with physical activity levels measured either by one or the other method are reported (i.e., listed variables do not cover all the variables investigated in each study). PA, physical activity; M, boys only; F, girls only; SES, socioeconomic status.

1 REFERENCES

- 2 1. Erlichman J, Kerbey AL, James WP: Physical activity and its impact on health outcomes. Paper 1: The
3 impact of physical activity on cardiovascular disease and all-cause mortality: an historical perspective.
4 *Obes Rev* 2002, 3(4):257-271.
- 5 2. Erlichman J, Kerbey AL, James WP: Physical activity and its impact on health outcomes. Paper 2:
6 Prevention of unhealthy weight gain and obesity by physical activity: an analysis of the evidence.
7 *Obes Rev* 2002, 3(4):273-287.
- 8 3. Strong WB, Malina RM, Blimkie CJ, Daniels SR, Dishman RK, Gutin B, Hergenroeder AC, Must A,
9 Nixon PA, Pivarnik JM *et al*: Evidence based physical activity for school-age youth. *J Pediatr* 2005,
10 146:732-737.
- 11 4. WHO: *Global Strategy on Diet, Physical Activity and Health*. Geneva: World Health Organization;
12 2004.
- 13 5. Children and young people - the importance of physical activity [<http://www.chnheart.org>]
- 14 6. Goran MI, Reynolds KD, Lindquist CH: Role of physical activity in the prevention of obesity in
15 children. *Int J Obes Relat Metab Disord* 1999, 23 Suppl 3:S18-33.
- 16 7. Kohn M, Booth M: The worldwide epidemic of obesity in adolescents. *Adolesc Med* 2003, 14(1):1-9.
- 17 8. Lobstein T, Baur L, Uauy R: Obesity in children and young people: a crisis in public health. *Obes Rev*
18 2004, 5 Suppl 1:4-104.
- 19 9. Steinbeck KS: The importance of physical activity in the prevention of overweight and obesity in
20 childhood: a review and an opinion. *Obes Rev* 2001, 2(2):117-130.
- 21 10. Johnson RK: Changing eating and physical activity patterns of US children. *Proc Nutr Soc* 2000,
22 59(2):295-301.
- 23 11. Services USDoHaH: *Physical Activity and Health: a report of the Surgeon General*. Atlanta, GA:
24 Centers for Disease Control and Prevention; 1996.
- 25 12. WHO: *Health behaviour in school-aged children: A WHO cross-national study (HBSC)*. Geneva:
26 International Report World Health Organization Regional Office for Europe; 2000.
- 27 13. Sallis JF: Age-related decline in physical activity: a synthesis of human and animal studies. *Med Sci*
28 *Sports Exerc* 2000, 32(9):1598-1600.
- 29 14. Caspersen CJ, Pereira MA, Curran KM: Changes in physical activity patterns in the United States, by
30 sex and cross-sectional age. *Med Sci Sports Exerc* 2000, 32(9):1601-1609.
- 31 15. Aaron DJ, Storti KL, Robertson RJ, Kriska AM, LaPorte RE: Longitudinal study of the number and
32 choice of leisure time physical activities from mid to late adolescence: implications for school cur-
33 ricula and community recreation programs. *Arch Pediatr Adolesc Med* 2002, 156(11):1075-1080.
- 34 16. Malina RM: Tracking of physical activity and physical fitness across the lifespan. *Res Q Exerc Sport*
35 1996, 67(3 Suppl):S48-57.
- 36 17. Gordon-Larsen P, Nelson MC, Popkin BM: Longitudinal physical activity and sedentary behavior
37 trends: adolescence to adulthood. *Am J Prev Med* 2004, 27(4):277-283.
- 38 18. Telama R, Leskinen E, Yang X: Stability of habitual physical activity and sport participation: a longi-
39 tudinal tracking study. *Scand J Med Sci Sports* 1996, 6:371-378.
19. Twisk JW, Kemper HC, van Mechelen W: Tracking of activity and fitness and the relationship with
cardiovascular disease risk factors. *Med Sci Sports Exerc* 2000, 32(8):1455-1461.
- Raitakari OT, Porkka KVK, Taimela S, Telama R, Rasanen L, Viikari JSA: Effects of persistent physi-
cal activity and inactivity on coronary risk factors in children an young adults. The Cardiovascular
Risk in Young Finns Study. *Am J Epidemiol* 1994, 140(3):195-205.

21. WHO: *Obesity: Preventing and Managing the Global Epidemic - Report of a World Health Organization Consultation on Obesity*. Geneva; 1998.
22. Booth KM, Pinkston MM, Poston WS: Obesity and the built environment. *J Am Diet Assoc* 2005, 105(5 Suppl 1):S110-117.
23. Hill JO, Wyatt HR, Reed GW, Peters JC: Obesity and the environment: where do we go from here? *Science* 2003, 299(5608):853-855.
24. Hill JO, Peters JC: Environmental contributions to the obesity epidemic. *Science* 1998, 280(5368):1371-1374.
25. Jeffery RW, Utter J: The changing environment and population obesity in the United States. *Obes Res* 2003, 11 Suppl:12S-22S.
26. Peters JC, Wyatt HR, Donahoo WT, Hill JO: From instinct to intellect: the challenge of maintaining healthy weight in the modern world. *Obes Rev* 2002, 3(2):69-74.
27. Booth ML, Macaskill P, Lazarus R, Baur LA: Sociodemographic distribution of measures of body fatness among children and adolescents in New South Wales, Australia. *Int J Obes Relat Metab Disord* 1999, 23(5):456-462.
28. Dowda M, Ainsworth BE, Addy CL, Saunders R, Riner W: Environmental influences, physical activity, and weight status in 8- to 16-year-olds. *Arch Pediatr Adolesc Med* 2001, 155(6):711-717.
29. Strauss RS, Knight J: Influence of the home environment on the development of obesity in children. *Pediatrics* 1999, 103(6):e85.
30. Timperio A, Crawford D, Telford A, Salmon J: Perceptions about the local neighborhood and walking and cycling among children. *Prev Med* 2004, 38(1):39-47.
31. Trost SG, Kerr LM, Ward DS, Pate RR: Physical activity and determinants of physical activity in obese and non-obese children. *Int J Obes Relat Metab Disord* 2001, 25(6):822-829.
32. Ritchie LD, Welk G, Styne D, Gerstein DE, Crawford PB: Family environment and pediatric overweight: what is a parent to do? *J Am Diet Assoc* 2005, 105(5 Suppl 1):S70-79.
33. Ball K, Crawford D: The obesity epidemic: Contextual influences on physical activity and body weight. 2003, 6(4):377-378.
34. Baranowski T, Mendlein J, Resnicow K, Frank E, Cullen KW, Baranowski J: Physical activity and nutrition in children and youth: an overview of obesity prevention. *Prev Med* 2000, 31:S1-S10.
35. Egger G, Swinburn B: An "ecological" approach to the obesity pandemic. *Bmj* 1997, 315(7106):477-480.
36. French SA, Story M, Jeffery RW: Environmental Influences on eating and physical activity. *Annu Rev Public Health* 2001, 22:309-335.
37. Nestle M, Jacobson MF: Halting the obesity epidemic: a public health policy approach. *Public Health Rep* 2000, 115(1):12-24.
38. Story M, Neumark-Sztainer D, French S: Individual and environmental influences on adolescent eating behaviors. *J Am Diet Assoc* 2002, 102(3 Suppl):S40-51.
39. Nutbeam D, Aar L, Catford J: Understanding childrens' health behaviour: the implications for health promotion for young people. *Soc Sci Med* 1989, 29(3):317-325.
40. Kohl III HW, Hobbs KE: Development of physical activity behaviors among children and adolescents. *Pediatrics* 1998, 101(3 Pt 2):549-554.
41. Richter KP, Harris JO, Paine-Andrews A, Fawcett SB, Schmid TL, Lankenau BH, HJohnston J: Measuring the health environment for physical activity and nutritions among youth: a review of the literature and applications for community initiatives. *Prev Med* 2000, 31:S98-S111.

42. Sallis JF, Simons-Morton BG, Stone EJ, Corbin CB, Epstein LH, Faucette N, Iannotti RJ, Killen JD, Klesges RC, Petray CK *et al*: Determinants of physical activity and interventions in youth. *Med Sci Sports Exerc* 1992, 24(6 Suppl):S248-257.
43. Brug J, Oenema A, Ferreira I: Theory, evidence and Intervention Mapping to improve behavioral nutrition and physical activity interventions. *Int J Behav Nutr Phys Act* 2005, 2:2.
44. Baranowski T, Cullen KW, Nicklas T, Thompson D, Baranowski J: Are current health behavioral change models helpful in guiding prevention of weight gain efforts? *Obes Res* 2003, 11 Suppl:23S-43S.
45. Sallis JF, Prochaska JJ, Taylor WC: A review of correlates of physical activity of children and adolescents. *Med Sci Sports Exerc* 2000, 32(5):963-975.
46. Howley ET: Type of activity: resistance, aerobic and leisure versus occupational physical activity. *Med Sci Sports Exerc* 2001, 33(6 Suppl):S364-369; discussion S419-320.
47. Biddle SJ, Gorely T, Marshall SJ, Murdey I, Cameron N: Physical activity and sedentary behaviours in youth: issues and controversies. *J R Soc Health* 2004, 124(1):29-33.
48. Gordon-Larsen P, McMurray RG, Popkin BM: Determinants of adolescent physical activity and inactivity patterns. *Pediatrics* 2000, 105(6):E83.
49. Lindquist CH, Reynolds KD, Goran MI: Sociocultural determinants of physical activity among children. *Prev Med* 1999, 29(4):305-312.
50. Owen N, Leslie E, Salmon J, Fotheringham MJ: Environmental determinants of physical activity and sedentary behavior. *Exerc Sport Sci Rev* 2000, 28(4):153-158.
51. Schmitz KH, Lytle LA, Phillips GA, Murray DM, Birnbaum AS, Kubik MY: Psychosocial correlates of physical activity and sedentary leisure habits in young adolescents: the Teens Eating for Energy and Nutrition at School study. *Prev Med* 2002, 34(2):266-278.
52. Gorely T, Marshall SJ, Biddle SJ: Couch kids: correlates of television viewing among youth. *Int J Behav Med* 2004, 11(3):152-163.
53. Flay BR, Petraitis J: The theory of triadic influence: a new theory of health behavior with implications for interventions. *Advances in Medical Sociology* 1994, 4:4-19.
54. Kumanyika S, Jeffery RW, Morabia A, Ritenbaugh C, Antipatis VJ: Obesity prevention: the case for action. *Int J Obes Relat Metab Disord* 2002, 26(3):425-436.
55. Swinburn B, Egger G, Raza F: Dissecting obesogenic environments: the development and application of a framework for identifying and prioritizing environmental interventions for obesity. *Prev Med* 1999, 29(6 Pt 1):563-570.
56. Cooper H: *Synthesizing research: a guide for literature reviews*. 3rd ed edition. London: Sage; 1998.
57. Pocock SJ, Collier TJ, Dandreo KJ, de Stavola BL, Goldman MB, Kalish LA, Kasten LE, McCormack VA: Issues in the reporting of epidemiological studies: a survey of recent practice. *Bmj* 2004, 329(7471):883.
58. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, Moher D, Becker BJ, Sipe TA, Thacker SB: Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000, 283(15):2008-2012.
59. Anderssen N, Jacobs DR, Jr., Aas H, Jakobsen R: Do adolescents and parents report each other's physical activity accurately? *Scand J Med Sci Sports* 1995, 5(5):302-307.
60. Norton DE, Froelicher ES, Waters CM, Carrieri-Kohlman V: Parental influence on models of primary prevention of cardiovascular disease in children. *Eur J Cardiovasc Nurs* 2003, 2(4):311-322.
61. Nader PR, Sallis JF, Patterson TL, Abramson IS, Rupp JW, Senn KL, Atkins CJ, Roppe BE, Morris JA, Wallace JP *et al*: A family approach to cardiovascular risk reduction: results from the San Diego Family Health Project. *Health Educ Q* 1989, 16(2):229-244.

62. Luepker RV, Perry CL, McKinlay SM, Nader PR, Parcel GS, Stone EJ, Webber LS, Elder JP, Feldman HA, Johnson CC *et al*: Outcomes of a field trial to improve children's dietary patterns and physical activity. The Child and Adolescent Trial for Cardiovascular Health. CATCH collaborative group. *JAMA* 1996, 275(10):768-776.
63. Nader PR, Sellers DE, Johnson CC, Perry CL, Stone EJ, Cook KC, Bebchuk J, Luepker RV: The effect of adult participation in a school-based family intervention to improve Children's diet and physical activity: the Child and Adolescent Trial for Cardiovascular Health. *Prev Med* 1996, 25(4):455-464.
64. Perry CL, Luepker RV, Murray DM, Kurth C, Mullis R, Crockett S, Jacobs DR, Jr.: Parent involvement with children's health promotion: the Minnesota Home Team. *Am J Public Health* 1988, 78(9):1156-1160.
65. Perry CL, Luepker RV, Murray DM, Hearn MD, Halper A, Dudovitz B, Maile MC, Smyth M: Parent involvement with children's health promotion: a one-year follow-up of the Minnesota home team. *Health Educ Q* 1989, 16(2):171-180.
66. Bradley RH, Corwyn RF: Socioeconomic status and child development. *Annu Rev Psychol* 2002, 53:371-399.
67. Evans GW: The environment of childhood poverty. *Am Psychol* 2004, 59(2):77-92.
68. Shropshire J, Carrol B: Family variables and children's physical activity: influence of parental exercise and socio-economic status. *Sport Educ Soc* 1997, 2(1):95-116.
69. Timperio A, Salmon J, Ball K: Evidence-based strategies to promote physical activity among children, adolescents and young adults: review and update. *J Sci Med Sport* 2004, 7(1 Suppl):20-29.
70. Matson-Koffman DM, Brownstein JN, Neiner JA, Greaney ML: A site-specific literature review of policy and environmental interventions that promote physical activity and nutrition for cardiovascular health: what works? *Am J Health Promot* 2005, 19(3):167-193.
71. Stone EJ, McKenzie TL, Welk GJ, Booth ML: Effects of physical activity interventions in youth. Review and synthesis. *Am J Prev Med* 1998, 15(4):298-315.
72. Prevention. CfDCA: Guidelines for school and community programs to promote lifelong physical activity among young people. *MMWR* 1997, 46:(No. RR-6).
73. Story M: School-based approaches for preventing and treating obesity. *Int J Obes Relat Metab Disord* 1999, 23 Suppl 2:S43-51.
74. Wechsler H, Devereaux RS, Davis M, Collins J: Using the school environment to promote physical activity and healthy eating. *Prev Med* 2000, 31:S121-S137.
75. McKenzie TL, Feldman H, Woods SE, Romero KA, Dahlstrom V, Stone EJ, Strikmiller PK, Williston JM, Harsha DW: Children's activity levels and lesson context during third-grade physical education. *Res Q Exerc Sport* 1995, 66(3):184-193.
76. Zask A, van Beurden E, Barnett L, Brooks LO, Dietrich UC: Active school playgrounds - myth or reality? Results of the "Move it groove it" project. *Prev Med* 2001, 33:402-408.
77. McKenzie TL, Marshall SJ, Sallis JF, Conway TL: Student activity levels, lesson context, and teacher behavior during middle school physical education. *Res Q Exerc Sport* 2000, 71(3):249-259.
78. McKenzie TL, Marshall SJ, Sallis JF, Conway TL: Leisure-time physical activity in school environments: an observational study using SOPLAY. *Prev Med* 2000, 30(1):70-77.
79. Sallis JF, Conway TL, Prochaska JJ, McKenzie TL, Marshall SJ, Brown M: The association of school environments with youth physical activity. *Am J Public Health* 2001, 91(4):618-620.
80. Simons-Morton BG, Taylor WC, Snider SA, Huang IW: The physical activity of fifth-grade students during physical education classes. *Am J Public Health* 1993, 83(2):262-264.

- 1 81. Simons-Morton BG, Taylor WC, Snider SA, Huang IW, Fulton JE: Observed levels of elementary and
2 middle school children's physical activity during physical education classes. *Prev Med* 1994, 23(4):437-
3 441.
- 4 82. McKenzie TL: Promoting physical activity in youth: focus on middle school environments. *Quest*
5 2001, 33:326-334.
- 6 83. Kristjansdottir G, Vilhjalmsson R: Sociodemographic differences in patterns of sedentary and physi-
7 cally active behavior in older children and adolescents. *Acta Paediatr* 2001, 90(4):429-435.
- 8 84. King AC, Jeffery RW, Fridinger F, Dusenbury L, Provence S, Hedlund SA, Spangler K: Environmental
9 and policy approaches to cardiovascular disease prevention through physical activity: issues and
10 opportunities. *Health Educ Q* 1995, 22(4):499-511.
- 11 85. Sallis JF, Johnson MF, Calfas KJ, Caparosa S, Nichols JF: Assessing perceived physical environmental
12 variables that may influence physical activity. *Res Q Exerc Sport* 1997, 68(4):345-351.
- 13 86. Braza M, Shoemaker W, Seeley A: Neighborhood design and rates of walking and biking to elemen-
14 tary school in 34 California communities. *Am J Health Promot* 2004, 19(2):128-136.
- 15 87. Duncan SC, Duncan TE, Strycker LA, Chaumeton NR: Neighborhood physical activity opportunity:
16 a multilevel contextual model. *Res Q Exerc Sport* 2002, 73(4):457-463.
- 17 88. Pate RR: Physical activity assessment in children and adolescents. *Crit Rev Food Sci Nutr* 1993, 33(4-
18 5):321-326.
- 19 89. Welk GJ, Corbin CB, Dale D: Measurement issues in the assessment of physical activity in children.
20 *Res Q Exerc Sport* 2000, 71(2 Suppl):S59-73.
- 21 90. Schutz Y, Weinsier RL, Hunter GR: Assessment of free-living physical activity in humans: an over-
22 view of currently available and proposed new measures. *Obes Res* 2001, 9(6):368-379.
- 23 91. Janz KF, Witt J, Mahoney LT: The stability of children's physical activity as measured by accelerometry
24 and self-report. *Med Sci Sports Exerc* 1995, 27(9):1326-1332.
- 25 92. Kohl III HW, Fulton JE, Caspersen CJ: Assessment of physical activity among children and adoles-
26 cents: a review and synthesis. *Prev Med* 2000, 31(2):S54-S76.
- 27 93. Dishman RK, Darracott CR, Lambert LT: Failure to generalize determinants of self-reported physical
28 activity to a motion sensor. *Med Sci Sports Exerc* 1992, 24(8):904-910.
- 29 94. Kirtland KA, Porter DE, Addy CL, Neet MJ, Williams JE, Sharpe PA, Neff LJ, Kimsey CD, Jr., Ain-
30 sworth BE: Environmental measures of physical activity supports: perception versus reality. *Am J*
31 *Prev Med* 2003, 24(4):323-331.
- 32 95. Sallis JF, Hovell MF, Hofstetter CR, Elder JP, Hackley M, Caspersen CJ, Powell KE: Distance between
33 homes and exercise facilities related to frequency of exercise among San Diego residents. *Public*
34 *Health Rep* 1990, 105(2):179-185.
- 35 96. Hoehner CM, Brennan Ramirez LK, Elliott MB, Handy SL, Brownson RC: Perceived and objective
36 environmental measures and physical activity among urban adults. *Am J Prev Med* 2005, 28(2 Suppl
37 2):105-116.
- 38 97. Owen N, Humpel N, Leslie E, Bauman A, Sallis JF: Understanding environmental influences on
39 walking: Review and research agenda. *Am J Prev Med* 2004, 27(1):67-76.
98. Lewis BA, Marcus BH, Pate RR, Dunn AL: Psychosocial mediators of physical activity behavior
among adults and children. *Am J Prev Med* 2002, 23(2 Suppl):26-35.
99. Bargh J, Chartrand T: The unbearable automaticity of being. *Am Psychol* 1999, 54:462-479.
100. Bauman AE, Sallis JF, Dzawaltowski DA, Owen N: Toward a better understanding of the influences
on physical activity: the role of determinants, correlates, causal variables, mediators, moderators, and
confounders. *Am J Prev Med* 2002, 23(2 Suppl):5-14.

1 **BIBLIOGRAPHY**

2 * indicates studies included in the previous review of Sallis et al. (2000).

- 3 1.* Aarnio M, Kujala UM, Kaprio J. Associations of health-related behaviors, school type and health
4 status to physical activity patterns in 16 year old boys and girls. *Scand J Soc Med* 1997; 25:156-167.
- 5 2. Aarnio M, Winter T, Kujala U, Kaprio J. Associations of health related behaviour, social relationships,
6 and health status with persistent physical activity and inactivity: a study of Finnish adolescent twins.
7 *Br J Sports Med* 2002; 36:360-364.
- 8 3. Aarnio M, Winter T, Kujala UM, Kaprio J. Familial aggregation of leisure-time physical activity -- a
9 three generation study. *Int J Sports Med* 1997; 18:549-556.
- 10 4.* Aaron DJ, Kriska AM, Dearwater SR, Anderson RL, Olsen TL, Cauley JA, Laporte RE. The epidemi-
11 ology of leisure physical activity in an adolescent population. *Med Sci Sports Exerc* 1993; 25:847-853.
- 12 5. Adkins S, Sherwood NE, Story M, Davis M. Physical activity among African-American girls: the role
13 of parents and the home environment. *Obes Res* 2004; 12:38S-45S.
- 14 6. Allison KR, Dwyer JJ, Makin S. Perceived barriers to physical activity among high school students.
15 *Prev Med* 1999; 28:608-615.
- 16 7. Andersen LB, Schelin B. Physical activity and performance in a random sample of adolescents at-
17 tending school in Denmark. *Scand J Med Sci Sports* 1994; 4:13-18.
- 18 8.* Anderssen N, Wold B. Parental and peer influences on leisure-time physical activity in young adoles-
19 cents. *Res Q Exerc Sport* 1992; 63:341-348.
- 20 9. Anthsel KM, Anderman EM. Social influences on sports participation during adolescence. *J Res Dev*
21 *Educ* 2000; 33:85-94.
- 22 10.* Baranowski T, Thompson WO, DuRant RH, Baranowski J, Puhl J. Observations on physical activity
23 in physical locations: age, gender, ethnicity, and month effects. *Res Q Exerc Sport* 1993; 64:127-133.
- 24 11. Barnett TA, O'Loughlin J, Paradis G. One- and two-year predictors of decline in physical activity
25 among inner-city schoolchildren. *Am J Prev Med* 2002; 23:121-128.
- 26 12. Baxter-Jones AD, Maffulli N. Parental influence on sport participation in elite young athletes. *J Sports*
27 *Med Phys Fitness* 2003; 43:250-255.
- 28 13. Beech BM, Kumanyika SK, Baranowski T, Davis M, Robinson TN, Sherwood NE, Taylor WC, Relyea
29 G, Zhou A, Pratt C, Owens A, Thompson NS. Parental cultural perspectives in relation to weight-
30 related behaviors and concerns of African-American girls. *Obes Res* 2004; 12 Suppl:7S-19S.
- 31 14.* Biddle S, Goudas M. Analysis of children's physical activity and its association with adult encourage-
32 ment and social cognitive variables. *J Sch Health* 1996; 66:75-78.
- 33 15. Bogaert N, Steinbeck KS, Baur LA, Brock K, Birmingham MA. Food, activity and family - environ-
34 mental vs. biochemical predictors of weight gain in children. *Eur J Clin Nutr* 2003; 57:1242-1249.
- 35 16. Bungum T, Dowda M, Weston A, Trost SG, Pate RR. Correlates of physical activity in male and
36 female youth. *Pediatr Exerc Sci* 2000; 12:71-79.
- 37 17.* Bungum TJ, Vincent ML. Determinants of physical activity among female adolescents. *Am J Prev*
38 *Med* 1997; 13:115-122.
- 39 18.* Butcher J. Socialization of adolescent girls into physical activity. *Adolescence* 1983; 18:753-766.
19. Carlin JB, Stevenson MR, Roberts I, Bennett CM, Gelman A, Nolan T. Walking to school and traffic
exposure in Australian children. *Aust N Z J Public Health* 1997; 21:286-292.
- 20.* Crocker PR, Bailey DA, Faulkner RA, Kowalski KC, McGrath R. Measuring general levels of physical
activity: preliminary evidence for the Physical Activity Questionnaire for Older Children. *Med Sci*
Sports Exerc 1997; 29:1344-1349.

- 1 21. Damore DT. Preschool and school age activities: comparison of urban and suburban populations. *J*
- 2 *Community Health* 2002; 27:203-211.
- 3 22. Davison KK, Cutting TM, Birch LL. Parents' activity-related parenting practices predict girls' physical
- 4 activity. *Med Sci Sports Exerc* 2003; 35:1589-1595.
- 5 23. Deflandre A, Lorant J, Gavarry O, Falgairette G. Determinants of physical activity and physical and
- 6 sports activities in French school children. *Percept Mot Skills* 2001; 92:399-414.
- 7 24. Deflandre A, Lorant J, Gavarry O, Falgairette G. Physical activity and sport involvement in French
- 8 high school students. *Percept Mot Skills* 2001; 92:107-120.
- 9 25.* Dempsey JM, Kimiecik JC, Horn TS. Parental influence on children's moderate to vigorous physical
- 10 activity participation: an expectancy-value approach. *Pediatr Exerc Sci* 1993; 5:151-167.
- 11 26.* DiLorenzo TM, Stucky-Ropp RC, Vander Wal JS, Gotham HJ. Determinants of exercise among
- 12 children. II. A longitudinal analysis. *Prev Med* 1998; 27:470-477.
- 13 27. Dollman J, Norton K, Tucker G. Anthropometry, fitness and physical activity of urban and rural
- 14 south Australian children. *Pediatr Exerc Sci* 2002; 14:297-312.
- 15 28. Dowda M, Pate RR, Trost SG, Almeida M, Sirard JR. Influences of preschool policies and practices on
- 16 children's physical activity. *J Community Health* 2004; 29:183-196.
- 17 29. Duncan SC, Duncan TE, Strycker LA, Chaumeton NR. A multilevel analysis of sibling physical activ-
- 18 ity. *J Sport Exerc Psychol* 2004; 26:57-68.
- 19 30. Dunton GF, Jamner MS, Cooper DM. Assessing the perceived environment among minimally active
- 20 adolescent girls: validity and relations to physical activity outcomes. *Am J Health Promot* 2003; 18:70-
- 21 73.
- 22 31.* DuRant RH, Baranowski T, Johnson M, Thompson WO. The relationship among television watching,
- 23 physical activity, and body composition of young children. *Pediatrics* 1994; 94:449-455.
- 24 32.* Epstein LH, Paluch RA, Coleman KJ, Vito D, Anderson K. Determinants of physical activity in obese
- 25 children assessed by accelerometer and self-report. *Med Sci Sports Exerc* 1996; 28:1157-1164.
- 26 33. Fein AJ, Plotnikoff RC, Wild T, Spence JC. Perceived environment and physical activity in youth. *Int*
- 27 *J Behav Med* 2004; 11:135-142.
- 28 34. Feldman DE, Barnett T, Shrier I, Rossignol M, Abenham L. Is physical activity differentially associ-
- 29 ated with different types of sedentary pursuits? *Arch Pediatr Adolesc Med* 2003; 157:797-802.
- 30 35. Felton GM, Dowda M, Ward DS, Dishman RK, Trost SG, Saunders R, Pate RR. Differences in physical
- 31 activity between black and white girls living in rural and urban areas. *J Sch Health* 2002; 72:250-255.
- 32 36.* Ferguson KJ, Yesalis CE, Pomrehn PR, Kirkpatrick MB. Attitudes, knowledge, and beliefs as predic-
- 33 tors of exercise intent and behavior in schoolchildren. *J Sch Health* 1989; 59:112-115.
- 34 37. Finn K, Johannsen N, Specker B. Factors associated with physical activity in preschool children. *J*
- 35 *Pediatr* 2002; 140:81-85.
- 36 38. Fogelholm M, Nuutinen O, Pasanen M, Myohanen E, Saatela T. Parent-child relationship of physical
- 37 activity patterns and obesity. *Int J Obes Relat Metab Disord* 1999; 23:1262-1268.
- 38 39.* Freedson PS, Evenson S. Familial aggregation in physical activity. *Res Q Exerc Sport* 1991; 62:384-389.
- 40.* Fuchs R, Powell KE, Semmer NK, Dwyer JH, Lippert P, Hoffmeister H. Patterns of physical activity
- 41 among German adolescents: the Berlin-Bremen Study. *Prev Med* 1988; 17:746-763.
- 42.* Garcia AW, Broda MA, Frenn M, Coviak C, Pender NJ, Ronis DL. Gender and developmental dif-
- ferences in exercise beliefs among youth and prediction of their exercise behavior. *J Sch Health* 1995;
- 65:213-219.
- 42.* Garcia AW, Pender NJ, Antonakos CL, Ronis DL. Changes in physical activity beliefs and behaviors
- of boys and girls across the transition to junior high school. *J Adolesc Health* 1998; 22:394-402.

43. Garton AF, Harvey R, Price C. Influence of perceived family environment on adolescent leisure participation. *Aust J Psychol* 2004; 56:18-24.
- 44.* Gentle P, Caves R, Armstrong N, Balding J, Kirby B. High and low exercisers among 14- and 15-year-old children. *J Public Health Med* 1994; 16:186-194.
45. Gillander Gadin K, Hammarstrom A. Can school-related factors predict future health behavior among young adolescents? *Public Health* 2002; 116:22-29.
46. Gilmer MJ, Harrell JS, Miles MS, Hepworth JT. Youth characteristics and contextual variables influencing physical activity in young adolescents of parents with premature coronary heart disease. *J Pediatr Nurs* 2003; 18:159-168.
- 47.* Godin G, Shephard RJ. Normative beliefs of school children concerning regular exercise. *J Sch Health* 1984; 54:443-445.
- 48.* Godin G, Shephard RJ. Psychosocial factors influencing intentions to exercise of young students from grades 7 to 9. *Res Q Exerc Sport* 1986; 57:41-52.
- 49.* Godin G, Shephard RJ, Colantonio A. Children's perception of parental exercise: influence of sex and age. *Percept Mot Skills* 1986; 62:511-516.
50. Gomez JE, Johnson BA, Selva M, Sallis JF. Violent crime and outdoor physical activity among inner-city youth. *Prev Med* 2004; 39:876-881.
51. Goran ML, Nagy TR, Gower BA, Mazariegos M, Solomons N, Hood V, Johnson R. Influence of sex, seasonality, ethnicity, and geographic location on the components of total energy expenditure in young children: implications for energy requirements. *Am J Clin Nutr* 1998; 68:675-682.
52. Gordon-Larsen P, Harris KM, Ward DS, Popkin BM. Acculturation and overweight-related behaviors among Hispanic immigrants to the US: the National Longitudinal Study of Adolescent Health. *Soc Sci Med* 2003; 57:2023-2034.
53. Gordon-Larsen P, McMurray RG, Popkin BM. Determinants of adolescent physical activity and inactivity patterns. *Pediatrics* 2000; 105:E83.
- 54.* Gottlieb NH, Chen MS. Sociocultural correlates of childhood sporting activities: their implications for heart health. *Soc Sci Med* 1985; 21:533-539.
55. Grunbaum JA, Lowry R, Kann L. Prevalence of health-related behaviors among alternative high school students as compared with students attending regular high schools. *J Adolesc Health* 2001; 29:337-343.
- 56.* Guillaume M, Lapidus L, Bjorntorp P, Lambert A. Physical activity, obesity, and cardiovascular risk factors in children. The Belgian Luxembourg Child Study II. *Obes Res* 1997; 5:549-556.
57. Harrell JS, Gansky SA, Bradley CB, McMurray RG. Leisure time activities of elementary school children. *Nurs Res* 1997; 46:246-253.
58. Harten N, Olds T. Patterns of active transport in 11-12 year old Australian children. *Aust N Z J Public Health* 2004; 28:167-172.
59. Heath GW, Pratt M, Warren CW, Kann L. Physical activity patterns in American high school students. Results from the 1990 Youth Risk Behavior Survey. *Arch Pediatr Adolesc Med* 1994; 148:1131-1136.
60. Higgins JW, Gaul C, Gibbons S, Van Gyn G. Factors influencing physical activity levels among Canadian youth. *Can J Public Health* 2003; 94:45-51.
61. Hoefler WR, McKenzie TL, Sallis JF, Marshall SJ, Conway TL. Parental provision of transportation for adolescent physical activity. *Am J Prev Med* 2001; 21:48-51.
62. Hofstetter RC, Hovell ME, Sallis JF, Zakarian J, Beirich H, Mulvihill M, Emerson J. Exposure to sports mass media and physical activity characteristics among ethnically diverse adolescents. *Med Exerc Nutr Health* 1995; 4:234-242.

- 63.* Hovell MF, Kolody B, Sallis JF, Black DR. Parent support, physical activity, and correlates of adiposity in nine year olds: an exploratory study. *J Health Educ* 1996; 27:126-129.
64. Hussey J, Gormley J, Bell C. Physical activity in Dublin children aged 7-9 years. *Br J Sports Med* 2001; 35:268-272; discussion 273.
65. Huurre T, Aro H, Rahlkonen O. Well-being and health behaviour by parental socioeconomic status: a follow-up study of adolescents aged 16 until age 32 years. *Soc Psychiatry Psychiatr Epidemiol* 2003; 38:249-255.
66. Johansson B, Drott P. Informal parental traffic education and children's bicycling behaviour. *Ups J Med Sci* 2001; 106:133-144.
67. Karvonen S, Rimpela AH. Urban small area variation in adolescents' health behaviour. *Soc Sci Med* 1997; 45:1089-1098.
68. Kimiecik JC, Horn TS. Parental beliefs and children's moderate-to-vigorous physical activity. *Res Q Exerc Sport* 1998; 69:163-175.
- 69.* Kimiecik JC, Horn TS, Shurin CS. Relationships among children's beliefs, perceptions of their parents' beliefs, and their moderate-to-vigorous physical activity. *Res Q Exerc Sport* 1996; 67:324-336.
- 70.* Klesges RC, Eck LH, Hanson CL, Haddock C, et al. Effects of obesity, social interactions, and physical environment on physical activity in preschoolers. *Health Psychol* 1990; 9:435-449.
- 71.* Klesges RC, Malott JM, Boschee PF, Weber JM. The effects of parental influences on children's food intake, physical activity, and relative weight. *Int J Eat Disord* 1986; 5:335-346.
72. Kristjansdottir G, Vilhjalmsson R. Sociodemographic differences in patterns of sedentary and physically active behavior in older children and adolescents. *Acta Paediatr* 2001; 90:429-435.
73. Lasheras L, Aznar S, Merino B, Lopez EG. Factors associated with physical activity among Spanish youth through the National Health Survey. *Prev Med* 2001; 32:455-464.
74. Lee RE, Cubbin C. Neighborhood context and youth cardiovascular health behaviors. *Am J Public Health* 2002; 92:428-436.
75. Lewko JH, Ewing ME. Sex differences and parental influence in sport involvement of children. *J Sport Psychol* 1980; 2:62-68.
76. Lindquist CH, Reynolds KD, Goran MI. Sociocultural determinants of physical activity among children. *Prev Med* 1999; 29:305-312.
- 77.* Lowry R, Kann L, Collins JL, Kolbe LJ. The effect of socioeconomic status on chronic disease risk behaviors among US adolescents. *JAMA* 1996; 276:792-797.
78. Macintosh D. Socio-economic, educational and status characteristics of Ontario interschool athletes. *Can J Appl Sport Sci* 1982; 7:272-283.
79. McGuire MT, Hannan PJ, Neumark-Sztainer D, Cossrow NH, Story M. Parental correlates of physical activity in a racially/ethnically diverse adolescent sample. *J Adolesc Health* 2002; 30:253-261.
80. McGuire MT, Neumark-Sztainer DR, Story M. Correlates of time spent in physical activity and television viewing in a multi-racial sample of adolescents. *Pediatr Exerc Sci* 2002; 14:75-86.
- 81.* McKenzie TL, Sallis JF, Nader PR, Broyles SL, Nelson JA. Anglo- and Mexican-American preschoolers at home and at recess: activity patterns and environmental influences. *J Dev Behav Pediatr* 1992; 13:173-180.
- 82.* McKenzie TL, Sallis JF, Nader PR, Patterson TL, Elder JP, Berry CC, Rupp JW, Atkins CJ, Buono MJ, Nelson JA. BEACHES: an observational system for assessing children's eating and physical activity behaviors and associated events. *J Appl Behav Anal* 1991; 24:141-151.
83. McLellan L, Rissel C, Donnelly N, Bauman A. Health behaviour and the school environment in New South Wales, Australia. *Soc Sci Med* 1999; 49:611-619.

- 84.* McMurray RG, Bradley CB, Harrell JS, Bernthal PR, Frauman AC, Bangdiwala SI. Parental influences on childhood fitness and activity patterns. *Res Q Exerc Sport* 1993; 64:249-255.
85. McMurray RG, Harrell JS, Bangdiwala SI, Deng S. Cardiovascular disease risk factors and obesity of rural and urban elementary school children. *J Rural Health* 1999; 15:365-374.
86. McMurray RG, Harrell JS, Bangdiwala SI, Gansky SA. Biologic and environmental factors influencing the aerobic power of children. *Med Exerc Nutr Health* 1995; 4:243-250.
87. Mellin AE, Neumark-Sztainer D, Story M, Ireland M, Resnick MD. Unhealthy behaviors and psychosocial difficulties among overweight adolescents: the potential impact of familial factors. *J Adolesc Health* 2002; 31:145-153.
88. Molnar BE, Gortmaker SL, Bull FC, Buka SL. Unsafe to play? Neighborhood disorder and lack of safety predict reduced physical activity among urban children and adolescents. *Am J Health Promot* 2004; 18:378-386.
- 89.* Moore LL, Lombardi DA, White MJ, Campbell JL, Oliveria SA, Ellison RC. Influence of parents' physical activity levels on activity levels of young children. *J Pediatr* 1991; 118:215-219.
90. Morgan CF, McKenzie TL, Sallis JF, Broyles SL, Zive MM, Nader PR. Personal, social, and environmental correlates of physical activity in a bi-ethnic sample of adolescents. *Pediatr Exerc Sci* 2003; 15:288-301.
91. Motl RW, Dishman RK, Ward DS, Saunders RP, Dowda M, Felton G, Pate RR. Examining social-cognitive determinants of intention and physical activity among black and white adolescent girls using structural equation modeling. *Health Psychol* 2002; 21:459-467.
92. Murphey DA, Lamonda KH, Carney JK, Duncan P. Relationships of a brief measure of youth assets to health-promoting and risk behaviors. *J Adolesc Health* 2004; 34:184-191.
93. Neumark-Sztainer D, Story M, Hannan PJ, Tharp T, Rex J. Factors associated with changes in physical activity: a cohort study of inactive adolescent girls. *Arch Pediatr Adolesc Med* 2003; 157:803-810.
94. Nutbeam D, Aar L, Catford J. Understanding childrens' health behaviour: the implications for health promotion for young people. *Soc Sci Med* 1989; 29:317-325.
95. O'Loughlin J, Paradis G, Kishchuk N, Barnett T, Renaud L. Prevalence and correlates of physical activity behaviors among elementary schoolchildren in multiethnic, low income, inner-city neighborhoods in Montreal, Canada. *Ann Epidemiol* 1999; 9:397-407.
96. Pate RR, Pfeiffer KA, Trost SG, Ziegler P, Dowda M. Physical activity among children attending preschools. *Pediatrics* 2004; 114:1258-1263.
- 97.* Pate RR, Trost SG, Felton GM, Ward DS, Dowda M, Saunders R. Correlates of physical activity behavior in rural youth. *Res Q Exerc Sport* 1997; 68:241-248.
- 98.* Pérusse L, Leblanc C, Bouchard C. Familial resemblance in lifestyle components: results from the Canada Fitness Survey. *Can J Public Health* 1988; 79:201-205.
- 99.* Pérusse L, Tremblay A, Leblanc C, Bouchard C. Genetic and environmental influences on level of habitual physical activity and exercise participation. *Am J Epidemiol* 1989; 129:1012-1022.
- 100.* Poest CA, Williams JR, Witt DD, Atwood ME. Physical activity patterns of preschool children. *Early Child Res Q* 1989; 4:367-376.
101. Prochaska JJ, Rodgers MW, Sallis JF. Association of parent and peer support with adolescent physical activity. *Res Q Exerc Sport* 2002; 73:206-210.
- 102.* Reynolds KD, Killen JD, Bryson SW, Maron DJ, Taylor CB, Maccoby N, Farquhar JW. Psychosocial predictors of physical activity in adolescents. *Prev Med* 1990; 19:541-551.
103. Robinson CH, Thomas SP. The Interaction Model of Client Health Behavior as a conceptual guide in the explanation of children's health behaviors. *Public Health Nurs* 2004; 21:73-84.

104. Romero AJ, Robinson TN, Kraemer HC, Erickson SJ, Haydel KF, Mendoza F, Killen JD. Are perceived neighborhood hazards a barrier to physical activity in children? *Arch Pediatr Adolesc Med* 2001; 155:1143-1148.
105. Rossow I, Rise J. Concordance of parental and adolescent health behaviors. *Soc Sci Med* 1994; 38:1299-1305.
106. Runyan SM, Stadler DD, Bainbridge CN, Miller SC, Moyer-Mileur LJ. Familial resemblance of bone mineralization, calcium intake, and physical activity in early-adolescent daughters, their mothers, and maternal grandmothers. *J Am Diet Assoc* 2003; 103:1320-1325.
- 107.* Sallis JF, Alcaraz JE, McKenzie TL, Hovell MF. Predictors of change in children's physical activity over 20 months: Variations by gender and level of adiposity. *Am J Prev Med* 1999; 16:222-229.
- 108.* Sallis JF, Alcaraz JE, McKenzie TL, Hovell MF, Kolody B, Nader PR. Parental behavior in relation to physical activity and fitness in 9-year-old children. *Am J Dis Child* 1992; 146:1383-1388.
- 109.* Sallis JF, Nader PR, Broyles SL, Berry CC, Elder JP, McKenzie TL, Nelson JA. Correlates of physical activity at home in Mexican-American and Anglo-American preschool children. *Health Psychol* 1993; 12:390-398.
- 110.* Sallis JF, Patterson TL, Buono MJ, Atkins CJ, Nader PR. Aggregation of physical activity habits in Mexican-American and Anglo families. *J Behav Med* 1988; 11:31-41.
- 111.* Sallis JF, Patterson TL, McKenzie TL, Nader PR. Family variables and physical activity in preschool children. *J Dev Behav Pediatr* 1988; 9:57-61.
112. Sallis JF, Prochaska JJ, Taylor WC, Hill JO, Geraci JC. Correlates of physical activity in a national sample of girls and boys in grades 4 through 12. *Health Psychol* 1999; 18:410-415.
113. Sallis JF, Taylor WC, Dowda M, Freedson PS, Pate RR. Correlates of vigorous physical activity for children in grades 1 through 12: Comparing parent-reported and objectively measured physical activity. *Pediatr Exerc Sci* 2002; 14:30-44.
114. Saunders RP, Motl RW, Dowda M, Dishman RK, Pate RR. Comparison of social variables for understanding physical activity in adolescent girls. *Am J Health Behav* 2004; 28:426-436.
115. Saunders RP, Pate RR, Felton G, Dowda M, Weinrich MC, Ward DS, Parsons MA, Baranowski T. Development of questionnaires to measure psychosocial influences on children's physical activity. *Prev Med* 1997; 26:241-247.
116. Saxena R, Borzekowski DL, Rickert VI. Physical activity levels among urban adolescent females. *J Pediatr Adolesc Gynecol* 2002; 15:279-284.
117. Schmitz KH, Lytle LA, Phillips GA, Murray DM, Birnbaum AS, Kubik MY. Psychosocial correlates of physical activity and sedentary leisure habits in young adolescents: the Teens Eating for Energy and Nutrition at School study. *Prev Med* 2002; 34:266-278.
- 118.* Shephard RJ, Jequier JC, Lavallee H, La Barre R, Rajic M. Habitual physical activity: effects of sex, milieu, season and required activity. *J Sports Med Phys Fitness* 1980; 20:55-66.
119. Shropshire J, Carrol B. Family variables and children's physical activity: influence of parental exercise and socio-economic status. *Sport Educ Soc* 1997; 2:95-116.
120. Simons-Morton BG, McKenzie TJ, Stone E, Mitchell P, Osganian V, Strikmiller PK, Ehlinger S, Cribb P, Nader PR. Physical activity in a multiethnic population of third graders in four states. *Am J Public Health* 1997; 87:45-50.
121. Starfield B, Riley AW, Witt WP, Robertson J. Social class gradients in health during adolescence. *J Epidemiol Community Health* 2002; 56:354-361.
122. Starfield B, Robertson J, Riley AW. Social class gradients and health in childhood. *Ambul Pediatr* 2002; 2:238-246.

123. Strauss RS, Rodzilsky D, Burack G, Colin M. Psychosocial correlates of physical activity in healthy children. *Arch Pediatr Adolesc Med* 2001; 155:897-902.
- 124.* Stucky-Ropp RC, DiLorenzo TM. Determinants of exercise in children. *Prev Med* 1993; 22:880-889.
- 125.* Tappe MK, Duda JL, Ehrnwald PM. Perceived barriers to exercise among adolescents. *J Sch Health* 1989; 59:153-155.
- 126.* Tappe MK, Duda JL, Menges-Ehrnwald P. Personal investment predictors of adolescent motivational orientation toward exercise. *Can J Sport Sci* 1990; 15:185-192.
127. Taveras EM, Rifas-Shiman SL, Field AE, Frazier AL, Colditz GA, Gillman MW. The influence of wanting to look like media figures on adolescent physical activity. *J Adolesc Health* 2004; 35:41-50.
- 128.* Terre L, Ghiselli W, Taloney L, DeSouza E. Demographics, affect, and adolescents' health behaviors. *Adolescence* 1992; 27:12-24.
- 129.* Theodorakis Y, Doganis G, Bagiatis K, Gouthas M. Preliminary study of the ability of reasoned action model in predicting exercise behaviour of young children. *Percept Mot Skills* 1991; 72:51-58.
130. Theodorakis Y, Papaioannou A, Karastogianidou K. Relations between family structure and students' health-related attitudes and behaviors. *Psychol Rep* 2004; 95:851-858.
131. Timperio A, Crawford D, Telford A, Salmon J. Perceptions about the local neighborhood and walking and cycling among children. *Prev Med* 2004; 38:39-47.
- 132.* Trost SG, Pate RR, Dowda M, Saunders R, Ward DS, Felton G. Gender differences in physical activity and determinants of physical activity in rural fifth grade children. *J Sch Health* 1996; 66:145-150.
- 133.* Trost SG, Pate RR, Saunders R, Ward DS, Dowda M, Felton G. A prospective study of the determinants of physical activity in rural fifth-grade children. *Prev Med* 1997; 26:257-263.
134. Trost SG, Pate RR, Ward DS, Saunders R, Riner W. Correlates of objectively measured physical activity in preadolescent youth. *Am J Prev Med* 1999; 17:120-126.
135. Trost SG, Sallis JF, Pate RR, Freedson PS, Taylor WC, Dowda M. Evaluating a model of parental influence on youth physical activity. *Am J Prev Med* 2003; 25:277-282.
136. Tuinstra J, Groothoff JW, van den Heuvel WJ, Post D. Socio-economic differences in health risk behavior in adolescence: do they exist? *Soc Sci Med* 1998; 47:67-74.
137. Unger JB, Reynolds K, Shakib S, Spruijt-Metz D, Sun P, Johnson CA. Acculturation, physical activity, and fast-food consumption among Asian-American and Hispanic adolescents. *J Community Health* 2004; 29:467-481.
138. Vermorel M, Vernet J, Bitar A, Fellmann N, Coudert J. Daily energy expenditure, activity patterns, and energy costs of the various activities in French 12-16-y-old adolescents in free living conditions. *Eur J Clin Nutr* 2002; 56:819-829.
139. Vilhjalmsón R. Effects of social support on self-assessed health in adolescence. *J Adolesc Health* 1994; 23:437-452.
140. Vilhjalmsón R, Kristjansdóttir G. Gender differences in physical activity in older children and adolescents: the central role of organized sport. *Soc Sci Med* 2003; 56:363-374.
141. Vilhjalmsón R, Thorlindsson T. Factors related to physical activity: a study of adolescents. *Soc Sci Med* 1998; 47:665-675.
142. Wagner A, Klein-Platat C, Arveiler D, Haan MC, Schlienger JL, Simon C. Parent-child physical activity relationships in 12-year old French students do not depend on family socioeconomic status. *Diabetes Metab* 2004; 30:359-366.
143. Wardle J, Jarvis M, Steggle N, Sutton S, Williamson S, Farrimond H, Cartwright M, Simon AE. Socioeconomic disparities in cancer-risk behaviors in adolescence: baseline results from the Health and Behaviour in Teenagers Study (HABITS). *Prev Med* 2003; 36:721-730.

- 1 144. Welk GJ, Wood K, Morss G. Parental influences on physical activity in children: An exploration of
2 potential mechanisms. *Pediatr Exerc Sci* 2003; 15:19-33.
- 3 145. Williams EA, Jenkins C, Nevill AM. Social area influences on leisure activity - an exploration of the
4 ACORN classification with reference to sport. *Leisure Studies* 1988; 7:81-94.
- 5 146. Wold B, Oygard L, Eder A, Smith C. Social reproduction of physical activity, Implications for health
6 promotion in young people. *Eur J Public Health* 1994; 4:163-168.
- 7 147. Woodfield L, Duncan M, Al-Nakeeb Y, Nevill A, Jenkins C. Sex, ethnic and socio-economic differ-
8 ences in children's physical activity. *Pediatr Exerc Sci* 2002; 14:277-285.
- 9 148. Yang X, Telama R, Laakso L. Parent's physical activity, socioeconomic status and education as predic-
10 tors of physical activity and sport among children and youths - a 12-year follow-up study. *Int Rev Soc*
11 *Sports* 1996; 31:273-291.
- 12 149.* Zakarian JM, Hovell MF, Hofstetter CR, Sallis JF, Keating KJ. Correlates of vigorous exercise in a
13 predominantly low SES and minority high school population. *Prev Med* 1994; 23:314-321.
- 14 150. Ziviani J, Scott J, Wadley D. Walking to school: incidental physical activity in the daily occupations of
15 Australian children. *Occup Ther Int* 2004; 11:1-11.
- 16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

Part III Socio-demographic correlates of energy balance- related behaviors

5 Gender, ethnic and school type differences in overweight and energy balance-related behaviors among Dutch adolescents

van der Horst K, Oenema A, te Velde SJ, Brug J. Gender, ethnic and school type differences in overweight and energy balance-related behaviors among Dutch adolescents.

International Journal of Pediatric Obesity 2009, May 15:1-10. [Epub ahead of print].

1 **ABSTRACT**

2

3 **Objective:** The aim of this study was to investigate gender, ethnic and school type differ-
4 ences in overweight and energy balance-related behaviors: snack, soft drink and breakfast
5 consumption, walking, bicycling, and playing sports during leisure time, active commuting
6 to school, television viewing and computer use among 12- 15- year-old adolescents.

7 **Methods:** Cross-sectional data on weight status and energy balance-related behaviors were
8 obtained from 1206 adolescents (2005-2006). Energy balance-related behaviors were self-
9 reported and body mass index was calculated from measured height and weight. Gender,
10 ethnic and school type differences in weight status and behaviors were examined with
11 multi-level logistic regression analyses.

12 **Results:** Overweight and unfavorable energy balance-related behaviors were more
13 likely among youth from non-Western ethnic backgrounds and those attending vocational
14 schools. Analyses stratified by ethnicity showed that girls from non-Western ethnic back-
15 grounds were more likely to be overweight (OR=1.5) and to report not bicycling (OR=2.4)
16 and watching more than two hours of television (OR=2.3) compared to boys from non-
17 Western ethnic backgrounds. Vocational students from Western ethnic backgrounds were
18 more likely to report high levels of soft drink consumption (OR=3.2), watching television
19 (OR=2.9) and computer use (OR=2.1) compared to higher-level education students from
20 Western ethnic backgrounds.

21 **Conclusions:** The study findings indicate important ethnic and educational differences in
22 overweight and energy balance-related behaviors. Future research should focus on what
23 kind of interventions work and for which target groups they work, taking demographic
24 variables such as gender, ethnicity, school type into account.

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

1 INTRODUCTION

2
3 To curb the obesity epidemic, it is important to identify and target adolescents at risk
4 for overweight and obesity. Because obesity persists into adulthood [1] and is associated
5 with severe health consequences [2], a detailed understanding of risk behaviors related to
6 the development of obesity is essential to developing preventive interventions. It is also
7 important to identify specific target groups of adolescents who are more at risk of becoming
8 obese by engaging in more (or specific) obesity-related risk behaviors. Being able to
9 distinguish specific target groups provides the opportunity to better tailor interventions
10 to the needs and perceptions of those most at risk [3]. Recent overviews have suggested a
11 range of specific energy balance-related behaviors (EBRB, i.e. behaviors that contribute to
12 energy intake or expenditure) that may contribute substantially to a higher or lower risk for
13 unnecessary weight gain [4-6]. Currently, there is insufficient insight into the occurrence
14 of a number of overweight risk behaviors among adolescents and whether it is possible to
15 distinguish specific subgroups that are more likely to engage in specific risk behaviors for
16 overweight and obesity.

17
18 Earlier evidence points out that the prevalence of overweight is considerably greater among
19 youth from racial or ethnic minority backgrounds [7-9]. In addition to genetic, economic
20 and environmental factors, ethnic disparities in overweight and obesity are likely due to
21 differences in EBRB [10, 11]. Studies conducted in different countries indicate that ethnic
22 minority groups participate less in physical activity, spend more time watching television [5,
23 12-14], are less likely to eat breakfast regularly [13], consume soft drinks and savory snacks
24 and visit fast-food restaurants more often [15], but also have higher fruit intakes [16]. The
25 higher rates of overweight among ethnic minority groups might be explained in part by
26 their lower educational levels. Educational level has also been found to be an independent
27 determinant of overweight [17] and to be associated with adolescent health behavior such
28 as physical activity [18]. Furthermore, differences have been found between boys and girls
29 in overweight and obesity and related risk behaviors, with girls being more likely to be
30 overweight [9, 19, 20] and to engage in less physical activity [14, 21].

31
32 Although most of the previous studies examined single behaviors, it is most likely that a
33 number of risk behaviors contribute to an increased risk for overweight. Therefore, we examined
34 how overweight and specific unfavorable EBRB (high soft drink intake, high snack
35 consumption, not eating breakfast on a daily basis, high amounts of television viewing and
36 computer use, little participation in sports, little walking and bicycling during leisure time,
37 and an absence of active commuting to school) vary by gender, ethnicity and school type. In
38 addition, we investigated possible interaction effects between gender, ethnicity and school
39 type, and performed stratified analyses when interaction effects were significant. Based on

1 the findings of previous studies, we hypothesized that girls, adolescents from non-Western
2 ethnic backgrounds, and adolescents attending vocational schools have a higher risk of
3 overweight and obesity and unfavorable EBRB compared to boys, adolescents from Western
4 ethnic backgrounds, and adolescents participating in higher-level education.

7 **METHODS**

9 Study design and sample selection

11 Data from the ENvironmental Determinants of Obesity in Rotterdam SchoolchildrEn
12 (ENDORSE) study were used. The ENDORSE study is a prospective two-year study among
13 adolescents aged 12 to 15, with assessments at baseline and a two-year follow-up. More de-
14 tails on this project are described elsewhere [22]. The Medical Ethics Committee of Erasmus
15 University Medical Center reviewed the proposal and issued a “declaration of no objection”
16 for the ENDORSE project.

17 After stratification according to the area in the city in which the schools are located,
18 seventeen school locations were randomly selected from 24 out of a total of 54 schools
19 that were willing to participate in the ENDORSE study. Stratification was done, to ensure a
20 range of physical and cultural environments. An average of five classes per school location
21 was randomly selected to participate in the study, and 1668 adolescents from these classes
22 were eligible for participation. In the baseline survey, 187 adolescents were absent during the
23 questionnaire assessment. Due to printing mistakes, it was necessary to delete 120 adoles-
24 cents, including one entire school. Respondents with missing data on ethnicity were deleted
25 from the sample (n=155). This meant that the study sample included 1206 participants (72%)
26 from 71 classes and 16 schools.

28 Procedure

30 The ENDORSE study collects data among adolescents in the first (12- to 13-year-olds) and
31 third (14- to 15-year-olds) years of secondary school. The school types varied from lower
32 vocational schools to higher-level secondary education. All data were gathered within the
33 ongoing health surveillance system of the local Municipal Health Service and as a part of the
34 government approved routine health examinations of the preventive youth health care [22].
35 Separate informed consent therefore was not requested. ENDORSE classroom question-
36 naires and anthropometrics were completed on a voluntary basis. Parents received written
37 information on these measurements and were free to object to participation of their child.
38 From October 2005 to May 2006, the students completed the ENDORSE questionnaire
39 in the classroom in the presence of a teacher and a trained research assistant during one

1 class period of approximately 50 minutes. Within a month after completing the ENDORSE
 2 questionnaire, two trained research assistants measured height and weight. The adolescents
 3 were asked to come into a private room one by one, where they were measured in street
 4 clothes without shoes and heavy clothes.

5 6 Measures

7 8 *Weight, height and body mass index*

9 Body height was measured using a Seca 225 mobile height rod with an accuracy of 0.1 cm. A
 10 calibrated electronic digital floor scale (Seca 888 class III with accuracy of 0.1 kg) was used to
 11 determine the body weight of the participants. Body mass index (BMI) was calculated from
 12 the measured height and weight (kg/m^2). BMI cut points for children/adolescents from the
 13 International Obesity TaskForce (IOTF) were used to define overweight and obesity [23].

14 15 *Energy balance-related behaviors*

16 The following EBRB were assessed: snack, soft drink and breakfast consumption, walking,
 17 bicycling and playing sports during leisure time, active commuting to school (walking and
 18 bicycling), television viewing and computer use.

19 Snacks were defined as sweet (candy, candy bars, chocolate, cake, cookies) and savory
 20 (fast food, pizza, fries, chips, nuts) snacks. Sweet snacks were assessed by two questions:
 21 “How many days a week do you usually eat sweet snacks or cookies?” and “On average, how
 22 many times a day do you eat sweet snacks or cookies?” The same two questions were asked
 23 for savory snacks. These questions were combined to compute a single score for the mean
 24 snack intake in times per day.

25 Soft drinks were defined as carbonated soft drinks, other non-carbonated sugar-sweetened
 26 drinks (water-based beverages that contain sugar) and sport drinks. The consumption of
 27 soft drinks was assessed by two questions: “How many days a week do you usually drink
 28 sugar-sweetened (not ‘light’ or ‘diet’) beverages?” and “If you drink sugar-sweetened bever-
 29 ages, how many glasses, cans, and/or bottles do you drink on average per day?” Total soft
 30 drink consumption was expressed in milliliters per day and therefore, serving sizes were
 31 transferred to a quantification in milliliters using the Dutch standard serving sizes (1 glass =
 32 200 ml, 1 can = 330 ml, 1 bottle = 500ml).

33 Breakfast consumption was assessed with two questions: “How often do you eat breakfast
 34 on school days?” and “How often do you eat breakfast on weekend days?” These questions
 35 were combined to compute a single score for breakfast consumption in days per week.

36 Ten-day test-retest reliability for snack consumption, soft drink consumption and break-
 37 fast consumption was $r=0.46$, $r=0.59$ and $r=0.65$ respectively.

38 An adapted version of the Activity QUEStionnaire for Adolescents & Adults (AQUAA)
 39 was used to assess physical activity (transportation, activities and sports during leisure time)

1 and sedentary behaviors (television viewing, computer use). The structure of the AQUAA
2 is obtained from the validated Short QUESTIONNAIRE to ASses Health-enhancing physical
3 activity (SQUASH) [24]. The frequency (days per week) and duration (hours and minutes)
4 of the activities were multiplied, and then divided by the total number of days to provide the
5 average minutes per day of the physical activity and sedentary sub-behaviors.

6 As most behaviors were not normally distributed the EBRB were dichotomized to express
7 these behaviors in “favorable” and “unfavorable” categories. If possible, categories were
8 distinguished based on participating in a behavior (favorable group) and not participating
9 in a behavior (unfavorable group) (bicycling during leisure time, playing sports and com-
10 muting to school). Variables that could not be dichotomized by engaging or not engaging
11 in behavior were dichotomized based on recommendations (television viewing, soft drink
12 consumption) or the median value in the data set (snacking, breakfast consumption, walk-
13 ing, computer use).

14 The questions, response options and the cut-off points for all EBRB are provided in the
15 appendix to this paper.

17 *Demographics*

18 Ethnicity was defined according to the definition used by Statistics Netherlands [25]. Ado-
19 lescents were considered to be from a Western ethnic background if both parents had been
20 born in an European country, North America, Oceania, Indonesia or Japan. Adolescents
21 were considered to be from a non-Western ethnic background if one or both parents had
22 been born in a non-Western country. School type was categorized into two levels: vocational
23 schools and higher-level secondary education. The schools provided this information. Age
24 was determined based on the date of the anthropometrical measurements and the date of
25 birth (provided by the schools).

27 *Data analyses*

28 Chi-square tests were used to test differences in gender, school type, overweight status and
29 EBRB between the participants who were included and not included in the analyses (i.e.
30 those with and without data on ethnicity). Chi-square tests were also used to test differences
31 in gender, ethnicity and school type between adolescents with missing values on the EBRB
32 and weight status and adolescents with reported EBRB and weight status. Respondents with
33 missing data on the behavior variables were not deleted from the sample, but were deleted
34 from the analysis. Because of this, the numbers of students in the analyses are different for
35 various outcome variables.

36 Categorical data were described using frequencies and percentages. To examine if gender,
37 ethnicity and school type are significant correlates of the EBRB and weight status, multi-level
38 logistic regression analyses were performed with the demographic factors as independent
39 variables.

1 Interaction effects between gender and ethnicity and school type and ethnicity were
 2 examined by adding interaction terms into the regression models. If the interactions had
 3 P values <0.1 , stratified analyses were conducted for ethnicity. Multi-level models with a
 4 three-level structure were used (child, class and school) in order to take into account that
 5 children were nested within classes and schools. The univariate analyses were conducted
 6 in SPSS version 15 and the multi-level analyses were performed using MlwiN version 2.02.

9 RESULTS

11 Adolescents from Western ethnic backgrounds and adolescents attending vocational
 12 schools had significantly more missing values on several EBRB compared to adolescents
 13 from Western ethnic backgrounds and adolescents attending higher-level education (data
 14 not shown).

15 Table 5.1 presents the frequency of demographics, weight status and EBRB in the study
 16 population. Among the respondents, 45.1% was female, the mean age was 14.1 years, 56.7%
 17 was attending vocational schools and 50.5% was from non-Western ethnic background.
 18 Overweight or obesity was present in 20.2% of the participating adolescents. Unfavorable
 19 sedentary and dietary behaviors such as consuming more than two glasses of soft drink per
 20 day (67.2%) were more often reported compared to unfavorable physical activity behaviors
 21 such as not playing sports (19.8%).

23 Gender differences

25 In the multivariate analyses (Table 5.2), we found girls to be more likely to report low
 26 breakfast consumption (OR=1.8), not bicycling during leisure time (OR=1.6), no sports
 27 participation (OR=3.0) and high television viewing (OR=1.8). Girls were less likely to report
 28 high soft drink consumption (OR=0.6). There were no differences between boys and girls
 29 in weight status.

31 Ethnic differences

33 The multivariate analyses (Table 5.2) demonstrated that adolescents from non-Western eth-
 34 nic backgrounds were more likely to be overweight or obese (OR=1.8), to not eat breakfast
 35 everyday (OR=1.9), to do no bicycling during leisure time (OR=3.2), to not participate in
 36 sports (OR=1.7), to use non-active modes of transportation to school (OR=1.6) and to spend
 37 more than two hours watching television (OR=2.4). A significant inverse association was
 38 found for walking during leisure time, indicating that adolescents from non-Western ethnic
 39 backgrounds were less likely to report low levels of walking (OR=0.6).

Table 5.1 Frequency of demographics, weight status and energy balance-related behaviors.

	Frequency in study population (%) (unless otherwise specified)
Demographic variables	
Mean age of respondents in years	14.1 (SD=1.2; range 10-17)
Girls	544 (45.1)
Non-Western ethnic background	609 (50.5)
Vocational education	684 (56.7)
Weight status	
Overweight (according to IOTF)	189 (15.7)
Obesity (according to IOTF)	54 (4.5)
Mean BMI	20.7 (SD=3.6; range 14.2-36.1)
Boys	20.2 (SD=3.6)
Girls	21.3 (SD=3.6)
Western ethnic background	20.0 (SD=3.1)
Non Western ethnic background	21.4 (SD=4.0)
Vocational education	21.1 (SD=3.8)
Higher-level education	20.2 (SD=3.3)
Dietary behaviours	
Soft drinks > 2 glasses/day	811 (67.2)
Snacks > 2 times/day	603 (50.0)
Breakfast 0-6 days/week	525 (43.5)
Physical activity	
Walking during leisure time < 60 min/day	531 (44.0)
Not bicycling during leisure time	344 (28.5)
Not playing sports	239 (19.8)
Non-active commuting to school	398 (33.0)
Sedentary behaviours	
Television viewing > 120 min/day	490 (40.6)
Computer use > 90 min/day	568 (47.1)

School type differences

The multivariate analyses (Table 5.2) showed that vocational students were more likely to be overweight or obese (OR=1.7) and to report high soft drink consumption (OR=2.0), high snack consumption (OR=1.5), no sports participation (OR=2.5) and to spend more than two hours watching television (OR=1.7). A significant inverse association was found for walking during leisure time, indicating that vocational students were less likely to report low levels of walking (OR=0.5).

Table 5.2 Results of multiple logistic regression analyses (odds ratios (OR) and 95% confidence intervals (95% CI)) with gender, ethnicity and school type as independent variables and energy-balance related behaviours as dependent variables

	Gender (girls)* OR (95% CI)	Ethnicity (non- Western)* OR (95% CI)	School type (vocational)* OR (95% CI)
Weight status			
Normal weight	1.00	1.00	1.00
Overweight and obesity	1.2 (0.91-1.65)	1.8 (1.29-2.39)	1.7 (1.19-2.33)
Dietary behaviours			
Soft drink consumption			
≤ 2 glasses	1.00	1.00	1.00
> 2 glasses	0.6 (0.43-0.76)	1.2 (0.88-1.64)	2.0 (1.19-3.22)
Breakfast consumption			
Every day	1.00	1.00	1.00
0-6 days/week	1.8 (1.30-2.36)	1.9 (1.35-2.58)	1.5 (0.88-2.39)
Snack consumption			
≤ 2 times/day	1.00	1.00	1.00
> 2 times/day	1.0 (0.80-1.33)	1.2 (0.91-1.52)	1.5 (1.16-2.00)
Physical activity			
Walking during leisure time			
≥60 min/day	1.00	1.00	1.00
<60 min/day	1.1 (0.79-1.39)	0.6 (0.46-0.85)	0.5 (0.32-0.79)
Bicycling			
Bicycling during leisure time	1.00	1.00	1.00
Not bicycling during leisure time	1.6 (1.14-2.12)	3.2 (2.29-4.58)	0.9 (0.55-1.34)
Sports			
Playing sports during leisure time	1.00	1.00	1.00
Not playing sports during leisure time	3.0 (1.98-4.48)	1.7 (1.11-2.62)	2.5 (1.37-4.43)
Commuting to school (walking/bicycling)			
Active commuting	1.00	1.00	1.00
Non-active commuting	0.9 (0.66-1.25)	1.6 (1.13-2.32)	1.4 (0.62-3.20)
Sedentary behaviours			
Television viewing			
≤ 2 hours per day	1.00	1.00	1.00
> 2 hours per day	1.8 (1.33-2.38)	2.4 (1.74-3.25)	1.7 (1.08-2.71)
Computer use			
≤ 90 minutes per day	1.00	1.00	1.00
> 90 minutes per day	0.8 (0.59-1.04)	1.3 (0.98-1.80)	1.4 (0.90-2.14)

Odds ratios in **bold** indicate a significant association. All analyses were adjusted for age.

* Reference groups were boys (gender), Western ethnic background (ethnicity) and higher-level education students (school type).

1 Interaction effects between ethnicity and gender

2
3 Interaction effects between ethnicity and gender were significant for weight status, soft
4 drink consumption, bicycling during leisure time, playing sports, television viewing and
5 computer use. The stratified analyses (Table 5.3) showed that compared to boys from non-
6 Western ethnic backgrounds, girls from non-Western ethnic backgrounds were more likely
7 to be overweight (OR=1.5), to do no bicycling during leisure time (OR=2.4) and to spend
8 more than two hours watching television (OR=2.3). These gender differences were not
9 significant for adolescents from Western ethnic backgrounds. Compared to boys, girls from
10 both non-Western (OR=3.5) and Western ethnic backgrounds (OR=2.0) were more likely to
11 report no sports participation. Compared to boys from Western ethnic backgrounds, girls
12 from Western ethnic backgrounds were less likely to report high soft drink consumption
13 (OR=0.4) and using the computer for more than 90 minutes (OR=0.6).

15 Interaction effects between ethnicity and school type

16
17 Interaction effects between ethnicity and school type were significant for soft drink
18 consumption, television viewing and computer use. The stratified analyses showed that
19 compared to high-level education students from Western ethnic backgrounds, vocational
20 students from Western ethnic backgrounds were more likely to report high soft drink con-
21 sumption (OR=3.2), more than two hours of television viewing (OR=2.9) and more than
22 90 minutes of computer use (OR=2.1). There were no significant school type differences for
23 adolescents from non-Western ethnic backgrounds.

26 DISCUSSION

27
28 This study examined gender, ethnic and school type differences in weight status and EBRB
29 among Dutch adolescents aged 12 to 15. As expected, we found girls to be more likely than
30 boys to engage in unfavorable behaviors (i.e. low breakfast consumption, not bicycling
31 during leisure time, not playing sports and spending more than two hours a day watching
32 television). Adolescents from non-Western ethnic backgrounds and vocational schools were
33 more likely to be overweight or obese and to engage in unfavorable EBRB. These results are
34 in accordance with studies from other countries in which girls were also found to be less
35 physically active [14, 19, 21] and minority groups were more likely to be overweight and
36 showed more unhealthy EBRB [5, 7-9, 12-15, 26] such as watching television [5, 12-14], low
37 breakfast consumption [13], soft drink and snack consumption [15]. That ethnic minority
38 groups have higher rates of overweight and unhealthy behaviors might partly be due to
39 their lower educational levels. However the examination of potential ethnicity by school

Table 5.3 Results of multiple logistic regression analyses with energy balance-related behaviours as dependent variables (odds ratios (OR) and 95% confidence intervals (95% CI)) stratified by ethnicity with gender and school type as independent variables

	Gender (girls)*		School type (vocational)*	
	Western OR (95% CI)	Non-Western OR (95% CI)	Western OR (95% CI)	Non-Western OR (95% CI)
Weight status				
Normal weight	1.00	1.00	1.00	1.00
Overweight	0.9 (0.55-1.51)	1.5 (1.05-2.26)	NA	NA
Dietary behaviours				
Soft drink consumption				
≤ 2 glasses	1.00	1.00	1.00	1.00
> 2 glasses	0.4 (0.28-0.64)	0.7 (0.50-1.08)	3.2 (1.63-6.34)	1.5 (0.92-2.30)
Breakfast consumption				
Every day	1.00	1.00	1.00	1.00
0-6 days/week	NA	NA	NA	NA
Snack consumption				
≤ 2 times/day	1.00	1.00	1.00	1.00
> 2 times/day	NA	NA	NA	NA
Physical activity				
Walking during leisure time				
≥60 min/day	1.00	1.00	1.00	1.00
<60 min/day	NA	NA	NA	NA
Bicycling				
Bicycling during leisure time	1.00	1.00	1.00	1.00
Not bicycling during leisure time	0.8 (0.45-1.25)	2.4 (1.64-3.45)	NA	NA
Sports				
Playing sports during leisure time	1.00	1.00	1.00	1.00
Not playing sports during leisure time	2.0 (1.07-3.77)	3.5 (2.20-5.67)	NA	NA
Commuting to school (walking/bicycling)				
Active commuting	1.00	1.00	1.00	1.00
Non-active commuting	NA	NA	NA	NA
Sedentary behaviours				
Television viewing				
≤ 2 hours per day	1.00	1.00	1.00	1.00
> 2 hours per day	1.1 (0.69-1.63)	2.3 (1.60-3.30)	2.9 (1.55-5.46)	1.3 (0.82-1.96)
Computer use				
≤ 90 minutes per day	1.00	1.00	1.00	1.00
> 90 minutes per day	0.6 (0.38-0.85)	0.9 (0.66-1.35)	2.1 (1.25-3.59)	1.2 (0.70-1.97)

NA = not applicable, no significant interaction by ethnicity. Odds ratios in **bold** indicate a significant association. All analyses were adjusted for age. * Reference groups were boys (gender) and higher-level education students (school type).

Appendix Items on the energy balance-related behaviours questionnaire

Items	Response categories
Sweet and savoury snacks (≤ 2 times/day; > 2 times/day)	
How many days a week do you usually eat sweet snacks or cookies?	8-point scale: from 0 = never to 7 = 7 days per week
On average, how many times a day do you eat sweet snacks or cookies?	10-point scale: from 1 = 1x to 10 = 10x or more
How many days a week do you usually eat savoury snacks?	8-point scale: from 0 = never to 7 = 7 days per week
On average, how many times a day do you eat savoury snacks?	10-point scale: from 1 = 1x to 10 = 10x or more
Soft drinks (≤ 2 glasses/day (400 ml); > 2 glasses/day)	
How many days a week do you usually drink sugar-sweetened (not "light" or "diet") beverages?	8-point scale: from 0 = never to 7 = every day
If you drink sugar-sweetened beverages, how many glasses, cans, and/or bottles do you drink on average per day?	8-point scale: from 0 = none to 7 glasses 8-point scale: from 0 = none to 7 cans 8-point scale: from 0 = none to 7 bottles
Breakfast consumption (every day; 0-6 days/week)	
How often do you eat breakfast on school days?	5-point scale: from 0 = I never eat breakfast on school days to 5 = 5 days
How often do you eat breakfast on weekend days?	3-point scale: 0 = I never eat breakfast on weekend days 1 = I eat breakfast on one weekend day (Saturday or Sunday) 2 = I eat breakfast on both weekend days (Saturday and Sunday)
Bicycling during leisure time (bicycling; not bicycling)	
How many days a week do you bicycle during leisure time? (include things like bicycling to the supermarket, sports club and movie theatre)	8-point scale: from 0 = never to 7 = 7 days per week
On a day that you bicycle, how long do you bicycle on average?	Open question (hours and minutes could be reported).
Walking during leisure time (≥ 60 min/day; < 60 min/day)	
How many days a week do you walk during leisure time? (include things like walking to the supermarket and sports club and walking the dog)	8-point scale: from 0 = never to 7 = 7 days per week
On a day that you walk, how long do you walk on average?	Open question (hours and minutes could be reported).
Active commuting to school (active commuting; non-active commuting)	
How many days a week do you walk from home to school?	6-point scale: from 0 = never to 5 = 5 days per week
How long does it take you to walk from home to school (one way only)?	Open question (minutes could be reported).
How many days a week do you bicycle from home to school?	6-point scale: from 0 = never to 5 = 5 days per week
How long does it take you to bicycle from home to school (one way only)?	Open question (minutes could be reported).
Sports during leisure time (playing sports; not playing any sports)	
Which sports did you play last week (at a sports club or with friends)?	Open question (three sports could be reported).
How many days did you play this sport last week?	7-point scale: from 1 = 1 day per week to 7 = every day. This scale could be filled in for the three sports listed in the preceding question.
On a day that you participate in this kind of sport, how long do you do this on average?	This open question could be filled in for the three sports listed in the first question (hours and minutes could be reported).
Watching television (≤ 120 min/day; > 120 min/day)	
How many days a week do you watch television?	8-point scale: from 0 = never to 7 = 7 days per week
On a day that you watch television, how long do you watch television on average?	Open question (hours and minutes could be reported).
Computer use (≤ 90 min/day; > 90 min/day)	
How many days per week do you use the computer (include things like the internet, games, X-box and PlayStation)?	8-point scale: from 0 = never to 7 = 7 days per week
On a day that you use the computer, how long do you use the computer on average?	Open question (hours and minutes could be reported).

1 type interaction effects and subsequent stratified analyses showed that differences according
2 to school type were not found among adolescents from non-Western ethnic backgrounds.
3 Cultural differences or level of acculturation [27, 28] might account for individual differ-
4 ences in unhealthy behaviors in non-Western ethnic groups. These findings indicate that
5 especially adolescents attending vocational schools and all adolescents from non-Western
6 ethnic backgrounds have to be targeted with interventions, since they are most likely to
7 engage in risk behaviors. Developing and implementing school based healthful diet and
8 physical activity promotion interventions that were specifically designed for lower voca-
9 tional schools is a promising strategy to prevent overweight and obesity [29, 30].

10
11 Examining gender by “ethnicity interaction effect” and subsequent stratified analyses
12 showed that girls from non-Western ethnic backgrounds were more likely to be overweight
13 or obese compared to boys from non-Western ethnic backgrounds and that they were more
14 likely to engage in risk behaviors. These gender differences were not found for adolescents
15 from Western ethnic backgrounds. This indicates that girls from non-Western ethnic back-
16 grounds in particular are an important target group. The same pattern of higher overweight
17 prevalence among non-Western female groups has also been found in the United States.
18 Ethnicity-overweight differences were greater among females, showing a higher overweight
19 prevalence among African-American girls compared to boys [9, 13]. We also found that
20 girls from non-Western ethnic backgrounds had a significantly higher risk of not bicycling,
21 not participating in sports and watching television for more than two hours compared to
22 boys from non-Western ethnic backgrounds. These similarities in patterns are interesting,
23 because the non-Western ethnic groups in the US are different from those in the Neth-
24 erlands. Whereas in the US ethnic minority groups are African American and Hispanic,
25 in the Netherlands the most important ethnic minority groups are Turkish, Moroccan,
26 Surinamese and Cape Verdean. Our findings are consistent with other studies that found
27 especially non-Western migrant women (Turkish and Moroccan) to be less physically active
28 [28, 31]. Hosper et al. (2007) also found that the prevalence’s for physical inactivity and
29 overweight of second-generation Turkish and Moroccan women seem to converge towards
30 the prevalence rates in the Dutch population [28]. Because it is unclear whether this accul-
31 turation process will occur for all risk behaviors and for all non-Western ethnic groups[27],
32 preventive interventions should still target these high-risk groups.

33
34 The following limitations should be taken into account when interpreting the results of
35 this study. The cross-sectional design of the study did not allow us to determine causal
36 effects. Longitudinal data are needed to see whether changes in weight status are associated
37 with changes in the differences in EBRB between boys and girls, Western and non-Western
38 adolescent and higher-level and lower-level education students. No test-retest data exists
39 for the physical activity measures and validation data is lacking for both dietary intake and

1 physical activity measures. The use of self-reported measures of EBRB could have caused
2 an overestimation of intakes and activities. The categorization of the data could also have
3 influenced the outcomes. In this study we made a distinction between adolescents from
4 Western and non-Western ethnic backgrounds. However, the group of adolescents from
5 non-Western ethnic backgrounds was quite diverse, including adolescents with parents born
6 in Turkey, Morocco, Cape Verde and Surinam. We were not able to examine the differences
7 between these groups.

8
9 Examining gender, ethnic and educational differences in overweight and risk behaviors for
10 overweight is important for target group segmentation and intervention development. In
11 this study, we observed the most differences in overweight and EBRB for ethnicity and
12 school type. Therefore, adolescents from non-Western ethnic backgrounds (girls in particu-
13 lar) and adolescents attending vocational schools (particularly those with Western ethnic
14 backgrounds) are important target groups for preventive interventions aimed at preventing
15 overweight and obesity. Interventions should focus on behaviors that are of specific impor-
16 tance to these high-risk groups. The effectiveness of interventions will probably increase
17 when they also take differences in individual cognitions, cultural influences and environ-
18 mental determinants of EBRB into account.

19 20 Conclusion

21
22 The results of this study showed that adolescents from non-Western ethnic backgrounds
23 and those attending vocational schools are important target groups for obesity preven-
24 tion. Given the relatively few published obesity-prevention and treatment studies that are
25 designed for specific educational or ethnic groups, it is important to promote the develop-
26 ment of culturally appropriate intervention strategies that are shown to be effective among
27 youth of diverse backgrounds. Future research should consider subgroups in the adolescent
28 population and focus on what kind of interventions work and for which target groups they
29 work taking demographic variables such as gender, ethnicity and school type into account.
30 Further qualitative and longitudinal research is needed to examine determinants of the
31 EBRB to better tailor interventions to the needs and perceptions of these specific target
32 groups.

33 34 35 ACKNOWLEDGEMENTS

36
37 Financial support for this study was provided by a grant from ZonMw, the Netherlands
38 Organization for Health Research and Development (grant ID no. 2100.0103). This study
39 was part of CEPHIR: the Center for Effective Public Health in the larger Rotterdam Area.

1 REFERENCES

- 2 1. Serdula MK, Ivery D, Coates RJ, Freedman DS, Williamson DF, Byers T: Do obese children become
3 obese adults? A review of the literature. *Prev Med* 1993, 22(2):167-177.
- 4 2. Must A, Jacques PF, Dallal GE, Bajema CJ, Dietz WH: Long-term morbidity and mortality of over-
5 weight adolescents. A follow-up of the Harvard Growth Study of 1922 to 1935. *N Engl J Med* 1992,
6 327(19):1350-1355.
- 7 3. Kreuter MW, Lukwago SN, Bucholtz RD, Clark EM, Sanders-Thompson V: Achieving cultural ap-
8 propriateness in health promotion programs: targeted and tailored approaches. *Health Educ Behav*
9 2003, 30(2):133-146.
- 10 4. Swinburn BA, Caterson I, Seidell JC, James WP: Diet, nutrition and the prevention of excess weight
11 gain and obesity. *Public health nutrition* 2004, 7(1A):123-146.
- 12 5. Gorely T, Marshall SJ, Biddle SJ: Couch kids: correlates of television viewing among youth. *Int J Behav*
13 *Med* 2004, 11(3):152-163.
- 14 6. Rennie KL, Johnson L, Jebb SA: Behavioural determinants of obesity. *Best Pract Res Clin Endocrinol*
15 *Metab* 2005, 19(3):343-358.
- 16 7. Fredriks AM, Van Buuren S, Sing RA, Wit JM, Verloove-Vanhorick SP: Alarming prevalences of
17 overweight and obesity for children of Turkish, Moroccan and Dutch origin in The Netherlands
18 according to international standards. *Acta Paediatr* 2005, 94(4):496-498.
- 19 8. Lobstein T, Frelut ML: Prevalence of overweight among children in Europe. *Obes Rev* 2003, 4(4):195-
20 200.
- 21 9. Gordon-Larsen P, Adair LS, Popkin BM: The relationship of ethnicity, socioeconomic factors, and
22 overweight in US adolescents. *Obes Res* 2003, 11(1):121-129.
- 23 10. Maffeis C: Aetiology of overweight and obesity in children and adolescents. *Eur J Pediatr* 2000, 159
24 Suppl 1:S35-44.
- 25 11. Olden K, White SL: Health-related disparities: influence of environmental factors. *Med Clin North*
26 *Am* 2005, 89(4):721-738.
- 27 12. Gordon-Larsen P, McMurray RG, Popkin BM: Adolescent physical activity and inactivity vary by
28 ethnicity: The National Longitudinal Study of Adolescent Health. *J Pediatr* 1999, 135(3):301-306.
- 29 13. Delva J, O'Malley PM, Johnston LD: Racial/ethnic and socioeconomic status differences in over-
30 weight and health-related behaviors among American students: national trends 1986-2003. *J Adolesc*
31 *Health* 2006, 39(4):536-545.
- 32 14. Sallis JF, Prochaska JJ, Taylor WC: A review of correlates of physical activity of children and adoles-
33 cents. *Med Sci Sports Exerc* 2000, 32(5):963-975.
- 34 15. Kuepper-Nybelen J, Lamerz A, Bruning N, Hebebrand J, Herpertz-Dahlmann B, Brenner H: Major
35 differences in prevalence of overweight according to nationality in preschool children living in
36 Germany: determinants and public health implications. *Arch Dis Child* 2005, 90(4):359-363.
- 37 16. te Velde SJ, Wind M, van Lenthe FJ, Klepp KI, Brug J: Differences in fruit and vegetable intake and
38 determinants of intakes between children of Dutch origin and non-Western ethnic minority children
39 in the Netherlands - a cross sectional study. *Int J Behav Nutr Phys Act* 2006, 3:31.
17. McMurray RG, Harrell JS, Deng S, Bradley CB, Cox LM, Bangdiwala SI: The influence of physical
activity, socioeconomic status, and ethnicity on the weight status of adolescents. *Obes Res* 2000,
8(2):130-139.
18. Westerstahl M, Barnekow-Bergkvist M, Jansson E: Low physical activity among adolescents in practi-
cal education. *Scand J Med Sci Sports* 2005, 15(5):287-297.

19. Wang Y, Liang H, Tussing L, Braunschweig C, Caballero B, Flay B: Obesity and related risk factors among low socio-economic status minority students in Chicago. *Public health nutrition* 2007, 10(9):927-938.
20. Shaw NJ, Crabtree NJ, Kibirige MS, Fordham JN: Ethnic and gender differences in body fat in British schoolchildren as measured by DXA. *Arch Dis Child* 2007, 92(10):872-875.
21. van der Horst K, Chin a Paw MJM, Twisk JWR, Van Mechelen W: A brief review on correlates of physical activity and sedentary behavior. *Med Sci Sports Exerc* 2007, 39(8):1241-1250.
22. van der Horst K, Oenema A, van de Looij-Jansen P, Brug J: The ENDORSE study: research into environmental determinants of obesity related behaviors in Rotterdam schoolchildren. *BMC public health* 2008, 8:142.
23. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH: Establishing a standard definition for child overweight and obesity worldwide: international survey. *Bmj* 2000, 320(7244):1240-1243.
24. Wendel-Vos GC, Schuit AJ, Saris WH, Kromhout D: Reproducibility and relative validity of the short questionnaire to assess health-enhancing physical activity. *J Clin Epidemiol* 2003, 56(12):1163-1169.
25. Statistics Netherlands [<http://statline.cbs.nl>]
26. Wang Y, Beydoun MA: The obesity epidemic in the United States--gender, age, socioeconomic, racial/ethnic, and geographic characteristics: a systematic review and meta-regression analysis. *Epidemiologic reviews* 2007, 29:6-28.
27. Hosper K, Klazinga NS, Stronks K: Acculturation does not necessarily lead to increased physical activity during leisure time: a cross-sectional study among Turkish young people in the Netherlands. *BMC public health* 2007, 7:230.
28. Hosper K, Nierkens V, Nicolaou M, Stronks K: Behavioural risk factors in two generations of non-Western migrants: do trends converge towards the host population? *Eur J Epidemiol* 2007, 22(3):163-172.
29. Martens MK, Van Assema P, Paulussen TG, Van Breukelen G, Brug J: Krachtvoer: effect evaluation of a Dutch healthful diet promotion curriculum for lower vocational schools. *Public health nutrition* 2008, 11(3):271-278.
30. Singh AS, Chin APMJ, Brug J, van Mechelen W: Short-term effects of school-based weight gain prevention among adolescents. *Archives of pediatrics & adolescent medicine* 2007, 161(6):565-571.
31. Crespo CJ, Smit E, Andersen RE, Carter-Pokras O, Ainsworth BE: Race/ethnicity, social class and their relation to physical inactivity during leisure time: results from the Third National Health and Nutrition Examination Survey, 1988-1994. *Am J Prev Med* 2000, 18(1):46-53.

6 Socio-demographic factors as correlates of active commuting to school in Rotterdam, the Netherlands

Bere E, van der Horst K, Oenema A, Prins R, Brug J. Socio-demographic factors as correlates of active commuting to school in Rotterdam, the Netherlands.

Preventive Medicine 2008, 47: 412-416.

1 **ABSTRACT**

2

3 **Objective:** Report frequencies of adolescents' active commuting to school in an inner city
4 environment in the Netherlands, and to explore potential socio-demographic correlates of
5 active commuting to school.

6 **Methods:** Cross-sectional data were obtained from the ENDORSE-study (2005–2006)
7 including 1361 adolescents (response=82%), aged 12-15 from 16 schools in Rotterdam. Socio-
8 demographic variables were assessed by questionnaire, height and weight were measured
9 and distance to school was calculated based on route planner information. Multilevel
10 logistic regressions were performed to analyze the data.

11 **Results:** The proportions of participants categorized as walkers, cyclists, non-active
12 commuters were 12%, 35% and 34% respectively. With cyclists as the reference category,
13 adolescents of non-Western ethnic background were more likely to be walkers (OR=4.1;
14 95%CI=2.1-8.2) and non-active commuters (OR=5.1; 95%CI=3.3-7.9), compared to native
15 Dutch adolescents. A further distance from home to school was inversely associated with
16 being a walker (OR=0.22; 95%CI=0.17-0.29) and being a cyclist (OR=0.83; 95%CI=0.79-0.86)
17 and positively associated with being a non-active commuter (OR=1.2; 95%CI=1.16-1.23).

18 **Conclusion:** Almost 50% of the adolescents reported to actively commute to school, and
19 mode of commuting was associated with ethnicity and distance. Further research is needed
20 to examine main barriers to active commuting among adolescents from non-Western ethnic
21 background.

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

1 INTRODUCTION

2
3 The prevalence of overweight and obesity among adolescents is increasing worldwide, as is
4 also the case in the Netherlands [1]. Evidence indicates that engaging in at least 60 minutes
5 of moderate intensity physical activity on preferably all days of the week, contributes to pre-
6 vention of overweight and obesity and to better health [2, 3]. Active commuting to school is
7 one of the daily activities that could be an important component of the daily-recommended
8 level of physical activity for adolescents. A meta-analytic review stated recently that active
9 commuting (among adults) was associated with an 11% reduction in cardiovascular risk [4].

10 There is a lack of research on trends in mode of transport to school over the past years.
11 In the US and Australia low and decreasing frequencies of active commuting to school have
12 been reported [5-8]. Different reasons have been suggested for these low and decreasing
13 levels of active commuting such as safety concerns, traffic, road-crossing, crime, conve-
14 nience to drop children off on way to work and environmental factors such as walkability
15 and distance to school [5, 9-14].

16 In the Netherlands between 1985 and 2005 no clear decreases in the total number of cy-
17 cling and walking trips and distances have been seen in the Dutch population [15]. Between
18 2004 and 2006, 12-16 year old adolescents reported to cycle approximately 6 km/per day and
19 to walk approximately 0.33 km/day [16]. The built environment in cities in the Netherlands
20 appears to be good for cycling compared to cities in other countries. The Netherlands have
21 a long tradition of cycling, which has resulted in a cycling-friendly infrastructure mak-
22 ing it more convenient and safer to cycle than in other countries. However, no study has
23 reported frequencies of active commuting to school in the Netherlands and few studies
24 in general have reported socio-demographic determinants of active commuting to school.
25 In the Netherlands, a large number of adolescents from non-Dutch ethnic backgrounds
26 live in the larger cities and we expect that differences between cultures exist for the mode
27 of commuting to school. Better insight in socio-demographic factors associated with ac-
28 tive commuting to school will enable tailoring interventions aimed at the prevention of
29 overweight to the needs of specific risk groups. Therefore, the aim of the present study was
30 to report frequencies of adolescents' active commuting to school in Rotterdam, the second-
31 largest city in the Netherlands, and to explore potential socio-demographic correlates of
32 active commuting to school.

33 34 35 METHODS

36
37 The present study is part of the ENDORSE (Environmental Determinants of Obesity among
38 Rotterdam SchoolchildrEn) project on identification of important individual and environ-
39 mental determinants of adolescent behaviors related to overweight and obesity. The EN-

1 DORSE study is an integral part of the ongoing health surveillance system of the Municipal
2 Health Service in the Rotterdam area (Youth Monitor Rotterdam, YMR). The Medical Ethics
3 Committee of the Erasmus University Medical Center approved the ENDORSE project.

4 5 Procedure and sample 6

7 The YMR and ENDORSE studies collected data in school year 2005/2006 among adoles-
8 cents in the first (12-13 year olds) and third year of secondary school (14-15 year olds). A
9 total of 24 schools from the 56 schools participating in the YMR were willing to participate
10 also in the ENDORSE study. After stratification according to location in the city, seventeen
11 school locations were randomly selected. On average five classes per school location were
12 selected at random to participate and a total of 1668 adolescents were eligible to participate
13 in the ENDORSE study. Between October 2005 and May 2006, the adolescents completed
14 the YMR and the ENDORSE questionnaires. Within a month after completion of the EN-
15 DORSE questionnaire, two trained research assistants measured height and weight.

16 During administration of the ENDORSE survey 187 adolescents were absent. Due
17 to printing mistakes, it was necessary to delete 120 adolescents, including one complete
18 school from the sample. Therefore, the study sample includes 1361 participants (82%), from
19 71 classes and 16 schools; 55% boys, 6.5% non-Dutch Western ethnic background and 53%
20 non-Western ethnic background. Mean age was 14.1 years (SD=1.2; range 11.0-17.6).

21 22 Measures 23

24 Commuting to school was measured by three questions: How many days a week do you
25 travel to school; (1) walking, (2) cycling, (3) by public transport or car. Response categories
26 were never, one day/week, two days/week, three days/week, four days/week, and five days/
27 week. The three items were combined to one variable with four categories: (1) walking 3
28 days/week or more (WALKERS), (2) cycling three days/week or more (CYCLISTS), (3) non-
29 active commuting three days/week or more (NON-ACTIVE COMMUTERS), (4) pupils
30 where the sum of the three answers counted up to less than or more than five days/week
31 (PUPILS NOT CATEGORIZED INTO MODE OF COMMUTING).

32 Sex, school level (vocational or university preparatory high school) and date of birth were
33 provided by the schools. Age was determined on the date of the anthropometrical measure-
34 ments. Employment of parents was assessed in the YMR questionnaire by two questions
35 asking whether their mother and father had paid work or not (0=mother or/and father
36 have NOT paid work, 1= both mother and father have paid work). Ethnicity was assessed in
37 the YMR questionnaire by two questions asking in which country their mother and father
38 had been born. Ethnicity was defined upon the definition used by Statistics Netherlands
39 [17]. The pupils were considered to be native Dutch if both parents had been born in the

1 Netherlands, the pupils were considered to be from Western ethnic background if one
 2 or both parents had been born in another European country, North America, Oceania,
 3 Indonesia or Japan. Adolescents with one or both parents born in a non-Western country
 4 were considered as from non-Western ethnic background. Body Mass Index was calculated
 5 from the measurements of height and weight, carried out by research assistants. Age and
 6 sex specific cut off points were used to categorize adolescents in categories of normal weight
 7 and overweight or obese [18]. Distance from home to school was calculated from pupil's
 8 reports of home address postal codes. The home address postal codes and the exact school
 9 addresses were entered into the route planner www.routenet.nl (in March 2007). The length
 10 of the optimal route for cars was derived from this service and entered into the data set
 11 for each pupil. Distances over 45 km were regarded as outliers, and therefore, pupils living
 12 further than 45km from school were not included in the analyses (6 pupils).

14 Statistics

16 Descriptive analyses of commuting to school in relation to the potential determinants
 17 were conducted using SPSS version 14. Multilevel logistic regression analyses, taking the
 18 clustering of pupils within schools into account, were performed with walking, cycling or
 19 non-active commuting to school as dependent variables, using MLwiN version 2.02. Walk-
 20 ing, cycling and non-active commuters were first compared to the rest of the sample (e.g.
 21 walkers, were compared to non-walkers (i.e. cyclists, non-active commuters and pupils not
 22 categorized into mode of commuting)), and then walkers and non-active commuters were
 23 compared to cyclists. All regression models included sex, work status of parents, ethnicity,
 24 weight status, age, distance from home to school and school level (high school or voca-
 25 tional). Three dummy variables were created and included in the analyses in order to keep
 26 adolescents with missing values on one or more of the following variables in the models: (1)
 27 Work status: quite a few adolescents (n=244) reported not to know whether their parents
 28 had paid work or not, or reported to have no parent or no contact with their mother and/
 29 or father. (2) Weight status: due to absence, anthropometrical measurements were lacking
 30 for 143 adolescents. (3) Ethnicity and work status: due to absence, 133 adolescents did not
 31 participate in the YMR survey. Odds ratios (OR) with confidence interval (95%) are given
 32 for each independent variable.

35 RESULTS

37 Table 6.1 shows descriptive characteristics of the sample and how the different potential
 38 correlates were bivariately related to commuting to school. The proportions of participants
 39 categorized as walkers, cyclists, non-active commuters and 'pupils not categorized into mode

Table 6.1 Description of the proposed determinants and the unadjusted relationship between these factors and commuting to school among adolescents in Rotterdam, the Netherlands, school year 2005/2006 (proportions of total sample, or mean, with 95% CI)

	WALKERS (n=168)		CYCLIST (n=471)		NON-ACTIVE COMMUTERS (n=465)		PUPILS NOT CATEGORISED INTO COMMUTING MODE (n=257)	
	N	Proportion (95% CI)	Proportion (95% CI)	Proportion (95% CI)	Proportion (95% CI)	Proportion (95% CI)	Proportion (95% CI)	
All	1361	12% (11, 14)	35% (32, 37)	34%	19%			
Sex								
Boys	752	10% (8, 12)	37% (33, 40)	36%	18%			
Girls	609	16% (13, 18)	32% (28, 36)	32%	21%			
Work status parents								
Both parents have work	611	9% (6, 11)	44% (40, 48)	31%	16%			
Not two working parents	338	18% (14, 22)	24% (20, 29)	37%	21%			
Ethnicity								
Native Dutch	512	4% (2, 6)	54% (49, 58)	28%	14%			
Western ethnicity	75	5% (0, 11)	40% (29, 51)	36%	19%			
Non-Western ethnicity	614	20% (17, 23)	18% (15, 21)	39%	23%			
Weight status								
Normal weight	919	11% (9, 13)	39% (36, 42)	32%	17%			
Overweight or obese	299	16% (12, 20)	26% (21, 31)	37%	21%			
Age (years, mean)	1361	14.5 (14.3, 14.7)	14.0 (13.9, 14.1)	14.1	14.1			

	WALKERS (n=168)	CYCLIST (n=471)	NON-ACTIVE COMMUTERS (n=465)	PUPILS NOT CATEGORISED INTO COMMUTING MODE (n= 257)
Distance to school				
Mean, km	1.4 (1.3, 1.6)	4.8 (4.5, 5.0)	9.7 (8.9, 10.4)	6.9 (6.0, 7.8)
Type of school				
High school	11% (9, 14)	42% (38, 46)	33% (29, 37)	14% (11, 17)
Vocational school	13% (11, 15)	30% (27, 33)	35% (32, 38)	22% (19, 25)

1 of commuting' were 12%, 35%, 34% and 19% respectively. The majority within the walking,
 2 cycling and non-actively commuting categories, respectively, reported to walk (88%), cycle
 3 (92%) or non-actively commute (95%) all five school days/week (data not shown). Mean
 4 distances from home to school were 1.4 km, 4.8 km, 9.7 km, and 6.9 km respectively for the
 5 walkers, the cyclists, the non-active commuters and the pupils not categorized into mode of
 6 commuting. The proportion of walkers, cyclists and non-active commuters living less than
 7 three km away from school were 97%, 30% and 10% respectively. Of the cyclists and the non-
 8 active commuters 96% and 70% lived within ten km from school. Fewer adolescents from
 9 non-Western (47%) and Western (non-Dutch) ethnic background (67%) reported to have
 10 bikes at home than native Dutch adolescents (83%). Adolescents with two working parents
 11 (73%) and with at least one parent not working (53%) reported to have bikes at home.

12
 13 Comparing walkers, cyclists and non-active commuters respectively to the remaining
 14 sample (including also the pupils not categorized into mode of commuting) (Table 6.2);
 15 adolescents from non-Western ethnic background (OR=2.0; 95%CI=1.0-4.0) and older ado-
 16 lescents (OR=1.3; 95%CI=1.0-1.7) were more likely to be walkers, while adolescents living
 17 further away from school (OR=0.22; 95%CI=0.17-0.29) were less likely to be walkers. Those
 18 having a parent without paid work (OR=0.6; 95%CI=0.4-0.9), from Western (OR=0.5;
 19 95%CI=0.3-0.8) and non-Western ethnic background (OR=0.3; 95%CI=0.2-0.4), as well
 20 as those living further away from school (OR=0.83; 95%CI=0.79-0.86) were less likely to
 21 be cyclists. Adolescents from Western (OR=2.3; 95%CI=1.3-4.2) and non-Western ethnic
 22 background (OR=3.0; 95%CI=2.1-4.4), as well as those living further away from school
 23 (OR=1.20; 95%CI=1.16-1.23) were more likely to be non-active commuters.

24
 25 Comparing walkers to cyclists (Table 6.3); adolescents from non-Western ethnic background
 26 (OR=4.1; 95%CI=2.1-8.2) were more likely to be a walker than a cyclist, while adolescents
 27 living further away from school (OR = 0.3; 95%CI=0.3-0.4) were less likely to be a walker
 28 than a cyclist. Comparing non-active commuters to cyclists; those having at least one parent
 29 without paid work (OR=1.7; 95%CI=1.1-2.6), being from Western (OR=2.6; 95%CI=1.3-5.2)
 30 and non-Western ethnic background (OR=5.1; 95%CI=3.3-7.9) and those living further away
 31 from school (OR=1.4; 95%CI=1.3-1.4) were all more likely to be non-active commuters than
 32 cyclists.

33 34 35 DISCUSSION

36
 37 Almost half (47%) of the sample do actively commute to school most school days. In the
 38 present study, differences in mode of commuting to school were found between adolescents
 39 from Dutch, non-Western and Western ethnic backgrounds. Cycling was the dominant

Table 6.2 Odds ratios for being a walker, cyclist or non-active commuter among adolescents in Rotterdam, the Netherlands, school year 2005/2006 (reference is “all other adolescents”)

	WALKER		CYCLIST		NON-ACTIVE COMMUTER	
	OR	(95%CI)	OR	(95%CI)	OR	(95%CI)
Girls vs. boys	0.9	(0.5, 1.4)	0.8	(0.6, 1.0)	1.0	(0.7, 1.3)
Not two working parents vs. both working	0.8	(0.4, 1.5)	0.6	(0.4, 0.9)	1.2	(0.9, 1.7)
Western ethnicity vs. native Dutch	0.7	(0.2, 2.7)	0.5	(0.3, 0.8)	2.3	(1.3, 4.2)
Non-western ethnicity vs. native Dutch	2.0	(1.0, 4.0)	0.3	(0.2, 0.4)	3.0	(2.1, 4.4)
Overweight vs. non-overweight	1.1	(0.6, 1.9)	0.8	(0.6, 1.1)	1.1	(0.8, 1.6)
Age (years)	1.3	(1.0, 1.7)	0.9	(0.8, 1.0)	1.1	(0.9, 1.2)
Distance (km)	0.22	(0.17, 0.29)	0.83	(0.79, 0.86)	1.20	(1.16, 1.23)
Vocational vs. high school	1.2	(0.6, 2.7)	0.6	(0.3, 1.2)	0.9	(0.5, 1.7)

Table 6.3 Odds ratios for being a non-active commuter (compared to cyclists) and for being a walker (compared to cyclist) among adolescents in Rotterdam, the Netherlands, school year 2005/2006

	WALKER vs. CYCLIST		NON-ACTIVE COMMUTER vs. CYCLIST	
	OR	(95% CI)	OR	(95% CI)
Girls vs. boys	0.9	(0.6, 1.6)	1.2	(0.9, 1.7)
Not two working parents vs. both working	1.1	(0.5, 2.1)	1.7	(1.1, 2.6)
Western ethnicity vs. native Dutch	1.0	(0.3, 4.1)	2.6	(1.3, 5.2)
Non-western ethnicity vs. native Dutch	4.1	(2.1, 8.2)	5.1	(3.3, 7.9)
Overweight vs. non-overweight	1.2	(0.6, 2.2)	1.3	(0.9, 2.0)
Age (years)	1.3	(1.0, 1.7)	1.1	(0.9, 1.3)
Distance (km)	0.32	(0.25, 0.43)	1.36	(1.28, 1.44)
Vocational vs. high school	1.7	(0.7, 4.4)	1.3	(0.5, 3.4)

mode of transport among the native Dutch adolescents, and 54% travel to school by bike at least three days per week. Non-active commuting was the dominant mode of transport among adolescents from non-Western ethnicity (39%), followed by walking (20%). The findings are in line with a study from de Bruijn et al. [19], which found native Dutch adolescents (mean age 14.8 years) to be nearly three times as likely to use a bicycle for general transportation as adolescents from other ethnic backgrounds. This difference may be explained by a difference in culture. In the Netherlands the bicycle is traditionally an important mode of transport and most Dutch families do have bikes [19]. In the present

1 study, lower proportions of immigrants (47% of non-Western and 67% of Western ethnicity)
2 reported to have bikes at home than native Dutch (83%).

3
4 Cycling seems to be a more prominent transportation mode among adolescents of higher
5 socio-economic position in Rotterdam. Adolescents with at least one parent without a paid
6 job were less likely to be cyclists, and more likely to be non-active commuters. There was
7 also a disparity in the parental work status measure on having bikes at home; 73% vs. 53%
8 respectively among the “two working parents” and the “at least one parent not working”
9 groups. Higher SES groups have previously been reported to cycle more often to school in
10 Australia; living in a high SES area increased the odds for walking and cycling to school [14].
11 However, in the USA and Portugal opposite results have been reported; adolescents from
12 lower socio-economic positions were more likely to walk or cycle to school [5, 20].

13 Bivariately an association between weight status and cycling to school was observed
14 (Table 1). However, no significant associations between mode of commuting to school and
15 weight status were seen in the multivariate analyses. Similarly, other studies did not find
16 clear associations between active commuting to school and overweight or BMI [21-23].
17 However, it has been reported that increased walking or cycling distance was significantly
18 associated with lower fat mass [22]. Although some studies failed to find an association with
19 overweight, active commuting to school is still an opportunity to increase physical activity
20 levels, and therefore contribute to a healthy lifestyle [24].

21
22 Few evidence-based indications have been reported in the international literature about
23 how far we can expect adolescents to walk and/or cycle to school. Studies have reported that
24 most adolescent ‘walkers’ live within a distance of 2.5 km from school [5, 12]. Colabianchi
25 et al. [25] found that girls think of an ‘easy’ walking distance as 15 minutes (approximately
26 0.75 miles/1.2 km). In Australia, parents reported 1.6 km as an appropriate walking distance
27 for 10-12 year old children [26], which is rather similar to the median walking distance
28 for the walkers in the present study (1.3km). However, walking to and from school, a total
29 of 1-2 km probably does not lead to significant increases in energy expenditure and will
30 probably not have an impact on weights status. The current study indicates that adolescents
31 can commute rather long distances in the Netherlands, at least up to 3km of walking and
32 10km of cycling (one way), and actively commuting such distances might make a difference
33 for obesity prevention. Further research is needed to examine individual and environmental
34 barriers to active commuting especially for adolescents from non-Western ethnicities who
35 live within cycling and walking distance to school to tailor interventions better to the needs
36 and perceptions of this target group. The cultural differences in transportation to school
37 might also indicate that other strategies for increasing physical activity are needed for
38 adolescents from non-Western ethnic background.

1 Study limitations

2
3 There are some limitations of the present study. Only one Dutch city was included in the
4 study and since cycling in the Netherlands is so typical for the native-Dutch population,
5 generalization of the findings to other countries might be difficult. No test/retest or valida-
6 tion data exist for the commuting to school measure. The measure can neither differentiate
7 public transportation commuters from car commuters, which would be an interesting
8 comparison since public transportation commuters do more physical activity than car
9 commuters [27]. Using public transportation to work has also been negatively associated
10 with overweight and obesity among both Swedish [28] and Australian [29] men. The SES
11 measures used in this paper are rather simple and including more proper measures for
12 SES (family educational level, type of work, and income) might explain more of the large
13 ethnicity disparities observed. Ethnicity is clearly not as homogenous as the three classified
14 groups. Stratifying the results on more specific ethnicities would be interesting, and this
15 clearly is an issue for future research. Distance to school was calculated as the optimal route
16 for cars, and not necessarily reflecting the true walking or cycling distance. The strength of
17 the study is that the analyses were adjusted for an objective measure of distance to school,
18 and that it reports walking and cycling rates from a cycling country, which might serve as a
19 good example for other countries.

21 Conclusion

22
23 Almost half of the adolescents living in an inner city environment in the Netherlands
24 actively commuted to school on most school days, and mode of commuting was strongly as-
25 sociated with ethnicity. Adolescents from non-Dutch ethnicities and from lower SES groups
26 are important target groups for the promotion of active commuting to school. However,
27 further research is needed to examine determinants of active and inactive commuting to
28 school to better tailor interventions to the needs and perceptions of these target groups.

31 ACKNOWLEDGMENTS

32
33 This study was financially supported by grants from ZonMw, The Netherlands Organization
34 for Health Research and Development (grant ID no 2100.0103). Elling Bere had a post doc
35 grant from the Norwegian Research Council, and spent one year (2006/07) at Department
36 of Public Health, Erasmus MC Rotterdam. This study was part of CEPHIR: the Center for
37 Evidence-based Public Health In the Rotterdam area.

1 REFERENCES

- 2 1. Schokker DF, Visscher TL, Nooyens AC, van Baak MA, Seidell JC: Prevalence of overweight and
3 obesity in the Netherlands. *Obes Rev* 2007, 8(2):101-108.
- 4 2. Diet, nutrition and the prevention of chronic diseases. *World Health Organ Tech Rep Ser* 2003, 916:i-
5 viii, 1-149, backcover.
- 6 3. Strong WB, Malina RM, Blimkie CJ, Daniels SR, Dishman RK, Gutin B, Hergenroeder AC, Must A,
7 Nixon PA, Pivarnik JM *et al*: Evidence based physical activity for school-age youth. *J Pediatr* 2005,
8 146(6):732-737.
- 9 4. Hamer M, Chida Y: Active commuting and cardiovascular risk: a meta-analytic review. *Prev Med*
10 2008, 46(1):9-13.
- 11 5. McDonald NC: Active transportation to school: trends among U.S. schoolchildren, 1969-2001. *Am J*
12 *Prev Med* 2007, 32(6):509-516.
- 13 6. Salmon J, Timperio A: Prevalence, trends and environmental influences on child and youth physical
14 activity. *Med Sport Sci* 2007, 50:183-199.
- 15 7. Salmon J, Timperio A, Cleland V, Venn A: Trends in children's physical activity and weight status in
16 high and low socio-economic status areas of Melbourne, Victoria, 1985-2001. *Aust NZ J Public Health*
17 2005, 29(4):337-342.
- 18 8. van der Ploeg HP, Merom D, Corpuz G, Bauman AE: Trends in Australian children traveling to
19 school 1971-2003: burning petrol or carbohydrates? *Prev Med* 2008, 46(1):60-62.
- 20 9. Bringolf-Isler B, Grize L, Mader U, Ruch N, Sennhauser FH, Braun-Fahrlander C: Personal and
21 environmental factors associated with active commuting to school in Switzerland. *Prev Med* 2008,
22 46(1):67-73.
- 23 10. Carver A, Timperio A, Crawford D: Playing it safe: the influence of neighbourhood safety on chil-
24 dren's physical activity. A review. *Health Place* 2008, 14(2):217-227.
- 25 11. Kerr J, Rosenberg D, Sallis JF, Saelens BE, Frank LD, Conway TL: Active commuting to school: As-
26 sociations with environment and parental concerns. *Med Sci Sports Exerc* 2006, 38(4):787-794.
- 27 12. Nelson NM, Foley E, O'Gorman DJ, Moyna NM, Woods CB: Active commuting to school: How far is
28 too far? *Int J Behav Nutr Phys Act* 2008, 5:1.
- 29 13. Sjolie AN, Thuen F: School journeys and leisure activities in rural and urban adolescents in Norway.
30 *Health Promot Int* 2002, 17(1):21-30.
- 31 14. Timperio A, Ball K, Salmon J, Roberts R, Giles-Corti B, Simmons D, Baur LA, Crawford D: Personal,
32 family, social, and environmental correlates of active commuting to school. *Am J Prev Med* 2006,
33 30(1):45-51.
- 34 15. [<http://www.statline.cbs.nl>]
- 35 16. The Ministry of Transport PWaWM, Public Works and Water Management (Rijkswaterstaat): Mobil-
36 ity Survey Netherlands. Published online at <http://www.mobiliteitsonderzoeknederland.nl>.
- 37 17. Netherlands S: Hoe doet het CBS dat nou? Standaard definitie allochtonen. Voorburg; 2000.
- 38 18. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH: Establishing a standard definition for child overweight
39 and obesity worldwide: international survey. *BMJ* 2000, 320(7244):1240-1243.
19. de Bruijn GJ, Kremers SP, Schaalma H, van Mechelen W, Brug J: Determinants of adolescent bicycle
use for transportation and snacking behavior. *Prev Med* 2005, 40(6):658-667.
20. Mota J, Gomes H, Almeida M, Ribeiro JC, Carvalho J, Santos MP: Active versus passive transporta-
tion to school-differences in screen time, socio-economic position and perceived environmental
characteristics in adolescent girls. *Ann Hum Biol* 2007, 34(3):273-282.

- 1 21. Heelan KA, Donnelly JE, Jacobsen DJ, Mayo MS, Washburn R, Greene L: Active commuting to and
2 from school and BMI in elementary school children-preliminary data. *Child Care Health Dev* 2005,
3 31(3):341-349.
- 4 22. Landsberg B, Plachta-Danielzik S, Much D, Johannsen M, Lange D, Muller MJ: Associations between
5 active commuting to school, fat mass and lifestyle factors in adolescents: the Kiel Obesity Prevention
6 Study (KOPS). *Eur J Clin Nutr* 2008, 62(6):739-747.
- 7 23. Rosenberg DE, Sallis JF, Conway TL, Cain KL, McKenzie TL: Active transportation to school over 2
8 years in relation to weight status and physical activity. *Obesity (Silver Spring)* 2006, 14(10):1771-1776.
- 9 24. Alexander LM, Inchley J, Todd J, Currie D, Cooper AR, Currie C: The broader impact of walking to
10 school among adolescents: seven day accelerometry based study. *BMJ* 2005, 331(7524):1061-1062.
- 11 25. Colabianchi N, Dowda M, Pfeiffer KA, Porter DE, Almeida MJ, Pate RR: Towards an understanding
12 of salient neighborhood boundaries: adolescent reports of an easy walking distance and convenient
13 driving distance. *Int J Behav Nutr Phys Act* 2007, 4:66.
- 14 26. Timperio A, Crawford D, Telford A, Salmon J: Perceptions about the local neighborhood and walk-
15 ing and cycling among children. *Prev Med* 2004, 38(1):39-47.
- 16 27. Edwards RD: Public transit, obesity, and medical costs: assessing the magnitudes. *Prev Med* 2008,
17 46(1):14-21.
- 18 28. Lindstrom M: Means of transportation to work and overweight and obesity: a population-based
19 study in southern Sweden. *Prev Med* 2008, 46(1):22-28.
- 20 29. Wen LM, Rissel C: Inverse associations between cycling to work, public transport, and overweight
21 and obesity: findings from a population based study in Australia. *Prev Med* 2008, 46(1):29-32.
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39

Part IV Individual and environmental correlates of energy balance-related behaviors

7 The school food environment: associations with adolescent soft drink and snack consumption

Van der Horst K, Timperio A, Crawford D, Roberts R, Brug J, Oenema A.
The school food environment: associations with adolescent soft drink and
snack consumption.

American Journal of Preventive Medicine 2008, 35(3): 217-23.

1 **ABSTRACT**

2

3 **Background:** Because students may purchase food and drinks in and around their schools,
4 the school food environment may be important for obesity-related eating behaviors such as
5 soft drink and snack consumption. However, research exploring the associations between
6 school environments and specific eating behaviors is sparse.

7 **Methods:** Associations of the availability of canteen food and drinks, the presence of food
8 stores around schools, and individual cognitions (attitudes, norms, modeling, perceived
9 behavioral control, and intentions) with soft drink and snack consumption were examined
10 in a cross-sectional study (2005–2006) among 1293 adolescents aged 12–15 years. Soft
11 drink and snack consumption and related cognitions were assessed with self-administered
12 questionnaires. The presence of food stores and the distance to the nearest food store were
13 calculated within a 500-meter buffer around each school. Data on the availability of soft
14 drinks and snacks in school canteens were gathered by observation. In 2007, multilevel
15 regression models were run to analyze associations and mediation pathways between cogni-
16 tions, environmental factors, and behaviors.

17 **Results:** Adolescents' attitudes, subjective norms, parental and peer modeling, and inten-
18 tions were positively associated with soft drink and snack consumption. There was an
19 inverse association between the distance to the nearest store and the number of small food
20 stores with soft drink consumption. These effects were mediated partly by cognitions.

21 **Conclusions:** This study provided little evidence for associations of environmental factors
22 in the school environment with soft drink and snack consumption. Individual cognitions
23 appeared to be stronger correlates of intake than physical school-environmental factors.
24 Longitudinal research is needed to confirm these findings.

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

1 INTRODUCTION

2
3 Obesity is a major problem in many countries, and its prevalence is increasing [1, 2]. Dietary
4 patterns such as the consumption of fast food, snacks, and soft drinks may contribute to the
5 development of overweight and obesity through the foods' high energy density and large
6 portion sizes [3-9].

7 The theory of planned behavior (TPB) has proven to be useful in understanding cor-
8 relates or determinants of soft drink and snack consumption [10-13]. According to the TPB,
9 behavior can be predicted from the intention to perform behavior that is determined by
10 attitudes, subjective norm, modeling, and perceived behavioral control [14-16]. However,
11 obesogenic dietary behaviors may also be influenced by the environmental opportunities to
12 eat food [17-19]. In social-ecologic models, (e.g., the Environmental Research framework for
13 weight Gain prevention [the EnRG framework]), it is proposed that environmental factors
14 indirectly influence behavior via the individual's cognitions [20]. Environments that offer
15 appealing opportunities for unhealthy foods may result in positive cognitions regarding the
16 consumption of these unhealthy foods, resulting in higher intake of them.

17
18 Schools may be an important setting for obesity-prevention interventions, as many schools
19 provide extensive facilities for selling food and drinks [21, 22]. During breaks, adolescents
20 may also purchase food items in the immediate area around the school. Relatively few stud-
21 ies [18, 23, 24] are available that examine environmental factors in a school setting. These
22 studies found that the number of snack vending machines was associated with student snack
23 purchases and lower fruit intake, and that in schools where soft drink machines were turned
24 off during lunch time, adolescents purchased fewer soft drinks [23, 24]. Many fast-food
25 restaurants are located within walking distance of a school, and an open-campus policy
26 during lunchtime was found to be associated with a higher likelihood of students' eating
27 lunch at a fast-food restaurant [24-27].

28
29 The present study expands on the limited literature that explores the role of school food
30 environments in influencing the dietary behaviors of youth. The overall hypothesis for this
31 study was that a greater availability of soft drinks and snacks at school and in the school
32 neighborhood as well as positive cognitions would lead to a higher intake of soft drinks and
33 snacks. A second hypothesis was that environmental factors influence behavior via cogni-
34 tions, that is, that cognitive factors mediate the association between environmental factors
35 and behavior. This study specifically aimed to (1) examine the associations between school
36 food availability and food stores in the school neighborhood with soft drink and snack
37 consumption, (2) examine the associations between cognitions and soft drink and snack
38 consumption, and (3) examine whether the effect of environmental factors on soft drink and
39 snack consumption is partly mediated by cognitions (mediation effect).

1 METHODS

2

3 Study Design and Sample Selection

4

5 The Environmental Determinants of Obesity in Rotterdam SchoolchildrEn (ENDORSE)
6 study is a prospective 2-year study among adolescents (aged 12–13 years) in the first year and
7 third year (aged 14–15 years) of secondary school [28]. In 2005–2006, 24 of 56 schools in
8 Rotterdam participated in the ENDORSE study. After stratification according to city region,
9 17 schools were selected. From them, a total of 78 classes (1668 adolescents) were randomly
10 selected to participate in the ENDORSE study. Because of absence, printing mistakes, and a
11 school location outside the municipal border, it was necessary to omit 375 respondents from
12 the sample. The study sample therefore included 1293 participants (78% of those eligible)
13 from 66 classes and 15 schools. The Medical Ethics Committee of the Erasmus University
14 Medical Center reviewed the proposal and gave a declaration of no objection for the EN-
15 DORSE project.

16

17 Procedure

18

19 The ENDORSE study was announced through a letter to parents. Parents could refuse to
20 allow the participation of their child(ren). From October 2005 to May 2006, the adoles-
21 cents completed a questionnaire in a lesson of approximately 50 minutes in the presence
22 of a teacher and a trained research assistant. Observations of the school canteens were
23 performed in the same time period.

24

25 Measures

26

27 *Sugar-sweetened soft drink and snack consumption*

28 Soft drinks were defined as carbonated drinks, other noncarbonated sugar-sweetened
29 drinks (water-based beverages that contain sugar), and sport drinks. The consumption of
30 soft drinks was assessed by two questions: *How many days a week do you usually drink*
31 *sugar-sweetened (not “light” or “diet”) beverages?* and *If you drink sugar-sweetened beverages,*
32 *how many glasses, cans, and/or bottles do you drink on average per day?* Total soft drink
33 consumption was expressed in liters per day, and was calculated from the two questions
34 according to Dutch standard serving sizes (1 glass=200 ml, 1 can=330 ml, 1 bottle=500 ml).
35 The soft drink consumption variable was normally distributed, allowing linear regression
36 analysis.

37

38 Snacks were classified as *sweet* (candy, candy bars, chocolate, cake, biscuits) and *savory*
39 (fast-food, pizza, fries, chips, nuts). Sweet-snack consumption was assessed by two ques-

1 tions: *How many days a week do you usually eat sweet snacks or cookies?* and *On average, how*
2 *many times a day do you eat sweet snacks or cookies?* Two similar questions were asked for
3 savory snacks. These questions were combined so that a single score could be computed for
4 the mean snack intake in times per day. As this variable was not normally distributed, snack
5 consumption was dichotomized by means of a median split, distinguishing adolescents
6 consuming two or fewer times per day (reference group) from those consuming more than
7 two times per day.

8 *Personal factors*

9 Cognitions specific to soft drink and snack consumption (attitude, subjective [parental]
10 norm, modeling, perceived behavioral control, and intention) were assessed according to
11 the TRB, using a 5-point bipolar scale [14]. Attitude was assessed with two items that asked
12 if the adolescent considered the behavior as good or bad and as pleasant or unpleasant.
13 The two items were collapsed in a single attitude variable by calculating the mean item
14 score (Cronbach's $\alpha=0.74$ soft drink, 0.61 snack consumption). Parental norm was assessed
15 with one item: *My parents consider consuming soft drinks/snacks as good/bad*. Modeling was
16 assessed with two items that asked if parents and friends consume a lot of or very little/very
17 few soft drinks/snacks. Perceived behavior control was assessed with two items that asked
18 how easy or difficult it is to consume soft drinks/snacks, and then asking if the decision
19 to consume soft drinks/snacks is completely or not completely under the control of the
20 adolescent. The two items were collapsed into one variable by calculating the mean item
21 score (Cronbach's $\alpha=0.61$ soft drink, 0.65 snack consumption). The intention to consume
22 soft drinks, snacks, or both was assessed with a single item that asked about the adolescent's
23 intention to consume soft drinks/snacks in the next 6 months. Because of a skewed distribu-
24 tion, responses to all variables were dichotomized to indicate agreement with the statement
25 (i.e., very good or good=1) or otherwise (i.e., neither good/bad, bad, very bad=0).

26 *School food environment*

27
28 Two observers audited each school. An audit instrument was developed to assess the avail-
29 ability of food in the schools (see Appendix). The instrument was reviewed by experts and
30 pilot-tested. It included observations of the different types of soft drinks/snacks that were
31 available from vending machines and at the canteen counter. Eight availability variables
32 were created: the availability of sugar-sweetened soft drinks in vending machines (1) and
33 at the canteen counter (2); the total availability of low-calorie drinks (3); the availability of
34 energy-dense snacks in vending machines (4) and at the canteen counter (5); the availability
35 of low-energy snacks in vending machines (6) and at the canteen counter (7); and the total
36 availability of fruit and vegetables at school (8).

1 Products from one brand with different flavors (e.g. Coke and Coke with vanilla flavor) were
2 counted as two distinct products. The availability variables were re-coded into variables
3 with two or three categories. Where possible, a group with no products available was cre-
4 ated; otherwise, the variables were categorized into tertiles or dichotomized, depending on
5 the variation in the data.
6

7 *Local neighborhood environment around school*

8 The neighborhood around a school was defined as a crow-fly buffer of 500 meters (stores
9 that could be accessed within a lunch break of 30 minutes). Each school address was geo-
10 coded using ArcView Version 3.3, and 500-meter buffers were created around each school.
11 The municipality of Rotterdam supplied cadastral data as well as road and road-attribute
12 information. Records from Locatus, a company that provides information on stores in The
13 Netherlands, were used to identify the locations of the food establishments surrounding
14 each school. These locations were geocoded, and the availability of stores (the total number
15 within 500 meters) was computed for five types of food establishments: (1) fast-food outlets;
16 (2) large supermarkets; (3) small food stores (small supermarkets, ethnic-food stores, news
17 agencies, stores at petrol stations); (4) bakeries; and (5) fruit/vegetable stores. The distance to
18 the nearest food store was calculated using the street network (walking route). All variables
19 were re-coded to categories based on tertiles (small food stores, bakeries, fruit/vegetable
20 stores, distance); the median value (fast-food outlets); and the possibility of distinguishing
21 a no-availability category (large supermarkets).
22

23 Demographics

24
25 Age was derived from date of birth and date of measurement. Ethnicity was defined accord-
26 ing to the definition used by the Netherlands Statistics. Adolescents were considered to be
27 from a Western ethnic background if both parents had been born in the Netherlands; in
28 another European country; or in North America, Oceania, Indonesia, or Japan. Adolescents
29 with one or both parents born in a non-Western country were considered to be from a non-
30 Western ethnic background. Schools provided the school-level information (higher-level
31 secondary education or vocational training).
32

33 Analyses

34
35 Respondents with missing data on relevant cognitive variables and soft drink or snack
36 consumption were deleted from the sample, resulting in study samples of 1174 and 1139
37 adolescents for soft drink and snack consumption, respectively. In 2007, multilevel linear
38 (soft drinks) and logistic (snack) regression analyses were performed, using MLwiN version
39

1 2.02. A three-level structure was used to take into account that adolescents were nested
2 within classes and classes were nested within schools [29].

3
4 First, a 3-level, random-intercept model was fitted without any explanatory variables to
5 examine the significance of the between-school and between-class variance (Model 0). A
6 significant variance would indicate that the individual behavior clustered within schools
7 and/or classes [30]. Second, demographic factors were included in Model 1. To control
8 for possible confounding, these factors were included in all other models that were fitted.
9 Third, school-canteen factors were added (Model 2) as well as school-neighborhood factors
10 (Model 3). Fourth, a model was fitted with individual cognitions (Model 4). To examine
11 individual cognitions as mediators of the associations between environmental factors and
12 behavior, the four-step procedure indicated by Baron and Kenny [31] was used. Mediation
13 can be established if (1) environmental factors are associated with the outcome behavior
14 (Models 2 and 3); (2) the individual cognitions are associated with the outcome behavior
15 (Model 4); (3) the environmental variables are associated with the individual cognitions
16 (model not presented); and (4) the association between the environmental factor and the
17 outcome behavior decreases when controlling for the mediators (Mediation Model).
18

19 20 **RESULTS**

21 22 **School Environment and Participant Characteristics**

23
24 Four of the 15 schools sold fruit/vegetables, and two schools had low-energy snacks avail-
25 able in their vending machines. A small food store was the closest store for five schools,
26 while only one school had a fruit/vegetable store as the closest store. The mean number
27 of food establishments within 500 meters of schools was 16.7 (range=1–45), and consisted
28 mostly of small food stores (M=6.7, range=0–21), followed by fast food outlets (M=4.7,
29 range=1–12); bakeries (M=2.9, range=0–8); fruit/vegetable stores (M=1.3, range=0–4); and
30 large supermarkets (M=1.2, range=0–4). The mean street-network distance from a school to
31 the nearest food establishment was 284 meters (range=89–619) (data not presented).
32

33 The demographic, cognitive, and behavioral characteristics of the sample are shown in Table
34 7.1. Adolescents reported drinking an average of 1 liter (SD=0.97) of soft drink per day, and
35 51.5% reported consuming more than two portions of snacks per day. The adolescents had
36 particularly positive cognitions regarding soft drink consumption.
37
38
39

Table 7.1 Demographic, cognitive, and behavioral characteristics of the study population

	Soft drink (n=1174) ^a	Snack (n=1139)
Demographics		
Age (mean, SD), years	14.1 (1.2)	14.1 (1.2)
Gender (boys), %	52.9	53.6
Ethnicity (Dutch + Western immigrants), %	49.2	50.2
School level (high school), %	46.1	46.8
Individual cognitions		
Attitude (<i>I think consuming ... is good and pleasant</i>), %		
Disagree/unsure	26.7	43.3
Agree	73.3	56.7
Parental subjective norm (<i>If I consume..., my parents think it's good</i>), %		
Disagree/unsure	54.2	78.4
Agree	45.8	21.6
Parental modeling (<i>My parents consume a lot of...</i>), %		
Disagree/unsure	79.5	92.0
Agree	20.5	8.0
Friends modeling (<i>My friends consume a lot of...</i>), %		
Disagree/unsure	32.8	35.2
Agree	67.2	64.8
Perceived behavioral control (<i>I am able to determine my own consumption, and I think it is easy for me to consume ...</i>), %		
Disagree/unsure	9.3	22.1
Agree	90.7	77.9
Intention (<i>I intend to consume ... in the coming 6 months</i>), %		
Disagree/unsure	20.0	35.3
Agree	80.0	64.7
Behavior		
Soft drink consumption (mean, SD), liters	1.05 (0.97)	—
Snack consumption, %		
≤2 pieces/day	—	48.5
>2 pieces/day	—	51.5

^aThe number of respondents is different for soft drink and snack consumption as the respondents with missing data on a behavior were excluded from the sample.

1 Soft Drink Consumption

2
3 Table 7.2 shows the multivariate associations of the potential correlates with soft drink
4 consumption. Significant between-school variance was found in the null model, which
5 was explained by the individual-level demographics of the students (when controlling for
6 demographics, there remained no between-school variance).

7
8 Gender, ethnicity, and school level were significantly associated with soft drink consump-
9 tion. School canteen-availability factors were not associated with soft drink intake. An
10 intermediate distance to the nearest store of 200–300 meters and the number of small food
11 stores were inversely associated with soft drink consumption. Attitude, parental norm,
12 modeling from friends and parents, and intention were positively associated with soft drink
13 consumption. No significant association was found for perceived behavioral control.

14
15 In the mediation analyses, significant inverse associations were found between cognitions
16 and the distance to the nearest store and the number of small food stores, with ORs ranging
17 between 0.6 and 0.7 (results not presented). The association between environmental factors
18 and soft drink consumption decreased, with percentages ranging from 19% to 48% after
19 controlling for the significant cognitive variables, indicating that more small food stores and
20 a 200–300 meter distance to the nearest shop decreased the positive cognitions toward soft
21 drink consumption, resulting in lower intake.

22 23 Snack Consumption

24
25 No significant between-school and between-class variance was found for snack consump-
26 tion. Vocational-training students were more likely to have high snack intake compared to
27 those attending higher-level education (OR=1.5; 95% CI=1.1, 2.1). No significant associations
28 were found for school-canteen and school-neighborhood factors. Adolescents with positive
29 scores on attitude (OR=1.6; 95% CI=1.2, 2.1); modeling parents (OR=2.0; 95% CI=1.2, 3.5);
30 modeling friends (OR=1.5; 95% CI=1.1, 2.0); parental norm (OR=1.4; 95% CI=1.0, 2.0); and
31 intention (OR=2.5; 95% CI=1.8, 3.4) were more likely to report high snack intake compared
32 to adolescents with more negative cognitions toward snack consumption.

33 34 35 DISCUSSION

36
37 This is one of the first studies to systematically examine the association of self-reported
38 cognitive factors and objectively measured school-environment factors with soft drink
39 and snack consumption among adolescents. As in other studies [10–13], significant positive

Table 7.2 School and class differences in soft drink consumption,^a and the effect of individual cognitions, school-canteen availability, and school-neighborhood factors on soft drink consumption in liters per day (unstandardized regression coefficients)^b

	Model 0 ^c	Model 1	Model 2	Model 3	Model 4	Mediation
Random effects						
Between-school variance	0.074*	0.029	0.006	0.000	0.015	0.000
Between-class variance	0.029	0.013	0.017	0.008	0.009	0.007
Demographics						
Gender (girls)		-0.231**	-0.247**	-0.245**	-0.177***	-0.180**
Age		0.008	0.022	-0.005	-0.017	-0.026
Ethnicity (non-Western)		0.162*	0.141*	0.157*	0.168***	0.179***
School level (vocational)		0.282***	0.561**	0.355**	0.270***	0.258**
School canteen						
Sugar-sweetened soft drink counter (0=ref)						
Medium			-0.373			
High			-0.168			
Sugar-sweetened soft-drink vending (0=ref)						
Medium			-0.258			
High			-0.066			
Low-calorie drinks (low=ref)						
Medium			-0.086			
High			0.010			
School neighborhood						
Supermarket (0=ref)				0.077		
Fast food (low=ref)				-0.055		
Small food stores (low=ref)						
Medium				-0.322***		-0.167*
High				-0.259*		-0.211*
Distance to nearest store (<200m=ref)						
200–300 meters				-0.376**		-0.246***
>300 meters				-0.098		-0.015
Individual cognitions						
Attitude					0.324**	0.352**
Modeling parents					0.294**	0.320**
Modeling friends					0.190**	
Parental norm					0.203**	0.214**
Perceived behavioral control					0.038	
Intention					0.424**	0.426**

Note: Beta's in **bold** indicate a significant association.

^aSchool and class differences are indicated by the between-school and between-class variance. Significance was calculated with the Wald statistic following a chi-square distribution with 1 df. A significant variance would indicate that the individual behavior clusters within schools and/or classes. ^bUnstandardized regression coefficients express the likelihood of soft drink consumption in liters per day. ^cAll models are adjusted for between-school and -class variance and all variables that were included in the specific model.

* $p < 0.05$; ** $p < 0.001$; *** $p < 0.01$

1 associations were found for individual cognitions for both behaviors. This may indicate
2 that cognitions are important factors to target in interventions, that is, by means of health
3 education techniques.

4
5 Only small associations between environmental factors and intake were found. No associa-
6 tions were found for the availability of products in school canteens, and associations with
7 school-neighborhood factors were found only for soft drink consumption. The associations
8 found did not clearly confirm the hypothesis that a higher availability of snack food and soft
9 drink in the school environment would be associated with higher intake of such items. The
10 association between the distance to the nearest food store and soft drink consumption was
11 inconsistent, as no significant inverse association was found for food stores located farther
12 than 300 meters away from a school. This might indicate that adolescents consider 300
13 meters too far to walk and do not visit these stores to buy drinks. Another study found that
14 residing closer to a fast-food restaurant was associated with increased high-fat vegetable
15 intake (e.g., fried potatoes) by adolescents [32]. However, the current study did not find
16 this association for snack intake, indicating that the distance to food stores may not be
17 important for all dietary behaviors. The inverse association between the number of small
18 food stores and soft drink consumption was unexpected, and the opposite of the hypothesis
19 that the presence of more food stores would have a positive effect on intake. The inverse
20 association that was found might indicate that the presence of a greater range of food stores
21 close to schools provides a larger variety of food and drinks from which student can choose,
22 including more healthful options; this may account for the inverse association, but more
23 research is necessary.

24
25 The mediation effect that was found provides some evidence for the hypothesis that environ-
26 mental factors influence soft drink consumption via the cognitions, as proposed by Kremers
27 and colleagues [20] in their EnRG framework. A possible reason why associations between
28 school-environment factors and intake were not detected might be that only crude measures
29 of the complex constructs of proximity and availability were used, without taking into ac-
30 count traffic safety, food prices, policy, and social factors [24, 33, 34]. The use of a broad
31 crow-fly buffer instead of a network buffer (a boundary based on potential walking routes)
32 may also be a reason for the lack of findings. In addition, all adolescents may have enough
33 access to soft drinks and snacks, so that the minor differences in the availability of products
34 in the school environment are not a limiting factor. This is expressed in the nonsignificant
35 between-school variance for snack consumption and in the between-school difference for
36 soft drink consumption that was, in large part, accounted for by differences in demographic
37 factors. Furthermore, intake was assessed based on the average intake per day instead of
38 the intake at school. Other studies found that adolescents consume soft drinks and snacks
39 mainly in other settings—for instance, at home and in fast-food restaurants [35, 36].

1 The results of the present study must be interpreted in the light of several limitations. The
2 cross-sectional design of the study did not allow the determination of causal effects. The use
3 of self-reported measures of intake is a well-known source of potential bias. The categoriza-
4 tion of the in-school availability measures could have led to nonsensitive categorization.
5 Testing the validity of the audit instrument was not possible, because a gold standard does
6 not exist for assessing environmental factors in the school environment. The strength of this
7 study is the combination of individual and objective environmental measures. However,
8 there are also limitations concerning GIS data, as the number of food stores in the com-
9 mercial database might be an under- or over-representation of the actual number of stores.

11 Conclusion

12
13 This study provided little evidence for associations of environmental factors in the school
14 environment with soft drink and snack consumption, while finding clear positive associa-
15 tions between cognitions and soft drink and snack intake. This indicates that such cogni-
16 tions, rather than environmental factors, should be the primary target for interventions.
17 However, the inverse associations between environmental factors and soft drink intake
18 might indicate that the environment can also exert a positive influence on dietary behaviors
19 and cognitions. As this is one of the first studies to examine these factors in the school
20 environment, longitudinal and experimental studies are needed to draw firmer conclusions.

23 ACKNOWLEDGEMENTS

24
25 This study was financially supported by a grant from ZonMw, The Netherlands Organization
26 for Health Research and Development (2100.0103). AT and DC are supported by fellowships
27 from the Victorian Health Promotion Foundation, Australia.

1 **Appendix.** Audit instrument for school canteen availability of drinks and snacks

2	Yes	# of products	Category
3	Availability of drinks in vending machines		
4	<input type="checkbox"/>		Sugar-sweetened soft drinks
5	<input type="checkbox"/>		Sugar-sweetened soft drinks
6	<input type="checkbox"/>		Low-calorie drinks
7	<input type="checkbox"/>		Low-calorie drinks
8	<input type="checkbox"/>		Sugar-sweetened soft drinks
9	<input type="checkbox"/>		Low-calorie drinks
10	Availability of drinks at canteen counter		
10	<input type="checkbox"/>		Sugar-sweetened soft drinks
11	<input type="checkbox"/>		Sugar-sweetened soft drinks
12	<input type="checkbox"/>		Low-calorie drinks
13	<input type="checkbox"/>		Low-calorie drinks
14	<input type="checkbox"/>		Sugar-sweetened soft drinks
15	<input type="checkbox"/>		Low-calorie drinks
16	Availability of snacks in vending machines		
17	<input type="checkbox"/>		Energy-dense snacks
18	<input type="checkbox"/>		Energy-dense snacks
19	<input type="checkbox"/>		Energy-dense snacks
20	<input type="checkbox"/>		Energy-dense snacks
21	<input type="checkbox"/>		Energy-dense snacks
22	<input type="checkbox"/>		Low-energy snacks
23	<input type="checkbox"/>		Low-energy snacks
24	<input type="checkbox"/>		Energy-dense snacks
25	<input type="checkbox"/>		Low-energy snacks
26	Availability of snacks at canteen counters		
27	<input type="checkbox"/>		Energy-dense snacks
28	<input type="checkbox"/>		Energy-dense snacks
29	<input type="checkbox"/>		Energy-dense snacks
30	<input type="checkbox"/>		Energy-dense snacks
31	<input type="checkbox"/>		Energy-dense snacks
32	<input type="checkbox"/>		Energy-dense snacks
33	<input type="checkbox"/>		Low-energy snacks
34	<input type="checkbox"/>		Low-energy snacks
35	<input type="checkbox"/>		Energy-dense snacks
36	<input type="checkbox"/>		Energy-dense snacks
37	<input type="checkbox"/>		Energy-dense snacks
38	<input type="checkbox"/>		Low-energy snacks
39	<input type="checkbox"/>		Low-energy snacks

1 REFERENCES

- 2 1. Hedley AA, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM: Prevalence of overweight
3 and obesity among US children, adolescents, and adults, 1999-2002. *JAMA* 2004, 291(23):2847-2850.
- 4 2. Lobstein T, Frelut ML: Prevalence of overweight among children in Europe. *Obes Rev* 2003, 4(4):195-
5 200.
- 6 3. Swinburn BA, Caterson I, Seidell JC, James WP: Diet, nutrition and the prevention of excess weight
7 gain and obesity. *Public Health Nutr* 2004, 7(1A):123-146.
- 8 4. Moreno LA, Rodriguez G: Dietary risk factors for development of childhood obesity. *Curr Opin Clin
9 Nutr Metab Care* 2007, 10(3):336-341.
- 10 5. Bowman SA, Gortmaker SL, Ebbeling CB, Pereira MA, Ludwig DS: Effects of fast-food consumption
11 on energy intake and diet quality among children in a national household survey. *Pediatrics* 2004,
12 113(1 Pt 1):112-118.
- 13 6. Paeratakul S, Ferdinand DP, Champagne CM, Ryan DH, Bray GA: Fast-food consumption among US
14 adults and children: dietary and nutrient intake profile. *J Am Diet Assoc* 2003, 103(10):1332-1338.
- 15 7. Frary CD, Johnson RK, Wang MQ: Children and adolescents' choices of foods and beverages high
16 in added sugars are associated with intakes of key nutrients and food groups. *J Adolesc Health* 2004,
17 34(1):56-63.
- 18 8. Harnack L, Stang J, Story M: Soft drink consumption among US children and adolescents: nutritional
19 consequences. *J Am Diet Assoc* 1999, 99(4):436-441.
- 20 9. Sanigorski AM, Bell AC, Swinburn BA: Association of key foods and beverages with obesity in
21 Australian schoolchildren. *Public Health Nutr* 2007, 10(2):152-157.
- 22 10. Grimm GC, Harnack L, Story M: Factors associated with soft drink consumption in school-aged
23 children. *J Am Diet Assoc* 2004, 104(8):1244-1249.
- 24 11. de Bruijn GJ, Kremers SP, Schaalma H, van Mechelen W, Brug J: Determinants of adolescent bicycle
25 use for transportation and snacking behavior. *Prev Med* 2005, 40(6):658-667.
- 26 12. Kassem NO, Lee JW: Understanding soft drink consumption among male adolescents using the
27 theory of planned behavior. *J Behav Med* 2004, 27(3):273-296.
- 28 13. Kassem NO, Lee JW, Modeste NN, Johnston PK: Understanding soft drink consumption among
29 female adolescents using the Theory of Planned Behavior. *Health Educ Res* 2003, 18(3):278-291.
- 30 14. Ajzen I: *Attitudes, personality, and behavior*: Homewood, IL, US: Dorsey Press; 1988.
- 31 15. Brug J, Lechner L, De Vries H: Psychosocial determinants of fruit and vegetable consumption. *Ap-
32 petite* 1995, 25(3):285-296.
- 33 16. Vries HD, Backbier E, Kok G, Dijkstra M: The impact of social influences in the context of attitude,
34 self-efficacy, intention, and previous behavior as predictors of smoking onset. *Journal of applied social
35 psychology* 1995, 25(3):237-257.
- 36 17. Swinburn B, Egger G, Raza F: Dissecting obesogenic environments: the development and application
37 of a framework for identifying and prioritizing environmental interventions for obesity. *Prev Med*
38 1999, 29(6 Pt 1):563-570.
- 39 18. van der Horst K, Oenema A, Ferreira I, Wendel-Vos W, Giskes K, van Lenthe F, Brug J: A systematic
review of environmental correlates of obesity-related dietary behaviors in youth. *Health Educ Res*
2007, 22(2):203-226.
19. Ball K, Timperio AF, Crawford DA: Understanding environmental influences on nutrition and physi-
cal activity behaviors: where should we look and what should we count? *Int J Behav Nutr Phys Act*
2006, 3:33.

20. Kremers SP, de Bruijn GJ, Visscher TL, van Mechelen W, de Vries NK, Brug J: Environmental influences on energy balance-related behaviors: a dual-process view. *Int J Behav Nutr Phys Act* 2006, 3:9.
21. Johnston LD, Delva J, O'Malley PM: Soft drink availability, contracts, and revenues in American secondary schools. *American journal of preventive medicine* 2007, 33(4 Suppl):S209-225.
22. O'Toole TP, Anderson S, Miller C, Guthrie J: Nutrition services and foods and beverages available at school: results from the School Health Policies and Programs Study 2006. *The Journal of school health* 2007, 77(8):500-521.
23. Kubik MY, Lytle LA, Hannan PJ, Perry CL, Story M: The association of the school food environment with dietary behaviors of young adolescents. *Am J Public Health* 2003, 93(7):1168-1173.
24. Neumark-Sztainer D, French SA, Hannan PJ, Story M, Fulkerson JA: School lunch and snacking patterns among high school students: associations with school food environment and policies. *Int J Behav Nutr Phys Act* 2005, 2(1):14.
25. Kipke MD, Iverson E, Moore D, Booker C, Ruelas V, Peters AL, Kaufman F: Food and park environments: neighborhood-level risks for childhood obesity in east Los Angeles. *J Adolesc Health* 2007, 40(4):325-333.
26. Austin SB, Melly SJ, Sanchez BN, Patel A, Buka S, Gortmaker SL: Clustering of fast-food restaurants around schools: a novel application of spatial statistics to the study of food environments. *Am J Public Health* 2005, 95(9):1575-1581.
27. Zenk SN, Powell LM: US secondary schools and food outlets. *Health Place* 2007.
28. van der Horst K, A O, van de Looij PM, H B: The ENDORSE study: Research into environmental determinants of obesity related behaviors in Rotterdam Schoolchildren. *BMC Public Health* to be resubmitted.
29. Twisk JWR: *Applied Multilevel Analysis: A Practical Guide for Medical Researchers*. 1 edition: Cambridge University Press; 2006.
30. Merlo J, Chaix B, Yang M, Lynch J, Rastam L: A brief conceptual tutorial of multilevel analysis in social epidemiology: linking the statistical concept of clustering to the idea of contextual phenomenon. *J Epidemiol Community Health* 2005, 59(6):443-449.
31. Baron RM, Kenny DA: The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol* 1986, 51(6):1173-1182.
32. Jago R, Baranowski T, Baranowski JC, Cullen KW, Thompson D: Distance to food stores & adolescent male fruit and vegetable consumption: mediation effects. *Int J Behav Nutr Phys Act* 2007, 4(1):35.
33. Hilbert A, Rief W, Braehler E: What determines public support of obesity prevention? *J Epidemiol Community Health* 2007, 61(7):585-590.
34. Nollen NL, Befort CA, Snow P, Daley CM, Ellerbeck EF, Ahluwalia JS: The school food environment and adolescent obesity: qualitative insights from high school principals and food service personnel. *Int J Behav Nutr Phys Act* 2007, 4:18.
35. Savage G, Macfarlane A, Ball K, Worsley A, Crawford D: Snacking behaviours of adolescents and their association with skipping meals. *Int J Behav Nutr Phys Act* 2007, 4(1):36.
36. French SA, Lin BH, Guthrie JF: National trends in soft drink consumption among children and adolescents age 6 to 17 years: prevalence, amounts, and sources, 1977/1978 to 1994/1998. *J Am Diet Assoc* 2003, 103(10):1326-1331.

8 Do individual cognitions mediate the association of socio-cultural and physical environmental factors with adolescent sports participation?

Van der Horst K, Oenema A, te Velde SJ, Brug J. Do individual cognitions mediate the association of socio-cultural and physical environmental factors with adolescent sports participation?

Public Health Nutrition (submitted).

1 **ABSTRACT**

2

3 **Objective:** To examine the associations of perceived physical environmental factors (avail-
4 ability of physical activity attributes at home, physical activity facilities in the neighbour-
5 hood, neighbourhood pleasantness and safety) and social environmental factors (parental
6 sports behaviour and parental rule regarding sports participation) with adolescent leisure
7 time sports participation, and to explore whether the associations found were mediated by
8 individual cognitions as derived from the Theory of Planned Behaviour (TPB).

9 **Design:** Cross-sectional

10 **Setting:** Adolescents from 17 schools in Rotterdam, the Netherlands, completed a ques-
11 tionnaire during school hours that included self-report measures of leisure time sports
12 participation, the perceived physical environmental factors and TPB variables. Information
13 about parental sports behaviour and parental rule was obtained from a questionnaire that
14 was completed by one parent of the adolescents.

15 **Subjects:** Data was collected from 584 adolescent – parent combinations.

16 **Results:** Data was analyzed with multilevel logistic regression analyses. Availability of
17 physical activity attributes at home (OR = 1.26), parents' sports behaviour (OR = 2.03) and
18 parental rule (OR = 1.64) were associated with a higher likelihood of adolescents' leisure
19 time sports participation. These associations were partly mediated by attitude and intention.

20 **Conclusions:** Adolescents were more likely to engage in leisure time sports when PA at-
21 tributes were available at home, when parents participated in sports activities and had a
22 rule about their offspring participation in sport activities. These associations were partly
23 mediated by attitude and intention. These results suggest that parents can importantly
24 promote sports participation among their offspring by making sports activities accessible
25 and a family routine.

26

27

28

29

30

31

32

33

34

35

36

37

38

39

1 INTRODUCTION

2
3 Insufficient physical activity (PA) is a risk factor for a range of chronic conditions including
4 obesity, among adults as well as adolescents [1, 2]. Most adolescents do not meet the recom-
5 mended minimum levels of engaging in at least 60 minutes of moderate to vigorous inten-
6 sity PA each day [3-6]. Adolescents are a particularly important target group to improve PA
7 levels since physically active adolescents are more likely to become active adults [7]. To be
8 able to increase PA levels among adolescents, it is important to develop interventions that
9 target the most important determinants of PA.

10
11 In addition to individual cognitions such as attitude, subjective norm, perceived behavioural
12 control and intention, as derived from the Theory of Planned Behaviour (TPB) [8] physical
13 and social environmental factors may be important determinants of PA behaviour. Kremers
14 and colleagues, in their Environmental Research framework for weight Gain prevention
15 (EnRG) [9] suggest that environmental factors may have a direct and an indirect association
16 with behaviour. The direct association reflects a more automatic and unconscious effect of
17 the environment on behaviours. The indirect influence suggests that environmental factors
18 influence PA via the individual cognitions, e.g. environments that offer appealing and easily
19 accessible opportunities for PA may result in more positive attitudes, perceived behavioural
20 control and intentions toward leisure time PA, which may result in higher PA levels. The
21 TPB also assumes that the impact of various external variables such as physical and social
22 environmental factors on behaviour is mediated by attitude, subjective norm, perceived
23 behavioural control and intention.

24
25 Physical environmental factors such as the availability and accessibility of PA opportuni-
26 ties have received most attention in exploring environmental determinants of PA [9-11].
27 However, a recent review indicated that the evidence for the role of social environmental
28 factors is stronger [12-14].

29 Earlier studies have found that among adults the association of perceived neighbourhood
30 with walking was mediated by attitude [15] and that associations of perceived neighbour-
31 hood aesthetics with walking were mediated by attitude and intention [16]. De Bruijn
32 and colleagues found that the association of environmental aesthetics and distance to PA
33 facilities on PA among adolescents was mediated by intention to be physically active [17].
34 Motl and colleagues found that the association of equipment accessibility with adolescent
35 girls PA was mediated by self-efficacy [18]. These previous studies indicate that some TPB
36 variables may be more likely to serve as a mediators in environment – behaviour relation-
37 ships than others [17, 19], with the strongest evidence for attitudes as a potential mediating
38 variable [15-17, 19, 20]. Most previous studies have investigated mediation pathways for
39 physical environmental factors.

1 The aim of the present study was to examine the associations between physical environ-
2 mental factors (availability of PA attributes at home, PA facilities in the neighbourhood,
3 neighbourhood pleasantness and safety) and social environmental factors (parents own
4 sports behaviour and parental rule about sport participation) with adolescents' leisure
5 time sports participation and to explore whether these associations are mediated by TPB
6 variables (figure 1).

7

8

9 **METHODS**

10

11 **Study design and sample selection**

12

13 Baseline data from the ENvironmental Determinants of Obesity in Rotterdam Schoolchil-
14 drEn (ENDORSE) study were used [21] for which data were collected among adolescents in
15 the first (12- to 13-year-olds) and third (14- to 15-year-olds) years of secondary school. The
16 Medical Ethics Committee of Erasmus University Medical Center declared no objection to
17 the project. Schools located in the Rotterdam area that participate in the Youth Monitor
18 Rotterdam (YMR) (N = 56) were invited for participation in the ENDORSE study. Sub-
19 sequently, a random sample of 17 school locations was drawn from the pool of 24 schools
20 that were willing to participate. On average, five classes per school location were randomly
21 selected to participate in the study and 1668 adolescents and their parents were eligible for
22 participation. In the baseline survey, 187 adolescents were absent during the questionnaire
23 assessment. Due to questionnaire printing mistakes, 120 records, including those from one
24 entire school had to be removed. Response rate for the parent questionnaire was 43%, result-
25 ing in 584 adolescent – parent combinations. There was no data available on the parents and
26 adolescents that did not participate in the study and examining response bias was therefore
27 not possible. Compared to the data from the total adolescent sample, in the sample used for
28 this study adolescents with a non-Western ethnic background (36.6% compared to 50.5%)
29 and attending vocational schools (49.1% compared to 56.7%) were underrepresented.

30

31 **Procedure**

32

33 Parents received a letter announcing and explaining the ENDORSE study and could refuse
34 participation of their child(ren) by sending a note to the adolescent's teacher. Between October
35 2005 and May 2006, the students completed the ENDORSE questionnaire in the classroom in
36 the presence of a teacher and a trained research assistant within one school hour (50 minutes).
37 The adolescents were handed a questionnaire with a pre-addressed and stamped envelope for
38 completion by one of their parents. To increase the participation rates, five I-pods were raffled
39 amongst the parent respondents and two reminders were send to the parents.

1 Measures

3 *Leisure time sports participation*

4 The relevant questions from the Activity QUestionnaire for Adolescents and Adults
5 (AQUAA) were used to assess leisure time sports activities [22]. The test-retest reliability
6 for this questionnaire was moderate (intraclass correlations 0.30-0.59) and validity with
7 accelerometer data was low (Spearman correlation coefficient = 0.21 for vigorous activities)
8 [22]. No validity data are available on sports behavior. First, adolescents were asked to write
9 in text boxes in a pre-structured format up to three sports activities that they had engaged
10 in, in the past week. Adolescents were asked to write down organized and unorganized
11 sports they engaged in. Second, they had to tick on how many days in the past week (1 to
12 7) they had engaged in this activity. Third, they had to indicate how long on average they
13 participated in this activity per occasion, in an open answering format with text boxes to
14 indicate hours and minutes. The frequency and duration of the activities were multiplied,
15 and then divided by the total number of days to provide the average minutes per day spent
16 doing leisure time sports activities. As this variable was highly skewed and the distribution
17 could not be improved through transformation, two categories were created: engaging in
18 leisure time sports activities for less than 30 minutes per day (coded as 0), or 30 minutes or
19 more per day (coded as 1). This cut-off point for sports behaviour was based on the Dutch
20 physical activity recommendation that children and adolescents should engage in moderate
21 to vigorous physical activity at least 30 minutes per day and practice sports at least 3 days
22 a week.

24 *TPB variables*

25 Attitude, subjective parental norm, perceived behaviour control and intention were spe-
26 cifically assessed in relation to participation in leisure time sports activities. All questions
27 could be answered on five-point bipolar answering scales. Attitude was assessed with two
28 items asking if the adolescent considers sports and PA in leisure time as very good (+2)
29 or very bad (-2) and as very pleasant (+2) or very unpleasant (-2). The mean item score
30 (Cronbach's alpha = 0.79, Intraclass correlation = 0.66) for these items was calculated [35].
31 Subjective norm was assessed with one item: 'if I engage in sports and PA in leisure time,
32 my parents consider that as very good (+2) – very bad (-2)'. Perceived behaviour control was
33 assessed with one item asking how easy or difficult it is to engage in sports and PA in leisure
34 time with an answering scale ranging from very easy (+2) to very difficult (-2). Intention to
35 perform the behaviour was assessed with a single item: 'Do you intend to engage in sports
36 and PA in leisure time in the next six months?' with an answering scale ranging from Yes, I
37 certainly do (+2) to no, I certainly do not (-2).

1 *Physical environmental variables*

2 We assessed availability of PA opportunities at home by providing a list of ten sport 'at-
3 tributes' (i.e. bicycle, dog, home trainer /treadmill, running shoes, stationary aerobic
4 equipment, step aerobics, skates, balls, racquets, jumping rope). This list was translated
5 from Sallis et al. [23]. The adolescents could tick which of these were available in their
6 home. A score of PA attributes available at home was calculated by adding up the "yes"
7 responses to these questions. Perceived neighbourhood pleasantness was assessed with two
8 questions: "I think my neighbourhood provides a pleasant living environment", and "I think
9 my neighbourhood is attractive", that could be answered on 5-point scales ranging from
10 totally agree (+2) to totally disagree (-2). The mean item score (Cronbachs alpha = 0.78,
11 Intraclass correlation = 0.64) of these two items was calculated. Perceived neighbourhood
12 safety was assessed with four questions: "There is a lot of traffic in my neighbourhood", "It is
13 unsafe to bicycle in my neighbourhood", "I feel safe when I am in my neighbourhood", and
14 "It is unsafe to be outside in my neighbourhood", using the same five point answering scale
15 format as neighbourhood attractiveness. The mean item score (Cronbachs alpha = 0.64) of
16 these four items was calculated. Perceived availability of PA facilities in the neighbourhood
17 was assessed by asking the adolescents to indicate whether or not (yes/no answering format)
18 there were parks, sports clubs, sports/playing grounds present in the neighbourhood where
19 they lived. The yes responses were summed to form one score for these four items.

20

21 *Social environmental variables*

22 Parental rule about PA was assessed with one question in the questionnaire for parents: "is
23 it a rule in your household that your child has to participate in sport activities?" in a yes/
24 no answering format. Parents own sports behaviour was assessed in the questionnaire for
25 parents with two questions assessing frequency and duration, using the relevant questions
26 from the SQUASH questionnaire [24]. The Spearman correlation for overall reproducibility
27 of the SQUASH was 0.58 (95% CI = 0.36-0.74), and correlations for the reproducibility of
28 leisure time sport was 0.90. Spearman's correlation coefficient between activity monitor
29 readings and the total activity score was 0.45 (95% CI = 0.17-0.66) [24]. Frequency was
30 assessed with: "How many days per week do you engage in sports activities?" on a 7-point
31 scale from 1 = 1 day per week to 7 = every day. The duration was assessed with "On a day
32 that you participate in sports activities, how long do you do this on average?" and hours and
33 minutes could be reported. The frequency and duration of the activities were multiplied,
34 and then divided by the total number of days to provide the average minutes per day. As
35 this variable was highly skewed, two categories were created: engaging in sports activities
36 (coded as 1) and not engaging in sports activities.

37

38

39

1 Demographics

2 To establish ethnicity according to the Statistics Netherlands definition, adolescents were
3 asked to report in which country their parents had been born [25]. Adolescents were consid-
4 ered to be from a Western ethnic background if both parents had been born in a European
5 country, North America, Oceania, Indonesia or Japan. Based on the socio-economic and
6 cultural position of immigrants in the Netherlands from Oceania, Japan and Indonesia (a
7 former colony of the Netherlands), children from these immigrants were also included in
8 the Western ethnic group. Adolescents were considered to be from a non-Western ethnic
9 background if one or both parents had been born in other countries. The school type the
10 adolescents attended was categorized into two levels: vocational schools and higher-level
11 secondary education (pre-academic). The schools provided educational level information.
12 Age was determined based on the date of the measurements and the date of birth that were
13 provided by the schools.

15 Data analyses

17 Possible multi-collinearity problems were examined with bivariate correlations and not
18 encountered; all inter-correlations between predictors were below 0.6.

19 Mediation analyses according to suggestions of MacKinnon (2008) were used to identify
20 total effects, direct effects and mediated effects in the associations of physical environmental
21 factors (availability of PA attributes at home, availability of PA facilities in the neighbour-
22 hood, perceived neighbourhood pleasantness and safety) and socio-cultural environmental
23 factors (parents sports behaviour and parental rule about sport participation) with ado-
24 lescents' leisure time sports participation [26]. To do so, we explored associations between
25 the environmental variables and TPB variables with multivariate linear regression analyses
26 (step 1, path a in figure 1). Next, we examined if the potential mediators from the TPB were
27 associated with leisure time sports, after adjustment for the environmental variables (step 2,
28 path b in figure 1). The total effect of physical environmental factors and socio-cultural envi-
29 ronmental factors on adolescents' leisure time sports participation (step 3, path c), and after
30 adjustment for the possible mediator, the direct effect of environmental variables on sports
31 participation (step 4, path c'), were examined in various models. As suggested by MacKin-
32 non [26] and also outlined by Cerin and MacKinnon [27], a significant association between
33 environmental variables (predictor variables) and sports participation (outcome variable)
34 is not a requirement for mediation to occur, since absence of an overall relationship may
35 be due to suppression effects. Therefore also non-significant environmental factors were
36 included in the mediation analyses. Steps 2-4 were examined by means of multi-level multi-
37 variate logistic regression analyses. All analyses were adjusted for gender, age, ethnicity and
38 school level, as these are possible confounding factors. All analyses were performed with
39 MLwiN version 2.02. A three-level structure was used to take into account that adolescents




Figure 8.1 Conceptual model of the direct and indirect association of physical and socio-cultural environmental factors

were nested within classes and classes within schools [28]. Because of the dichotomous outcome variables there are different scales across the (logistic) regression analyses that makes it incorrect to use the 'difference-of-coefficients estimate' as an estimate of the mediation effect [27, 29]. One solution to overcome this difference in scaling is to standardize the regression coefficients before mediation is estimated [29, 30]. The standardized coefficients were subsequently used to estimate the proportion mediated ($(c_{\text{standardized}} - c'_{\text{standardized}}) / c_{\text{standardized}}$) and were additionally entered in the Sobel test [31] to formally test the mediation effect.

RESULTS

Sixty percent of the adolescents reported to participate in leisure time sports for more than 30 minutes per day. The adolescents reported positive cognitions regarding leisure time sports participation. On average adolescents reported to have four of the listed PA attributes available at home (range 0-10) and to have three PA facilities available in their neighborhood (range 0-4), and reported positive perceptions of neighborhood safety (mean = 0.64, SD = 0.76) and pleasantness (mean = 0.76, SD = 0.99). A majority of the parents reported that it was a rule in the household that the adolescent had to do some kind of sports (66%) and 59% of the parents participated in leisure time sports activities themselves (Table 8.1).

Associations between environmental factors and TPB variables

Multivariate analyses showed that most of the physical and social environmental variables were significantly positively associated with TPB variables (Table 8.2). No associations were found for neighborhood facilities with attitude, perceived behavior control and intention. PA attributes at home and neighborhood safety were not associated with perceived behavior control. Parents' sport behavior was not associated with subjective parental norm.

Table 8.1. Characteristics of the study population (N=584)

Variable	Percentage / Mean; SD	Range	Cronbachs alpha / Intraclass correlation
Behavior			
Sports participation (>= 30 min/week)	59.8%		
Demographics			
Gender (girls)	45.2%		
Ethnicity (non-Western)	36.6%		
Age	M=13.91; SD=1.13	11.9 – 17.6	
Educational level (lower)	49.1%		
Individual cognitions			
Attitude	M=1.29; SD=0.64	-2 – 2	0.79 / 0.66
Parental norm	M=1.50; SD=0.61	-2 – 2	
Perceived behavior control	M=1.12; SD=0.81	-2 – 2	
Intention	M=1.57; SD=0.79	-2 – 2	
Physical environment			
PA attributes at home	M=3.97; SD=2.08	0 – 10	
Neighborhood safety	M=0.64; SD=0.76	-2 – 2	0.64
Neighborhood pleasantness	M=0.76; SD=0.99	-2 – 2	0.78 / 0.64
Neighborhood facilities	M=3.26; SD=0.95	0 – 4	
Socio-cultural environment			
Parental rule to play sports (yes)	65.8%		
Parent behavior (parent does practice sports)	59.4%		

Associations of TPB variables with adolescents' leisure time sports participation

To establish a mediation effect, the potential mediators must be associated with the outcome variable after adjustment for de independent, environmental variables [26]. Multivariate analyses (Table 8.3) showed that attitude and intention were found to be significantly associated with a higher likelihood of participating in leisure time sports after adjustment for physical environmental variables (attitude OR = 2.30; intention OR = 2.10) and after adjustment for social environmental variables (attitude OR = 2.24; intention OR = 2.03).

Mediating effects of TPB variables

As neighborhood facilities were not associated with attitude and intention, this factor was not included in the mediation models (Table 8.4). The association of PA attributes at home with a higher likelihood of participating in leisure time sports was partly mediated by attitude (17.4%) and intention (21.6%), as indicated by the significant Sobel test results. The

Table 8.2 Results of multivariate linear regression analyses (unstandardized regression coefficients) of physical and social environmental variables with TPB variables as dependent variables, adjusted for age, gender, ethnicity, school type and clustering within classes and schools

	Attitude	Parental norm	Perceived behavioral control	Intention
PA attributes at home	0.039**	0.042**	0.026	0.056***
Neighborhood safety	0.146***	0.120***	0.016	0.186***
Neighborhood pleasantness	0.168***	0.090***	0.105**	0.115***
Neighborhood facilities	0.035	0.059**	0.028	0.049
Parental rule	0.141*	0.121*	0.174*	0.166*
Parent behavior	0.135*	-0.001	0.185**	0.180**

*p<0.05; **p<0.01, ***p<0.001

Table 8.3 Results of multivariate logistic regression analysis (odds ratios) examining the association between potential mediators (TPB variables) and leisure time sports participation (≥ 30 min), adjusting for the physical (model 1) and social (model 2) environmental factors^a

	Model 1		Model 2	
	OR	95% CI	OR	95% CI
Physical environmental factors				
PA attributes at home	1.20	1.07-1.35		
Neighborhood safety	1.22	0.86-1.73		
Neighborhood pleasantness	0.89	0.68-1.16		
PA facilities in the neighborhood	1.02	0.83-1.23		
<i>Socio-cultural environmental factors</i>				
Parental rule			1.37	0.85-2.20
Parents sports behavior			1.79	1.14-2.80
TPB variables				
Attitude	2.30	1.46-3.61	2.24	1.48-3.39
Parental norm	1.21	0.82-1.80	1.29	0.87-1.92
Perceived behavioral control	1.00	0.76-1.33	0.96	0.72-1.27
Intention	2.10	1.47-3.02	2.03	1.42-2.91

^a Multivariate logistic regression analyses adjusted for age, gender, ethnicity, school type and clustering within classes and schools

association between neighborhood safety and sports participation was significantly mediated by intention; however, both direct and indirect associations were not significant. The direction of the association between neighborhood pleasantness and sports participation changed after adjustment for attitude, which suggests an inconsistent mediation model.

The associations of parental rule and parents' sports behavior with a higher likelihood of participating in leisure time sports were also partly mediated by attitude and intention with

Table 8.4 Results of logistic regression analyses to examine the mediation of the association between physical environmental factors with leisure time sports participation (≥ 30 min) by attitude (model 2) and intention (model 3) ^a

	Model 1		Model 2		Sobel test	Proportion mediated ^b	Model 3		Sobel test	Proportion mediated ^b
	OR	95% CI	OR	95% CI			OR	95% CI		
Physical environmental factors										
PA attributes at home	1.26	1.13-1.42	1.24	1.10-1.39	2.54 **	17.4	1.22	1.09-1.37	2.86**	21.6
Neighborhood safety	1.28	0.93-1.77	1.32	0.94-1.86	3.40 ***	-3.4 ^c	1.16	0.83-1.63	3.37***	44.3
Neighborhood pleasantness	1.06	0.83-1.36	0.87	0.66-1.13	4.40***	312.2 ^d	1.01	0.78-1.31	2.92**	82.2
TPB variables										
Attitude			3.44	2.33-5.08						
Intention							2.71	1.93-3.80		

^a Multivariate logistic regression analyses adjusted for age, gender, ethnicity, school type and clustering within classes and schools

^b as calculated with the standardized coefficients (see methods section)

^{c,d} Negative values and values $> 100\%$ indicate inconsistent mediation models and the results cannot be interpreted.

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

percentages ranging between 20.1 and 31.7 (Table 8.5). The associations of PA attributes at home and parent behavior with leisure time sports remained statistically significant, while the association of parental rule lost significance in the mediation models.

DISCUSSION

In this study, associations of socio-cultural and physical environmental factors with adolescents' leisure time sports participation were examined and it was explored if these associations were mediated by individual cognitions such as attitudes and intentions. Results showed that parents' sport behaviour, parental rule about sports behaviour and availability of PA attributes at home were associated with a higher likelihood that adolescents engaged in sports behaviour. We cannot draw conclusions upon the findings that resulted from an inconsistent mediation model [32]. The inconsistent model is a result of the fact that the direct association between neighbourhood pleasantness and sports participation was weak but positive, while the indirect association was also weak and non significant, but negative.

Table 8.5 Results of logistic regression analyses to examine the mediation of the association between social environmental factors with leisure time sports participation (≥ 30 min) by attitude (model 2) and intention (model 3) ^a

	Model 1		Model 2		Sobel test	Proportion mediated ^b	Model 3		Sobel test	Proportion mediated ^b
	OR	95% CI	OR	95% CI	z score	%	OR	95% CI	z score	%
Social environmental factors										
Parental rule	1.64	1.05-2.56	1.51	0.95-2.42	2.20*	23.1	1.45	0.91-2.78	2.43*	31.7
Parent behavior	2.03	1.26-3.25	1.85	1.19-2.87	2.29*	20.1	1.80	1.16-2.78	2.13*	23.3
TPB variables										
Attitude			3.38	2.32-4.91						
Intention							2.77	2.00-3.85		

^a Multivariate logistic regression analyses adjusted for age, gender, ethnicity, school type and clustering within classes and schools

^b as calculated with the standardized coefficients (see methods section)

* $p < 0.05$

Evidence was found for partial mediation of social and physical environmental factors by attitude and intention. However, also a direct significant association remained of PA attributes at home and parental behaviour with leisure time sports participation. These findings are in accordance with those of earlier studies where attitudes were found to be strong mediators of the association between physical environmental factors and PA [15-17, 19, 20]. The results suggests that both direct, automatic influences of the environment and more reasoned cognitive processes are important in adolescents' sports participation as suggested in the EnRG framework [9] However, as we found rather strong associations of attitude and intention with leisure time sports participation in this study, adolescents leisure time sports behaviour seems to be, at least partly, also the result of a more reasoned, deliberate process, that is not influenced by the environmental factors considered in the present study. This does make sense, since sport activities are less likely to be part of routine habits, more likely to need to be planned in advance and to be dependent on explicit positive motivation, than, for example, daily activities. The present results support this hypothesis to a certain extend, as we found significant but small correlations between environmental factors and cognitions. This association of cognitions, independent from the environmental factors is not clearly stated in the EnRG framework that focuses on cognitions as mediators of environmental influences. Two of the four cognitions included in the present study, i.e. parental subjective norm and perceived behavioural control, were not found to be associated with

1 sports behaviour. This might indicate that not all cognitions as suggested by the Theory
2 of Planned Behaviour are important mediators or play a role in the suggested reasoned
3 process for this particular behaviour in this population group. On the other hand, the non-
4 significant results might be caused by the limited assessment of these constructs with only
5 one or two items.

6
7 In accordance with other studies, social factors seem to be more strongly associated with
8 physical activity behaviour than physical environmental factors [12-14]. The direct associa-
9 tions of parents' sports behaviour and the availability of PA equipment at home, support
10 earlier evidence that parental example and support (for instance through providing good
11 sports equipment at home), are important for PA promotion. The range of social factors
12 considered in the present study was narrow, and factors such as social networks, friends'
13 support and behaviour and the perceived behaviour of parents should be included in future
14 studies to provide further insight into the specific aspects of the social environment that are
15 most important for adolescents PA behaviours [18, 33]. Next to this, future research should
16 address the moderating effects of socio-demographic factors mentioned in the EnRG
17 framework as there are clear gender differences in sports participation and correlates of
18 physical activity might be different for boys and girls. For example, mothers' physical activ-
19 ity appears to be more often associated with girls' rather than boys' physical activity [12].

20
21 One possible limitation of this study was that we used perceptions of the environment
22 instead of more objective measures of the physical environment. Perceived environmental
23 factors are, of course, also cognitive representations (i.e. of environmental factors), and
24 different mediating pathways may be apparent with more objective assessments of the
25 environment. Evidence points out that perceived and objective environmental factors are
26 different constructs [34] and that perceptions of the environment only partly depend on
27 what is objectively available in the environment [35]. Studies exploring TPB variables as
28 well as perceived environmental factors as mediators of the associations between objective
29 environmental characteristics and PA behaviour may help to unravel the interplay between
30 individual and environmental factors in influencing energy balance-related behaviours as
31 proposed in the EnRG framework.

32
33 The following limitations should be taken into account when interpreting the results of this
34 study.

35 The cross-sectional design of the study did not allow us to determine causal effects and
36 is an important limitation in research examining mediation pathways. Physically active
37 adolescent might be more aware of physical activity equipment in their environment and
38 they might select more or less the neighbourhood they are active in by having a specific
39 definition about how large the neighbourhood is. Having positive cognitions towards sports

1 might shape the adolescents environment. For instance, adolescents might influence their
2 parents by promoting sports activities and asking for more equipment. Next to this, the
3 sample size of this study was restricted because of the rather low response rate for the parent
4 questionnaire. The fact that adolescents from non-Western ethnic background and lower
5 educational level were underrepresented in the sample suggests selection bias. Several limi-
6 tations relate to the measurement instruments used in the study. First, adolescents sports
7 behaviour was based on self-report and in a validation study with use of accelerometers was
8 shown that the questionnaire had limited validity and that adolescents over-reported their
9 activity levels [22]. Second, the TPB variables were assessed with only one or two items
10 leading to limited reliability. The TPB variables, particularly perceived behaviour control,
11 might have been not robust enough to demonstrate associations and to show up as a media-
12 tor. Environmental constructs were often measured with only one ore two items with only
13 moderate reliability. In explorative research more effort needs to be done to construct better
14 scales that examine all aspects of the perceived environmental factors. More qualitative
15 research is needed to improve existing measurement instruments and scales. Next to this,
16 only a limited set of perceived environmental variables was used in this study. Especially
17 other social environmental influences such as encouragement of parents and friends might
18 be important in explaining physical activity behaviours [12, 14, 36]. Parents' sports behaviour
19 was also assessed with other questions compared to adolescents' sports participation and
20 other cut-off points were used. This could also have affected the associations found. Third,
21 adolescents with overweight or lack of physical activity might have given social desirable
22 answers on sports behaviour and on the theory of planned behaviour items as well, which
23 could have influenced the associations between cognitions and behaviour.

24 Nevertheless, this explorative study contributes to the structured examination of the as-
25 sociations between environmental factors and physical activity and the suggested mediation
26 by TPB variables as supposed by the EnRG framework.

27 28 Conclusion

29
30 Dutch adolescents were more likely to engage in leisure time sports when PA attributes
31 were available at home, when parents participated in sports activities and when parents had
32 the rule in their household that the child has to play a sport. These associations were partly
33 mediated by attitude and intention. This indicates that parents are important actors in shap-
34 ing the environmental factors of interest by making sports activities accessible and a family
35 routine. Therefore, not only adolescents, but also parents should be targeted in interven-
36 tions aiming to improve PA among adolescents. However, the cross-sectional design of this
37 study should be taken into account and the findings have to be verified in longitudinal and
38 experimental studies. Effort needs to be done to construct better measurement instruments
39 and scales to examine perceived environmental factors.

1 Acknowledgements

2
3 This study was part of CEPHIR: the Center for Effective Public Health in the larger Rot-
4 terdam Area. Financial support for this study was provided by grants from ZonMw, the
5 Netherlands Organization for Health Research and Development (grant ID no. 2100.0103)
6 and The World Cancer Research Fund - WCRF-NL (Grant number; 2007/47). This paper
7 has been facilitated by the EU funded HOPE project: “Health-promotion through Obe-
8 sity Prevention across Europe (the Commission of the European Communities, SP5A-
9 CT-2006-044128). The study does not necessarily reflect the Commission’s views and in no
10 way anticipates the Commission’s future policy in this area’.

1 REFERENCES

- 2 1. Boreham C, Riddoch C: The physical activity, fitness and health of children. *J Sports Sci* 2001,
3 19(12):915-929.
- 4 2. Wareham NJ, van Sluijs EM, Ekelund U: Physical activity and obesity prevention: a review of the
5 current evidence. *The Proceedings of the Nutrition Society* 2005, 64(2):229-247.
- 6 3. Lampert T, Mensink GB, Romahn N, Woll A: [Physical activity among children and adolescents
7 in Germany. Results of the German Health Interview and Examination Survey for Children and
8 Adolescents (KiGGS)] Körperlich-sportliche Aktivität von Kindern und Jugendlichen in Deutsch-
9 land. Ergebnisse des Kinder- und Jugendgesundheits surveys (KiGGS). *Bundesgesundheitsblatt*
10 *Gesundheitsforschung Gesundheitsschutz* 2007, 50(5-6):634-642.
- 11 4. Scully M, Dixon H, White V, Beckmann K: Dietary, physical activity and sedentary behaviour among
12 Australian secondary students in 2005. *Health Promot Int* 2007, 22(3):236-245.
- 13 5. Tammelin T, Ekelund U, Remes J, Nayha S: Physical activity and sedentary behaviors among Finnish
14 youth. *Med Sci Sports Exerc* 2007, 39(7):1067-1074.
- 15 6. Prevention CfDcA: Youth Risk Behavior Surveillance - United States, 2005. *MMWR Surveill Summ*
16 2006, 55(SS-5):1-112.
- 17 7. Telama R, Yang X, Viikari J, Valimaki I, Wanne O, Raitakari O: Physical activity from childhood to
18 adulthood: a 21-year tracking study. *Am J Prev Med* 2005, 28(3):267-273.
- 19 8. Ajzen I: The Theory of Planned Behavior. *Organizational Behavior and Human Decision Processes*
20 1991, 50:179-211.
- 21 9. Kremers SP, De Bruijn GJ, Visscher TL, Van Mechelen W, De Vries NK, Brug J: Environmental influ-
22 ences on energy balance-related behaviors: A dual-process view. *Int J Behav Nutr Phys Act* 2006,
23 3(1):9.
- 24 10. Sallis JF, Owen N: Ecological models of health behavior. In *Health behavior and health education*. 3
25 edition. Edited by Glanz K, Rimer BK, Lewis FM. San Francisco: Jossey-Bass.
- 26 11. Stokols D: Establishing and maintaining healthy environments. Toward a social ecology of health
27 promotion. *Am Psychol* 1992, 47(1):6-22.
- 28 12. Ferreira I, van der Horst K, Wendel-Vos W, Kremers S, van Lenthe F, Brug J: Environmental cor-
29 relates of physical activity in youth - A review and update. *Obesity Reviews* 2007, 8(2):129-154.
- 30 13. Sallis JF, Prochaska JJ, Taylor WC: A review of correlates of physical activity of children and adoles-
31 cents. *Med Sci Sports Exerc* 2000, 32(5):963-975.
- 32 14. Van Der Horst K, Paw MJ, Twisk JW, Van Mechelen W: A brief review on correlates of physical
33 activity and sedentariness in youth. *Med Sci Sports Exerc* 2007, 39(8):1241-1250.
- 34 15. Rhodes RE, Brown SG, McIntyre CA: Integrating the perceived neighborhood environment and the
35 theory of planned behavior when predicting walking in a Canadian adult sample. *Am J Health Promot*
36 2006, 21(2):110-118.
- 37 16. Rhodes RE, Courneya KS, Blanchard CM, Plotnikoff RC: Prediction of leisure-time walking: an
38 integration of social cognitive, perceived environmental, and personality factors. *Int J Behav Nutr*
39 *Phys Act* 2007, 4:51.
17. de Bruijn GJ, Kremers SP, Lensvelt-Mulders G, de Vries H, van Mechelen W, Brug J: Modeling in-
dividual and physical environmental factors with adolescent physical activity. *Am J Prev Med* 2006,
30(6):507-512.
18. Motl RW, Dishman RK, Saunders RP, Dowda M, Pate RR: Perceptions of physical and social environ-
ment variables and self-efficacy as correlates of self-reported physical activity among adolescent girls.
Journal of pediatric psychology 2007, 32(1):6-12.

19. Van der Horst K, Timperio A, Crawford D, Roberts R, Brug J, Oenema A: The school food environment: associations with adolescent soft drink and snack consumption. *American Journal of Preventive Medicine* in press.
20. van der Horst K, Kremers S, Ferreira I, Singh A, Oenema A, Brug J: Perceived parenting style and practices and sugar-sweetened beverage consumption in adolescents. *Health Educ Res* 2007, 22:295-304.
21. van der Horst K, Oenema A, van de Looij-Jansen P, Brug J: The ENDORSE study: research into environmental determinants of obesity related behaviors in Rotterdam schoolchildren. *BMC Public Health* 2008, 8(1):142.
22. Slootmaker SM, Schuit AJ, Chinapaw MJ, Seidell JC, van Mechelen W: Disagreement in physical activity assessed by accelerometer and self-report in subgroups of age, gender, education and weight status. *Int J Behav Nutr Phys Act* 2009, 6:17.
23. Sallis JF, Johnson ME, Calfas KJ, Caparosa S, Nichols JF: Assessing perceived physical environmental variables that may influence physical activity. *Res Q Exerc Sport* 1997, 68(4):345-351.
24. Wendel-Vos GC, Schuit AJ, Saris WH, Kromhout D: Reproducibility and relative validity of the short questionnaire to assess health-enhancing physical activity. *J Clin Epidemiol* 2003, 56(12):1163-1169.
25. Statistics Netherlands [<http://statline.cbs.nl>]
26. Mackinnon DP: *Introduction to statistical mediation analysis*: Routledge Academic; 2008.
27. Cerin E, Mackinnon DP: A commentary on current practice in mediating variable analyses in behavioural nutrition and physical activity. *Public health nutrition* 2008:1-7.
28. Twisk JWR: *Applied Multilevel Analysis: A Practical Guide*: Cambridge University Press; 2006.
29. MacKinnon DP, Lockwood CM, Brown CH, Wang W, Hoffman JM: The intermediate endpoint effect in logistic and probit regression. *Clinical trials (London, England)* 2007, 4(5):499-513.
30. Jasti S, Dudley WN, Goldwater E: SAS macros for testing statistical mediation in data with binary mediators or outcomes. *Nursing research* 2008, 57(2):118-122.
31. Baron RM, Kenny DA: The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol* 1986, 51(6):1173-1182.
32. MacKinnon DP, Krull JL, Lockwood CM: Equivalence of the mediation, confounding and suppression effect. *Prev Sci* 2000, 1(4):173-181.
33. Brug J, Kremers SP, van Lenthe F, Ball K, Crawford D: Environmental determinants of healthy eating: in need of theory and evidence. *Proceedings of the Nutrition Society* in press.
34. McGinn AP, Evenson KR, Herring AH, Huston SL, Rodriguez DA: Exploring associations between physical activity and perceived and objective measures of the built environment. *J Urban Health* 2007, 84(2):162-184.
35. Scott MM, Evenson KR, Cohen DA, Cox CE: Comparing perceived and objectively measured access to recreational facilities as predictors of physical activity in adolescent girls. *J Urban Health* 2007, 84(3):346-359.
36. Hohepa M, Scragg R, Schofield G, Kolt GS, Schaaf D: Social support for youth physical activity: Importance of siblings, parents, friends and school support across a segmented school day. *Int J Behav Nutr Phys Act* 2007, 4:54.

9 Perceived parenting style and practices and sugar-sweetened beverage consumption by adolescents

van der Horst K, Kremers S, Ferreira I, Singh A, Oenema A, Brug J.
Perceived parenting style and practices and sugar-sweetened beverage consumption by adolescents.

Health Education Research 2007, 22: 295-304.

1 ABSTRACT

2

3 **Purpose:** To investigate whether perceived parenting practices and parenting style dimen-
4 sions (strictness and involvement) are associated with adolescents' consumption of sugar-
5 sweetened beverages.

6 **Methods:** In this cross-sectional study, secondary-school students ($n = 383$, mean age 13.5)
7 completed a self-administered questionnaire on their consumption of sugar-sweetened
8 beverages, attitude, social influences, self-efficacy, habit strength, food-related parenting
9 practices, and the general parenting style dimensions of 'strictness' and 'involvement'. Data
10 were analyzed using multiple linear regression analyses.

11 **Results:** More restrictive parenting practices were associated with lower consumption of
12 sugar-sweetened beverages ($\beta = -38.0$ ml, 95% CI = $-48.1; -28.0$). This association was highly
13 mediated (about 55%) by attitude, self-efficacy, and modeling from parents. Nevertheless, a
14 significant direct effect remained ($\beta = -17.1$ ml, 95% CI = $-27.2; -6.90$). Interactions between
15 perceived parenting style and parenting practices showed that the association between
16 parenting practices and sugar-sweetened beverage consumption was stronger among ado-
17 lescents who perceived their parents as being moderately strict and highly involved.

18 **Conclusions:** Parents influence their children's sugar-sweetened beverage consumption and
19 should therefore be involved in interventions aimed at changing dietary behaviors. Inter-
20 ventions aimed at the promotion of healthy parenting practices will improve when they are
21 tailored to the general parenting style of the participants.

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

1 INTRODUCTION

2
3 The prevalence of overweight among adolescents has increased rapidly over the last decades
4 [1, 2]. Overweight is caused by an imbalance between energy intake and energy expenditure.
5 Various behaviors, such as watching television [3-5], fast food consumption [6, 7], and con-
6 sumption of large serving portions [8, 9] have been identified as determinants of a positive
7 energy balance. Additionally, consumption of sugar-sweetened soft drinks has been found
8 to be positively associated with adolescent obesity [10-13]. Effective promotion of healthful
9 eating requires a detailed understanding of the factors influencing dietary behavior. This
10 is especially important for sugar-sweetened drinks, given the large increase in soft drink
11 consumption in adolescents in recent years and the increase in soft drink consumption
12 throughout adolescence [14]. Between 1977 and 1998, the consumption of carbonated soft
13 drinks by school-aged children in the United States (aged 6-17 years) has increased from
14 5 fluid oz. (148 ml) to 12 fluid oz. (355 ml) per day [15], contributing 8% to the total daily
15 energy intake of adolescents (10.3% and 8.6% in overweight males and females respectively)
16 [16, 17]. In Dutch adolescents (aged 13-18 years), similar trends have been reported between
17 1987 and 1998, with sugar sweetened carbonated and non-carbonated soft drink consump-
18 tion increasing by 50.2% (317 ml - 476 ml) and 32.5% (212 ml - 281 ml) per day for boys and
19 girls, respectively [18].

20
21 Research indicates that parents play an important role in the eating behavior of adolescents
22 [19]. Parents influence the availability of soft drinks at home, but can also exert their in-
23 fluence through food related parenting practices [20-24]. Parenting practices are directly
24 related to specific behaviors of their children, such as the consumption of soft drinks, and
25 parents use different parenting practices for different behaviors. Studies on food-related
26 parenting practices have reported contradictory results: on the one hand, the results of
27 some studies indicate that strict parenting practices may increase children's preference
28 for (and the intake of) the restricted foods [23, 25, 26], whereas an other study suggests
29 that adolescents have a healthier diet and consume less soft drinks when they report more
30 food related rules in their family [27]. These mixed results suggest that additional factors
31 play a role. For instance Darling & Steinberg [28] postulated that parenting style modifies
32 the association between parenting practices and adolescent behavior (figure 1). Parent-
33 ing style refers to general patterns of parenting and the emotional climate in which the
34 parents' behaviors are expressed. In contrast to parenting practices, parenting style refers
35 to parent-child interactions in general, whereas parenting practices are related to specific
36 behaviors, and are reflected in things like food rules [28]. Thus, parenting practices operate
37 in the context of parenting style. Parenting styles are classified according to two dimen-
38 sions of parental behavior: 'strictness' or parental control, and 'involvement' or parental
39 warmth and acceptance [29]. Food-related parenting practices might have a different effect




Figure 9.1 Moderating contextual model of parenting style

on adolescents' behavior depending on the parenting style of their parents. Few studies have proposed this interaction effect between parenting style and parental child-feeding practices [19, 30]. Elaborating on the model of Darling & Steinberg, we aimed to examine the interactive nature of parenting style dimensions in more detail.

Cognitive variables, such as attitude, social influence, self-efficacy or habit strength may also be predictors of the consumption of sugar-sweetened beverages [21, 22]. Whether behavior-specific cognitions and habit strength can explain the link between parenting practices and adolescents' sugar-sweetened beverage consumption is not known, but most social cognitive theories assume that environmental influences are mediated by behavior-specific cognitions.

In this cross-sectional study, we investigated (a) whether behavior-specific cognitions from the Attitude, Social influence, self-Efficacy (ASE) model and habit strength are associated with sugar-sweetened beverage consumption by adolescents, (b) whether perceived parenting practices are associated with adolescent sugar-sweetened beverage consumption and whether cognitions and habit strength explain such an association, (c) possible interactions between perceived parenting style dimensions and perceived parenting practices. We combined the contextual model of parenting style [28] with the ASE model [31] and habit strength [32] to investigate sugar-sweetened beverage consumption by adolescents.

METHODS

Study population and procedure

This study was part of the Dutch Obesity Intervention in Teenagers. The medical ethical committee of the VU University Medical Center granted ethical approval for this study.

1 Data were collected at Dutch secondary schools in May and June 2003. The subjects were
 2 383 adolescents from 16 first and second grades of five secondary schools. The question-
 3 naires were completed in the classroom. No refusals to complete the questionnaire were
 4 reported. The mean age (SD; range) of the respondents was 13.5 years (0.62; 12 to 17); 211
 5 (55.1%) were female; and 55 (14.4%) were of recent immigrant origin, defined as one or both
 6 parents born abroad.

8 Measurements

9
 10 The questionnaire was based on other validated questionnaires that assessed dietary intakes
 11 and behavior-specific cognitions, habit strength and parenting variables in adolescent popu-
 12 lations [30-34]. The self-administered questionnaire was pre-tested for clarity and length,
 13 by means of cognitive interviewing among four adolescents not participating in the study.
 14

15 *Outcome measure: Sugar-sweetened beverage consumption*

16 Sugar-sweetened beverages were defined as carbonated soft drinks, other non-carbonated
 17 sugar-sweetened drinks (water-based beverages that contain sugar), and so-called sport
 18 drinks. The consumption of sugar-sweetened beverages was assessed by two questions: 'On
 19 how many days a week do you drink sugar-sweetened (not 'light' or 'diet') beverages?', with
 20 answering categories ranging from zero to seven days per week, and 'On days that you drink
 21 sugar-sweetened beverages, how many glasses, cans, and/or bottles do you drink?', with the
 22 amount to be filled in by hand. Total sugar-sweetened beverage consumption was expressed
 23 in milliliters per day and calculated from these two questions according to Dutch standard
 24 serving sizes (1 glass = 200 ml; 1 can = 330 ml; 1 bottle = 500 ml). Reported consumption of
 25 more than 4 l per day (n = 5) was recoded as 4 l.
 26

27 *Behavior-specific cognitions*

28 Cognitions specific to sugar-sweetened beverage consumption, i.e. attitude, subjective
 29 norm, social modeling, social pressure and self-efficacy were assessed according to the ASE
 30 Model [31]. All cognitions were assessed by two questions on a five-point bipolar scale. The
 31 internal consistency of the constructs was assessed using Cronbach's alpha (α). In the case
 32 of Cronbach's alpha > 0.6 , the items were combined in one scale by calculating the mean
 33 item score [35].

34 Attitude was assessed using the statements 'I think it is good to drink a lot of sugar-
 35 sweetened beverages' and 'I think it is pleasant to drink a lot of sugar-sweetened beverages'.
 36 Answering categories ranged from: 'I completely agree' (2) to 'I completely disagree' (-2)
 37 ($\alpha=0.65$). Social influences were assessed by three constructs: subjective norms, modeling
 38 and social pressure. Subjective norm was assessed by: 'My friends think that I should drink
 39 sugar-sweetened beverages' and 'My parents think that I should drink sugar-sweetened

1 beverages, with answering categories ranging from ‘Yes, definitely’ (+2) to ‘No, definitely
 2 not’ (-2) ($\alpha=0.50$). Modeling was assessed by: ‘Do your friends drink sugar-sweetened bev-
 3 erages?’ and ‘Do your parents drink sugar-sweetened beverages?’, with answering categories
 4 ranging from ‘Yes, a lot’ (+2) to ‘No, very little’ (-2) ($\alpha=0.34$). Social pressure was assessed
 5 by two questions: ‘Do your parents encourage you to drink sugar-sweetened beverages?’,
 6 and ‘Do your friends encourage you to drink sugar-sweetened beverages?’, with answering
 7 categories ranging from ‘Yes, a lot’ (+2) to ‘No, very little’ (-2) ($\alpha=0.73$). Self-efficacy was
 8 assessed by asking ‘Do you think you are able to drink less sugar-sweetened beverages?’,
 9 and ‘Does drinking less sugar-sweetened beverages seem difficult to you?’, with answering
 10 categories ranging from ‘Yes, definitely’ (+2) to ‘No, definitely not’ (-2) ($\alpha=0.72$).

11 *Habit strength*

12 We assessed habit strength of sugar-sweetened beverage consumption by means of the Self
 13 Report Habit Index [32]. This questionnaire assesses three features of habitual behavior:
 14 (1) the extent to which a behavior is automatic, e.g. ‘drinking sugar-sweetened beverages is
 15 something I do without thinking’, (2) the repeated character of the behavior, e.g. ‘drinking
 16 sugar-sweetened beverages is something I have been doing for a long time’, and (3) the sense
 17 of identity the behavior reflects, e.g. ‘drinking sugar-sweetened beverages, that’s typically
 18 “me”’. These three features were assessed by twelve questions on a five-point bipolar scale,
 19 ranging from ‘I completely agree’ (+2) to ‘I completely disagree’ (-2). An overall score for
 20 habit strength was constructed by summing the item scores ($\alpha = .88$).

21 *Perceived parenting practices and parenting style dimensions*

22 Based on the parent–child food control questionnaire developed by Cullen et al.[33], we as-
 23 sessed perceived parenting practices using nine items. Four questions (identical for fathers
 24 and mothers) assessed specific practices regarding sugar-sweetened beverage consumption
 25 (e.g. ‘My father/mother tells me *how much* sugar-sweetened beverages I am allowed to
 26 consume’, ‘My father/mother tells me *which* sugar-sweetened beverages I am allowed to
 27 consume’). An additional item assessed the availability of soft drink in the home environ-
 28 ment: ‘My mother always has my favorite sugar-sweetened beverage available at home’. All
 29 parenting items were measured on five-point bipolar scales ranging from completely agree
 30 (+2) to completely disagree (-2). A single score was computed by summing the scores on
 31 these items ($\alpha = .86$), in such a way that a higher score reflects more restrictive parenting
 32 practices.

33 We assessed two parenting style dimensions, perceived strictness and perceived involve-
 34 ment, according to den Exter and colleagues [34, 36]. Strictness was assessed by seven items,
 35 e.g.: ‘My parents know exactly where I am most afternoons after school’ and ‘At what time do
 36 you have to be at home at night on weekdays?’. Involvement was assessed by ten items, e.g.:
 37 ‘My parents make time to talk to me’ and ‘When I get a poor grade in school, my parents
 38
 39

1 encourage me to do better'. Composite scores were computed for involvement ($\alpha = .83$) and
2 strictness ($\alpha = .77$) by summing the scores on these items. Higher scores meant perceiving
3 parents as more involved or stricter. The two dimensions of strictness and involvement can
4 be used to define four parenting styles: authoritarian, authoritative, indulgent and neglect-
5 ful, by dichotomizing the scores on both dimensions. In this study, however, we used the
6 two continuous dimensions of strictness and involvement.

7 8 Data analyses

9
10 Missing data on the cognitions, habit strength and parenting variables were imputed using
11 the median value of all respondents without missing values. The missing data on sugar-
12 sweetened beverage consumption were replaced by the group mean. The highest frequency
13 of missing values was 10 (2.6%), for a parenting practice item. Multi-collinearity problems
14 were not encountered; all inter-correlations between predictors were below 0.5 [37].

15 In all conducted analyses, we used multiple linear regression analyses to examine whether
16 the associations between the determinants of interest and sugar-sweetened beverage con-
17 sumption differed with age, sex, and ethnicity. Since no significant interactions were found
18 (all $p > 0.05$), data are presented for the whole sample with adjustments for these variables
19 as potential confounders.

20 The first set of multiple linear regression analyses examined whether cognitions and habit
21 strength were associated with sugar-sweetened beverage consumption. A second set of mul-
22 tiple regression analyses investigated whether perceived parenting practices were associated
23 with sugar-sweetened beverage consumption, and the possible mediating effects of the
24 behavior-specific cognitions. It used the following requirements for establishing mediation
25 effects: (a) perceived parenting practices must be associated with sugar-sweetened beverage
26 consumption, (b) the potential mediators must be associated with sugar-sweetened beverage
27 consumption, and (c) the mediators must cause a significant reduction in the association
28 between perceived parenting practices and sugar-sweetened beverage consumption, after
29 controlling for the mediator [38]. A p-value below 0.05 was considered to be significant. A
30 Sobel test was conducted [39] to examine whether the strength of the association between
31 perceived parenting practices and sugar-sweetened beverage consumption (given by the
32 regression coefficient) decreased significantly after a potential mediator was added to the
33 model. Finally, we investigated the interaction between perceived strictness and perceived
34 parenting practices, and between perceived involvement and perceived parenting practices.
35 To this end, interaction terms between parenting practices and the strictness and involve-
36 ment dimensions were added to the regression model testing the association between
37 parenting practices and sugar-sweetened beverage consumption. If the interactions had
38 p-values below 0.1, stratified analyses were conducted for the quartiles of strictness and
39 involvement.

1 RESULTS

2
3 Table 9.1 shows the mean scores of the studied variables for boys and for girls. A significant
4 difference between boys and girls in perceived strictness, attitude, and self-efficacy was
5 found. Overall, respondents reported to perceive their parents using restrictive parenting
6 practices regarding sugar-sweetened beverage consumption and they perceived the parent-
7 ing style of their parents as low in strictness and high in involvement.

9 Behavior-specific cognitions, habit strength, and associations with sugar-sweetened 10 beverage consumption

11
12 In the first set of regression analyses (adjusting for age, gender and ethnicity), all cognitions,
13 except the social norm of friends, were significantly associated with sugar-sweetened bever-
14 age consumption (Table 9.2). When all significant cognitions were included in the regres-
15 sion model, only attitude, self-efficacy, and modeling from parents remained significantly
16 associated with sugar-sweetened beverage consumption. Habit strength was also associated
17 with sugar-sweetened beverage consumption ($\beta=35.4$, 95%CI=28.8; 42.1).

19 Mediation of the association between perceived parenting practices and sugar- 20 sweetened beverage consumption

21
22 Perceiving more restrictive parenting practices was associated with less consumption of
23 sugar-sweetened beverages (Table 9.3, model 1). Further adjustments for habit strength
24 and cognitions significantly reduced the strength of this association, which nevertheless
25 remained significant. Habit strength explained 44% of the association between perceived
26 parenting practices and sugar-sweetened beverage consumption, as can be inferred from
27 the reduction of the unstandardized regression coefficient from -38.0 ml/day to -21.3 ml/
28 day. Among the cognitions, attitude was the strongest mediator (22.7%; β -38.0 to β -29.4),
29 followed by modeling from parents (17.6%; β -38.0 to β -31.3) and self-efficacy (9.9%; β -38.0
30 to β -34.3) (Table 9.3). The cognitions, age, gender, ethnicity, and habit strength together
31 (model 6) explained 55.0% of the association between perceived parenting practices and
32 sugar-sweetened beverage consumption (β changed from -38.0 to β -17.1 ml/day).

34 Interaction between perceived parenting style dimensions and parenting practices

35
36 We further examined whether the perceived dimensions of parenting style (involvement
37 and strictness) modified the associations between parenting practices and sugar-sweetened
38 beverage consumption. P-values of interaction terms between perceived parenting prac-
39 tices and strictness ($p=0.065$), and between perceived parenting practices and involvement

Table 9.1 General characteristics of the study population.

Variable (scale range)	Mean (SD)		P-value*
	Boys (n=172)	Girls (n=211)	T-test
Age	13.4 (.542)	13.5 (.679)	NS
Ethnicity (% native)	86.6	84.8	NS
Sugar-sweetened beverage consumption (ml /day)	809 (854)	674 (677)	NS
Parenting practices (-18; 18)	-7.0 (7.25)	-7.1 (7.15)	NS
Strictness (-14; 14)	2.0 (5.42)	3.8 (4.50)	.000
Involvement (-20; 20)	10.5 (5.68)	11.1 (5.78)	NS
Habit strength (-24; 24)	-2.8 (10.2)	-3.5 (10.0)	NS
Attitude (-2; 2)	0.39 (.883)	0.12 (.904)	.003
Self-efficacy (-2; 2)	0.54 (1.19)	0.82 (1.03)	.013
Modeling from parents (-2; 2)	-0.23 (.887)	-0.34 (.950)	NS
Modeling from friends (-2; 2)	0.90 (.726)	0.90 (.654)	NS
Social norm of parents (-2; 2)	-0.20 (.942)	-0.28 (.978)	NS
Social norm of friends (-2; 2)	-0.08 (.812)	-0.13 (.779)	NS
Social pressure (-2; 2)	-1.2 (.903)	-1.3 (.724)	NS

* P-values represent the differences between genders

Table 9.2 Associations between cognitions and sugar-sweetened beverage consumption.

Cognitions	'Univariate' ^a		Multivariate ^b	
	β^c	95% CI	β^c	95% CI
Attitude	282.7 ^f	201.6; 363.7	189.3 ^f	105.8; 272.8
Self-efficacy	-204.2 ^f	-269.7; -138.7	-128.2 ^f	-194.1; -62.3
Modeling from parents	269.8 ^f	191.6; 347.9	191.2 ^f	110.6; 271.8
Modeling from friends	117.7 ^d	7.5; 228.0	-28.1	-132.7; 76.4
Social norm of parents	103.9 ^e	24.8; 183.0	-12.2	-88.7; 64.2
Social pressure	161.8 ^c	68.0; 255.5	36.1	-56.3; 128.5
Social norm of friends	17.3	-78.8; 113.4	-	-

Abbreviation: 95% CI = 95% confidence interval

^a Univariate = model adjusted for age, gender, and ethnicity.

^b Multivariate = model further adjusted for cognitions.

^c β (unstandardized coefficient) indicates the change in soft drink consumption (in ml) for a 1 unit increase in the ASE variable.

^d $p < .05$, ^e $p < .01$, ^f $p < .001$

($p = 0.069$), were below 0.1. Further stratified analyses revealed that the association between perceived parenting practices and sugar-sweetened beverage consumption varied by different quartiles of strictness and involvement (Table 9.4): parenting practices were most effective (i.e. associated with less sugar-sweetened beverage consumption) in the second and third quartiles of strictness and in the highest quartile of involvement.

Table 9.3 Mediation of the association between parenting practices and daily sugar-sweetened beverage consumption

Model	β (ml/day)	95% CI	R ²	Sobel test z-score	Percentage of the total effect that is mediated
1	-38.0 ^c	-48.1; -28.0	.154	NA	NA
2	-21.3 ^c	-31.4; -11.2	.280	-5.88 ^c	44.0
3	-29.4 ^c	-39.8; -19.0	.203	-3.97 ^c	22.7
4	-34.3 ^c	-44.0; -24.5	.217	-2.44 ^a	9.9
5	-31.3 ^c	-41.3; -21.3	.241	-3.63 ^c	17.6
6	-17.1 ^b	-27.2; -6.90	.325	NA	55.0

Abbreviation: NA, not applicable

Model 1: adjusted for age, gender, and ethnicity

Model 2: model 1 + adjusted for habit strength

Model 3: model 1 + adjusted for attitude

Model 4: model 1 + adjusted for self-efficacy

Model 5: model 1 + adjusted for modeling from parents

Model 6: model 1 + adjusted for habit strength, attitude, self-efficacy, and modeling from parents

^a $p < .05$, ^b $p < .01$, ^c $p < .001$

Table 9.4 Associations between parenting practices and sugar-sweetened beverage consumption, stratified by the quartiles of strictness and involvement.

Quartiles ^a	Strictness			Involvement		
	β^b (ml/day)	95% CI	R ²	β^b (ml/day)	95% CI	R ²
1 (lowest)	-11	-36; 14	.500	-9	-30; 13	.397
2	-27 ^c	-49; -5	.404	-18	-40; 3	.395
3	-35 ^d	-55; -14	.319	-15	-38; 8	.325
4 (highest)	-15	-32; 3	.182	-28 ^d	-46; -10	.433

^a Ranges for strictness (-14; 14) per quartile: (1) -14;-1, (2) 0;3, (3) 4;6, (4) 7;14, Ranges for involvement (-20; 20) per quartile: (1) -12;7, (2) 8;11, (3) 12;15, (4) 16;20

^b Unstandardized beta, adjusted for age, sex, ethnicity, habit strength, attitude, modeling from parents, and self-efficacy

^c $p < .05$

^d $p < .01$

DISCUSSION

The present study investigated the association between perceived parenting practices and sugar-sweetened beverage consumption by adolescents on the one hand, and the influence of perceived parenting style dimensions on this association on the other. Perceiving more restrictive parenting practices was found to be associated with less soft drink consumption, which is in agreement with the findings of the study by De Bourdeaudhuij & Van Oost (2000). [27]. However, findings from studies among younger children suggest that strict

1 parental practices can in fact increase children's preferences for, and intake of, the restricted
2 foods [23, 25, 26]. These contrasting outcomes may have been caused by the difference
3 between parenting practices that are used in childhood and in adolescence. For instance,
4 parents might use pressure to get their young children to eat, or may restrict access to foods.
5 For adolescents, parents might use clearly defined rules about the times when a certain food
6 can be eaten and how much of a certain food they can eat.

7
8 In our study, the model with age, gender, ethnicity, habit strength, modeling from parents,
9 attitude, and self-efficacy explained 55% of the relationship between perceived parenting
10 practices and sugar-sweetened beverage consumption, the largest contribution being
11 that by habit strength (44%). Nevertheless, perceived parenting practices had a direct
12 association with sugar-sweetened beverage consumption unmediated by cognitions and
13 habit strength. Sugar-sweetened beverage consumption may thus not always be reasoned
14 behavior, and this finding has some theoretically important implications. Potential distal
15 determinants of intakes in the social, cultural or physical environment may increase our
16 understanding of sugar-sweetened beverage consumption in adolescents. In addition, since
17 perceived parenting practices were still independently associated with the consumption of
18 sugar-sweetened beverages, other factors may also be involved in the association between
19 perceived parenting practices and sugar-sweetened beverage consumption by adolescents,
20 for instance the influence of taste preferences [20]. Parents shape their children's eating
21 environment in different ways. Parental feeding practices in early childhood, for instance
22 using foods as a reward or to comfort [40], exposure to foods [41], and parental control
23 of how much and what children eat [42] can influence a child's taste preferences which
24 may persist into adolescence. In addition, the amount and diversity of sugar-sweetened
25 beverages parents make available and accessible at home can influence the amount of such
26 beverages adolescents consume [15, 20]. It has been suggested that the exposure to soft drink
27 advertising may lead to a higher consumption of soft drinks during TV watching [20, 43],
28 an activity which constitutes a considerable part of adolescents' leisure time activity. Finally,
29 the availability of soft drink vending machines in the adolescents' immediate environment
30 (e.g. schools) could also contribute to a higher consumption of soft drinks [15, 20, 44].

31
32 We also explored whether the association between perceived parenting practices (specific
33 rules about sugar-sweetened beverage consumption) and sugar-sweetened beverage con-
34 sumption by adolescents was different depending on the perceived parenting style of their
35 parents. We therefore examined if the dimensions of parenting style strictness (parental
36 control) and involvement (parental warmth and acceptance) modified the association be-
37 tween parenting practices and the consumption of such beverages. The results indicated that
38 the effect of parenting practices in this respect was most pronounced in those families with
39 a highly involved or moderately strict parenting style. Kremers et al. (2003) also found that

1 adolescents raised in a family with authoritative parenting style (highly strict and highly
2 involved) showed the most favorable consumption of fruits and vegetables [30]. In the high-
3 est quartile of strictness, we found no significant association between perceived parenting
4 practices and sugar-sweetened beverage consumption. This indicates that if parents use a
5 very strict parenting style, parenting practices relating to sugar-sweetened beverage con-
6 sumption may not have an additional direct limiting effect on their children's consumption
7 of these beverages. The strongest association between perceived parenting practices and
8 sugar-sweetened beverage consumption was found in the highest quartile of involvement
9 indicating a stronger direct effect of parenting practices on adolescent behavior, in the case
10 of involved parents.

11
12 The results of our study and that by Kremers et al. (2003) are not entirely comparable. We
13 used the continuous measures of the two perceived parenting style dimensions instead of
14 the four categories of parenting style. There were several reasons why we chose to use the
15 two dimensions of strictness and involvement instead of the four parenting style categories.
16 First, the skewed distribution on these dimensions would have caused misclassification
17 when dichotomized into categories. Second, dichotomization itself is quite arbitrary since
18 there are no 'universal' cut-off values for those dimensions and the cut-offs will therefore
19 vary for different populations. Third, classification of two continuous variables into four
20 categorical parenting styles means loss of information. Since we were interested in the role
21 of various types and dimensions of parenting we chose to include the two dimensions as
22 continuous variables and investigate the existence of interactions.

23
24 To our knowledge, this is the first study to indicate the role of parenting styles as an environ-
25 mental context factor that can influence the effectiveness of food-related parenting practices
26 in terms of adolescents' consumption of sugar-sweetened beverages. As such, it contributes
27 to theory development of the influence of parents on adolescents' dietary behaviors. In
28 contrast to assumptions that underlie theories such as the Theory of Planned Behavior
29 [45] the results indicate that sugar-sweetened beverage consumption may not always be
30 reasoned or planned [46]. Additionally, contextual variables may moderate the associations
31 between cognitive variables and intake levels. Notably, Bandura's Social Cognitive Theory
32 [47] does include the reciprocal interaction of person, environment and behavior. Such a
33 theoretical approach may guide future research aimed at examining potential moderators of
34 the environment – behavior relationship.

35 There are several limitations to the interpretation of the results of this study. Since the
36 design of this study was cross-sectional, inferences regarding cause and effect must be made
37 with caution and will not be conclusive. Parenting practices could determine, but also be a
38 result of children's behavior regarding sugar-sweetened beverage consumption (and indeed
39 that of other food items). Another limitation is that the schools and classes included in this

1 study were not randomly selected. In addition, the study population included few children
2 from the various ethnic minorities in the Netherlands, which made it impossible to examine
3 the potential role ethnic background in sugar-sweetened beverage consumption. We used
4 adolescents' reports of parenting practices and parenting style dimensions. Therefore, it
5 could be that parental reports of practices and style would differ from their children's per-
6 ceptions. In addition, what adolescents perceive as 'strict' and 'involved' may vary among
7 individuals. Obtaining data from multiple sources (adolescents, parents and siblings) would
8 probably result in data that are more valid. Finally, the assessment of sugar-sweetened bev-
9 erage consumption relied on self-report and was not supported by any objective assessment.
10 Assessment of validity and reliability data were not available for this measure. Although
11 the measurement instrument used in this study was designed to be as clear as possible, it
12 is not known whether adolescents under-reported or over-reported their sugar-sweetened
13 beverage consumption. Validation studies on measures of sugar-sweetened beverage con-
14 sumption are clearly needed, and might be undertaken as part of future research to improve
15 the assessment of this behavior.

16
17 Despite these limitations, our findings emphasize the importance of parental rules and
18 the interaction between these rules and parenting style dimensions for sugar-sweetened
19 beverage consumption by adolescents. The central role parents can play on the primary
20 prevention of obesity-related behaviors was clearly illustrated: a one 'unit' decrease on the
21 parenting practice scale accounted for an increase of 38 ml per day in sugar-sweetened
22 beverage consumption. Small increases in energy intake, not accompanied by concomitant
23 increasing energy expenditure, will substantially contribute to weight gain.

24 Parents are thus important intermediates for changing dietary behaviors of adolescents
25 and should therefore be involved in interventions aimed at changing dietary behaviors, and
26 reducing overweight. The present study showed that interventions aimed at the promotion
27 of healthy parenting practices will improve when they are tailored to the general parenting
28 style of the participants.

1 REFERENCES

- 2 1. Hedley AA, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM: Prevalence of overweight
3 and obesity among US children, adolescents, and adults, 1999-2002. *JAMA* 2004, 291(23):2847-2850.
- 4 2. Lobstein T, Frelut ML: Prevalence of overweight among children in Europe. *Obes Rev* 2003, 4(4):195-
5 200.
- 6 3. Andersen RE, Crespo CJ, Bartlett SJ, Cheskin LJ, Pratt M: Relationship of physical activity and televi-
7 sion watching with body weight and level of fatness among children: results from the Third National
8 Health and Nutrition Examination Survey. *JAMA* 1998, 279(12):938-942.
- 9 4. Eisenmann JC, Bartee RT, Wang MQ: Physical activity, TV viewing, and weight in U.S. youth: 1999
10 Youth Risk Behavior Survey. *Obes Res* 2002, 10(5):379-385.
- 11 5. Berkey CS, Rockett HR, Field AE, Gillman MW, Frazier AL, Camargo CA, Jr., Colditz GA: Activity,
12 dietary intake, and weight changes in a longitudinal study of preadolescent and adolescent boys and
13 girls. *Pediatrics* 2000, 105(4):E56.
- 14 6. Diet, nutrition and the prevention of chronic diseases. *World Health Organ Tech Rep Ser* 2003, 916:i-
15 viii, 1-149, backcover.
- 16 7. Levitsky DA, Halbmaier CA, Mrdjenovic G: The freshman weight gain: a model for the study of the
17 epidemic of obesity. *Int J Obes Relat Metab Disord* 2004:1-8.
- 18 8. Kral TV, Rolls BJ: Energy density and portion size: their independent and combined effects on energy
19 intake. *Physiol Behav* 2004, 82(1):131-138.
- 20 9. McConahy KL, Smiciklas-Wright H, Mitchell DC, Picciano MF: Portion size of common foods
21 predicts energy intake among preschool-aged children. *J Am Diet Assoc* 2004, 104(6):975-979.
- 22 10. Mrdjenovic G, Levitsky DA: Nutritional and energetic consequences of sweetened drink consump-
23 tion in 6- to 13-year-old children. *J Pediatr* 2003, 142(6):604-610.
- 24 11. Ludwig DS, Peterson KE, Gortmaker SL: Relation between consumption of sugar-sweetened drinks
25 and childhood obesity: a prospective, observational analysis. *Lancet* 2001, 357(9255):505-508.
- 26 12. James J, Thomas P, Cavan D, Kerr D: Preventing childhood obesity by reducing consumption of
27 carbonated drinks: cluster randomised controlled trial. *Bmj* 2004, 328(7450):1237.
- 28 13. Giammattei J, Blix G, Marshak HH, Wollitzer AO, Pettitt DJ: Television watching and soft drink
29 consumption: associations with obesity in 11- to 13-year-old schoolchildren. *Arch Pediatr Adolesc Med*
30 2003, 157(9):882-886.
- 31 14. Rampersaud GC, Bailey LB, Kauwell GP: National survey beverage consumption data for children
32 and adolescents indicate the need to encourage a shift toward more nutritive beverages. *J Am Diet*
33 *Assoc* 2003, 103(1):97-100.
- 34 15. French SA, Lin BH, Guthrie JF: National trends in soft drink consumption among children and
35 adolescents age 6 to 17 years: prevalence, amounts, and sources, 1977/1978 to 1994/1998. *J Am Diet*
36 *Assoc* 2003, 103(10):1326-1331.
- 37 16. Troiano RP, Briefel RR, Carroll MD, Bialostosky K: Energy and fat intakes of children and adolescents
38 in the united states: data from the national health and nutrition examination surveys. *Am J Clin Nutr*
39 2000, 72(5 Suppl):1343S-1353S.
17. Briefel RR, Johnson CL: Secular trends in dietary intake in the United States. 2004, 24:401-431.
18. Gezondheidsraad: Enkele belangrijke ontwikkelingen in de voedselconsumptie. Den Haag: Gezond-
heidsraad. Commissie Trends Voedselconsumptie; 2002.
19. Golan M, Crow S: Parents are key players in the prevention and treatment of weight-related problems.
Nutr Rev 2004, 62(1):39-50.

20. Grimm GC, Harnack L, Story M: Factors associated with soft drink consumption in school-aged children. *J Am Diet Assoc* 2004, 104(8):1244-1249.
21. Kassem NO, Lee JW: Understanding soft drink consumption among male adolescents using the theory of planned behavior. *J Behav Med* 2004, 27(3):273-296.
22. Kassem NO, Lee JW, Modeste NN, Johnston PK: Understanding soft drink consumption among female adolescents using the Theory of Planned Behavior. *Health Educ Res* 2003, 18(3):278-291.
23. Birch LL, Fisher JO: Mothers' child-feeding practices influence daughters' eating and weight. *Am J Clin Nutr* 2000, 71(5):1054-1061.
24. Robinson TN, Kiernan M, Matheson DM, Haydel KF: Is parental control over children's eating associated with childhood obesity? Results from a population-based sample of third graders. *Obes Res* 2001, 9(5):306-312.
25. Fisher JO, Birch LL: Restricting access to palatable foods affects children's behavioral response, food selection, and intake. *Am J Clin Nutr* 1999, 69(6):1264-1272.
26. Brown R, Ogdan J: Children's eating attitudes and behaviour: a study of the modelling and control theories of parental influence. *Health Educ Res* 2004, 19(3):261-271.
27. De Bourdeaudhuij I, Van Oost, P.: Personal and family determinants of dietary behaviour in adolescents and their parents. *Psychology and Health* 2000, 15:751-770.
28. Darling N, Steinberg L: Parenting style as context: An integrative model. *Psychological Bulletin* 1993, 113(3):487-496.
29. Maccoby EE, Martin JA: Socialization in the context of the family: parent-child interaction. In *Handbook of Child Psychology Personality and Social Development. Volume 4*. Edited by Hetherington EM. New York: Wiley; 1983:1-110.
30. Kremers SP, Brug J, de Vries H, Engels RC: Parenting style and adolescent fruit consumption. *Appetite* 2003, 41(1):43-50.
31. de Vries H, Dijkstra M, Kuhlman P: Self-efficacy: The third factor besides attitude and subjective norm as a predictor of behavioural intentions. *Health Educ Res* 1988, 3(3):273-282.
32. Verplanken B, Orbell S: Reflections on past behavior: A self-report index of habit strength. *J Appl Soc Psychol* 2003, 33(6):1313-1330.
33. Cullen KW, Baranowski T, Rittenberry L, Cosart C, Hebert D, de Moor C: Child-reported family and peer influences on fruit, juice and vegetable consumption: reliability and validity of measures. *Health Educ Res* 2001, 16(2):187-200.
34. Steinberg L, Elmen JD, Mounts NS: Authoritative parenting, psychosocial maturity, and academic success among adolescents. *Child Dev* 1989, 60(6):1424-1436.
35. Nunnally JC: *Psychometric theory*. 2 edition. New York: McGraw Hill; 1978.
36. den Exter Blokland EAW, Engels RCME, Finkenauer C: Parenting styles, self-control and male juvenile delinquency: The mediation role of self-control. In *Prevention and control of aggression and the impacts on its victims*. Edited by Martinez M. Dordrecht/New York: Kluwer/Plenum Press; 2001:201-207.
37. Kleinbaum DG, Kupper LL, Muller KE: *Applied regression analysis and other multivariable methods*. Boston: PWS-KENT Publishing Company; 1978.
38. Baron RM, Kenny DA: The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol* 1986, 51(6):1173-1182.
39. MacKinnon DP, Dwyer JH: Estimating mediated effects in prevention studies. *Evaluation Review*, 17(2):144-158.
40. Birch LL: Psychological influences on the childhood diet. *J Nutr* 1998, 128(2 Suppl):407S-410S.

- 1 41. Liem DG, de Graaf C: Sweet and sour preferences in young children and adults: role of repeated
2 exposure. *Physiol Behav* 2004, 83(3):421-429.
- 3 42. Fisher JO, Mitchell DC, Smiciklas-Wright H, Birch LL: Parental influences on young girls' fruit and
4 vegetable, micronutrient, and fat intakes. *J Am Diet Assoc* 2002, 102(1):58-64.
- 5 43. Van Den Bulck J, Van Mierlo J: Energy intake associated with television viewing in adolescents, a
6 cross sectional study. *Appetite* 2004, 43(2):181-184.
- 7 44. Vereecken CA, Bobelijm K, Maes L: School food policy at primary and secondary schools in Belgium-
8 Flanders: does it influence young people's food habits? *Eur J Clin Nutr* 2005, 59(2):271-277.
- 9 45. Ajzen I: *Attitudes, personality, and behavior*: Homewood, IL, US: Dorsey Press; 1988.
- 10 46. Kremers SP, De Bruijn GJ, Visscher TL, Van Mechelen W, De Vries NK, Brug J: Environmental influ-
11 ences on energy balance-related behaviors: A dual-process view. *Int J Behav Nutr Phys Act* 2006,
12 3(1):9.
- 13 47. Bandura A: *Social foundations of thought and action: a social cognitive theory*.: Englewood Cliffs, NJ:
14 Prentice-Hall; 1986.
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39

10 General Discussion

1 ABSTRACT

2

3 The studies described in this thesis aimed at gaining insight into important individual and
4 environmental correlates of energy balance-related behaviors among adolescents. The aims
5 of this thesis were based on the EnRG framework [1] and were to (I) examine individual
6 and environmental correlates of energy balance-related behaviors and (II) explore to what
7 extent the association between environmental factors and energy balance-related behaviors
8 is mediated by individual cognitions. In this chapter the main findings of this thesis are
9 summarized. Next, methodological issues are discussed, followed by an integration of
10 findings. Finally implications for research and practice will be formulated to inform future
11 research and intervention development.

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

10.1 MAIN FINDINGS

Environmental correlates of energy balance-related behaviors: reviews of the literature

Two systematic reviews (chapters 3 and 4) were conducted to get an overview of the existing evidence-base regarding important environmental correlates of energy balance-related behaviors and to inform the development of measurement instruments. The ANGELO framework was used in these reviews to categorize the environmental factors [2]. The review on dietary behaviors revealed that consistent evidence was found for associations between parent and sibling intakes with adolescent's energy and fat intake, and between parents' educational level and adolescent's fruit and vegetable intake. In the review on adolescents' physical activity behaviors, support from significant others, mother's education level, family income, non-vocational school attendance, and low crime incidence were found to be associated with higher physical activity. Several gaps in the available evidence of associations between environmental factors and energy balance-related behaviors were identified, such as the lack of high-quality studies and study replications. Many potentially relevant environmental factors have not been studied at all [3] and the available research focused mainly on factors in the home environment such as parental influences. Only a limited number of studies assessing environmental correlates of snack and soft drink intakes were retrieved, while these two behaviors may be of specific importance in obesity prevention [4, 5].

Demographic correlates of energy balance-related behaviors

The second part of this thesis reports on two studies in which associations between demographic correlates and overweight and energy balance-related behaviors were explored. Chapter 5 reports on gender, ethnic and educational differences in overweight and energy balance-related behaviors. The study findings indicate important ethnic and educational differences in overweight and energy balance-related behaviors. In line with previous findings, youth from non-Western ethnic backgrounds, especially girls and those attending vocational schools were more likely to be overweight and to engage in unfavorable energy balance-related behaviors [6, 7].

Chapter 6 reports on adolescents' active commuting to school and in this chapter we explored potential socio-demographic correlates of active commuting to school. Almost half of the adolescents reported to actively commute to school. Adolescents of non-Western ethnic background were more likely to be walkers and non-active commuters than cyclists compared to native Dutch adolescents. A further distance from home to school was inversely associated with being a walker or cyclist and positively associated with being a non-active commuter.

1 Individual and environmental correlates of energy balance-related behavior

2
3 The final part of this thesis explored possible mediating effects of individual cognitions.
4 According to the EnRG framework [1] we examined the direct association of environmental
5 correlates on energy balance-related behaviors, but also the suggested mediating effect of
6 individual cognitions between environmental factors and energy balance-related behaviors.

7 In chapter 7 associations of the availability of foods/drinks in school canteens, the pres-
8 ence of food stores around schools and individual cognitions with soft drink and snack
9 consumption were examined. Mediation of the environment – behavior relationship by
10 individual cognitions was also examined. This study indicated that individual cognitions
11 appeared to be stronger correlates of intakes than physical school environmental factors
12 and little evidence for associations of environmental factors in the school environment with
13 soft drink and snack consumption was found. There was an inverse association between dis-
14 tance to the nearest store and the number of small food stores with soft drink consumption
15 and these associations were partly mediated by cognitions. Therefore, this study provides
16 some evidence for the hypothesis that environmental factors are associated with soft drink
17 consumption via the cognitions, as proposed in the EnRG framework.

18 The possible mediation effect of cognitions was also examined in chapter eight and in this
19 study significant mediation effects were also found for the individual cognitions. A direct
20 association of environmental factors on sports participation was found for availability of
21 physical activity attributes at home, parents sports behavior and parental rule about sports
22 participation. These associations were partly mediated by attitude and intention. This study
23 provided evidence for the indirect effect of environmental factors on adolescent sports
24 behavior

25 In chapter 9, we used another dataset to investigate the possible mediating role of cogni-
26 tions in further detail. In this study we examined whether cognitions and habit strength
27 mediated the association between parenting practices and sugar-sweetened beverage
28 intake and possible moderating effects of the parenting style dimensions “strictness” and
29 “involvement”, because Darling and Steinberg [8] postulated that parenting style modifies
30 the association between parenting practices and adolescent behavior. The results of this
31 study indicated that more restrictive parenting practices were associated with lower sugar-
32 sweetened beverage intake. Mediation was found for the cognitions attitude, self-efficacy
33 and modeling. Nevertheless, a significant direct association between environmental par-
34 enting practices and intake also remained. A possible moderating effect of parenting style
35 was found, showing that the association between parenting practices and sugar-sweetened
36 beverage consumption was stronger among adolescents who perceived their parents as be-
37 ing moderately strict and highly involved.

10.2 METHODOLOGICAL ISSUES

The studies presented in this thesis have several limitations and the results and conclusions should be interpreted in the light of these limitations. In this section, considerations concerning the study design, sampling and subjects, the assessment of energy balance-related behaviors and individual and environmental factors are discussed.

Cross sectional study design

The studies in this thesis were based on the baseline data collection from the ENDORSE project as the longitudinal data collection took place in 2007/2008 and was not yet available. Therefore, causal conclusions cannot be drawn and caution is needed interpreting the findings reported in the studies. For instance, we found parental influences, such as parenting practices to have an important association with energy balance-related behaviors. However parenting practices could also be a result of the adolescents' behavior or weight status [9, 10]. Reciprocal determinism, where the causal relationships are bi-directional also exists for the associations found between perceptions of the environment and physical activity behavior. Physically active adolescent might be more aware of physical activity equipment in their environment. This makes discussion of traditional "causal" pathways more complex [11].

The cross sectional design of the study also made the investigation of mediating effects more difficult as mediation effects refer to causal mechanisms. We used the method suggested by Baron & Kenny [12, 13] which specifies a series of tests of the links in a causal chain to investigate mediation effects in a cross sectional design. Despite its widespread use, the Baron-Kenny method has some limitations. For instance, it does not provide a direct estimate of the size of the indirect (mediated) effect, and the Baron-Kenny approach has low statistical power in studies with a relatively small sample size, even in the presence of large mediation effects [14, 15]. Nevertheless, the Baron-Kenny approach is useful in understanding mediating effects of environmental factors on energy balance-related behaviors as proposed in the EnRG framework because it specifically tests the direct and indirect associations of environmental factors on energy balance-related behavior. The mediation analyses that were conducted for this thesis should preferably be repeated with a longitudinal design. A cross-sectional approach to mediation can generate substantially biased estimates even under the ideal conditions when mediation is complete [16]. However, a cross-sectional study design is efficient for exploring and identifying environment – behavior associations in a relatively new research field. With the longitudinal data from the ENDORSE project, the results from various studies can be verified and further examined.

1 Sampling and subjects

2

3 *City*

4 The data collection process for the ENDORSE study took place in close cooperation with
5 the Municipal Health Service Rotterdam area and was an integral part of their ongoing
6 health surveillance, the Youth Monitor Rotterdam. Therefore, all participating schools were
7 located in Rotterdam. Rotterdam is a city with a relatively high proportion of people from
8 lower socio-economic positions and foreign ethnicity compared to the rest of the Nether-
9 lands. The results can thus not be generalized. The study described in chapter 9 was based
10 on another dataset. This study was part of the Dutch Obesity Intervention in Teenagers [17].
11 Data was collected at Dutch secondary schools in May and June 2003. These schools were
12 located in the eastern part of the Netherlands in smaller cities compared to Rotterdam. The
13 subjects were 383 adolescents from 16 first and second grades of five secondary schools.
14 The ethnic background of this study population was representative for the Dutch general
15 adolescent population.

16

17 *Schools*

18 The study results must also be interpreted with the possibility of selection bias at the school
19 level. Only schools that already participated in the Youth Monitor Rotterdam were con-
20 tacted to participate in the study. Of these 54 schools, 24 were interested in participating.
21 These schools might be more motivated and already involved in promoting healthy lifestyles
22 than the other schools in Rotterdam. Subsequently, a random sample of 17 school locations
23 was drawn from the pool of schools that were willing to participate, after stratification of the
24 schools according to four city areas in which they were located. Stratification was done, to
25 ensure variation in physical and cultural environments. No schools located in the western
26 part of the city with more deprived neighborhoods were willing to participate in the study.

27

28 *Students*

29 No inclusion criteria were defined for participating in the ENDORSE study. Per school ap-
30 proximately five classes were selected to participate in the study. All adolescents in one class
31 participated in the study, unless they or their parents indicated that they were not willing to
32 participate or if they were absent on the day of the assessment. We did not obtain data from
33 the adolescents that refused to participate, and thus we were not able to determine to what
34 extent selection bias at this level may have occurred. However it is likely that overweight
35 and obese adolescents were less likely to participate in the anthropometrical measurements.
36 There were no significant differences in the number of missing values on weight status for
37 gender, ethnicity and educational level.

38

39

1 *Parents*

2 To study the effects of parental behavior and parental rules on sports behavior (chapter
3 8) we used data from the parent questionnaire. However, the response rate for the parent
4 questionnaire was quite low. Within the sample of full parent-adolescent combinations
5 respondents from non-Western ethnic background were underrepresented, contrary to
6 the other studies in which the full adolescent sample was used. This under-representation
7 might have occurred because all materials were in the Dutch language. As the parents were
8 mostly first generation immigrants, language problems could have been the main problem
9 for answering and returning the questionnaire.

11 Measurement of energy balance-related behaviors

13 *Validity and reliability*

14 Measuring dietary intake and physical activity behaviors by self-reports is a major problem
15 as the assessment relies on a child's recall of behaviors. No really valid questionnaires are
16 available and the use of these self-reported measures of energy balance-related behaviors
17 could have caused an overestimation or underestimation of behaviors. Therefore, the qual-
18 ity of the measurement instrument should be taken into account in evaluating the results
19 of this study.

20 Food frequency questions to assess food intake are often used for epidemiological studies
21 since they are relatively easy to administer and less expensive than other methods such
22 as 24-hour recall and food records, but food frequency questionnaires tend to over- and
23 underestimate energy and nutrient intakes [18]. A validation study to test the validity of the
24 food frequency questions used in this study was not undertaken. However validation meth-
25 ods are also imperfect, since a gold standard does not exist for assessing dietary behaviors
26 and developing a good food frequency questionnaire is therefore very difficult. The dietary
27 intake measures used in the ENDORSE project were developed by adapting validated Dutch
28 questionnaires on dietary intake to the ENDORSE study population and dietary behaviors
29 [17, 19, 20]. The test-retest reliability for the dietary intake variables was considered to be
30 reasonably good (soft drink $r=0.59$, breakfast consumption $r=0.65$; sweet & savory snack
31 consumption $r=0.46$).

32 Also the assessment of physical activity and sedentary behaviors in adolescents is dif-
33 ficult. In our study we used the Activity Questionnaire for Adults & Adolescents (AQuAA)
34 [21], which is a short questionnaire to assess physical activity at school and during leisure
35 time, active transportation to school and sedentary behaviors in leisure time. The AQuAA
36 refers to activities in the past week (7-day recall). The test-retest reproducibility was fair to
37 moderate for this questionnaire; with intra-class correlations ranging from 0.46 to 0.59. The
38 validation study with use of accelerometers showed that adolescents over-reported their
39 activity levels and validity of this measure was low [21]. Examination of the self-reported

1 amounts of time adolescents spent on the various physical activity and sedentary behav-
2 iors in the ENDORSE dataset made clear that the overestimation of physical activity and
3 sedentary behaviors was also present in the ENDORSE data. The average minutes per day
4 spent on most of the behaviors was highly skewed, with some adolescents reporting very
5 high amounts of physical activity and sedentary behaviors. We did not use more objective
6 measurements such as accelerometers. Accelerometer data give an indication of overall
7 physical activity and for the ENDORSE study we investigated various physical activity
8 sub-behaviors such as walking in leisure time and active commuting to school. Another
9 important weakness of accelerometers is that they are insensitive for many forms of activity,
10 including cycling [22].

11 In spite of the limitations discussed, questionnaires are often used to assess behavior and
12 they are considered to be easy and inexpensive in use. However, the quality of the mea-
13 surement instruments should be taken into account in evaluating the results of this study.
14 The development of valid and reliable questionnaires to examine energy balance-related
15 behaviors needs more attention in future research.

16 *Context specific assessment of behaviors*

17 As described in the ANGELO framework, energy balance-related behaviors can occur in a
18 wide range of behavioral settings, such as homes, schools, restaurants, and neighborhoods
19 [2, 23, 24]. People may behave differently in different settings and it is important to incor-
20 porate the behavioral setting in the assessment of energy balance-related behaviors. For
21 example, physical environmental correlates are likely to be different for physical activity at
22 school and physical activity at home [24]. Research has begun to focus on specific behaviors
23 such as walking to school or walking for recreation instead of a generic measure of walk-
24 ing, but still most research focuses on context-free behavioral outcomes. This is also the
25 case for the ENDORSE study as in chapter 7 environmental correlates were assessed at the
26 school level, but the behaviors of interest, soft drink and snack consumption were assessed
27 in general measures as the average intake per day. If adolescents consume most soft drinks
28 and snacks in other settings than at school, for instance at home, this might explain why no
29 associations were found in this study.

30 The context specific approach may underestimate the association between environmental
31 and behavioral variables [25]. However a context-specific approach to the assessment of
32 energy balance-related behaviors does have its own limitations. Questionnaires will be too
33 long and give insight in determinants of behaviors in specific situations while the results of
34 this kind of research is often used to inform interventions aimed at changing the behavior
35 regardless the context in which the behavior takes place.

1 Measurement of individual and environmental correlates

2
3 The following measurement instruments were used in the ENDORSE project: adolescent
4 and parent questionnaires, interviews with school representatives and canteen managers,
5 audits of the school environment, census data collection and adolescent body measure-
6 ments. In this paragraph, the use of questionnaires, the audit instrument and information
7 from Geographic Information Systems (GIS) to assess individual and environmental cor-
8 relates will be discussed.

9 10 *Assessment of individual correlates*

11 The assessment of individual correlates of energy balance-related behaviors was based on
12 the Theory of Planned Behavior (TPB). According to the TPB the assessment of cogni-
13 tions should be action, target, time and context specific [26, 27]. The questions used in the
14 ENDORSE study did not meet all of these criteria. For instance the questions on sports and
15 physical activity were specific on action (sports and physical activity) and the context (in
16 leisure time), but did not address the target and time in the question. Similar to measure-
17 ments of behavior, pursuing such specificity would have made the questionnaire even longer,
18 making administration within one school hour impossible. Besides, it would have made the
19 questions too complicated and long for the study population and therefore could have led
20 to incomplete or unreliable data. Therefore, we also assessed the cognitions through direct
21 measures (good-bad, pleasant-unpleasant, easy-difficult) instead of underlying beliefs [28].

22 Some attempts to test the reliability and validity of the TPB questions were undertaken.
23 The questions were formulated as much as possible according to generally accepted in-
24 structions provided from the TPB. In the developing phase, we pre-tested the adolescent
25 questionnaire among ten adolescents by means of cognitive interviewing to examine the
26 questions on clarity and comprehensibility. Subsequently, the questionnaire was completed
27 twice by 89 schoolchildren (aged 13–14) ten-days apart to assess the test-re-test reliability
28 and other psychometrics of the questionnaire. Items with low reliability were adjusted or
29 deleted from the questionnaire.

30 31 *Assessment of environmental correlates*

32 In the ENDORSE project, environmental correlates were assessed with questionnaires, by
33 audits of the schools and objective Geographic Information System (GIS) data. Potential
34 important environmental correlates were categorized according to the ANGELO frame-
35 work. In the ENDORSE study physical, socio-cultural, economic and policy correlates were
36 examined at the micro level (home, school and neighborhood level), and combinations
37 of perceived and objectively measured environmental correlates were used. Assessing
38 environmental correlates is difficult as there is not much known about which correlates are
39 important and how these correlates need to be measured. As it is impossible to assess all

1 potential important environmental factors in one study, several considerations should be
2 taken into account. First, one should decide to what extent the complexity of the environ-
3 ment could be assessed. Second, one can assess environmental factors in a subjective way
4 with questionnaires or more objectively with an audit instrument or census data. However
5 the development of a valid audit instrument is difficult because of the limited available
6 literature and instruments. Third, when we speak of assessing environmental factors in the
7 neighborhood around school or homes, it is important to define what is considered as a
8 neighborhood. These points will be further explained in the following paragraphs.

9 10 Complexity of environmental factors 11

12 A problem in environmental research is which aspects of the environment we should focus
13 on. Most research has primarily focused on the availability of facilities and the attractiveness
14 of neighborhoods. However, other or more complex measures of the environment might be
15 necessary to assess environmental influences on behavior [24]. For example, in chapter 7 we
16 focused on the school food environment using crude measures such as counting the avail-
17 ability of food products, the number of food shops and the distance to the nearest food shop.
18 However, the nearest food shop might be desirable in that it is nearby school, but could have
19 also undesirable characteristics for adolescents we did not assess, such as high prices, low
20 availability of preferred foods and drinks, and inconvenient opening hours, so that the shop
21 is never used. In chapter 8 crude measures of availability of physical activity equipment and
22 facilities were used, while the quality of the facilities might also be important. These more
23 complex aspects of environmental factors were not taken into account in this study.

24 25 Perceived versus objectively measured environmental factors 26

27 The development of objective measures of environmental factors is an important direc-
28 tion for research, as well as studies that compare self-reported perceptions with objective
29 environmental factors. Studies that report the use of objective environmental measures to
30 assess the food environment are limited. Therefore, we used objective measures to assess
31 the availability of products in the school canteens and the distance and number of shops in
32 the school neighborhood (Chapter 7). However, the perceived availability of products might
33 also be important in explaining dietary behaviors [29]. In chapter eight we used perceived
34 environmental measures such as perceived neighborhood safety and pleasantness. As we
35 did not find significant associations with sports behavior it might be that more objective
36 factors might play a role. Currently there is limited evidence about whether “objective” or
37 “perceived” environmental factors are more strongly associated with energy balance-related
38 behaviors. Previous studies have found associations between objective environmental mea-
39 sures with adolescents physical activity [30-32], active commuting to school [33, 34], fruit

1 and vegetable consumption [35, 36] and snack and soft drink consumption and purchases
2 [37]. Studies that examined associations of perceived environmental factors and physical
3 activity found also significant associations [38-40]. In a study in which perceived and objec-
4 tive environmental factors were compared, only the perceived environment was related to
5 adolescent girls physical activity and perceived and objectively measured environmental
6 factors were associated with each other [39]. In a recent study of Prins et al. [41] no agree-
7 ment between the perceived and objective environmental factors was found, which is in line
8 with studies conducted among adults [42-44]. This indicates that perceived and objective
9 environmental factors are different constructs [44] and recent findings suggest that percep-
10 tions of the environment may depend on what is objectively available in the environment
11 [39, 41]. It is therefore very relevant to combine the assessment of objective and perceived
12 environmental factors in one study to establish their influence on each other and on energy
13 balance-related behaviors.

15 Validity of the audit instrument and GIS data

17 Another limitation of the ENDORSE study was that the audit instrument to assess physical
18 environmental correlates in the school environment was not tested on validity. Testing the
19 validity of the audit instrument was not possible since a gold standard does not exist for as-
20 sessing environmental factors in the school environment. Most existing instruments focus
21 on the assessment of facilities for physical activities, aesthetics, traffic safety and food shops
22 in the neighborhood instead of the school environment. It was also not possible to compare
23 the results of the school canteen observations with product lists or sales records because
24 these records were not always available.

25 In chapter 7 GIS data was used to examine the number of food stores around schools.
26 However, the data in these kinds of databases might be an under- or over-representation of
27 the actual number of stores, depending on the quality of the databases.

29 Definition of neighborhood

31 A problem that arises with assessing environmental factors in the school and home neigh-
32 borhood is how neighborhoods are defined. Perceptions of the neighborhood environment
33 are highly individual [45] and it is still unclear what people define as their “neighborhood”
34 [24, 46]. To examine the perceptions of the neighborhood environment, we did not specify
35 what we meant by “your neighborhood” in the questionnaire. The definition adolescent have
36 of “their neighborhood” may depend on other environmental characteristics than distance,
37 such as access to vehicles and public transport [46], the kind of facility one is traveling to [47]
38 or living in rural or urban neighborhoods [48], making the area that adolescents perceive as
39 their neighborhood quite variable [46]. It is therefore important to assess perceptions of the

1 neighborhood by explicit items such as “Are there any parks within a 10 minute walk from
2 school” instead of “Are there any parks in your school neighborhood”.

3 We used a neighborhood boundary of 500 meters from school to examine the food en-
4 vironment. However, there is little agreement in the literature as to what might constitute
5 a good boundary from school or home. The choice for a neighborhood boundary might
6 depend on the behavior of interest, the likelihood of the behavior to occur close to school
7 or home and the age of the study population. Other problems with defining boundaries
8 is that key factors located just outside the defined boundary are missed and that factors
9 located inside the boundary are examined even if they are not used by the subjects because
10 of the presence of main barriers such as busy roads. Further research should therefore not
11 only assess facilities within the defined boundary but also combine these measures with the
12 assessment of the use of facilities within and outside the neighborhood boundary.

13 14 15 **10.3 INTEGRATION OF STUDY FINDINGS** 16

17 This thesis provides important information on the associations between environmental
18 factors and energy balance-related behaviors as proposed in the EnRG framework. The
19 research questions for this thesis were:


20 I What are important individual and environmental correlates of energy balance-related
21 behaviors?

22 II To what extent is the association between environmental factors and energy balance-
23 related behaviors mediated by individual cognitions?

24 The associations as supposed by the EnRG framework (Figure 10.1) were used to facilitate
25 the interpretation of the findings.

26
27 **What are important individual correlates of energy balance-related behaviors?**
28

29 This part of research question I relates to the association between cognitive factors and
30 energy-balance related behaviors. Cognitive factors have been found to be associated with
31 various health behaviors in many studies. This study contributes to the consistent evidence
32 for a strong association between cognitive factors from the TPB and energy balance-related
33 behaviors. Especially attitude was found to be a consistent correlate for various behaviors
34 such as soft drink consumption, snack consumption and leisure time sports participation.
35 However, perceived behavioral control was not found to be a correlate of soft drink and
36 snack consumption and leisure time sports participation. An explanation for this might be
37 that most adolescents perceive the dietary and physical activities as very easy to perform.
38
39



17 **Figure 10.1** Environmental Research framework for weight Gain prevention [1].

18
19

20 Next to the changeable cognitive correlates we also examined demographic factors as individual correlates of energy balance-related behaviors. Ethnicity and the school type adolescents attended (vocational or higher education) were found to be important individual correlates of energy balance-related behaviors. Demographic factors might be important moderators of the environment – behavior relationship [1] as the environment – behavior association might be different for various demographic sub-groups of the population [49-53]. As there were relatively few published obesity-prevention and treatment studies that are designed for specific educational or ethnic groups, the results of these studies can be used to distinguish specific risk groups and target groups for preventive interventions.

29

30 What are important environmental correlates of energy balance-related behaviors?

31

32 This part of research question I relates to the direct association between environmental factors and energy-balance related behaviors. Literature reviews have shown that evidence on environmental factors was inconsistent or lacking for some energy balance-related behaviors such as snack and soft drink consumption [54, 55]. Therefore, the ENDORSE study focused on these gaps in the literature and mainly examined factors in the school and neighborhood environment to replicate studies and build evidence on environmental correlates of energy balance-related behaviors. In Table 10.1 an overview according to the ANGLO framework is given of the environment – behavior relationships that were investigated in this thesis.

39

Table 10.1 Investigated environment – behavior relationships, categorized according to the ANGELO framework

Types	Levels	Home	School	Neighborhood	
Physical		Availability of sport attributes – leisure time sports	Availability of snacks and soft drinks and drinks in school canteens – soft drink and snack consumption	Distance from home to school – mode of commuting	
			Availability of snacks and soft drinks and drinks in vending machines – soft drink and snack consumption	Availability of food stores in the school neighborhood – soft drink and snack consumption	
				Distance to the nearest food store – soft drink and snack consumption	
				Neighborhood safety – leisure time sports	
				Neighborhood pleasantness – leisure time sports	
				Neighborhood facilities – leisure time sports	
	<hr/>				
	Economic				
	<hr/>				
	Political				
	<hr/>				
	Socio-cultural		Parents behavior – leisure time sports		
			Parenting practices – leisure time sports and soft drink consumption		

In line with results from the reviews, few significant associations between physical environmental factors and energy balance-related behaviors were found. We found availability of sports attributes associated with leisure time sports; the distance from home to school associated with the mode of commuting, and the distance to the nearest shop and the number of small food stores with soft drink consumption. Not all of these associations were very clear to explain, for example the inverse association found between the presence of small food shops nearby school and soft drink consumption.

In accordance with the review studies, our further studies suggest that social factors more strongly associated with physical activity and dietary behaviors than physical environmental factors [54, 55] as we found parenting practices or rules associated with soft drink consumption and leisure time sports and parents own sports behavior was associated with adolescents' leisure time sports. The range of social factors considered in this thesis was

1 narrow, and factors such as social networks, social capital, friends' support and behavior
2 should be included in future studies to provide further insight into the specific aspects of
3 the social environment that are most important for adolescents behaviors [3, 56].

4 The weak evidence found in the various studies should however not be interpreted as
5 an absence of a direct environment – energy balance-related behavior relationship. The
6 research field of environmental correlates of energy balance-related behavior is relatively
7 new and in a developing phase. The environment is very large which means that there are
8 many possible environmental factors that can be important for energy balance-related
9 behaviors. Most research has focused on just a part of the potentially relevant environment.
10 Therefore, future studies should focus on all aspects of the environment, also on economic
11 and political factors such as pocket money and school policies to reveal to what extent
12 environmental factors are associated with energy balance-related behaviors. Next to this,
13 the development of valid and reliable measures of the environment is necessary. As the
14 research field develops, more and more different instruments and definitions are used to
15 assess environmental factors. This hinders the comparison of results between studies and
16 makes the systematic development of theory for the relation of environment with energy
17 balance-related behaviors difficult.

18
19 To what extent is the association between environmental factors and energy balance-
20 related behaviors mediated by individual cognitions?

21
22 The second aim of this thesis was to investigate mediating factors to gain insight into the
23 mechanisms that underlie environment – energy balance-related behaviors relationships.
24 In this thesis cognitions partly mediated the association of the distance to the nearest shop
25 and the number of small food shops (chapter 7) and of parenting practices (chapter 9) with
26 adolescents soft drink consumption. Attitude and intention partly mediated the associa-
27 tions of physical activity attributes at home, parents' sport behavior and parental rule about
28 sports participation with leisure time sports (chapter 8). This mediating effect of cognitions
29 was also found in other studies. In a study of Jago et al. was found that the association
30 between distance to a small food store and low fat vegetable consumption was mediated by
31 low fat vegetable preferences [57]. De Bruijn and colleagues found that the association of
32 environmental aesthetics and distance to PA facilities on PA among adolescents was medi-
33 ated by intention to be physically active [58]. Motl and colleagues found that the association
34 of equipment accessibility with adolescent girls PA was mediated by self-efficacy [56]. In
35 these studies it is shown that not all cognitions from the Theory of Planned Behavior serve
36 as a mediator. The studies described above indicate that some TPB variables may be more
37 likely to serve as mediators in environment – behavior relationships than others [58, 59],
38 with the strongest evidence for attitude as a potential mediating variable [58-62]. Next to an
39 indirect association of environmental factors, in all studies on mediation, the environmen-

tal factors remained significantly associated with the outcome behavior after controlling for the mediating effect of cognitions. This is also shown in other studies that have combined environmental and cognitive factors [63-65]. According to the EnRG framework this partly unmediated environmental effect is an important explanatory mechanism and it can be argued that energy balance-related behaviors result partly from automatic and unconscious processes [66, 67].

As the EnRG-framework is specifically developed to generate hypotheses regarding when, how and for whom environmental factors might be influential, more research should focus on these questions instead of examining which environmental factors are important for energy balance-related behaviors. The studies in this thesis show that environmental factors can have both an indirect and direct association with energy balance-related behaviors. However, new questions also arise such as whether other mediating factors than cognitions for example preferences and environmental barriers play a role in the indirect association between environment and behavior. Baranowski et al. [68], proposed that there can be a range of mediating processes and cascading sequences of mediating processes. For instance, a child's self-efficacy for asking to be active after school will affect the likelihood that the child will ask to be active after school at home, which increases the likelihood that a parent will play with the child at home. Most previous studies have investigated mediation pathways for physical environmental factors. However, the studies in this thesis indicate that also social factors such as parents own behavior and parenting practices show an indirect association with energy balance-related behaviors. As the body of evidence on possible mediators grows, important questions for further research are whether mediating effects differ for different target groups and if some energy balance-related behaviors are more under the influence of automatic processes than others.

10.4 IMPLICATIONS OF THE STUDY FINDINGS

Findings from the studies reported in this thesis indicate that both individual and environmental factors are important for energy balance-related behaviors. However more research is needed to examine the relative influence and mechanisms behind these influences. Therefore, the results presented in this thesis have several implications for future research, theory and practice to assist further research and the development of obesity prevention interventions, which are described in this paragraph.

Implications for research

The research field of examining associations between environmental factors and energy balance-related behaviors is still in its infancy and needs further development and maturation.

1 tion. More insight is needed in how specific aspects of the environment can be assessed for
2 specific behaviors, what the best measurement instruments are, how the environment can
3 be defined for different target groups, and how gathered data can be used in analyses.

4 First, a large variety of potential environmental determinants of energy balance-related
5 behaviors have been studied, with few replicated studies for environment – behavior com-
6 binations. Further examination of potential relevant environmental factors is needed to
7 provide public health practitioners with recommendations for intervention development
8 and to improve theory and models on environmental determinants of health behaviors.
9 Research should not only be restricted to just a part of the potentially relevant environment,
10 but on all aspects listed in the ANGELO framework. Currently there is limited evidence
11 about whether “objective” or “perceived” environmental factors are more strongly associ-
12 ated with energy balance-related behaviors. Objective measures are generally regarded as
13 being superior to subjective self-reports. However, people may perceive their environments
14 differently even if they live in the same “objective” environment. The assessment of objective
15 and perceived factors should be combined in one study to establish their influence on each
16 other and on behavior to gain insight in which kind of determinants should be targeted in
17 preventive interventions: perceptions of the environment, the objective environment or a
18 combination of both.

19 Second, most studies on environmental determinant still apply relatively weak study
20 designs and measurement instruments. Longitudinal and intervention research is needed
21 to gain better insight into the relative importance of individual cognitive determinants
22 and environmental (physical, socio-cultural, economic, political) determinants for energy
23 balance-related behaviors. With these studies it can be examined if and how changes in the
24 environment lead to changes in behavior.

25 Third, valid and reliable measurement instruments to assess objective and subjective
26 environmental factors should be developed. More effort should be put into the develop-
27 ment of validated instruments that assess perceived environmental factors. Therefore, also
28 qualitative research is needed to examine how the target group perceives their neighborhood
29 taking the behavior of interest into account. More and more detailed objective measures of
30 environmental factors are available, for example, those documented in geographic infor-
31 mation systems (GIS). The use of GIS can be helpful to examine objective environmental
32 factors as environmental factors within defined boundaries can be examined. Next to this
33 research should also focus on developing good validated and reliable measurement instru-
34 ments for assessing dietary and physical activity behaviors as in adolescents these behaviors
35 are often over reported.

36 Finally, advanced research techniques need to be used to identify mediating and moderat-
37 ing factors of the environment – behavior relationship. It is important to examine which
38 cognitive factors mediate the environment – behavior relationship and if there are other
39 important mediators such as environmental barriers or parental influences.

1 Implications for theory development

2

3 The EnRG framework we used in this study was developed as a tool to help disentangle the
4 role of obesogenic environments. It is specifically directed at generating questions regarding
5 when, how and for whom environmental factors may be influential [1]. In this framework,
6 environmental influences are supposed to influence behavior both directly and indirectly.
7 In this thesis, individual cognitions were important correlates of soft drink and snack
8 consumption and sports participation but also mediators of the environment – behavior
9 relationship confirming the supposed indirect influence of the environment on behavior
10 in the EnRG framework. This indicates that socio-ecological models in which individual
11 cognitions and environmental factors are combined need specificity in the hypothesized
12 pathways between individual and environmental factors. To further develop theory on envi-
13 ronment – behavior relationships the mediation effects should be examined in longitudinal
14 research and for other environment – behavior relationships. Also other possible mediators
15 such as barriers and preferences should be examined. Future research is also needed to
16 examine if mediating effects differ for different target groups and if some energy balance-
17 related behaviors are more under the influence of automatic processes than others.

18 The direct association of environmental factors with energy balance-related behaviors
19 indicates also that many behaviors are more or less ‘automatic’. Environmental features can
20 prompt behavioral choices, without mediation by conscious decision making and theoretic-
21 al models are needed that focus also on environmentally cued habitual behavioral patterns
22 [69].

23

24 Recommendations for practice

25

26 The findings and implications lead to the following recommendations for practice and
27 intervention development.

28

29 *Intervention approach*

30 The partly mediated association between environmental factors and energy balance-related
31 behaviors indicate that both a health education approach and a health protection approach
32 by changing the environment might be appropriate strategies for behavior change interven-
33 tions [3]. Both intervention approaches are important, but since individual cognitions are
34 important correlates of energy balance-related behaviors, these factors remain important to
35 be targeted in interventions.

36

37 *Intervention ingredients*

38 Parental influences, including their own behavior and parenting rules seem to be important
39 factors for sports participation, soft drink and snack consumption. It is therefore recom-

1 mended to not only aim interventions at adolescents but also at the parents. Parents should
2 be made aware of their important role in promoting healthy dietary and physical activity be-
3 haviors. Possibly it is important to start to target parents at an early stage in the development
4 of the child, to teach parents to set clear rules and give a good example. The importance of
5 parents for energy balance-related behaviors also means that intervention should not only
6 take place at schools but also in the home environment. The results of this thesis do not
7 give a clear view on which factors in the school environment should be changed to promote
8 healthy energy balance-related behaviors. However, schools should promote healthy behav-
9 ior and healthy food should be made available and accessible in school.

11 *Target groups*

12 Adolescents at lower educational levels and adolescents with a non-Western ethnic back-
13 ground, mainly girls, engage in the most risk behaviors for overweight. Interventions are
14 needed that are effective in these groups. Therefore, studies to develop and evaluate inter-
15 ventions aimed at the promotion of healthy dietary and physical activity behaviors should
16 study the effects for these different target groups.

19 **10.5 GENERAL CONCLUSION**

21 There is only little evidence for associations between objective or perceived physical envi-
22 ronmental factors and energy balance-related behaviors. The evidence for social factors,
23 mainly parental factors is stronger and these factors might be more important for energy
24 balance-related behaviors among adolescents. Parents can importantly promote healthy en-
25 ergy balance-related behaviors among their offspring by shaping the environment by setting
26 clear rules, setting a good example and creating opportunities for the behavior. Next to this
27 we can conclude that environmental factors can have both an indirect and direct association
28 with energy balance-related behaviors. Especially attitudes and intentions are likely to be
29 important mediators of environment – behavior relationships. However, new questions also
30 arise, such as if other mediating factors than individual cognitions such as preferences and
31 environmental barriers play a role in the indirect association between environment and
32 behavior. More research is also needed on the investigation of mediating effects for different
33 target groups and if some energy balance-related behaviors are more under the influence of
34 automatic processes than others.

1 REFERENCES

- 2 1. Kremers SP, De Bruijn GJ, Visscher TL, Van Mechelen W, De Vries NK, Brug J: Environmental influ-
3 ences on energy balance-related behaviors: A dual-process view. *Int J Behav Nutr Phys Act* 2006,
4 3(1):9.
- 5 2. Swinburn B, Egger G, Raza F: Dissecting obesogenic environments: the development and application
6 of a framework for identifying and prioritizing environmental interventions for obesity. *Prev Med*
7 1999, 29(6 Pt 1):563-570.
- 8 3. Brug J, Kremers SP, van Lenthe F, Ball K, Crawford D: Environmental determinants of healthy eating:
9 in need of theory and evidence. *Pro Nutr Soc* 2008, 67(3):307-316.
- 10 4. Rennie KL, Johnson L, Jebb SA: Behavioural determinants of obesity. *Best Pract Res Clin Endocrinol*
11 *Metab* 2005, 19(3):343-358.
- 12 5. Swinburn BA, Caterson I, Seidell JC, James WP: Diet, nutrition and the prevention of excess weight
13 gain and obesity. *Public Health Nutr* 2004, 7(1A):123-146.
- 14 6. Crespo CJ, Smit E, Andersen RE, Carter-Pokras O, Ainsworth BE: Race/ethnicity, social class and
15 their relation to physical inactivity during leisure time: results from the Third National Health and
16 Nutrition Examination Survey, 1988-1994. *Am J Prev Med* 2000, 18(1):46-53.
- 17 7. Hosper K, Nierkens V, Nicolaou M, Stronks K: Behavioural risk factors in two generations of non-
18 Western migrants: do trends converge towards the host population? *Eur J Epidemiol* 2007, 22(3):163-
19 172.
- 20 8. Darling N, Steinberg L: Parenting style as context: An integrative model. *Psychological Bulletin* 1993,
21 113(3):487-496.
- 22 9. Birch LL, Fisher JO: Mothers' child-feeding practices influence daughters' eating and weight. *Am J*
23 *Clin Nutr* 2000, 71(5):1054-1061.
- 24 10. Costanzo PR, Woody, E.Z.: Domain-specific parenting styles and their impact on the child's develop-
25 ment op particular deviance: The example of obesity proneness. *J Soc Clin Psychol* 1985, 3(4):425-445.
- 26 11. Bauman AE, Sallis JF, Dzewaltowski DA, Owen N: Toward a better understanding of the influences
27 on physical activity: the role of determinants, correlates, causal variables, mediators, moderators, and
28 confounders. *Am J Prev Med* 2002, 23(2 Suppl):5-14.
- 29 12. Baron RM, Kenny DA: The moderator-mediator variable distinction in social psychological research:
30 conceptual, strategic, and statistical considerations. *J Pers Soc Psychol* 1986, 51(6):1173-1182.
- 31 13. Cerin E, Mackinnon DP: A commentary on current practice in mediating variable analyses in behav-
32 ioural nutrition and physical activity. *Public Health Nutr* 2008:1-7.
- 33 14. Cerin E, Taylor LM, Leslie E, Owen N: Small-scale randomized controlled trials need more powerful
34 methods of mediational analysis than the Baron-Kenny method. *J Clin Epidemiol* 2006, 59(5):457-
35 464.
- 36 15. MacKinnon DP, Lockwood CM, Hoffman JM, West SG, Sheets V: A comparison of methods to test
37 mediation and other intervening variable effects. *Psychol Methods* 2002, 7(1):83-104.
- 38 16. Maxwell SE, Cole DA: Bias in cross-sectional analyses of longitudinal mediation. *Psychol Methods*
39 2007, 12(1):23-44.
17. Singh AS, Chin APMJ, Kremers SP, Visscher TL, Brug J, van Mechelen W: Design of the Dutch Obesity
Intervention in Teenagers (NRG-DOiT): systematic development, implementation and evaluation of
a school-based intervention aimed at the prevention of excessive weight gain in adolescents. *BMC*
public health 2006, 6:304.
18. McPherson S, Hoelscher DM, Alexander M, Scanlon KS, Serdula MK: Dietary assessment methods
among school-aged children: validity and reliability. *Prev Med* 2000, 31:S11-S33.

19. Bogers RP, Van Assema P, Kester AD, Westerterp KR, Dagnelie PC: Reproducibility, validity, and responsiveness to change of a short questionnaire for measuring fruit and vegetable intake. *Am J Epidemiol* 2004, 159(9):900-909.
20. Van Assema P, Brug J, Ronda G, Steenhuis I, Oenema A: A short dutch questionnaire to measure fruit and vegetable intake: relative validity among adults and adolescents. *Nutrition and health (Berkhamsted, Hertfordshire)* 2002, 16(2):85-106.
21. Chin A Paw MJM, Sloomaker SM, Schuit AJ, van Zuidam M, Van Mechelen W: Test-retest reliability and concurrent validity of the Activity Questionnaire for Adults and Adolescents (AQuAA). Submitted
22. Trost SG: Objective measurement of physical activity in youth: current issues, future directions. *Exerc Sport Sci Rev* 2001, 29(1):32-36.
23. Sallis JF, Johnson ME, Calfas KJ, Caparosa S, Nichols JF: Assessing perceived physical environmental variables that may influence physical activity. *Res Q Exerc Sport* 1997, 68(4):345-351.
24. Giles-Corti B, Timperio A, Bull F, Pikora T: Understanding physical activity environmental correlates: increased specificity for ecological models. *Exerc Sport Sci Rev* 2005, 33(4):175-181.
25. Humpel N, Owen N, Leslie E: Environmental factors associated with adults' participation in physical activity: a review. *Am J Prev Med* 2002, 22(3):188-199.
26. Ajzen I, Fishbein M: *Understanding attitudes and predicting social behavior*. Englewood Cliffs NJ: Prentice-Hall; 1980.
27. Fishbein M, Ajzen I: *Belief, attitude, intention and behavior*. Don Mills (NY): Addison-Wesley; 1975.
28. Kremers SP, Visscher TL, Seidell JC, van Mechelen W, Brug J: Cognitive determinants of energy balance-related behaviours: measurement issues. *Sports medicine (Auckland, NZ)* 2005, 35(11):923-933.
29. Wiecha JL, Finkelstein D, Troped PJ, Fragala M, Peterson KE: School vending machine use and fast-food restaurant use are associated with sugar-sweetened beverage intake in youth. *J Am Diet Assoc* 2006, 106(10):1624-1630.
30. Gordon-Larsen P, Nelson MC, Page P, Popkin BM: Inequality in the built environment underlies key health disparities in physical activity and obesity. *Pediatrics* 2006, 117(2):417-424.
31. Cohen DA, Ashwood JS, Scott MM, Overton A, Evenson KR, Staten LK, Porter D, McKenzie TL, Catellier D: Public parks and physical activity among adolescent girls. *Pediatrics* 2006, 118(5):e1381-1389.
32. Powell LM, Chaloupka FJ, Slater SJ, Johnston LD, O'Malley PM: The availability of local-area commercial physical activity-related facilities and physical activity among adolescents. *Am J Prev Med* 2007, 33(4 Suppl):S292-300.
33. Timperio A, Ball K, Salmon J, Roberts R, Giles-Corti B, Simmons D, Baur LA, Crawford D: Personal, family, social, and environmental correlates of active commuting to school. *Am J Prev Med* 2006, 30(1):45-51.
34. Kerr J, Rosenberg D, Sallis JF, Saelens BE, Frank LD, Conway TL: Active commuting to school: Associations with environment and parental concerns. *Med Sci Sports Exerc* 2006, 38(4):787-794.
35. Timperio A, Ball K, Roberts R, Campbell K, Andrianopoulos N, Crawford D: Children's fruit and vegetable intake: Associations with the neighbourhood food environment. *Prev Med* 2008, 46(4):331-335.
36. Kubik MY, Lytle LA, Hannan PJ, Perry CL, Story M: The association of the school food environment with dietary behaviors of young adolescents. *Am J Public Health* 2003, 93(7):1168-1173.
37. Neumark-Sztainer D, French SA, Hannan PJ, Story M, Fulkerson JA: School lunch and snacking patterns among high school students: associations with school food environment and policies. *Int J Behav Nutr Phys Act* 2005, 2(1):14.

- 1 38. Mota J, Almeida M, Santos P, Ribeiro JC: Perceived Neighborhood Environments and physical activity in adolescents. *Prev Med* 2005, 41(5-6):834-836.
- 2 39. Scott MM, Evenson KR, Cohen DA, Cox CE: Comparing perceived and objectively measured access
3 to recreational facilities as predictors of physical activity in adolescent girls. *J Urban Health* 2007,
4 84(3):346-359.
- 5 40. Evenson KR, Scott MM, Cohen DA, Voorhees CC: Girls' perception of neighborhood factors on
6 physical activity, sedentary behavior, and BMI. *Obesity* 2007, 15(2):430-445.
- 7 41. Prins R, Oenema A, van der Horst K, Brug J: Objective and perceived availability of physical activity
8 opportunities: differences in associations with Physical Activity behavior among adolescents. Submitted
- 9 42. Boehmer TK, Hoehner CM, Deshpande AD, Brennan Ramirez LK, Brownson RC: Perceived and
10 observed neighborhood indicators of obesity among urban adults. *Int J Obes (Lond)* 2007, 31(6):968-
11 977.
- 12 43. Kirtland KA, Porter DE, Addy CL, Neet MJ, Williams JE, Sharpe PA, Neff LJ, Kimsey CD, Ainsworth
13 BE: Environmental measures of physical activity supports. Perception versus reality. *Am J Prev Med*
14 2003, 24(4):323-331.
- 15 44. McGinn AP, Evenson KR, Herring AH, Huston SL, Rodriguez DA: Exploring associations between
16 physical activity and perceived and objective measures of the built environment. *J Urban Health* 2007,
17 84(2):162-184.
- 18 45. Hume C, Salmon J, Ball K: Children's perceptions of their home and neighborhood environments,
19 and their association with objectively measured physical activity: a qualitative and quantitative study.
20 *Health Educ Res* 2005, 20(1):1-13.
- 21 46. Ball K, Timperio AF, Crawford DA: Understanding environmental influences on nutrition and physical
22 activity behaviors: where should we look and what should we count? *Int J Behav Nutr Phys Act*
23 2006, 3:33.
- 24 47. McCormack GR, Giles-Corti B, Bulsara M, Pikora TJ: Correlates of distances traveled to use recreational
25 facilities for physical activity behaviors. *Int J Behav Nutr Phys Act* 2006, 3:18.
- 26 48. Colabianchi N, Dowda M, Pfeiffer KA, Porter DE, Almeida MJ, Pate RR: Towards an understanding
27 of salient neighborhood boundaries: adolescent reports of an easy walking distance and convenient
28 driving distance. *Int J Behav Nutr Phys Act* 2007, 4(1):66.
- 29 49. Giles-Corti B, Donovan RJ: Socioeconomic status differences in recreational perceived activity levels
30 and real and perceived access to a supportive physical environment. *Prev Med* 2002, 35:601-611.
- 31 50. Suminski RR, Poston WS, Petosa RL, Stevens E, Katzenmoyer LM: Features of the neighborhood
32 environment and walking by U.S. adults. *Am J Prev Med* 2005, 28(2):149-155.
- 33 51. Humpel N, Marshall AL, Leslie E, Bauman A, Owen N: Changes in neighborhood walking are related
34 to changes in perceptions of environmental attributes. *Ann Behav Med* 2004, 27(1):60-67.
- 35 52. Humpel N, Owen N, Iverson D, Leslie E, Bauman A: Perceived environment attributes, residential
36 location, and walking for particular purposes. *Am J Prev Med* 2004, 26(2):119-125.
- 37 53. Garcia Bengoechea E, Spence JC, McGannon KR: Gender differences in perceived environmental
38 correlates of physical activity. *Int J Behav Nutr Phys Act* 2005, 2:12.
- 39 54. Ferreira I, van der Horst K, Wendel-Vos W, Kremers S, van Lenthe F, Brug J: Environmental correlates of physical activity in youth - A review and update. *Obesity Reviews* 2007, 8(2):129-154.
55. van der Horst K, Oenema A, Ferreira I, Wendel-Vos W, Giskes K, van Lenthe F, Brug J: A systematic review of environmental correlates of obesity-related dietary behaviors in youth. *Health Educ Res* 2007, 22:203-226.

- 1 56. Motl RW, Dishman RK, Saunders RP, Dowda M, Pate RR: Perceptions of physical and social environ-
2 ment variables and self-efficacy as correlates of self-reported physical activity among adolescent girls.
3 *Journal of pediatric psychology* 2007, 32(1):6-12.
- 4 57. Jago R, Baranowski T, Baranowski JC, Cullen KW, Thompson D: Distance to food stores & adolescent
5 male fruit and vegetable consumption: mediation effects. *Int J Behav Nutr Phys Act* 2007, 4(1):35.
- 6 58. de Bruijn GJ, Kremers SP, Lensvelt-Mulders G, de Vries H, van Mechelen W, Brug J: Modeling in-
7 dividual and physical environmental factors with adolescent physical activity. *Am J Prev Med* 2006,
8 30(6):507-512.
- 9 59. Van der Horst K, Timperio A, Crawford D, Roberts R, Brug J, Oenema A: The school food environ-
10 ment: associations with adolescent soft drink and snack consumption. *American Journal of Prev Med*
11 2008, 35(3): 217-223.
- 12 60. Rhodes RE, Brown SG, McIntyre CA: Integrating the perceived neighborhood environment and the
13 theory of planned behavior when predicting walking in a Canadian adult sample. *Am J Health Promot*
14 2006, 21(2):110-118.
- 15 61. Rhodes RE, Courneya KS, Blanchard CM, Plotnikoff RC: Prediction of leisure-time walking: an
16 integration of social cognitive, perceived environmental, and personality factors. *Int J Behav Nutr*
17 *Phys Act* 2007, 4:51.
- 18 62. van der Horst K, Kremers S, Ferreira I, Singh A, Oenema A, Brug J: Perceived parenting style and
19 practices and sugar-sweetened beverage consumption in adolescents. *Health Educ Res* 2007, 22:295-
20 304.
- 21 63. de Bruijn GJ, Kremers SP, Schaalma H, van Mechelen W, Brug J: Determinants of adolescent bicycle
22 use for transportation and snacking behavior. *Prev Med* 2005, 40(6):658-667.
- 23 64. Lewis BA, Marcus BH, Pate RR, Dunn AL: Psychosocial mediators of physical activity behavior
24 among adults and children. *Am J Prev Med* 2002, 23(2 Suppl):26-35.
- 25 65. De Bourdeaudhuij I, Sallis J: Relative contribution of psychosocial variables to the explanation of
26 physical activity in three population-based adult samples. *Prev Med* 2002, 34(2):279-288.
- 27 66. Aarts H, Dijksterhuis A: The automatic activation of goal-directed behaviour: The case of travel habit.
28 *J Environm Psychol* 2000, 20(1):75-82.
- 29 67. Chartrand T-L: The Role of Conscious Awareness in Consumer Behavior. *J Consumer Psychol* 2005,
30 15:203-210.
- 31 68. Baranowski T, Anderson C, Carmack C: Mediating variable framework in physical activity interven-
32 tions. How are we doing? How might we do better? *Am J Prev Med* 1998, 15(4):266-297.
- 33 69. Owen N, Humpel N, Leslie E, Bauman A, Sallis JF: Understanding environmental influences on
34 walking: Review and research agenda. *Am J Prev Med* 2004, 27(1):67-76.
- 35
- 36
- 37
- 38
- 39

Summary

Over the past decades the numbers of children and adults being overweight or obese have increased so rapidly, that overweight and obesity are among the most important and challenging public health problems. It is therefore important to prevent overweight in all age groups. Children and adolescents may however be especially important groups to target. Obesity at young age is associated with a higher likelihood of the development of chronic diseases at an early age or later in life. Furthermore, overweight or obese children and adolescents are more likely to become overweight or obese adults.

It is already known that weight gain is the result of a positive energy balance in which the energy intake (via diet) exceeds energy expenditure (mainly via physical activity). So, to be able to successfully prevent overweight among children and adolescents, it is important to know which specific dietary and physical activity behaviors (energy balance-related behaviors - EBRB) contribute most to weight gain among children and adolescents.

Another important element for successful prevention of overweight is a detailed understanding of the factors that determine these EBRB. Individual determinants of behavior such as knowledge, attitudes, social influences, and motivation have been found to be relevant in earlier research. However, it is also likely that physical and social environmental factors such as availability of food products, opportunities to be active and parents can influence dietary and physical activity behaviors of children and adolescents.

To study this, and the interplay between potential personal and environmental determinants of behavioral nutrition as well as physical activity among young adolescents, the ENDORSE project (ENvironmental Determinants of Obesity in Rotterdam SchoolchildrEn) was initiated in 2004. The main goals of the ENDORSE project were: (1) to identify which energy balance-related behaviors are associated with overweight and obesity; (2) to examine important individual and environmental correlates of these energy balance-related behaviors; (3) to investigate the associations with and the interactions between these correlates and energy balance-related behaviors; and (4) to formulate objectives to be targeted in interventions aimed at the prevention of overweight in adolescents aged 12-15 years. This thesis presents a series of studies in which these questions are addressed.

In the general introduction (**Chapter 1**), the background, aims and theoretical framework used in the ENDORSE study are presented.

In **Chapter 2**, the study protocol of the ENDORSE study is described together with the design and the measurement instruments. Data were collected among adolescents in the first (12- to 13-year-olds) and third (14- to 15-year-olds) years of secondary school. Seventeen schools participated in the ENDORSE study and 1668 adolescents and their parents were selected to participate in the study. In this chapter the results of the explorative research on

1 the identification of risk behaviors for overweight and obesity are described as well. Based
2 on a review of the literature and the opinion of consulted experts, the following EBRB were
3 selected to be addressed in the ENDORSE study: active transport to school, leisure time
4 activities, sports, watching television, computer use, soft drink consumption, sweet snack
5 consumption, savory snack consumption and breakfast consumption.
6

7 Research into environmental influences on behavior is relatively new. Therefore, we started
8 by searching literature on existing evidence on environmental correlates of energy balance-
9 related behaviors (**Chapters 3 and 4**). Evidence was found for associations between parents'
10 intake and the intakes of their children, especially for energy and fat intake. If parents had
11 higher educational levels, their adolescent's had a higher fruit and vegetable intake. Social
12 support, mother's education level, family income, non-vocational school attendance, and
13 low crime incidence were associated with higher physical activity among adolescents. Both
14 reviews showed that there were only a limited number of studies available that examined
15 physical environmental factors, such as the availability of food products or facilities to be
16 physically active. Although many studies were school-based studies, mainly home and
17 neighborhood environmental factors were examined in these studies. School environmental
18 factors were often left out of consideration.

19 The Chapters 5 and 6 focus on identifying important socio-demographic factors.

20 In **Chapter 5** differences in energy balance-related behaviors and overweight according
21 to gender, ethnicity and school-level are studied. The results of this study showed that
22 youth from a non-Western ethnic background and those attending lower-level secondary
23 education were more likely to be overweight and to engage in unfavorable energy balance-
24 related behaviors than youth from a Western ethnic background and higher-level secondary
25 education.

26 In **Chapter 6** these differences are examined for commuting to school. Adolescents of
27 a non-Western ethnic background were more likely to walk to school and use non-active
28 transportation (public transport, scooter) compared to native Dutch adolescents, who were
29 more likely to cycle to school. Adolescents were less likely to walk or cycle to school if
30 they lived further away from school, in which case they were more likely to use non-active
31 transportation.
32

33 The studies that are presented in chapters 7-9 examine the association of individual and en-
34 vironmental factors with energy balance-related behaviors. Moreover these studies explore
35 *how* the environment may influence energy balance-related behaviors.

36 The study in **Chapter 7** examines whether the availability of snacks and drinks in school
37 canteens and the presence of food stores in the school neighborhood were associated with
38 a higher soft drink and snack consumption. The results showed that if the distance to the
39 nearest food store was greater than 200 meters, and if there were more small food stores in

1 the school neighborhood, the adolescents drank less soft drinks. The availability of school
2 canteen products and the availability food stores in the proximity of the school were not
3 associated with snack consumption.

4 **Chapter 8** focuses on whether the availability of sports facilities, neighborhood factors
5 (availability of facilities, safety and pleasantness) and parental factors were associated with
6 adolescent sports participation. The results showed that adolescents participated more in
7 leisure time sports if they had more sports equipment at home, if their parents participated
8 more in sports and if their parents endorsed the rule that their child should participate in
9 sports. The influence of these environmental factors was not only direct (i.e., more equip-
10 ment was associated with higher sports participation), but also indirect via personal factors.
11 Adolescents with more sports equipment at home, with parents doing sports and the rule
12 to participate in sports, had also a more positive attitude and intention towards sports, and
13 were in turn also more likely to participate in sports.

14 **Chapter 9** reports on a study about potential parenting influences on adolescent soft
15 drink consumption. The results of this study indicated that more restrictive parenting
16 practices (rules) were associated with less soft drink consumption. It appeared that the
17 working mechanism was both direct (i.e., strict rules were associated with lower soft drink
18 consumption) and indirect. Having stricter rules about consuming soft drinks at home was
19 associated with a more negative attitude and cognitions towards soft drink consumption.
20 These more negative cognitions in turn were associated with lower soft-drink consumption.
21 Next to this, it was found that the association between parenting practices and soft drink
22 consumption was stronger among adolescents who perceived the parenting style of their
23 parents as being moderately strict and highly involved (authoritative parenting style).

24
25 In the general conclusion (**Chapter 10**) the findings of all studies are integrated and conclu-
26 sions and recommendations for practice and future research are given.

27 There is only little evidence for associations between physical environmental factors and
28 energy balance-related behaviors. The evidence for socio-cultural factors and in particular
29 parental influences is stronger and might be more important for the prevention of overweight
30 among adolescents. Parents can promote healthy energy balance-related behaviors by giving
31 the right example, by setting clear rules and by creating possibilities for healthy behaviors.

32 Environmental factors can have both an indirect (via cognitions) and a direct (automatic,
33 unconscious) association with energy balance-related behaviors. Future research should ex-
34 amine if some energy balance-related behaviors are more under the influence of automatic
35 processes than others.

36 Adolescents attending lower level education and adolescents from a non-Western ethnic
37 background are important target groups for the prevention of overweight. Interventions that
38 target these groups should be developed and investigated. Next to this, parents should be
39 made aware of their important role in the promotion of healthy behaviors of their children.

1 Samenvatting

2
3 Het aantal mensen dat in Nederland en andere Westerse landen aan overgewicht of obesitas
4 lijdt, is de laatste decennia zo sterk gestegen dat overgewicht en obesitas inmiddels behoren
5 tot de belangrijkste volksgezondheid problemen. Daarom is het van belang overgewicht in
6 alle leeftijdsgroepen te voorkómen. Met name kinderen en adolescenten zijn daarbij een
7 belangrijke doelgroep. Obesitas op jonge leeftijd gaat gepaard met een verhoogde kans op
8 het ontstaan van chronische ziekten op jonge en latere leeftijd. Daarnaast hebben kinderen
9 en jongeren met obesitas een grotere kans om ook overgewicht of obesitas te hebben als ze
10 volwassen zijn.

11 Het is al bekend dat gewichtsstijging het gevolg is van een positieve energiebalans waarbij
12 de energie-inname (via de voeding) groter is dan het energieverbruik (met name via licha-
13 melijke activiteit). Voor de succesvolle preventie van overgewicht is het belangrijk te weten
14 welke specifieke energiebalans gerelateerde gedragingen bijdragen aan gewichtsstijging bij
15 kinderen en jongeren.

16
17 Een ander belangrijk element voor succesvolle preventie van overgewicht is het verkrijgen
18 van een gedetailleerd overzicht van de factoren die deze energiebalans gerelateerde gedra-
19 gingen beïnvloeden. Uit voorgaand onderzoek is gebleken dat persoonlijke determinanten
20 van gedrag zoals kennis, attitudes, sociale invloed en motivatie belangrijk zijn. Echter, ook
21 omgevingsfactoren zoals de beschikbaarheid van voedingsmiddelen, de mogelijkheden tot
22 lichamelijke activiteit en de ouders kunnen voedings- en beweeggedrag van kinderen en
23 jongeren beïnvloeden.

24 Om meer zicht te krijgen op het samenspel tussen de persoonlijke en omgevingsdeter-
25 minanten van voedings- en beweeggedrag bij jongeren, is in 2004 het ENDORSE project
26 (ENvironmental Determinants of Overweight in Rotterdam SchoolchildEn) geïnitieerd. De
27 doelen van het ENDORSE project waren (1) het identificeren welke energiebalans gerela-
28 teerde gedragingen gerelateerd zijn aan overgewicht, (2) het onderzoeken van individuele
29 en omgevingsfactoren van deze energiebalans gerelateerde gedragingen, (3) het onderzoe-
30 ken van het verband en het samenspel tussen deze factoren en energiebalans gerelateerde
31 gedragingen en (4) het formuleren van doelstellingen voor interventies die gericht zijn op
32 de preventie van overgewicht bij 12-15 jarige jongeren. In dit proefschrift worden studies
33 gepresenteerd die de resultaten van dit onderzoek beschrijven.

34
35 In de algemene introductie (**hoofdstuk 1**) worden de achtergrond, de doelen en het theore-
36 tische raamwerk van de ENDORSE studie beschreven.

37 In **hoofdstuk 2** wordt het ENDORSE onderzoeksprotocol beschreven samen met de
38 onderzoeksopzet en de meetinstrumenten. Het ENDORSE onderzoek vond plaats onder
39 jongeren in de eerste (12 tot 13 jaar oud) en derde (14 tot 15 jaar oud) klas van het voortgezet

1 onderwijs. Zeventien scholen deden mee aan de ENDORSE studie en 1668 jongeren en
2 hun ouders werden geselecteerd om deel te nemen. Daarnaast wordt in dit hoofdstuk het
3 resultaat van een verkennend onderzoek beschreven waarin de risicogedragingen voor
4 het ontstaan van overgewicht en obesitas werden verkend. Een literatuuronderzoek en
5 het raadplegen van experts resulteerde in de selectie van de energiebalans gerelateerde
6 gedragingen die in het ENDORSE project nader onderzocht zouden worden: transport naar
7 school, lichamelijke activiteit in de vrije tijd, sporten, televisie kijken, computer gebruik en
8 de consumptie van suikerhoudende frisdranken, tussendoortjes en ontbijt.

9
10 Onderzoek naar omgevingsinvloeden op gedrag is relatief nieuw. Daarom is er eerst een
11 literatuuronderzoek uitgevoerd om het beschikbare bewijs over de relatie tussen omge-
12 vingsfactoren en energiebalans gerelateerde gedragingen te inventariseren (**Hoofdstuk 3**
13 **en 4**). Met dit literatuuronderzoek werd consistent bewijs gevonden dat er een verband is
14 tussen de inname van energie en vet door ouders en de inname van energie en vet door hun
15 kinderen. Ook liet dit onderzoek zien dat bij een hoger opleidingsniveau van de ouders de
16 jongeren meer groenten en fruit eten.

17 Sociale steun, het opleidingsniveau van de moeder, gezinsinkomen, het volgen van een
18 hogere middelbare schoolopleiding en een lage criminaliteitsincidentie in de woonomge-
19 ving waren geassocieerd met meer lichamelijke activiteit bij jongeren. Het literatuuronder-
20 zoek liet verder zien dat er maar een beperkt aantal onderzoeken beschikbaar waren die
21 fysieke omgevingsdeterminanten onderzoeken. Ondanks dat veel studies op scholen zijn
22 uitgevoerd, werden voornamelijk factoren in de thuis- en woonomgeving onderzocht en
23 bleef de schoolomgeving vaak buiten beschouwing.

24
25 De hoofdstukken 5 en 6 richten zich op het identificeren van belangrijke sociaaldemogra-
26 fische factoren.

27 In **hoofdstuk 5** werd onderzocht of er verschillen zijn in energiebalans gerelateerde
28 gedragingen en overgewicht voor geslacht, etniciteit en schoolniveau. De resultaten van
29 deze studie lieten zien dat jongeren met een niet westerse achtergrond en jongeren van het
30 VMBO vaker overgewicht hadden en ongezondere energiebalans gerelateerde gedragingen
31 vertoonden.

32 In **hoofdstuk 6** werden deze verschillen onderzocht voor vervoer naar school. Jongeren
33 van een niet Westerse afkomst wandelden vaker naar school of gebruikten vaker niet actieve
34 transportmiddelen (openbaar vervoer, scooter) dan Nederlandse jongeren, die vaker de
35 fiets gebruikten. Jongeren gingen minder vaak fietsend of lopend naar school naarmate de
36 afstand tot de school groter was.

1 De studies in hoofdstuk 7-9 onderzoeken het verband tussen de individuele en omgevings-
 2 factoren met de energiebalans gerelateerde gedragingen. Daarnaast wordt in deze studies
 3 verkend *hoe* de omgevingsfactoren de energiebalans gerelateerde gedragingen beïnvloeden.

4 In **hoofdstuk 7** werd onderzocht of de beschikbaarheid van frisdranken en snacks in
 5 schoolkantines en de aanwezigheid van levensmiddelenwinkels rond scholen samenhangen
 6 met de inname van frisdrank en snacks. De resultaten lieten zien dat als de afstand van
 7 school tot de dichtstbijzijnde winkel meer dan 200 meter was en als er meer kleine winkels
 8 in de schoolomgeving waren, de jongeren minder frisdrank dronken. Er werd geen verband
 9 gevonden voor het aanbod in schoolkantines en de winkels in de schoolomgeving met
 10 snackconsumptie.

11 In **hoofdstuk 8** onderzochten we of de beschikbaarheid van sportfaciliteiten, buurtfacto-
 12 ren, en de invloed van ouders verband hielden met het sportgedrag van jongeren. De
 13 jongeren sportten meer in de vrije tijd als ze thuis meer sportfaciliteiten hadden, als de ou-
 14 ders zelf meer sportten en als de ouders de regel hadden dat hun kind aan sport moet doen.
 15 De relatie van deze omgevingsfactoren was niet alleen direct (b.v. meer faciliteiten hielden
 16 verband met meer sporten), maar ook indirect, via de persoonlijke factoren. Jongeren met
 17 meer sportfaciliteiten thuis, ouders die sportten en met de regel dat ze aan sport moeten
 18 doen, hadden ook een positievere attitude en intentie tot sporten, en vervolgens ook een
 19 grotere kans om meer te sporten.

20 **Hoofdstuk 9** beschrijft een onderzoek naar de invloed van de ouders op het frisdrank-
 21 gebruik van jongeren. De resultaten van dit onderzoek lieten zien dat meer restrictieve
 22 regels over frisdrankgebruik waren geassocieerd met minder frisdrank consumptie. Het
 23 werkingsmechanisme leek ook hier niet alleen direct (restrictieve regels resulteerden in
 24 minder frisdrank consumptie) maar ook indirect. Jongeren die meer regels ondervonden,
 25 hadden ook een negatievere attitude en andere cognities ten aanzien van frisdranken, die
 26 vervolgens verband hielden met minder frisdrank consumptie. Daarnaast werd gevonden
 27 dat het verband tussen opvoedingsregels en frisdrankconsumptie sterker was bij jongeren
 28 die hun ouders als matig strikt en zeer betrokken ervaren (autoritatieve opvoedingsstijl).

29
 30 In de algemene conclusie (**hoofdstuk 10**) worden de resultaten uit de onderzoeken geïnte-
 31 greerd en conclusies en aanbevelingen gegeven voor de praktijk en verder onderzoek.
 32 De belangrijkste conclusies zijn dat er weinig bewijs is voor een verband tussen fysieke
 33 omgevingsfactoren en energiebalans gerelateerde gedragingen. Voor sociaal-culturele om-
 34 gevingsfactoren, met name de invloed van ouders, is het bewijs sterker en is mogelijk
 35 belangrijker voor de preventie van overgewicht bij jongeren. Ouders kunnen gezonde
 36 energiebalans gerelateerde gedragingen bevorderen, door het goede voorbeeld te geven,
 37 duidelijke regels te formuleren en mogelijkheden te creëren voor gezond gedrag.

38 Omgevingsfactoren kunnen op een indirecte (via individuele cognities) en directe
 39 (automatische en onbewuste) manier verband houden met energiebalans gerelateerde ge-

1 dragingen. In toekomstig onderzoek moet onderzocht worden in hoeverre de verschillende
2 energiebalans gerelateerde gedragingen door automatische processen beïnvloed worden.

3 Voor de preventie van overgewicht zijn met name jongeren van het VMBO en jongeren
4 met een niet-westerse achtergrond belangrijke doelgroepen. Interventies die gericht zijn
5 op deze groepen moeten ontwikkeld en onderzocht worden. Daarnaast moeten ouders
6 bewust gemaakt worden van hun belangrijke rol in het bevorderen van gezond gedrag bij
7 hun kinderen.

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

1 Dankwoord

2
3 Toen ik besloot om in Maastricht Gezondheidsvoorlichting te gaan studeren wist ik dat ik
4 hier verder mee wilde, maar het hoe en wat moest nog vorm krijgen. Nu, niet eens zo heel
5 veel jaren later, ligt dit proefschrift er. Stef, je enthousiasme tijdens mijn jaarwerkstuk en
6 afstudeeronderzoek was besmettelijk en doorslaggevend. Je overhandigde mij mijn diploma
7 in Maastricht en ook nu ben je er weer bij. Dit keer in de grote promotiecommissie. Wie
8 weet waar we elkaar de volgende keer treffen.

9
10 Bovenal wil ik Hans en Anke bedanken. Tenslotte hebben zij de grootste bijdrage geleverd
11 aan dit proefschrift. Hans, je dacht altijd in oplossingen en zelden in problemen. Dat zijn
12 voor een promotor goede kenmerken. Je zag overal kansen en mogelijkheden, zowel binnen
13 als buiten je functie. Ik hoop dat ik daar wat van heb opgepikt! Anke, het was eerst even
14 aftasten, maar we hebben altijd goed kunnen samenwerken. Je maakte wel veel werk van het
15 begeleiden,waardoor de papers altijd beter zijn geworden. Bedankt voor de vele hulp en
16 de fijne tijd samen.

17
18 Lottie & Nellie, mijn dappere onderzoeksassistenten! Gewapend met weegschaal, meetlat
19 en vragenlijsten gingen jullie op pad om pubers te meten en te wegen. Meestal was het leuk,
20 soms lastig. Met jullie enthousiasme, energie en blonde haren hebben jullie de leerlingen
21 en leraren voor jullie gewonnen. Zonder jullie had ik het niet gekund! Hartelijk dank voor
22 jullie inzet.

23
24 De samenwerking met de GGD Rotterdam is voor het ENDORSE onderzoek erg belangrijk
25 geweest. Ik kon gebruik maken van de structuur van “de Jeugdmonitor” waardoor veel scho-
26 len, leraren, leerlingen en ouders bereid waren om deel te nemen. Petra en Wilma bedankt!

27
28 Een leuke sfeer en veel koffiepauzes maken werken makkelijker en leuker. Kamergenotes
29 Birgitte, Willemieke en Wendy, bedankt voor alle gezelligheid. Paranimf Tinneke, wat heb-
30 ben wij veel koffie gedronken en gekletst. Het was altijd gezellig om het leven van alledag
31 met je te bespreken. Congresmaatje Carlijn; walvissenjacht in Boston, luxe kamer met
32 ontbijt in Olso en de Gondola van Banff... we kwamen tenslotte niet alleen voor het congres.
33 Appel Gert Jan, je favoriete koffiedrinkstagiaire is klaar! We hebben alleen nog steeds geen
34 paper met elkaar geschreven, dus... toch nog maar een koffie date? Saskia bedankt voor
35 de hulp met statistische problemen, het schrijven van papers en de fijne samenwerking.
36 Nannah, Merel, Marianne, Rick, Hein en andere (ex) DGG-ers: het was leuk om jullie als
37 collega's te hebben, bedankt voor de gezelligheid.

38 David and Anna, thanks for inviting me for a stay in Melbourne and the opportunity to
39 work at the Centre for Physical Activity and Nutrition Research. It was a great life and work

1 experience with a nice paper as a result. Jo, Karen, Abbie, Clare, Kylie and Rebecca, thank
2 you for making my stay so enjoyable!

3

4 Familie & vrienden, het 'gewoon mijn werk' is klaar. Steun bij mijn proefschrift heb ik niet
5 echt nodig gehad, dat liep eigenlijk op rolletjes. Maar er is veel gebeurd de afgelopen jaren,
6 leuke en verdrietige gebeurtenissen. Dank voor alle steun, maar vooral voor alle gezellige
7 momenten.

8

9 Pa, misschien komt het er nog eens van, maar de eerste Dr. dat ben ik! Ma, bedankt voor de
10 goede zorgen en gezelligheid! Het is altijd fijn om thuis te komen.

11

12 Bas, bedankt voor alles. Voor mij ben je de beste man die er is. Door jou is de combinatie
13 carrière & Femke perfect!

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

Curriculum Vitae

1
2
3 Klazine van der Horst is geboren op 20 december 1978 in Zwolle. Na de middelbare school
4 volgde ze de opleiding Voeding & Diëtetiek aan de Hogeschool van Arnhem en Nijmegen.
5 Na haar afstuderen in 2001 volgde ze de Master Gezondheidsvoorlichting van de opleiding
6 Gezondheidswetenschappen aan de universiteit van Maastricht. Tijdens en na haar afstu-
7 deeronderzoek werkte Klazine als onderzoeksmedewerker en junior onderzoeker bij het
8 EMGO instituut van het VU Medisch Centrum in Amsterdam. In augustus 2004 kon ze
9 beginnen met haar promotieonderzoek bij de afdeling Maatschappelijke Gezondheidszorg
10 van het Erasmus MC Rotterdam. Daar voltooide zij gelijktijdig de Master of Public Health
11 van het NIHES (2006). Vanaf mei 2008 werkte ze in Rotterdam als wetenschappelijk me-
12 dewerker verder aan de follow-up studie van het ENDORSE project. Sinds december 2008
13 werkt ze als postdoctoraal onderzoeker in Zwitserland aan de ETH te Zürich bij de afdeling
14 Consumer Behavior.

List of Publications

1
2
3
4 Hume C, van der Horst K, Brug J, Salmon J, Oenema A. Understanding the correlates of
5 adolescents' TV viewing: a social ecological approach. *International Journal of Pediatric*
6 *Obesity (in press)*.

7
8 van der Horst K, Oenema A, te Velde SJ, Brug J. Gender, ethnic and school type differences
9 in overweight and energy balance-related behaviours among Dutch adolescents. *Interna-*
10 *tional Journal of Pediatric Obesity* 2009, May 15:1-10. [Epub ahead of print].

11
12 DeJong CS, van Lenthe F, van der Horst K, Oenema A. Environmental and cognitive cor-
13 relates of adolescent breakfast consumption. *Preventive Medicine* 2009, 48(4): 372-377.

14
15 van der Horst K, Oenema A, van de Looij-Jansen P, Brug J. The ENDORSE study: Research
16 into environmental determinants of obesity related behaviors in Rotterdam Schoolchildren.
17 *BMC Public Health* 2008, 28;8: 142.

18
19 van der Horst K, Timperio A, Crawford D, Roberts R, Brug J, Oenema A. The school food
20 environment: associations with adolescent soft drink and snack consumption. *American*
21 *Journal of Preventive Medicine* 2008, 35(3): 217-23.

22
23 Bere E, van der Horst K, Oenema A, Prins R, Brug J. Socio-demographic factors as cor-
24 relates of active commuting to school in Rotterdam, the Netherlands. *Preventive Medicine*
25 2008, 47(4): 412-416.

26
27 van der Horst K, Oenema A, Ferreira I, Wendel-Vos W, Giskes K, Van Lenthe F, Brug J. A
28 review of environmental correlates of obesity-related dietary behaviors in youth. *Health*
29 *Education Research* 2007, 22: 203-226.

30
31 van der Horst K, Kremers S, Ferreira I, Singh A, Oenema A, Brug J. Perceived parenting
32 style and practices and sugar-sweetened beverage consumption by adolescents. *Health*
33 *Education Research* 2007, 22: 295-304

34
35 van der Horst K, Chin a Paw M, Twisk J, Van Mechelen W. A brief review on correlates of
36 physical activity and sedentariness in youth. *Medicine & Science in Sports & Exercise* 2007,
37 39(8): 1241-1250.

1 Ferreira I, van der Horst K, Wendel-Vos W, Kremers S, van Lenthe F, Brug J. Environmental
2 correlates of physical activity in youth - A review and update. *Obesity Reviews* 2007, 8(2):
3 129-154.

4
5 Kremers SPJ, van der Horst K, Brug J. Adolescent screen-viewing behaviour is associated
6 with consumption of sugar-sweetened beverages: the role of habit strength and perceived
7 parental norms. *Appetite* 2007, 48: 345-350.

8

9

10 **SUBMITTED**

11

12 van der Horst K, Oenema A, te Velde SJ, Brug J. Do individual cognitions mediate the
13 association of socio-cultural and physical environmental factors with adolescent sports
14 participation? *Public Health Nutrition* (submitted).

15

16 Prins R, Oenema A, van der Horst K, Brug J. Objective and perceived availability of physi-
17 cal activity opportunities: differences in associations with physical activity behavior among
18 adolescents. *International Journal for Behavioral Nutrition and Physical Activity* (submitted).

19

20 van der Horst K, Siegrist M, Orlow P, Giger, M. Residents' reasons for specialty choice:
21 Gender, time, patient and career aspects. *Medical Education* (submitted)

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

PhD Portfolio

Summary of PhD training and teaching activities

Name PhD student: Klazine van der Horst
Erasmus MC Department: Public Health

PhD period: 2004-2008
Promotor: Prof.dr.ir. J. Brug
Supervisor: Dr. A. Oenema

	Year	Workload (Hours/ ECTS)
1. PhD training		
General courses		
Master of Science in Health Sciences, specialization Public Health, NIHES, Erasmus MC Rotterdam	2006	70 ECTS
Specific courses		
Theory construction and statistical modelling, Faculty of Social and Behavioural Sciences, Utrecht University	2006	1 ECTS
Presentations		
Van der Horst K, Kremers S, Ferreira I, Singh A, Oenema A, Brug J. "Perceived parenting practices and parenting styles and adolescent soft drink consumption". Paper presented at the European Congress on Obesity, Athens, Greece (June 1-4)	2005	1 ECTS
Van der Horst K, Oenema A, Ferreira I, Brug J. "Environmental determinants of weight gain-related behaviours in youth". Paper presented at the sixth Conference on Psychology and Health, Kerkrade, the Netherlands (May 8-10)	2006	1 ECTS
Van der Horst K, Oenema A, Ferreira I, Wendel-Vos W, Giskes K, Van Lenthe F, Brug J. "Environmental correlates of obesity related behaviors in youth". Paper presented at the fifth conference of the International Society of Behavioral Nutrition and Physical Activity, Boston, USA (July 13-16)	2006	1 ECTS
Research Meeting, CPAN Deakin University, Melbourne. 'The ENDORSE Study'	2007	1 ECTS
Van der Horst K, Oenema A, Brug J. "Exploring environmental determinants of (in) activity in adolescents: the ENDORSE study". Paper presented at the European Health Psychology Society conference, Maastricht, the Netherlands. (August 15-18).	2007	1 ECTS
Van der Horst K, Oenema A, te Velde SJ, Brug J. "The influence of parenting styles and practices on adolescents' energy balance related behaviours in the Netherlands". Paper presented at the sixth conference of the International Society of Behavioral Nutrition and Physical Activity, Oslo, Norway (June 20-23)	2007	1 ECTS
Nederlands Congres Volksgezondheid, Groningen 'Frisdrank en snack consumptie bij jongeren: De invloed van voedingsmiddelen in schoolkantines & winkels in de schoolomgeving'	2008	1 ECTS
Van der Horst K, Oenema A, Brug J. "Are school physical activity policies and schoolyard facilities associated with sports and active commuting to school?" Paper presented at the seventh conference of the International Society of Behavioral Nutrition and Physical Activity, Banff, Canada. (May 21-24).	2008	1 ECTS

International Conferences

1			
2	Fourth conference of the International Society of Behavioral Nutrition and Physical Activity, Amsterdam, the Netherlands (July 16-18)	2004	1 ECTS
3	European Congress on Obesity, Athens, Greece (June 1-4)	2005	1 ECTS
4	Sixth Conference on Psychology and Health, Kerkrade, the Netherlands (May 8-10)	2006	1 ECTS
5	Fifth conference of the International Society of Behavioral Nutrition and Physical Activity, Boston, USA (July 13-16)	2006	1 ECTS
6			
7	Sixth conference of the International Society of Behavioral Nutrition and Physical Activity, Oslo, Norway (June 20-23)	2007	1 ECTS
8			
9	Seventh conference of the International Society of Behavioral Nutrition and Physical Activity, Banff, Canada. (May 21-24).	2008	1 ECTS
10			

2. Teaching activities

11			
12	Supervising Bachelor thesis	2007	15 hours
13	Curriculum Medical students, 2 nd year, Erasmus MC Rotterdam	2007	5 hours
14	Theme 2.2: 'Disorders in nutrition, metabolism and hormonal regulation'. Supervising students.		
15	Curriculum Medical students, 4 th year, Erasmus MC Rotterdam	2006	15 hours
16	Theme 4.2: 'The population as a patient'. Coordination and supervising students.	2007	20 hours
17			
18			
19			
20			
21			
22			
23			
24			
25			
26			
27			
28			
29			
30			
31			
32			
33			
34			
35			
36			
37			
38			
39			