

Hypertension

JOURNAL OF THE AMERICAN HEART ASSOCIATION



*Learn and Live*SM

Bradykinin Potentiation by Angiotensin-(1-7) and ACE Inhibitors Correlates With ACE C- and N-Domain Blockade

Beril Tom, René de Vries, Pramod R. Saxena and A.H. Jan Danser

Hypertension 2001;38;95-99

Hypertension is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75214

Copyright © 2001 American Heart Association. All rights reserved. Print ISSN: 0194-911X. Online ISSN: 1524-4563

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://hyper.ahajournals.org/cgi/content/full/38/1/95>

Subscriptions: Information about subscribing to Hypertension is online at <http://hyper.ahajournals.org/subscriptions/>

Permissions: Permissions & Rights Desk, Lippincott Williams & Wilkins, 351 West Camden Street, Baltimore, MD 21202-2436. Phone 410-5280-4050. Fax: 410-528-8550. Email: journalpermissions@lww.com

Reprints: Information about reprints can be found online at <http://www.lww.com/static/html/reprints.html>

Bradykinin Potentiation by Angiotensin-(1-7) and ACE Inhibitors Correlates With ACE C- and N-Domain Blockade

Beril Tom, René de Vries, Pramod R. Saxena, A.H. Jan Danser

Abstract—ACE inhibitors block B₂ receptor desensitization, thereby potentiating bradykinin beyond blocking its hydrolysis. Angiotensin (Ang)-(1-7) also acts as an ACE inhibitor and, in addition, may stimulate bradykinin release via angiotensin II type 2 receptors. In this study we compared the bradykinin-potentiating effects of Ang-(1-7), quinaprilat, and captopril. Porcine coronary arteries, obtained from 32 pigs, were mounted in organ baths, precontracted with prostaglandin F_{2α}, and exposed to quinaprilat, captopril, Ang-(1-7), and/or bradykinin. Bradykinin induced complete relaxation (pEC₅₀ = 8.11 ± 0.07, mean ± SEM), whereas quinaprilat, captopril, and Ang-(1-7) alone were without effect. Quinaprilat shifted the bradykinin curve to the left in a biphasic manner: a 5-fold shift at concentrations that specifically block the C-domain (0.1 to 1 nmol/L) and a 10-fold shift at concentrations that block both domains. Captopril and Ang-(1-7) monophasically shifted the bradykinin curve to the left, by a factor of 10 and 5, respectively. A 5-fold shift was also observed when Ang-(1-7) was combined with 0.1 nmol/L quinaprilat. Repeated exposure of porcine coronary arteries to 0.1 μmol/L bradykinin induced B₂ receptor desensitization. The addition of 10 μmol/L quinaprilat or Ang-(1-7) to the bath, at a time when bradykinin alone was no longer able to induce relaxation, fully restored the relaxant effects of bradykinin. Angiotensin II type 1 or 2 receptor blockade did not affect any of the observed effects of Ang-(1-7). In conclusion, Ang-(1-7), like quinaprilat and captopril, potentiates bradykinin by acting as an ACE inhibitor. Bradykinin potentiation is maximal when both the ACE C- and N-terminal domains are inhibited. The inhibitory effects of Ang-(1-7) are limited to the ACE C-domain, raising the possibility that Ang-(1-7) synergistically increases the blood pressure-lowering effects of N-domain-specific ACE inhibitors. (*Hypertension*. 2001;38:95-99.)

Key Words: angiotensin ■ bradykinin ■ angiotensin-converting enzyme inhibitors ■ receptors, bradykinin ■ coronary artery

Angiotensin (Ang)-(1-7) is a heptapeptide that is formed endogenously from both Ang I and Ang II.¹ In rats and dogs, Ang-(1-7) exerts direct vasodilatory effects via non-angiotensin II type 1 (AT₁), non-angiotensin II type 2 (AT₂) receptors, possibly by stimulating bradykinin and NO release.^{1,2} In contrast, in humans or pigs, no direct vasodilatory effects of Ang-(1-7) were observed,³⁻⁷ although Ang-(1-7) did antagonize the pressor effects of Ang II, suggesting that it may cause vasodilation indirectly, by acting as an AT₁ receptor antagonist.^{5,6} In addition, Ang-(1-7) potentiates bradykinin, either via an AT₂ receptor-dependent mechanism or through inhibition of ACE.^{3,5,8} The latter effect is not necessarily based on blockade of bradykinin hydrolysis, because recent studies have shown that ACE inhibitors, including Ang-(1-7), potentiate bradykinin by inhibiting desensitization of its receptor.⁹⁻¹¹ Somatic ACE has 2 homologous domains, each containing an active center. According to their position (N- or C-terminal), these domains are designated as the N- or C-domain, respectively. Interestingly, Ang-(1-7) inhibits the C-domain more potently than the N-domain (by 1 order of magnitude)⁸ and is cleaved to Ang-(1-5) by the N-domain.¹²

In the present study, we investigated the bradykinin-potentiating effects of Ang-(1-7) in porcine coronary arteries (PCAs), its dependency on ACE, and the possible involvement of AT₁ and/or AT₂ receptors. The effects of Ang-(1-7) were compared with those of quinaprilat and captopril, 2 ACE inhibitors with preference for the ACE C- and N-domains, respectively.^{13,14} We also verified the effect of Ang-(1-7) in human coronary arteries (HCAs).

Methods

Tissue Collection

HCAs were obtained from 4 "heart beating" organ donors (2 men and 2 women; age range, 14 to 38 years; mean ± SEM age, 23 ± 5 years) who died of noncardiac causes (1 subarachnoid bleeding, 3 head trauma) <24 hours before the heart was taken to the laboratory. Hearts were provided by the Rotterdam Heart Valve Bank (Bio Implant Services/Eurotransplant Foundation, Rotterdam, The Netherlands) after removal of the aortic and pulmonary valves for transplantation purposes. The study was approved by the joint ethics committee of Erasmus University Rotterdam and University Hospital Rotterdam. Immediately after circulatory arrest, the hearts were stored in an ice-cooled, sterile, organ-protecting solution. After

Received December 8, 2000; first decision January 8, 2001; revision accepted January 12, 2001.

From the Department of Pharmacology, Erasmus University Rotterdam, The Netherlands.

Correspondence to A.H.J. Danser, PhD, Department of Pharmacology, Room EE1418b, Erasmus University Rotterdam, Dr Molewaterplein 50, 3015 GE Rotterdam, The Netherlands. E-mail danser@farma.fgg.eur.nl

© 2001 American Heart Association, Inc.

Hypertension is available at <http://www.hypertensionaha.org>

arrival in the laboratory, the HCAs were removed and stored overnight in a cold, oxygenated Krebs' bicarbonate solution of the following composition (mmol/L): NaCl 118, KCl 4.7, CaCl₂ 2.5, MgSO₄ 1.2, KH₂PO₄ 1.2, NaHCO₃ 25, and glucose 8.3 (pH 7.4). Vessels were then cut into segments of ≈4 mm in length, suspended on stainless steel hooks in 15-mL organ baths containing Krebs' bicarbonate solution, aerated with 95% O₂/5% CO₂, and maintained at 37°C. Segments containing macroscopically visible atherosclerotic lesions were not used.

PCAs were obtained from 32 pigs age 2 to 3 months (Yorkshire×Landrace; weight, 10 to 15 kg). The pigs had been used in *in vivo* experiments studying the effects of α -adrenoceptor and serotonin receptor agonists and antagonists under pentobarbital (600 mg IV) anesthesia.¹⁵ The ethics committee of Erasmus University Rotterdam dealing with the use of animals for scientific experiments approved the protocol for this investigation. Hearts were explanted at the end of the experiment, and the coronary arteries were removed immediately and handled in the same way as HCAs.

Organ Bath Studies

All vessel segments were allowed to equilibrate for at least 30 minutes, and the organ bath fluid was refreshed every 15 minutes during this period. Changes in tissue contractile force were recorded with a Harvard isometric transducer. The vessel segments, stretched to a stable force of ≈15 mN, were exposed to 30 mmol/L K⁺ twice. The functional integrity of the endothelium was verified by observing relaxation to 1 nmol/L substance P after precontraction with 1 μ mol/L prostaglandin F_{2 α} (PGF_{2 α}).¹⁶ Subsequently, the tissue was exposed to 100 mmol/L K⁺ to determine the maximal contractile response to K⁺. The segments were then allowed to equilibrate in fresh organ bath fluid for 30 minutes. Thereafter, the following experiments were performed.

First, Ang-(1-7) concentration-response curves (CRCs) (0.1 nmol/L to 10 μ mol/L) were constructed in HCAs and PCAs, both at baseline and after precontraction with 10 μ mol/L PGF_{2 α} , in the absence or presence of the AT₁ receptor antagonist irbesartan (1 μ mol/L) or the AT₂ receptor antagonist PD 123,319 (1 μ mol/L).

Second, the effects of Ang-(1-7), quinaprilat, and captopril on bradykinin-induced vasorelaxation were studied in PCAs. Vessels were preincubated for 30 minutes in the absence or presence of Ang-(1-7) (10 pmol/L to 10 μ mol/L), quinaprilat (0.1 pmol/L to 0.1 mmol/L), captopril (0.1 pmol/L to 0.1 mmol/L), 10 μ mol/L Ang-(1-7)+1 μ mol/L irbesartan, 10 μ mol/L Ang-(1-7)+1 μ mol/L PD 123,319, 10 μ mol/L Ang-(1-7)+0.1 nmol/L quinaprilat, or 10 μ mol/L Ang-(1-7)+10 μ mol/L quinaprilat. Vessels were then precontracted with 10 μ mol/L PGF_{2 α} or 1 μ mol/L U46619, and CRCs to bradykinin (0.1 nmol/L to 1 μ mol/L) were constructed.

Third, the effect of quinaprilat and Ang-(1-7) on desensitized B₂ receptors was studied in PCAs. Vessels were preincubated for 30 minutes with or without 1 μ mol/L irbesartan or 1 μ mol/L PD 123,319. Vessels were then precontracted with 10 μ mol/L PGF_{2 α} and exposed 3 times to a concentration of bradykinin (0.1 μ mol/L) that is capable of inducing maximal relaxation. Each next exposure was started as soon as the effect of the previous exposure had disappeared, ie, after ≈15 minutes. After the third exposure, when bradykinin no longer exerted a vasodilatory effect, quinaprilat (10 μ mol/L) or Ang-(1-7) (10 μ mol/L) was added to the organ bath. Thereafter, when the effects of quinaprilat and Ang-(1-7) had disappeared, a fourth bradykinin dose (0.1 μ mol/L) was added to the organ bath.

Statistical Analysis

Data are given as mean±SEM and expressed as a percentage of the contraction in response to PGF_{2 α} or U46619. CRCs were analyzed by the logistic function described by de Lean et al¹⁷ to obtain pEC₅₀ (−¹⁰log EC₅₀) values, EC₅₀ representing the concentration at which 50% of the maximal relaxant effect has been reached. Statistical analysis was by ANOVA, followed by post hoc evaluation (according to Tukey or Dunnett where appropriate). Values of *P*<0.05 were considered significant.




Figure 1. Relaxations of PCAs, precontracted with 10 μ mol/L PGF_{2 α} or 1 μ mol/L U46619, to bradykinin in the absence (control) or presence of 10 μ mol/L Ang-(1-7), 10 μ mol/L quinaprilat, or 10 μ mol/L captopril (top). The effect of Ang-(1-7) on bradykinin-induced relaxations was also studied in the presence of 1 μ mol/L irbesartan or 1 μ mol/L PD 123,319 (bottom). Data (mean±SEM of 5 to 12 experiments) are expressed as a percentage of the contraction induced by PGF_{2 α} or U46619. Ang-(1-7), quinaprilat, and captopril significantly shifted the bradykinin CRC to the left (*P*<0.05 vs control for Ang-(1-7); *P*<0.01 vs control for the 2 ACE inhibitors), and the effect of Ang-(1-7) was not influenced (*P*=NS) by the presence of irbesartan or PD 123,319.

Results

Effect of Ang-(1-7) on HCAs and PCAs

Ang-(1-7), at concentrations up to 10 μ mol/L, did not exert a response in HCAs (*n*=4) and PCAs (*n*=5 to 12) at baseline or after precontraction with PGF_{2 α} or in the presence of irbesartan or PD 123,319.

Effects of Ang-(1-7), Quinaprilat, and Captopril on Bradykinin-Induced Relaxation in PCAs

Bradykinin caused complete relaxation of precontracted PCAs in a concentration dependent-manner (pEC₅₀=8.11±0.07; *n*=12; Figure 1, top panel). Quinaprilat (10 μ mol/L) and captopril (10 μ mol/L), like Ang-(1-7) at this concentration, did not exert any effect on precontracted PCAs. However, in the presence of 10 μ mol/L Ang-(1-7), the bradykinin CRC was shifted ≈5-fold to the left (pEC₅₀=8.72±0.09; *n*=12; *P*<0.05 versus control), whereas in the presence of 10 μ mol/L quinaprilat or 10 μ mol/L captopril the bradykinin CRC was shifted ≈10-fold to the left (pEC₅₀=9.03±0.21 and 8.91±0.09, respectively; *n*=8; *P*<0.01 versus control) (Figure 1, top panel). The effect of Ang-(1-7) on the bradykinin CRC was not affected by irbesartan or PD 123,319 (Figure 1, bottom panel).

To study the concentration dependency of the Ang-(1-7)- and ACE-inhibitor-induced leftward shifts, bradykinin CRCs




Figure 2. Change in $-\log(\text{EC}_{50})$ of the bradykinin CRC in the presence of increasing concentrations of quinaprilat (top), captopril (middle), or Ang-(1-7) (bottom). An increase in $-\log(\text{EC}_{50})$ represents a leftward shift of the bradykinin CRC. Data (mean \pm SEM of 6 to 8 experiments) were obtained in PCAs precontracted with 1 $\mu\text{mol/L}$ U46619. Con indicates control. Significant differences ($P < 0.05$) vs control were obtained for quinaprilat, captopril, and Ang-(1-7) at concentrations ≥ 0.1 , 100, and 1000 nmol/L, respectively.

were also constructed in the presence of a wide range of Ang-(1-7), quinaprilat, and captopril concentrations (Figure 2). The leftward shift caused by quinaprilat occurred in a biphasic manner, with a 5-fold shift at concentrations in the subnanomolar range and a 10-fold shift at concentrations > 1 nmol/L. This biphasic shift is in agreement with earlier studies¹³ demonstrating different quinaprilat binding affinities of the ACE C- ($K_d = 7$ pmol/L) and N-domains ($K_d = 1267$ pmol/L) and suggests that the 5- and 10-fold shifts represent C-domain inhibition and complete (ie, C- plus N-domain) ACE inhibition, respectively. The 10-fold leftward shift caused by captopril occurred monophasically (Figure 2), suggesting that captopril is bound with similar affinity by the 2 ACE domains. The 5-fold leftward shift caused by Ang-(1-7) also occurred monophasically (Figure 2), suggesting either that Ang-(1-7) inhibits 1 ACE domain only or that Ang-(1-7) potentiates bradykinin independently of its effects on ACE. To sort out the latter, the effect of 10 $\mu\text{mol/L}$ Ang-(1-7) on top of quinaprilat, either at a concentration that selectively inhibits the ACE C-domain (0.1 nmol/L) or at a concentration that inhibits both ACE domains (10 $\mu\text{mol/L}$), was studied. Ang-(1-7) affected neither the 5-fold leftward shift of the bradykinin CRC at 0.1 nmol/L quinaprilat ($\text{pEC}_{50} = 8.59 \pm 0.39$ and 8.68 ± 0.29 with and without Ang-




Figure 3. Relaxations of PCAs, precontracted with 10 $\mu\text{mol/L}$ $\text{PGF}_{2\alpha}$ or 1 $\mu\text{mol/L}$ U46619, to bradykinin in the absence (control, + symbols) or presence of 0.1 nmol/L quinaprilat (squares) or 10 $\mu\text{mol/L}$ quinaprilat (circles), with (closed symbols) or without (open symbols) 10 $\mu\text{mol/L}$ Ang-(1-7). Data (mean \pm SEM of 11 to 13 experiments) are expressed as a percentage of the contraction induced by $\text{PGF}_{2\alpha}$ or U46619. The addition of Ang-(1-7) on top of quinaprilat did not have a significant effect.

(1-7), respectively; $n = 13$; $P = \text{NS}$) nor the 10-fold shift at 10 $\mu\text{mol/L}$ quinaprilat ($\text{pEC}_{50} = 9.06 \pm 0.20$ and 9.03 ± 0.21 with and without Ang-(1-7), respectively; $n = 11$; $P = \text{NS}$) (Figure 3). Most likely, therefore, Ang-(1-7), at concentrations up to 10 $\mu\text{mol/L}$, acts as a selective inhibitor of the ACE C-domain.

Effect of Ang-(1-7) and Quinaprilat on Desensitized B_2 Receptors in PCAs

Repeated exposure of precontracted vessel segments to 0.1 $\mu\text{mol/L}$ bradykinin produced progressively smaller responses (Figure 4, top panel). The response to the third bradykinin dose was $< 50\%$ of the response to the first bradykinin dose. Quinaprilat or Ang-(1-7), added to the organ bath after the effect of the third bradykinin dose had disappeared, both at a concentration of 10 $\mu\text{mol/L}$, completely restored the relaxant effect of bradykinin ($90 \pm 10\%$ and $83 \pm 8\%$ relaxation, respectively; $n = 8$; $P = \text{NS}$). A fourth bradykinin dose, added after the effect of quinaprilat or Ang-(1-7) had disappeared, induced no further effect. The effect of Ang-(1-7) was not different in the presence of irbesartan or PD 123,319 (Figure 4, bottom panel).

Discussion

The present study shows that Ang-(1-7) potentiates the vasodilator effects of bradykinin in PCAs through inhibition of the ACE C-domain. AT_1 or AT_2 receptors are not involved in this effect, nor did Ang-(1-7) exert direct (ie, independently of bradykinin) relaxant effects in either HCAs or PCAs. Furthermore, the maximal potentiating effect of Ang-(1-7) was smaller than that of the ACE inhibitors quinaprilat and captopril, suggesting that full potentiation will only be obtained when both ACE domains are inhibited.

The concentrations of Ang-(1-7) required to obtain ACE C-domain inhibition (≥ 1 $\mu\text{mol/L}$) are in agreement with previously reported values.^{5,8} These concentrations exceed the in vivo tissue and plasma Ang-(1-7) concentrations in rats and humans by at least 4 orders of magnitude. Ang-(1-7)




Figure 4. Top, Original tracings of an experiment in which a PCA, after precontraction with 10 $\mu\text{mol/L}$ $\text{PGF}_{2\alpha}$, is subsequently exposed to 0.1 $\mu\text{mol/L}$ bradykinin (BK) (3 times), 10 $\mu\text{mol/L}$ quinaprilat (top), or 10 $\mu\text{mol/L}$ Ang-(1-7) (bottom), and 0.1 $\mu\text{mol/L}$ bradykinin. Bottom, Relaxations of PCAs, after precontraction with 10 $\mu\text{mol/L}$ $\text{PGF}_{2\alpha}$, to 3 consecutive bradykinin doses (0.1 $\mu\text{mol/L}$; BK1, BK2, BK3), 10 $\mu\text{mol/L}$ Ang-(1-7), and a fourth bradykinin dose (0.1 $\mu\text{mol/L}$; BK4) in the absence or presence of 1 $\mu\text{mol/L}$ irbesartan or 1 $\mu\text{mol/L}$ PD 123,319. Data (mean \pm SEM of 4 to 8 experiments) are expressed as a percentage of the contraction induced by $\text{PGF}_{2\alpha}$.

levels increase during ACE inhibition^{18–21} because ACE inhibitors block the ACE N-domain-mediated degradation of Ang-(1-7).^{1,8} However, the fact that most ACE inhibitors also block the ACE C-domain will mask any potential C-domain-blocking effect of the high Ang-(1-7) levels during ACE inhibition. Indeed, Ang-(1-7) did not enhance bradykinin-induced vasodilation in the forearm in human subjects treated with an ACE inhibitor.⁷ Therefore, the effects of Ang-(1-7) described in the present study may only be of physiological importance if sufficiently high concentrations of Ang-(1-7) are reached in the immediate vicinity of ACE (the levels measured per gram tissue do not exclude this possibility) and/or in combination with ACE inhibitors that specifically block the ACE N-domain.

Studies with the N-domain-specific substrate *N*-acetyl-ser-asp-lys-pro (AcSDKP) have revealed that the ACE inhibitor captopril inhibits AcSDKP hydrolysis 16 times more potently than Ang I hydrolysis.¹⁴ Because the ACE N- and C-domains exhibit similar catalytic activities toward Ang I,²² these data suggest that captopril, at low concentrations, preferentially inhibits the ACE N-domain. Using captopril, however, we were unable to make a clear distinction between the effects of ACE N- and C-domain inhibition on the relaxant effects of

bradykinin. In contrast, using the ACE inhibitor quinaprilat, which binds to the ACE C-domain with almost 200-fold greater affinity than to the ACE N-domain,¹³ we could make such a distinction: quinaprilat shifted the bradykinin CRC to the left in a biphasic manner, with an ≈ 5 -fold shift occurring at concentrations that selectively block the C-domain and an ≈ 10 -fold shift at concentrations that block both domains. Our inability to find a similar biphasic shift in the presence of increasing concentrations of captopril most likely relates to the limited selectivity of this ACE inhibitor for the ACE N-domain.

Ang-(1-7), like captopril, induced a monophasic leftward shift of the bradykinin CRC. At the highest Ang-(1-7) concentration (10 $\mu\text{mol/L}$) that was tested, however, the shift was only half of that caused by similar concentrations of quinaprilat or captopril. Adding quinaprilat, at a concentration that selectively blocks the ACE C-domain, to 10 $\mu\text{mol/L}$ Ang-(1-7) did not cause a further leftward shift, thereby demonstrating that Ang-(1-7), at least at this concentration, blocks the ACE C-domain only. Because a concentration of 10 $\mu\text{mol/L}$ is already many orders of magnitude above the measured concentrations of Ang-(1-7) in vivo,^{18–21} we did not evaluate whether even higher concentrations of Ang-(1-7) resulted in ACE N-domain inhibition.

Does ACE inhibition result in bradykinin potentiation by blocking its hydrolysis? Because bradykinin, like Ang I, is cleaved equally well by the 2 ACE domains,²² our finding that blockade of both domains results in a twice as large leftward shift as blockade of 1 domain appears to support this possibility. However, the long half-life of bradykinin in this preparation (30 minutes),³ as well as our previous finding in PCAs that the quinaprilat-induced leftward shift of the ACE-resistant bradykinin analogue [Hyp³-Tyr(Me)⁸]-bradykinin was not different from that of bradykinin,¹¹ argues against this concept. Moreover, under conditions of B₂ receptor desensitization (induced by repeated exposure to a concentration of bradykinin that causes maximal relaxation), when bradykinin itself was no longer active, both quinaprilat and Ang-(1-7) immediately restored the effect of bradykinin. Taken together, therefore, a more likely explanation for the Ang-(1-7)-, quinaprilat-, and captopril-induced bradykinin potentiation is the upregulation (or resensitization) of B₂ receptors that has recently been described in Chinese hamster ovary (CHO) cells transfected with the human B₂ receptor and human ACE.^{8,9,23,24} This effect only occurs when ACE and B₂ receptors are sterically closely associated, probably forming a heterodimer.²⁴ ACE inhibitors are believed to alter the heterodimer interaction, thereby promoting a conformation in the B₂ receptor that affects its sequestration and coupling to second messengers.²⁴ Our findings in intact PCAs show that ACE-B₂ receptor crosstalk is not limited to transfected CHO cells and, in addition, suggest that the conformational changes underlying bradykinin potentiation correlate directly with inhibition of the ACE C- and N-domains.

References

- Ferrario CM, Chappell MC, Tallant EA, Brosnihan KB, Diz DI. Counterregulatory actions of angiotensin-(1-7). *Hypertension*. 1997;30:535–541.

2. Brosnihan KB, Li P, Ferrario CM. Angiotensin-(1-7) dilates canine coronary arteries through kinins and nitric oxide. *Hypertension*. 1996;27:523–528.
3. Gorelik G, Carbini LA, Scicli AG. Angiotensin-(1-7) induces bradykinin-mediated relaxation in porcine coronary artery. *J Pharmacol Exp Ther*. 1998;286:403–410.
4. Pörsti I, Bara AT, Busse R, Hecker M. Corrigendum. *Br J Pharmacol*. 1996;117:231.
5. Roks AJM, van Geel PP, Pinto YM, Buikema H, Henning RH, de Zeeuw D, van Gilst WH. Angiotensin-(1-7) is a modulator of the human renin-angiotensin system. *Hypertension*. 1999;34:296–301.
6. Ueda S, Masumori-Maemoto S, Ashino K, Nagahara T, Gotoh E, Umemura S, Ishii M. Angiotensin-(1-7) attenuates vasoconstriction evoked by angiotensin II but not by noradrenaline in man. *Hypertension*. 2000;35:998–1001.
7. Davie AP, McMurray JJV. Effect of angiotensin-(1-7) and bradykinin in patients with heart failure treated with an ACE inhibitor. *Hypertension*. 1999;34:457–460.
8. Deddish PA, Marcic B, Jackman HL, Wang H-Z, Skidgel RA, Erdös EG. N-domain-specific substrate and C-domain inhibitors of angiotensin-converting enzyme: angiotensin-(1-7) and keto-ACE. *Hypertension*. 1998;31:912–917.
9. Minshall RD, Tan F, Nakamura F, Rabito SF, Becker RP, Marcic B, Erdös EG. Potentiation of the actions of bradykinin by angiotensin I-converting enzyme inhibitors: the role of expressed human bradykinin B₂ receptors and angiotensin-converting enzyme in CHO cells. *Circ Res*. 1997;81:848–856.
10. Benzing T, Fleming I, Blaukat A, Müller-Esterl W, Busse R. Angiotensin-converting enzyme inhibitor ramiprilat interferes with the sequestration of the B₂ kinin receptor within the plasma membrane of native endothelial cells. *Circulation*. 1999;99:2034–2040.
11. Danser AHJ, Tom B, de Vries R, Saxena PR. L-NAME-resistant bradykinin-induced relaxation in porcine coronary arteries is NO-dependent: effect of ACE inhibition. *Br J Pharmacol*. 2000;131:195–202.
12. Chappell MC, Pirro NT, Sykes A, Ferrario CM. Metabolism of angiotensin-(1-7) by angiotensin-converting enzyme. *Hypertension*. 1998;31:362–367.
13. Perich RB, Jackson B, Johnston CI. Structural constraints of inhibitors for binding at two active sites on somatic angiotensin converting enzyme. *Eur J Pharmacol*. 1994;266:201–211.
14. Michaud A, Williams TA, Chauvet M-T, Corvol P. Substrate dependence of angiotensin I-converting enzyme inhibition: captopril displays a partial selectivity for inhibition of *N*-acetyl-seryl-aspartyl-lysyl-proline hydrolysis compared with that of angiotensin I. *Mol Pharmacol*. 1997;51:1070–1076.
15. de Vries P, Willems EW, Heiligers JPC, Villalón CM, Saxena PR. Investigation of the role of 5-HT_{1B} and 5-HT_{1D} receptors in the sumatriptan-induced constriction of porcine carotid arteriovenous anastomoses. *Br J Pharmacol*. 1999;127:405–412.
16. MaassenVanDenBrink A, de Vries R, Saxena PR, Schalekamp MADH, Danser AHJ. Vasoconstriction by in situ formed angiotensin II: role of ACE and chymase. *Cardiovasc Res*. 1999;44:407–415.
17. de Lean A, Munson PJ, Rodbard D. Simultaneous analysis of families of sigmoidal curves: application to bioassay, radioligand assay, and physiological dose-response curves. *Am J Physiol*. 1978;235:E97–E102.
18. Campbell DJ, Kladis A, Duncan A-M. Nephrectomy, converting enzyme inhibition, and angiotensin peptides. *Hypertension*. 1993;22:513–522.
19. Lawrence AC, Evin G, Kladis A, Campbell DJ. An alternative strategy for the immunoassay of angiotensin peptides using amino-terminal-directed antisera: measurement of eight angiotensin peptides in human plasma. *J Hypertens*. 1990;8:715–724.
20. Campbell DJ, Duncan A-M, Kladis A. Angiotensin-converting enzyme inhibition modifies angiotensin but not kinin peptide levels in human atrial tissue. *Hypertension*. 1999;34:171–175.
21. Yamada K, Iyer SN, Chappell MC, Ganten D, Ferrario CM. Converting enzyme determines plasma clearance of angiotensin-(1-7). *Hypertension*. 1998;32:496–502.
22. Jaspard E, Wei L, Alhenc-Gélas F. Differences in the properties and enzymatic specificities of the two active sites of angiotensin I-converting enzyme (kininase II): studies with bradykinin and other natural peptides. *J Biol Chem*. 1993;268:9498–9503.
23. Marcic B, Deddish PA, Jackman HL, Erdös EG. Enhancement of bradykinin and resensitization of its B₂ receptor. *Hypertension*. 1999;33:835–843.
24. Marcic B, Deddish PA, Skidgel RA, Erdös EG, Minshall RD, Tan F. Replacement of the transmembrane anchor in angiotensin I-converting enzyme (ACE) with a glycosylphosphatidylinositol tail affects activation of the B₂ bradykinin receptor by ACE inhibitors. *J Biol Chem*. 2000;275:16110–16118.