

Water Beetles as Models in Ecology and Evolution

David T. Bilton^{1,*}, Ignacio Ribera² and Andrew Edward Z. Short³

¹ Marine Biology and Ecology Research Centre, School of Biological and Marine Sciences, University of Plymouth, Plymouth PL4 8AA, UK; email:

d.bilton@plymouth.ac.uk

²Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra), Passeig Marítim de la Barceloneta 37-49, 08003 Barcelona, Spain; email: ignacio.ribera@ibe.upf-csic.es

³Department of Ecology and Evolutionary Biology; and Division of Entomology, Biodiversity Institute, University of Kansas, Lawrence, Kansas 66045, USA; email:

aezshort@ku.edu

*Corresponding author

Keywords

Coleoptera, habitat shifts, model organisms, biogeography, sexual selection, indicator taxa

Abstract

Beetles have colonized water many times during their history, some of these events involving extensive evolutionary radiations and multiple transitions between media. With over 13,000 described species, they are one of the most diverse macroinvertebrate groups in most non-marine aquatic habitats, on all continents except Antarctica. A combination of wide geographical and ecological range, together with relatively accessible taxonomy make these insects an excellent model system for addressing a range of questions in ecology and evolution. Work on water beetles has recently made important contributions to fields as diverse as DNA taxonomy, macroecology, historical biogeography, sexual selection and conservation biology as well as predicting organismal responses to global change. Aquatic beetles have some of the best resolved phylogenies of any comparably diverse insect group, and this, coupled with recent advances in taxonomic and ecological knowledge are likely to drive an expansion of studies in the future.

INTRODUCTION

Beetles are an evolutionary success story *par excellence*, being by far the most speciose order of animals on earth. The Coleoptera are an old radiation, whose evolutionary origins may date back to the Permian or even Carboniferous (148, 161). Of the features that underpin the evolutionary success of beetles, the presence of elytra is probably the single most important trait facilitating their colonization of aquatic habitats as adults, a lifestyle almost unknown in other holometabolous insects. As well as protecting the hindwings, the elytra enclose a subelytral air store, fundamental to gas exchange in almost all water beetle adults (30). At least 23 beetle families, from three of the four extant suborders, are predominantly aquatic as adults, larvae or both (see **Figure 1**). Additionally, members of a number of other families are aquatic, or associated with water during at least one life-history stage, making the precise definition of a 'water beetle' somewhat subjective (75, 131). What is clear is that water beetles are an ecological guild rather than a clade, each aquatic colonization representing an independent transition between media, some giving rise to evolutionary radiations across the entire spectrum of inland waters. With more than 13,000 described species, water beetles are abundant and ecologically important in almost all non-marine aquatic habitats, from the smallest phytotelmata to larger lakes and rivers, on all continents except Antarctica. Their wide geographical and ecological range, together with a relatively stable and accessible taxonomy means that these insects are excellent models for addressing a range of ecological and evolutionary questions. Here we explore how water beetles have contributed to a range of disciplines, from macroecology to biogeography, sexual selection and conservation biology.

EVOLUTION AND MACROECOLOGY

Terrestrial-Aquatic Transitions

Amongst the insects, only Diptera have crossed the aquatic-terrestrial boundary as many times as Coleoptera (75). There are at least eight major transitions, and the actual number may be more than twice that (**Figure 1**) (131), with four of these resulting in radiations greater than 1,000 species. Not all aquatic beetles share the same ecology or evolutionary pathway to aquatic adaptation. In most, both the adult and larvae are aquatic. In others, only the larvae (Scirtidae) or adult (some Dryopidae and Helophoridae) live in water. Additionally, a number of water beetle lineages have experienced one or more secondary transitions back to terrestriality, particularly the Hydrophilidae and Hydraenidae (44, 103, 133, 144). The impact of these aquatic-terrestrial transitions on the evolutionary trajectories of lineages are not yet fully understood. In Hydrophilidae, habitat shifts were not themselves linked to changes in diversification rate, although increased habitat breadth may play a role (24). In other cases, transitions to terrestriality seem to be the result of unusual or idiosyncratic conditions, such as a rapid shift from streams to tree moss on Oceanic Islands (135) or to forest litter (144). Hygropetric habitats, recently revealed to support diverse water beetle assemblages (111, 134), may be important 'stepping-stones' between media.

From Freshwater to Salt

Water beetles, together with some Diptera and Heteroptera, are amongst the few insects able to tolerate hypersaline waters, with concentrations up to more than 200 gr/l. This tolerance has developed independently –and recurrently– in several lineages, mostly Hydrophilidae (e.g. *Berosus*, *Enochrus*, *Paracymus*), Hydraenidae (*Ochthebius*), and Dytiscidae (some species of *Nebrioporus*, *Hygrotus* and *Boreonectes*). Hypersaline habitats are usually coastal, most commonly saltmarshes or rockpools, but they may also be found inland, with saline streams forming one of the most unusual aquatic environments (87). The evolution of salinity tolerance has been studied in detail in the *Enochrus* subgenus *Lumetus*, in which the evolutionary origin of salinity tolerance was associated with geological periods of increased aridification (10). There were also direct and relatively fast transitions between freshwater and hypersaline habitats, suggesting species may have evolved salinity tolerance before actually occupying saline habitats. Salinity tolerance may thus represent an exaptation from an adaptation to aridity, favored by a similar physiological mechanism. This hypothesis has been corroborated by further studies of physiological tolerance to salinity and desiccation (25, 96–100). Data on the evolution of salinity tolerance in other groups of Coleoptera are scanty, but the association between the origin of salinity tolerance and periods of aridification seems to be recurrent (157). Another common pattern is the evolutionary conservation of habitat preferences. Despite multiple origins of salinity tolerance, species typical of saline habitats tend to cluster together in phylogenies more than expected given their morphological similarities (97, 121, 157).

Going Underground

Given the global extent of subterranean waters (54) it is not surprising that aquatic beetles have colonized these systems multiple times, across five families in various parts of the world (62, 91, 138, 158, 159). Prior to 2000, most publications on subterranean water beetles were isolated species descriptions. Since then it has been recognized that the subterranean dytiscid fauna of Australia is astoundingly rich, with around 100 species now known (37, 82). Australia's extensive subterranean environments, present for long periods of evolutionary time, may mean that this hypogean is uniquely diverse. However, other areas with subterranean faunas that share similar paleogeography, including parts of southern Africa and India remain largely uninvestigated (58). Most Australian taxa are associated with calcrete aquifers in Western Australia, where Miocene-Pliocene aridity (10–5 million years ago) has driven the colonization of groundwaters by previously epigeal beetles (82; 142). In all cases, described species are restricted to a single calcrete, these aquifers forming an archipelago of biologically isolated systems associated with paleodrainages (37, 82). Calcrete aquifers support one-three endemic dytiscids, their colonization representing independent experiments in evolution (79). Coexisting species are always morphologically different from each other, with a relatively consistent average body-size ratio of 1.6 between them, despite marked differences in absolute beetle body sizes across calcretes – these insects providing one of the few empirical examples of self-organizing limiting similarity in nature (129, 156). In most aquifers, species derive from different ancestral lineages (82) suggesting that speciation occurred in allopatry. Eleven calcretes are so far known to contain sister species, however, which may have diverged locally (79). Whether the speciation of such taxa was truly sympatric, remains unclear, as physicochemical conditions, including salinity, do vary across many individual calcretes (73), and population

genetic structure has been observed in a number of taxa within aquifers (59). In at least one case stable isotope analysis has revealed differential prey specialization across a triplet of sister species (26), an observation consistent with sympatric divergence through resource partitioning.

Most subterranean Australian taxa were originally described in new, entirely hypogean, genera on the basis of their derived morphology, although subsequent molecular analyses have shown that they instead nest within genera with surface-dwelling representatives, mostly *Paroster* and *Limbodessus* (12, 81). As with the evolution of salinity tolerance, transitions to the subterranean environment are concentrated in a few lineages, for unknown reasons; in the northern hemisphere, most stygobiont species are dytiscids of the subtribe Siettitina (91, 92). The diversification dynamics of Australian *Paroster* suggest an early burst of speciation, which, together with the high degree of morphological diversity seen in the genus, is consistent with an adaptive radiation in groundwaters (142). However, the overall speciation and extinction rates do not appear to differ between surface and underground lineages (142).

The regressive evolution of features, particularly eyes, in subterranean taxa, has long intrigued evolutionary biologists. There remains limited empirical evidence that eye regression is associated with the loss of protein-coding gene function, however. A comparison of Australian subterranean species with their fully eyed surface-dwelling relatives revealed that the eye pigment gene *cinnabar* behaves in a pseudogene-like manner in hypogean taxa (80). Two out of three species investigated further showed a complete loss of opsin transcription, consistent with neutral, regressive evolution. The remaining species retained transcription of a long-

wavelength opsin orthologue, despite living in an aphotic environment, which may indicate a novel pleiotropic role or an early stage of pseudogene development (139).

Macroecology and Range Size

Understanding the drivers of geographical range size forms an important research focus in both macroecology (52), and its more recent offshoot macrophysiology (53). Most multicellular organisms have relatively limited geographical ranges, and relative range sizes typically vary considerably within clades, the majority being restricted whilst a few are much more widespread (52). Various explanations have been put forward to account for such observations, including differences in evolutionary age (52) dispersal ability (18) and fundamental niche breadth (27). Few empirical studies have explored these questions within a comparative framework, however.

Deronectes diving beetles are one of the first and best-studied model systems in this regard, range sizes in these Palearctic stream-dwellers varying from point endemics to species present throughout most of Europe. Multiple regression models show that thermal range – the absolute number of degrees centigrade between a species mean upper and lower thermal limits – is a strong predictor of both latitudinal range extent and relative latitudinal position, and always more important than relative dispersal ability (33). This result remained when controlling for phylogeny, suggesting limited influence of evolutionary age – perhaps unsurprising in a clade whose ranges have been shaped by Pleistocene climatic events which post-date their evolutionary origin (49). Differences in thermal performance may themselves be linked to metabolic plasticity (35) and setal tracheal gill densities (78, 155) in the

genus, and in both *Deronectes* and *Ilybius* diving performance has also been observed to differ between widespread and restricted taxa (34).

A strong association between thermal biology and range parameters is also seen in the *Agabus brunneus* group (Dytiscidae) (31, 67), where differences in the expression of proteins involved in energy metabolism and molecular chaperones (including heat shock proteins) have been observed between widespread and more restricted congeners (64, 65); pointing to a possible mechanistic basis behind differences in whole-organism performance. Whether physiological features of widespread species arose prior to (and therefore facilitated) their range expansion, or appeared afterwards, as a result of local adaptation, remains incompletely understood. The fact that broad physiological tolerances have been observed in single populations (31, 33) together with the genetic and ecological uniformity of many widespread taxa and their relatively recent origins (49) point toward the former explanation, however. In the temperate latitudes occupied by these beetles, widespread species are those which have been successful at expanding their geographical ranges in the Holocene, meaning that the location of a species ice-age refugium may also influence its present range size and position (50, 51).

Going with the Flow?

One of the most readily observable habitat preferences in aquatic Coleoptera (and macroinvertebrates in general) is whether they occupy running (lotic) or standing (lentic) waters. Species are usually found in one of these two broad types, very few in both. These contrasting preferences occur at all taxonomic levels, between families (e.g. Elmidae are almost exclusively lotic), genera, or even closely related species

(e.g. *Ochthebius notabilis* (Hydraenidae) or *Nebrioporus ceresyi* (Dytiscidae) groups – see 2). Studies of Iberian aquatic Coleoptera first noted that lentic species had much larger geographical range sizes than their lotic relatives (116). These differences were related to the contrasting geological stability of the two habitat types: species in more geologically ephemeral lentic waters are forced to disperse when the habitat disappears, whilst in more long-lived lotic systems species can maintain local populations for longer. Consequently, lentic species were predicted to have higher dispersal abilities and inter-population gene flow, resulting in larger, more dynamic geographical ranges and slower evolutionary turnover. Lotic species, in contrast, would tend to have lower dispersal abilities, with higher persistence of local populations and reduced inter-population gene flow. Latitudinal diversity gradients of lentic and lotic species may also be expected to differ, as lotic species will be more dependent on historical factors and distance to glacial refugia, whereas lentic species will be closer to an equilibrium with current ecological and geographical conditions (109). It must be noted, however, that multiple factors influence dispersal ability in addition to habitat stability (18), so macroecological patterns related to habitat type will manifest as statistical trends, exceptions always being possible (e.g. 132). It is also obvious that the lotic-lentic divide is a simplification of the complexity of freshwater habitats, which could be further subdivided (e.g. 74). Despite these limitations, most of the predictions of (116) have proved accurate, mostly with data from aquatic Coleoptera but also other aquatic groups (e.g. 71, 84). Differences in geographical range between lotic and lentic species are not restricted to Western Europe (38, 70) and can be associated with dispersal ability rather than ecological tolerance (8). The *Enochrus bicolor* complex includes eight Palearctic species, all of which occupy saline habitats (11). Widespread lentic species had significantly larger wings, relative to body

size, than restricted lotic relatives, but running and standing water taxa differed little in thermal physiology (8).

The higher mobility of lentic species has apparently resulted in a faster recolonization of formerly glaciated areas in the Northern hemisphere, and a closer equilibrium with ecological conditions, whilst lotic species show a stronger dependence on latitude (115, 127). Differences in habitat stability are reflected in gene flow between populations (2, 84), although it is still not clear how these differences affect net diversification rates of lineages over longer evolutionary timescales (40, 41, 109, 112).

BIOGEOGRAPHY AND DIVERSIFICATION

Because most water beetle lineages arose prior to the breakup of Gondwana (**Figure 1**), they have been proven useful in providing a comparative model for the biogeographic consequences of Mesozoic vicariance events as well as more recent episodes of Cenozoic colonization and diversification. Australia-Oceania and the Western Palearctic have been the primary foci for recent regional-scale biogeography and diversification studies, with Madagascar and the Neotropics having been studied to a lesser degree. The Nearctic and Oriental regions have been largely overlooked to date.

Gondwanan Fragmentation

The separation of Africa and South America has been implicated in the diversification of several water beetle groups, including the dytiscid tribe Aciliini (28)

and the hydrophilid tribe Hydrophilini (141). The separation of Madagascar and India has been linked to the distributional pattern of hydrophilid cascade beetles in those regions (143). Gondwanan fragmentation may have played a role in the diversification of the Hydrobiuisini (Hydrophilidae) (150) although the lineage likely originated in Laurasia. Additionally, there are a number of water beetle lineages that show a classical austral disjunction between southern South America, Australia/New Zealand, and/or South Africa including the hydrophilid Cylominae (43), and Copelatinae (Dytiscidae) (22).

Africa and Madagascar

No studies have focused on the internal biogeography of continental Africa, although several have examined the phylogenetic placement of newly discovered endemic lineages such as South African *Aspidytes* (Aspidytidae) (111, 140) and *Capelatus* (Dytiscidae) (22). In contrast, Madagascar has been a particular focus for biogeographical and diversification studies. The current fauna has been shown to be a mix of ancient endemics (57), Cenozoic colonizers, as well as perhaps serving as a source of lineages back-dispersing to mainland Africa (29), a finding which should be tested with wider taxon sampling. A faunistic link between Madagascar and the South African Cape has been highlighted in dytiscids (110) and Hydraenidae (102).

The Neotropics

Our knowledge of the Neotropical fauna, and particularly that of tropical South America, has grown substantially in the last twenty years. The region harbors

substantial deep phylogenetic diversity and recent studies have begun to unravel its distribution and origins. In Platynectinae diving beetles, the Andean and Guiana/Brazilian Shield faunas represent separate colonizations, which have diversified independently (145, 149). This phylogenetic separation between lineages found in the Andes and eastern Shield regions is implied in a number of other groups, including Hydroscaphidae (134). .Additionally, the hypothesis that South America has served as a reservoir of diversity which has repeatedly dispersed northwards into Central America, North America, and the Caribbean has been supported in a variety of water beetles (39, 141, 28).

Australia and Oceania

The Cenozoic diversification of diving beetles across Australia and Oceania is one of the best-studied water beetle systems outside the Palearctic. Phylogenetic analyses of Australian Hydroporini reveal that eastern Australia was likely the ancestral source of the lineage when it began to diversify about 27 mya, and that the ongoing aridification of the continent has led to rampant extinction, as well as excursions underground (144, 146, 147). Recent radiations of Copelatinae (Dytiscidae) in New Guinea and New Caledonia are the result of repeated dispersal from Australian ancestors during the Miocene (14). There have been few incidents of back-dispersal or inter-island mixing, with the notable exception of an incredible “supertramp” species in the genus *Rhantus* (Dytiscidae), which originated in New Guinea and has dispersed as far as the Azores and New Zealand (13, 147).

Western Palearctic

There has been substantial progress in recent years towards understanding the origins of the Mediterranean – and more widely western Palearctic – water beetle fauna. Two general patterns emerge regarding the origin of Mediterranean endemics and the role of Pleistocene glacial cycles in shaping current faunas.

Virtually all Western Palearctic water beetles with restricted distributions are Mediterranean. The most ancient have been estimated to date from the Miocene, and these are mostly restricted to the Iberian and Anatolian peninsulas (49, 66, 114), in some cases forming two reciprocally monophyletic sister lineages (e.g. *Hydrochus* (Hydrochidae) and *Deronectes*). There are no ancient endemics of such genera in mainland Italy and most of the Balkans, resulting in an east-west disjunction that has been long recognized biogeographically. The likely explanation for this disjunction is that most of the Italian and Balkan peninsulas remained submerged until the Pliocene (106), meaning that species of these genera endemic to mainland Italy south of the Alps and southern Greece are of Plio-Pleistocene origin. In these areas there are, however, many endemics in taxa with an abundance of recent species, such as *Limnebius* and *Hydraena* (Hydraenidae) (118, 151, 152).

The availability of comprehensive phylogenies and phylogeographies of various groups of water beetles in Western Europe and the Mediterranean region has revealed the complex role of Pleistocene glaciations in shaping current faunas. There are examples of the classic pattern of recolonization of recently deglaciated areas from the Mediterranean peninsulas (26, 63), mostly from populations at the northern edge of southern refugia. This implies that southern endemic species remained localized (e.g. *Deronectes* (49) and some *Hydraena* (113, 151)). These southern endemics, in some cases of Pleistocene origin and sister to species with

northern distributions (e.g. 117), may not have undergone significant range movements during their entire evolutionary histories, never colonizing areas directly affected by glaciations (1, 3, 113). In central and northern Europe the fauna is dominated by widespread lentic species with good dispersal abilities (115, 127), or lotic species that expanded their ranges in a short temporal window with favorable conditions after the last glaciation (50, 113). These cycles of range expansion with subsequent fragmentation and speciation (similar to refuge speciation (95) or vicariance by niche conservatism (160)) may have acted as a “species pump”, contributing substantially to current diversity.

SEXUAL SELECTION IN WATER BEETLES

Sexual selection occurs when changes in trait frequency result from differential reproductive success between individuals. This includes intrasexual competition, as well as intersexual interactions such as mate choice (cryptic or otherwise) and forced matings (136). Sexual selection is implicated in the evolution of many complex traits, including insect genitalia (72, 137), and water beetle mating systems are complex and varied, but best known to date in Dytiscidae. As with most insect groups, male genitalia often provide the primary means of differentiating closely related water beetle taxa (e.g. 91, 102, 103), and in some cases reach striking levels of complexity, particularly in the Hydraenidae (see **Figure 2**), where the precise homology of some structures remains unclear (118). In *Limnebius*, changes in body size appear to driven trends in genital evolution; reduced body size in the subgenus *Bilimnius* being accompanied by shrinkage and simplification of male genitalia, and several independent increases in body size in *Limnebius* s. str. associated with

larger, more complex genitalia (118, 119). There is also evidence for Rensch's rule in this genus, where male body size is more evolutionarily labile than female (120). The fact that greater sexual size dimorphism in *Limnebius* is not associated with more strongly developed secondary sexual characters suggests that the increased variation in male body size may have more to do with the lack of constraints associated with egg development and reproduction, than with directional sexual selection (120).

Given differential gametic investment, the evolutionary interests of the two sexes often diverge, particularly when mating is relatively costly (136), leading to sexual conflict, which can drive evolutionary "arms-races" between males and females and result in extreme sexual dimorphism (101). Most male dytiscids have modified, sucker-like articulo-setae on the tarsi of their fore and middle legs (see **Figure 2**), which increase their ability to grasp females during mating (7). In contrast, females of many species have enhanced dorsal sculpture, which reduces male grasping ability during pairing (76). Such female sculpture results from either a modification of existing surface reticulation (e.g. 19, 21) or the evolution of novel structures, including ridges and furrows or macroscopic granules on the elytra and irregular sculpture and hair-filled pits on the pronotum, in areas where male tarsi attach during mating initiation (**Figure 2**). Such traits are common in the larger Dytiscinae, where pre-insemination sexual conflict dominates a sexual system characterized by long pairings, post-inseminatory mate-guarding (sometimes with the production of mating plugs), and vigorous attempts by females to dislodge males, particularly at the onset of pair formation (90). Although not quantified to date, such pairings are likely to impart a greater energetic cost to females than males, particularly since males restrict females' access to air when surfacing, in an apparent

attempt to manipulate mating success (7, 90). Pairing duration and mating behavior in other water beetles are poorly known, but sexual dimorphism consistent with sexual conflict has been reported in a number of other dytiscid groups (19, 21) and Haliplidae (107). Male attachment devices seen in Gyrinidae and some Hydrophilidae may also indicate sexual conflict, although to date these have not been investigated.

Some dytiscids are also intrasexually dimorphic, with two forms of female differing in their resistance traits; some being rough, others smooth, like males (89, 90). Differences in female resistance appear to drive the evolution of counter-modifications in male attachment devices, with non-random mating between male and female morphs leading to linkage disequilibrium between male and female traits and the coexistence of morphs through negative frequency dependent selection (60, 61). In some species there is marked geographical variation in the relative frequencies of rough and smooth female morphs, although the drivers of these distributions remain poorly understood (77). Sometimes, rough and smooth female populations are allo/parapatric, and associated with males differing in the extent of development of tarsal attachment suckers (17, 21). Differential mating success may drive observed changes in the geographical position of such contact zones (17).

In addition to pre-insemination conflict, the Dytiscidae show great variation in female reproductive tract morphology (88), particularly in the subfamily Hydroporinae. Female hydroporine tracts frequently feature long, convoluted spermathecal and fertilization ducts, as well as other modifications (88), all of which point to the occurrence of sperm selection by females (23, 93). Dytiscid sperm morphology is varied and complex, again particularly in the Hydroporinae, where complex sperm conjugations and heteromorphisms have been reported (**Figure 2**)

(68, 69). The evolution of sperm and female reproductive tract features are correlated across the family, consistent with strong sexual selection (68, 69). The role of morphologically complex spermatophores in this process (130) remains unknown, but these structures may play a role in sperm delivery and positioning as well as functioning as mating plugs.

BIODIVERSITY, CONSERVATION, AND GLOBAL CHANGE

Water beetles have great potential for biodiversity and conservation assessment of inland water habitats, a number of features making them an excellent indicator group. These include high species richness, wide ecological/habitat range, high functional diversity (reflecting multiple aquatic colonizations), relative ease of sampling (at least as adults), and the fact that they are relatively well known taxonomically and biogeographically (47, 75, 108, 131). Being well known taxonomically, they have featured heavily in attempts to explore the effectiveness of DNA taxonomy and species delineation methods (e.g. 15, 94). In addition, taxa vary considerably in both their degree of ecological specialization, and dispersal abilities, with some species being reliable indicators of water quality (48, 86) or long-term habitat stability (e.g. 45, 46). To date, most detailed applications have been in Europe (47, 55, 56, 104, 123), studies in other regions largely concerned with the diversity and conservation of the insects themselves. Beetles are also effective surrogates of wider macroinvertebrate diversity (20, 56, 122), reflecting patterns in both species richness and compositional similarity in the wider aquatic community. This surrogate effect has been widely used to address a range of conservation-related questions, particularly in southwest Europe. These include the setting of

regional conservation priorities (4), area selection (123), the effectiveness of protected area networks (5, 56), sampling bias in environmental datasets (124, 125) and species distribution models (126), the influence of surrounding land cover on aquatic assemblages (42) and the importance of conserving the evolutionary history of a group (6).

Recent studies of ecophysiology and geographical range size in water beetles have provided insights into relative vulnerability to global change, being examples of insect conservation physiology (9, 11). The distinction can be made between species with high persistence ability, and those more likely to shift distribution, these two groups requiring different conservation approaches (9). *Deronectes* diving beetles are weak dispersers (16), whose occupied ranges bear little resemblance to those predicted by species distribution models based on macroclimate (128). In these beetles, geographically restricted southern endemics are more vulnerable to climate warming than their widespread congeners, due to limited heat tolerance and thermal plasticity (32, 33). Global change is reducing the extent of suitable habitat in Mediterranean mountains (32), placing such taxa in double jeopardy. In reality, global change involves multiple stressors operating synergistically. In inland waters, these include increased temperatures and hypoxia (154), the latter resulting both from eutrophication and increased metabolic demand at high temperatures (153). Recent work suggests that gas exchange mechanism, and the degree to which individuals can regulate internal oxygen levels, are good predictors of vulnerability to the combined effects of rising temperature and hypoxia, plastron-breathing elmids being much more strongly affected than surface exchanging dytiscids (154).

FUTURE ISSUES AND DIRECTIONS

1. Water beetle families (particularly Dytiscidae, Hydrophilidae, Hydraenidae) will be amongst the first diverse insect groups for which phylogenies with an almost complete taxon sampling could be available, allowing us to accurately explore the diversification processes that have shaped the biodiversity of lineages and regions.

2. Genomic/transcriptomic data will soon allow us to resolve difficult nodes in phylogenies (e.g. the status of Hydradephaga) *and* to explore the mechanistic bases of morphological and physiological adaptations. How are convergences between lineages at the phenotypic scale (e.g. complex antennal modifications used for gas exchange in Hydrophiloidea and Hydraenidae, similarities in thermal physiology or adaptations to extreme salinity and the subterranean environment) reflected at the genomic level? Such approaches could also explore convergences in the sensory apparatus associated with aquatic colonization and the level at which these changes have occurred - deep in the reception mechanism, or only in the structures that receive the stimulus?

3. How general are the relationships between physiology and geographical range size revealed in Palearctic water beetles? There is an urgent need for similar comparative studies in tropical and southern temperate regions.

4. Most studies of water beetle sexual systems to date have focused on a limited number of diving beetle taxa. To understand the drivers of sexual conflict, further comparative studies are needed, both within the Dytiscidae and in other water beetle

families. The emergence of larger, more robust phylogenies means that such studies can be conducted within a sound evolutionary framework.

5. Water beetles are excellent surrogates of aquatic biodiversity. To date, however, their use as indicator taxa has been largely restricted to Europe. Ongoing improvements to taxonomic, ecological, and biogeographical knowledge in other regions will make the wider development of water beetles in habitat assessment a realistic possibility in the near future.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGEMENTS

We are grateful to Steven Routledge for supplying fresh *Acilius* specimens for imaging, and to Jiří Hájek and Ján Kodada for providing some of the habitus images used in Figure 1. Dawn Higginson kindly supplied the sperm images used in Figure 2. DTB is grateful to the Leverhulme Trust for initially funding work on water beetle macroecology. IR acknowledges funding from grant CGL2013-48950-C2-1-P (AEI/FEDER, UE).

LITERATURE CITED

1. Abellán P, Benetti CJ, Angus RB, Ribera I. 2011. A review of Quaternary range shifts in European aquatic Coleoptera. *Global Ecol. Biogeogr.* 20:87–100
2. Abellán P, Millán A, Ribera I. 2009. Parallel habitat-driven differences in the phylogeographical structure of two independent lineages of Mediterranean saline water beetles. *Mol. Ecol.* 18:3885–902
3. Abellán P, Ribera I. 2017. Using phylogenies to trace the geographic signal of speciation. *J. Biogeogr.* 44:2236–46
4. Abellán P, Sánchez-Fernández D, Velasco J, Millán A. 2005. Assessing conservation priorities for insects: status of water beetles in southeast Spain. *Biol. Conserv.* 121:79–90
5. Abellán P, Sánchez-Fernández D, Velasco J, Millán A. 2007. Effectiveness of protected area networks in representing freshwater biodiversity: the case of a Mediterranean river basin (SE Spain). *Aquat. Conserv.* 17:361–74

6. Abellán P, Sánchez-Fernández D, Picazo F, Millán A, Lobo JM, Ribera I. 2013. Preserving the evolutionary history of freshwater biota in Iberian National Parks. *Biol. Conserv.* 162:116–26
7. Aiken RB, Khan A. 1992. The adhesive strength of the palettes of males of a boreal water beetle, *Dytiscus alaskanus* J. Balfour Browne (Coleoptera: Dytiscidae). *Can. J. Zool.* 70:1321–24
8. Arribas P, Velasco J, Abellán P, Sánchez-Fernández D, Andújar C, Calosi P, Millán A, Ribera I, Bilton DT. 2012a. Dispersal ability rather than ecological tolerance drives differences in range size between lentic and lotic water beetles (Coleoptera: Hydrophilidae). *J. Biogeogr.* 39:984–94
9. Arribas P, Abellán P, Velasco J, Bilton DT, Millán A, Sánchez-Fernández D. 2012b. Evaluating drivers of vulnerability to climate change: a guide for insect conservation strategies. *Global Change Biol.* 18:2135–46
10. Arribas P, Andujar C, Abellán P, Velasco J, Millán A, Ribera I. 2014. Tempo and mode of the multiple origins of salinity tolerance in a water beetle lineage. *Mol. Ecol.* 23:360–73
11. Arribas P, Abellán P, Velasco J, Millán A, Sánchez-Fernández D. 2017. Conservation of insects in the face of Global Climate Change. In *Global Climate Change and Terrestrial Invertebrates*, ed. SN Johnson, TH Jones, pp. 349–67. Chichester: Wiley

12. Balke M, Ribera I. 2004. Jumping across Wallace's line: *Allodessus* and *Limbodessus* revisited (Coleoptera: Dytiscidae). *Austr. J. Entomol.* 43:114–28
13. Balke M, Ribera I, Hendrich L, Miller MA, Sagata K, Posman A, Vogler AP, Meier R. 2009. New Guinea highland origin of a widespread arthropod supertramp. *Proc. Roy. Soc. Lond. B. Bio.* 276:2359–67
14. Balke M, Pons J, Ribera I, Sagata K, Vogler AP. 2007. Infrequent and unidirectional colonization of hyperdiverse *Papuadytes* diving beetles in New Caledonia and New Guinea. *Molecular Phylogenetics and Evolution* 42:505–16
15. Bergsten J, Bilton DT, Fujisawa T, Elliott M, Monaghan MT, Balke M, Hendrich L, Geijer J, Herrmann J, Foster GN, Ribera I, Nilsson AN, Barraclough TG, Vogler AP. 2012. The Effect of Geographical Scale of Sampling on DNA Barcoding. *Syst. Biol.* 61:851–69
16. Bilton DT. 2014. Dispersal in Dytiscidae. In *Ecology, Systematics, and Natural History of Predaceous Diving Beetles (Coleoptera: Dytiscidae)*, ed. DA Yee, pp. 387–407. New York: Springer
17. Bilton DT, Foster GN. 2016. Observed shifts in the contact zone between two forms of the diving beetle *Hydroporus memnonius* are consistent with predictions from sexual conflict. *PeerJ* 4:e2089

18. Bilton DT, Freeland JR, Okamura B. 2001. Dispersal in freshwater invertebrates. *Annu. Rev. Ecol. Syst.* 32:159–81
19. Bilton DT, Hayward JWG, Roch, J, Foster, GN. 2016a. Sexual dimorphism and sexual conflict in the diving beetle *Agabus uliginosus* (L.) (Coleoptera: Dytiscidae). *Biol. J. Linn. Soc.* 119:1089–95
20. Bilton DT, McAbendroth L, Bedford A, Ramsay PM. 2006. How wide to cast the net? Cross-taxon congruence of species richness, community similarity and indicator taxa in ponds. *Freshwater Biol.* 51:578–90
21. Bilton DT, Thompson A, Foster GN. 2008. Inter- and intra-sexual dimorphism in the diving beetle *Hydroporus memnonius* Nicolai (Coleoptera: Dytiscidae). *Biol. J. Linn. Soc.* 94:685–97
22. Bilton DT, Turner C, Toussaint E, Balke M. 2015 *Capelatus prykei* gen. n., sp. n. (Coleoptera: Dytiscidae: Copelatinae) - a phylogenetically isolated diving beetle from the Western Cape of South Africa. *Syst. Entomol.* 40:520–31
23. Birkhead TR, Møller AP, Sutherland WJ. 1993. Why do females make it so difficult to fertilize their eggs *J. Theor. Biol.* 161:51–60
24. Bloom DD, Fikáček M, Short AEZ. 2014. Clade age and diversification rate variation determine species richness patterns in aquatic beetle lineages.

PLoS ONE 9: e98430

25. Botella-Cruz M, Carbonell JA, Pallarés S, Millán A, Velasco J. 2016. Plasticity of thermal limits in the aquatic saline beetle *Enochrus politus* (Küster 1849) (Coleoptera: Hydrophilidae) under changing environmental conditions. *Limnetica* 35:131–42
26. Bradford TM, Humphreys WF, Austin AD, Cooper SJ. 2014. Identification of trophic niches of subterranean diving beetles in a calcrete aquifer by DNA and stable isotope analyses *Mar. Freshwater Res.* 65:95–104
27. Brown JH. 1984. On the relationship between abundance and distribution of species. *Am. Nat.* 124:255–79
28. Bukontaite R, Miller KB, Bergsten J. 2014. The utility of CAD in recovering Gondwanan vicariance events and the evolutionary history of Aciliini (Coleoptera: Dytiscidae). *BMC Evol. Biol.* 14:5
29. Bukontaite R, Ranarilalaitiana T, Randriamihaja JH, Bergsten J. 2015. In or out-of-Madagascar?—Colonization patterns for large-bodied diving beetles (Coleoptera: Dytiscidae). *PloS ONE* 10:p.e0120777
30. Calosi P, Bilton DT, Spicer JI. 2007. The diving response of the diving beetle *Ilybius montanus* (Coleoptera: Dytiscidae): the effects of temperature and acidification. *J. Zool.* 273:289–97

31. Calosi P, Bilton DT, Spicer JI, Atfield A. 2008a. Thermal tolerance and geographic range size in the *Agabus brunneus* group of European diving beetles (Coleoptera: Dytiscidae). *J. Biogeogr.* 35:295–305
32. Calosi P, Bilton DT, Spicer JI. 2008b. Thermal tolerance, acclimatory capacity and vulnerability to global climate change. *Biol. Letters* 4:99–102
33. Calosi P, Bilton DT, Spicer JI, Votier S, Atfield A. 2010. What determines a species' geographical range? Thermal biology and latitudinal range size relationships in European diving beetles (Coleoptera: Dytiscidae). *J. Anim. Ecol.* 79: 194–204
34. Calosi P, Bilton DT, Spicer JI, Verberk WCEP, Atfield A, Garland TJr. 2012. The comparative biology of diving in two genera of European Dytiscidae (Coleoptera). *J. Evol. Biol.* 25:329–41
35. Cioffi R, Moody JA, Millán A, Billington R, Bilton DT. 2016. Physiological niche and geographical range in European diving beetles (Coleoptera: Dytiscidae). *Biol. Letters* 12:20160130
36. Coope GR. 1994. The response of insect faunas to glacial– interglacial climatic fluctuations. *Philos. T. Roy. Soc. B.* 344: 19–26
37. Cooper SJB, Hinze S, Leys R, Watts CHS, Humphreys WF 2002. Islands under the desert: molecular systematics and evolutionary origins of stygobitic water

beetles (Coleoptera: Dytiscidae) from central Western Australia. *Invertebr. Syst.* 16:589–98

38. Dehling M, Hof C, Brändle M, Brandl R. 2010. Habitat availability does not explain the species richness patterns of European lentic and lotic freshwater animals. *J. Biogeogr.* 37:1919–26
39. Deler-Hernández A, Sýkora V, Seidel M, Cala-Riquelme F, Fikáček M. 2017. Multiple origins of the *Phaenonotum* beetles in the Greater Antilles (Coleoptera: Hydrophilidae): phylogeny, biogeography and systematics. *Zool. J. Linn. Soc.* <https://doi.org/10.1093/zoolinnean/zlx071>.
40. Désamoré A, Laenen B, Miller KB, Bergsten J. 2018. Early burst in body size evolution in uncoupled from species diversification in diving beetles (Dytiscidae). *Mol. Ecol.* doi:10.1111/mec.14492
41. Dijkstra KDB, Monaghan MT, Pauls SU. 2014. Freshwater biodiversity and aquatic insect diversification. *Annu. Rev. Entomol.* 59:143–63
42. Eyre MD, Foster GN, Luff M, Staley JR. 2003. An investigation into the relationship between water beetle (Coleoptera) distribution and land cover in Scotland and northeast England. *J. Biogeogr.* 30:1835–49

43. Fikáček M, Minoshima YN, Newton AF. 2014. A Review of *Andotypus* and *Austrotypus* gen. nov., Rygmodine Genera with an Austral Disjunction (Hydrophilidae: Rygmodinae). *Ann. Zool.* 64:557–96
44. Fikacek M, Prokin AA, Angus RB, Ponomarenko AG, Yue Y, Ren D, Prokop J. 2012. Phylogeny and the fossil record of the Helophoridae reveal Jurassic origin of modern hydrophiloid lineages (Coleoptera: Polyphaga). *Syst. Entomol.* 37:420–47
45. Foster GN. 1993. Pingo fens, water beetles and site evaluation. *Antenna* 17:184–95
46. Foster GN, Bilton DT. 2014. The conservation of predaceous diving beetles: knowns, unknowns and anecdotes. In *Ecology, Systematics, and Natural History of Predaceous Diving Beetles (Coleoptera: Dytiscidae)*, ed. DA Yee, pp. 437–62 New York: Springer
47. Foster GN, Foster AP, Eyre MD, Bilton DT. 1989. Classification of water beetle assemblages in arable fenland and ranking of sites in relation to conservation value. *Freshwater Biol.* 22:343–54
48. García-Criado F, Fernández-Aláez M, Fernández-Aláez C. 2001. Hydraenidae and Elmidae assemblages (Coleoptera) from a Spanish river basin: good indicators of coal mining pollution? *Arch. Hydrobiol.* 150:641–60

49. García-Vázquez D, Bilton DT, Alonso R, Benetti CJ, Garrido J, Valladares LF, Ribera I. 2016. Reconstructing ancient Mediterranean crossroads in *Deronectes* diving beetles. *J. Biogeogr.* 43:1533–45
50. García-Vázquez D, Bilton DT, Foster GN, Ribera I. 2017. Pleistocene range shifts, refugia and the origin of widespread species in Western Palaearctic water beetles. *Mol. Phylogenet. Evol.* 114:122–36
51. García-Vázquez D, Ribera I. 2016 The origin of widespread species in a poor dispersing lineage (diving beetle genus *Deronectes*). *PeerJ.* 4:e2514
52. Gaston KJ. 2003. *The Structure and Dynamics of Geographic Ranges*. Oxford: Oxford University Press
53. Gaston KJ, Chown SL, Calosi P, Bernardo J, Bilton DT, Clarke A, Clusella-Trullas S, Ghalambor CK, Konarzewski M, Peck LS, Porter WP, Pörtner HO, Rezende EL, Schuller PM, Spicer JI, Stillman J, Terblanche JS, van Kleunen M. 2009. Macrophysiology: a conceptual reunification. *Am. Nat.* 174:595–612
54. Gleeson T, Wada Y, Bierkens MFP, van Beek LPH. 2012 Water balance of global aquifers revealed by groundwater footprint *Nature* 488:197-200
55. Guareschi S, Bilton DT, Velasco J, Millán A, Abellán P. 2015. How well do protected area networks support taxonomic and functional diversity in non-target taxa? The case of Iberian freshwaters. *Biol. Conserv.* 187:134–44

56. Guareschi S, Gutiérrez-Cánovas C, Picazo F, Sánchez-Fernández D, Abellán P, Velasco J, Millán A. 2012. Aquatic macroinvertebrate biodiversity: patterns and surrogates in mountainous Spanish national parks. *Aquat. Conserv.* 22:598–615
57. Gustafson G, Prokin AA, Bukontaite R, Bergsten J, Miller KB. 2017. Tip-dated phylogeny of whirligig beetles reveals ancient lineage surviving on Madagascar. *Sci. Rep.* 7:1–9
58. Guzik MT, Austin AD, Cooper SJB, Havey MS, Humphreys WF, Bradford T, Eberhard SM, King RA, Leys R, Muirhead KA, Tomlinson M. 2010. Is the Australian subterranean fauna uniquely diverse? *Invertebr. Syst.* 24:407–18
59. Guzik MT, Cooper S.J.B, Humphreys WF, Ong S, Kawakami T, Austin AD. 2011. Evidence for population fragmentation within a subterranean aquatic habitat in the Western Australian desert. *Heredity* 107:215–30
60. Härdling R, Bergsten J. 2006. Nonrandom mating preserves intrasexual polymorphism and stops population differentiation in sexual conflict. *Am. Nat.* 167:401–9
61. Härdling R, Karlsson K. 2009. The dynamics of sexually antagonistic coevolution and the complex influences of mating system and genetic correlation. *J. Theor. Biol.* 260:276–82

62. Hernando C, Aguilera P, Ribera I. 2001. *Limnius stygius* sp. nov., the first stygobiontic riffle beetle from the Palearctic Region (Coleoptera: Elmidae). *Entomol. Problems.* 32:69–72
63. Hewitt G. 2000. The genetic legacy of the Quaternary ice ages. *Nature* 405:907–13
64. Hidalgo-Galiana A, Monge M, Biron DG, Canals F, Ribera I, Cieslak A. 2014b. Reproducibility and consistency of proteomic experiments on natural populations of a non-model aquatic insect species. *PLoS ONE* 9:e104734
65. Hidalgo-Galiana A, Monge M, Biron DG, Canals F, Ribera I, Cieslak, A. 2016. Protein expression parallels thermal tolerance and ecologic changes in the diversification of a diving beetle species complex. *Heredity* 116:114–23
66. Hidalgo-Galiana A, Ribera I. 2011. Late Miocene diversification of the genus *Hydrochus* (Coleoptera, Hydrochidae) in the west Mediterranean area. *Mol. Phylogenet. Evol.* 59: 377–85
67. Hidalgo-Galiana A, Sánchez-Fernández D, Bilton DT, Cieslak A, Ribera, I. 2014a. Thermal niche evolution and geographic range expansion in a species complex of western Mediterranean diving beetles. *BMC Evol. Biol.* 14:187

68. Higginson DM, Miller KB, Segraves KA, Pitnick S. 2012a. Female reproductive tract form drives the evolution of complex sperm morphology. *P. Natl. A. Sci. Biol.* 109:4538–43
69. Higginson DM, Miller KB, Segraves KA, Pitnick S. 2012b. Convergence, recurrence and diversification of complex sperm traits in diving beetles (Dytiscidae). *Evolution* 66:1650-61
70. Hjalmarsson AE, Bergsten J, Monaghan MT. 2015. Dispersal is linked to habitat use in 59 species of water beetles (Coleoptera: Adepaga) on Madagascar. *Ecography* 38:732–9
71. Hof C, Brändle M, Brandl R 2006 Lentic odonates have larger and more northern ranges than lotic species. *J. Biogeogr.* 33:63–70
72. Hosken DJ, Stockley P. 2004. Sexual selection and genital evolution. *Trends Ecol. Evol.* 19:87–93
73. Humphreys WF, Watts CHS, Cooper SJB, Leijns R. 2009. Groundwater estuaries of salt lakes: buried pools of endemic biodiversity on the western plateau, Australia. *Hydrobiologia* 626:79–95
74. Iversen LL, Jacobsen D, Sand-Jensen K. 2016 Are latitudinal richness gradients in European freshwater species only structured according to dispersal and time? *Ecography* 39:1247–9

75. Jäch MA, Balke M. 2008. Global diversity of water beetles (Coleoptera) in freshwater. *Hydrobiologia* 595:419–42
76. Karlsson Green K, Kovalev A, Svensson EI, Gorb SN. 2013. Male clasping ability, female polymorphism and sexual conflict: fine-scale elytral morphology as a sexually antagonistic adaptation in female diving beetles. *J. Roy. Soc. Interface* 10:20130409
77. Karlsson Green K, Svensson EI, Bergsten J, Härdling R, Hansson B. 2014. The interplay between local ecology, divergent selection, and genetic drift in population divergence of a sexually antagonistic female trait. *Evolution* 68:1934–46
78. Kehl S, Dettner K. 2009. Surviving submerged—setal tracheal gills for gas exchange in adult rheophilic diving beetles. *J. Morphol.* 270:1348–55
79. Leijs R, van Nes EH, Watts CH, Cooper SJB, Humphreys WF, Hogendoorn K. 2012. Evolution of blind beetles in isolated aquifers: a test of alternative modes of speciation. *PLoS One* 7:e34260
80. Leys R, Cooper SJB, Strecker U, Wilkens H. 2005. Regressive evolution of an eye pigment gene in independently evolved eyeless subterranean diving beetles. *Biol. Letters* 1:496–9

81. Leys R, Watts CHS 2008. Systematics and evolution of the Australian subterranean hydroporine diving beetles (Dytiscidae), with notes on *Carabhydrus*. *Invertebr. Syst.* 22:217–25
82. Leys R, Watts CHS, Cooper SJB, Humphreys WF 2003. Evolution of subterranean diving beetles (Coleoptera: Dytiscidae: Hydroporini, Bidessini) in the arid zone of Australia. *Evolution* 57:2819–34
83. Lytle DA, Peckarsky BL. 2001. Spatial and temporal impacts of a diesel fuel spill on stream invertebrates. *Freshwater Biol.* 46:693–704
84. Marten A, Brändle M, Brandl R. 2006 Habitat type predicts genetic population differentiation in freshwater invertebrates. *Mol. Ecol.* 15: 2643–51
85. McKenna DD, Wild AL, Kanda K, Bellamy CL, Beutel RG, Caterino MS, Farnum CW, Hawks DC, Ivie MA, Jameson ML, Leschen RAB, Marvaldi AE, McHugh JV, Newton AF, Robertson JA, Thayer MK, Whiting MF, Lawrence JF, Slipinski A, Maddison DR, Farrell BD. 2015. The beetle tree of life reveals that Coleoptera survived end-Permian mass extinction to diversify during the Cretaceous terrestrial revolution. *Syst. Entomol.* 40:835–80
86. Miguelez D, Maze RA, Ansola G, Valladares LF. 2013. Aquatic Coleoptera and Hemiptera assemblages of a Cantabrian coastal stream (North Spain): composition, seasonal variation and environmental factors. *Limnetica* 32:47–

87. Millán A, Velasco J, Gutiérrez-Cánovas C, Arribas P, Picazo F, Sánchez-Fernández D, Abellán P. 2011. Mediterranean saline streams in southeast Spain: What do we know? *J. Arid Environ.* 75:1352–9
88. Miller, K.B. 2001. On the phylogeny of the family Dytiscidae Linnaeus (Insecta: Coleoptera) with an emphasis on the morphology of the female reproductive tract. *Insect Syst. Evol.* 32:45–92
89. Miller KB. 2003. The phylogeny of diving beetles (Coleoptera: Dytiscidae) and the evolution of sexual conflict. *Biol. J. Linn. Soc.* 79:359–88
90. Miller KB, Bergsten J. 2014. Predaceous diving beetle sexual systems. In *Ecology, systematics, and natural history of predaceous diving beetles (Coleoptera: Dytiscidae)*, ed. DA Yee, pp. 199–234. New York: Springer
91. Miller KB, Bergsten J. 2016. *Diving Beetles of the World*. Baltimore: Johns Hopkins University Press
92. Miller KB, Jean A, Alarie Y, Hardy N, Gibson R. 2013. Phylogenetic placement of North American subterranean diving beetles (Coleoptera: Dytiscidae). *Arthropod Syst. Phylo.* 71:75–90
93. Miller GT, Pitnick S. 2002. Sperm-female coevolution in *Drosophila*. *Science* 298:1230–3

94. Monaghan MT, Wild R, Elliot M, Fujisawa T, Balke M, Inward DJ, Lees DC, Ranaivosolo R, Eggleton P, Barraclough TG, Vogler AP. 2009. Accelerated species inventory on Madagascar using coalescent-based models of species delineation. *Syst. Biol.* 58:298–311
95. Moritz C, Patton JL, Schneider CJ, Smith TB. 2000. Diversification of rainforest faunas: an integrated molecular approach. *Annu. Rev. Ecol. Syst.* 31:533–63
96. Pallares S, Arribas P, Bilton DT, Millán A, Velasco J. 2015. The comparative osmoregulatory ability of two water beetle genera whose species span the fresh-hypersaline gradient in inland waters (Coleoptera: Dytiscidae, Hydrophilidae). *PLoS ONE* 10:e0124299
97. Pallares S, Arribas P, Bilton DT, Millán A, Velasco J, Ribera I. 2017a The chicken or the egg? Adaptation to desiccation and salinity tolerance in a lineage of water beetles. *Mol. Ecol.* 26:5614–28
98. Pallares S, Arribas P, Cespedes V, Millán A, Velasco J. 2012. Lethal and sublethal behavioural responses of saline water beetles to acute heat and osmotic stress. *Ecol. Entomol.* 37:508–20

99. Pallarés S, Botella-Cruz M, Arribas P, Millán A, Velasco J. 2017b. Aquatic insects in a multistress environment: cross-tolerance to salinity and desiccation. *J. Exp. Biol.* 220:1277–86
100. Pallarés S, Velasco J, Millán A, Bilton DT, Arribas P. 2017c. Aquatic insects dealing with dehydration: do desiccation resistance traits differ in species with contrasting habitat preferences? *PeerJ* 4:e2382
101. Parker GA. 2006. Sexual conflict over mating and fertilization: an overview. *Philos. T. Roy. Soc. B.* 361:235–59
102. Perkins PD. 2017. Hydraenidae of Madagascar (Insecta: Coleoptera). *Zootaxa* 4342:1–264
103. Perkins PD, Balfour-Browne J. 1994. A contribution to the taxonomy of aquatic and humicolous beetles of the family Hydraenidae in southern Africa. *Fieldiana Zool.* 77:1–159
104. Picazo F, Bilton DT, Moreno JL, Sánchez-Fernández D, Millán A. 2012. Water beetle biodiversity in Mediterranean standing waters: assemblage composition, environmental drivers and nestedness patterns. *Insect Divers. Conser.* 5:146–58
105. Ponomarenko AG, Prokin AA. 2015. Review of paleontological data on the evolution of aquatic beetles (Coleoptera). *Paleontolog. J.* 49:1383–412

106. Popov SV, Rogl F, Rozanov AY, Steininger FF, Shcherba IG, Kovac, M. 2004. Lithological-Paleogeographic maps of Paratethys. 10 maps Late Eocene to Pliocene. *Cour. Forsch. Sencken.* 250:1–46
107. Potts K. 2016. Secondary sexual characters in crawling water beetles (Coleoptera: Haliplidae): evidence for sexual conflict? *The Plymouth Student Scientist* 9:162–213
108. Ribera I. 2000. Biogeography and conservation of Iberian water beetles. *Biol. Conserv.* 92:131–50
109. Ribera I. 2008. Habitat constraints and the generation of diversity in freshwater macroinvertebrates. In *Aquatic Insects: Challenges to Populations* ed. J Lancaster, RA Briers, pp. 289–311 Wallingford: CAB International
110. Ribera I, Balke M. 2007. Recognition of a species-poor, geographically restricted but morphologically diverse Cape lineage of diving beetles (Coleoptera: Dytiscidae: Hyphydrini). *J. Biogeogr.* 34:1220–32
111. Ribera I, Beutel RG, Balke M, Vogler AP. 2002. Discovery of Aspidytidae, a new family of aquatic beetles. *Proc. Roy. Soc. Lond. B. Bio.* 269:2351–6

112. Ribera I, Barraclough TG, Vogler AP. 2001. The effect of habitat type on speciation rates and range movements in aquatic beetles: inferences from species-level phylogenies. *Mol. Ecol.* 10:721–35
113. Ribera I, Castro A, Diaz JA, Garrido J, Izquierdo A, Jäch MA, Valladares LF. 2011. The geography of speciation in narrow-range endemics of the '*Haenydra*' lineage (Coleoptera, Hydraenidae, *Hydraena*). *J. Biogeogr.* 38:502–16
114. Ribera I, Castro A, Hernando C. 2010 *Ochthebius* (*Enicocerus*) *aguilerai* sp.n. from central Spain, with a molecular phylogeny of the Western Palaearctic species of *Enicocerus* (Coleoptera, Hydraenidae). *Zootaxa* 2351:1–13
115. Ribera I, Foster GN, Vogler AP. 2003. Does habitat use explain large scale species richness patterns of aquatic beetles in Europe? *Ecography* 26:145–52
116. Ribera I, Vogler AP. 2000. Habitat type as a determinant of species range sizes: the example of lotic-lentic differences in aquatic Coleoptera. *Biol. J. Linn. Soc.* 71:33–52
117. Ribera I, Vogler AP. 2004. Speciation of Iberian diving beetles in Pleistocene refugia (Coleoptera, Dytiscidae). *Mol. Ecol.* 13:179–93

118. Rudoy A, Beutel RG, Ribera I. 2016. Evolution of the male genitalia in the genus *Limnebius* (Coleoptera, Hydraenidae). *Zool. J. Linn. Soc.* 178:97–127
119. Rudoy A, Ribera I. 2016. The macroevolution of size and complexity in insect male genitalia. *PeerJ* 4:e1882
120. Rudoy A, Ribera I. 2017. Evolution of sexual dimorphism and Rensch's rule in the beetle genus *Limnebius* (Hydraenidae) - is sexual selection opportunistic? *PeerJ* 5:e3060
121. Sabatelli S, Audisio P, Antonini G, Solano E, Martinoli A, Trizzino M. 2016. Molecular ecology and phylogenetics of the water beetle genus *Ochthebius* revealed multiple independent shifts to marine rockpools lifestyle. *Zool. Scr.* 45:175–86
122. Sánchez-Fernández D, Abellán P, Mellado A, Velasco J, Millán A. 2006. Are water beetles good indicators of biodiversity in Mediterranean aquatic ecosystems? The case of the Segura river basin (SE Spain). *Biodivers. Conserv.* 15:4507–20
123. Sánchez- Fernández D, Abellán P, Velasco J, Millán A. 2004. Selecting areas to protect the biodiversity of aquatic ecosystems in a semiarid Mediterranean region using water beetles. *Aquatic Conserv.* 14:465–79

124. Sánchez-Fernández D, Lobo JM, Abellán P, Ribera I, Millán A. 2008. Bias in freshwater biodiversity sampling: the case of Iberian water beetles. *Divers. Distrib.* 14:754–62
125. Sánchez-Fernández D, Lobo JM, Abellán P, Millán A. 2011a. How to identify future sampling areas when information is biased and scarce: an example using predictive models for species richness of Iberian water beetles. *J. Nat. Conserv.* 19:54–9
126. Sánchez-Fernández D, Lobo JM, Hernández-Manrique OL. 2011b. Species distribution models that do not incorporate global data misrepresent potential distributions: a case study using Iberian diving beetles. *Divers. Distrib.* 17:163–71
127. Sánchez-Fernández D, Lobo JM, Millán A, Ribera I. 2012a. Habitat type mediates equilibrium with climatic conditions in the distribution of Iberian diving beetles. *Global Ecol. Biogeogr.* 21:988–97
128. Sánchez-Fernández D, Aragón P, Bilton DT, Lobo JM. 2012b. Assessing the congruence of thermal niche estimations derived from distribution and physiological data. A test using diving beetles. *PLoS ONE* 7:e48163

129. Scheffer M, Vergnon R, van Nes EH, Cuppen JGM, Peeters ETHM, Leijds R, Nilsson AN. 2015. The Evolution of Functionally Redundant Species; Evidence from Beetles. *PLoS ONE* 10:e0137974
130. Shirt DB, Angus RB. 1992. A revision of the Nearctic water beetles related to *Potamonectes depressus* (Fabricius) (Coleoptera: Dytiscidae). *Colept. Bull.* 46:109–41
131. Short AEZ. 2018. Systematics of aquatic beetles (Coleoptera): current state and future directions. *Syst. Entomol.* 43:1–18
132. Short AEZ, Caterino MS. 2009. On the validity of habitat as a predictor of genetic structure in aquatic systems: a comparative study using California water beetles. *Mol. Ecol.* 18:403–14
133. Short AEZ, Fikáček M. 2013. Molecular Phylogeny, Evolution, and Classification of the Hydrophilidae (Coleoptera). *Syst. Entomol.* 38:723–52
134. Short AEZ, Joly LJ, García M, Wild A, Bloom DD, Maddison DR. 2015. Molecular phylogeny of the Hydroscaphidae (Coleoptera: Myxophaga) with description of a remarkable new lineage from the Guiana Shield. *Syst. Entomol.* 40:214–29

135. Short AEZ, Liebherr JK. 2007. Systematics and biology of the endemic water scavenger beetles of Hawaii (Coleoptera: Hydrophilidae: Hydrophilini). *Syst. Entomol.* 32:601–24
136. Shuker D, Simmons LW, eds. 2014. *The Evolution of Insect Mating Systems*. Oxford: Oxford University Press
137. Simmons LW. 2014. Sexual selection and genital evolution. *Austral Entomol.* 53:1–17
138. Spangler PJ, Decu V. 1998. Coleoptera aquatica In *Encyclopaedia Biospeologica*. Tom II, ed. C Juberthie, V Decu, pp. 1031–46. Société de Biospéologie Moulis: Bucarest
139. Tierney SM, Cooper SJB, Saint KM, Bertozzi T, Hyde J, Humphreys WF, Austin AD. 2015. Opsin transcripts of predatory diving beetles: A comparison of surface and subterranean photic niches. *Roy. Soc. Open Sci.* 2:140386
140. Toussaint EFA, Beutel RG, Morinière J, Jia F, Xu S, Michat MC, Zhou X, Bilton DT, Ribera I, Hájek J, Balke M. 2016c. Molecular phylogeny of the highly disjunct cliff water beetles from South Africa and China (Coleoptera: Aspidytidae). *Zool. J. Linn. Soc.* 176:537–46

141. Toussaint EFA, Bloom DD, Short AEZ. 2017b. Cretaceous West Gondwana vicariance shaped giant water scavenger beetle palaeobiogeography. *J. Biogeogr.* 44:1952–65
142. Toussaint EFA, Condamine FL, Hawlitschek O, Watts CHS, Porch N, Hendrich L, Balke M. 2015. Unveiling the diversification dynamics of Australasian predaceous diving beetles in the Cenozoic. *Syst. Biol.* 64:3–24
143. Toussaint EFA, Fikáček M, Short AEZ. 2016b. India-Madagascar vicariance explains cascade beetle biogeography. *Biol. J. Linn. Soc.* 118:982–91
144. Toussaint EFA, Hendrich L, Escalona H, Porch N, Balke M. 2016c. Evolutionary history of a secondary terrestrial Australian diving beetle (Coleoptera, Dytiscidae) reveals a lineage of high morphological and ecological plasticity. *Syst. Entomol.* 41:650–7
145. Toussaint EFA, Hendrich L, Hájek J, Michat M, Panjaita R, Short AEZ, Balke M. 2017c. Evolution of Pacific Rim diving beetles sheds light on Amphipacific biogeography. *Ecography* 40:500–10
146. Toussaint EFA, Hendrich L, Shaverdo H, Balke M. 2015. Mosaic patterns of diversification dynamics following the colonization of Melanesian islands. *Sci. Rep.* 5:16016

147. Toussaint EFA, Sagata K, Surbakti S, Hendrich L, Balke M. 2013. Australasian sky islands act as a diversity pump facilitating peripheral speciation and complex reversal from narrow endemic to widespread ecological supertramp. *Ecol. Evol.* 3:1031–49
148. Toussaint EFA, Seidel M, Arriaga-Varela E, Hájek J, Král D, Sekerka, L, Short AEZ, Fikáček M. 2017a. The peril of dating beetles. *Syst. Entomol.* 42:1–10
149. Toussaint EFA, Short AEZ. 2016. Miocenic Evolution of Brazilian Shield *Platynectes* Diving Beetles in Amazon Basin Paleodrainage Systems. *Annales Soc. Entomol. France* 52:185–91
150. Toussaint EFA, Short AEZ. 2017. Biogeographic mirages? Molecular evidence for dispersal-driven evolution in Hydrobiusini water scavenger beetles. *Syst. Entomol.* 42:692–702
151. Trizzino M, Audisio PA, Antonini G, Mancini E, Ribera I. 2011. Molecular phylogeny and diversification of the “*Haenydra*” lineage (Hydraenidae, genus *Hydraena*), a north-Mediterranean endemic-rich group of rheophilic Coleoptera. *Mol. Phylogen. Evol.* 61:772–83
152. Trizzino M, Jäch MA, Audisio P, Alonso R, Ribera I. 2013. A molecular phylogeny of the cosmopolitan hiperdiverse genus *Hydraena* Kugelann (Coleoptera, Hydraenidae). *Syst. Entomol.* 38:192–20

153. Verberk WCEP, Bilton DT, Calosi P, Spicer JI. 2011. Oxygen supply in aquatic ectotherms: Partial pressure and solubility together explain biodiversity and size patterns. *Ecology* 92:1565–72
154. Verberk WCEP, Bilton DT. 2013. Respiratory control in aquatic insects dictates their vulnerability to global warming. *Biol. Letters* 9:20130473
155. Verberk WCEP, Calosi P, Spicer JI, Kehl S, Bilton DT. 2018. Does plasticity in thermal tolerance trade off with inherent tolerance? The influence of setal tracheal gills on thermal tolerance and its plasticity in a group of European diving beetles. *J. Insect Physiol.* in press
156. Vergnon R, Leijs R, van Nes EH, Scheffer M. 2013. Repeated parallel evolution reveals limiting similarity in subterranean diving beetles. *Am. Nat.* 182:67–75
157. Villastrigo A, Fery H, Manuel M, Millán A, Ribera I. 2018. Evolution of salinity tolerance in the diving beetle tribe Hygrotini (Coleoptera, Dytiscidae). *Zool. Scr.* 47:63–71
158. Watts CHS, Humphreys WF. 2004. Thirteen new Dytiscidae (Coleoptera) of the genera *Boongurrus* Larson, *Tjirtudessus* Watts & Humphreys and *Nirripirti* Watts and Humphreys, from underground waters in Australia. *Transactions of the Royal Society of South Australia.* 128:99–129

159. Watts CHS, Humphreys WF. 2006. Twenty-six new Dytiscidae (Coleoptera) of the genera *Limbodessus* Guignot and *Nirripirti* Watts & Humphreys, from underground waters in Australia. *T. Roy. Soc. South Aust.* 130:123–85
160. Wiens JJ. 2004. Speciation and ecology revisited: phylogenetic niche conservatism and the origin of species. *Evolution* 58:193–7
161. Zhang S-Q, Che L-H, Li Y, Liang D, Pang H, Ślipiński A, Zhang P. 2018. Evolutionary history of Coleoptera revealed by extensive sampling of genes and species. *Nat. Commun.* 9:205

Reference annotations

3. Development of a novel approach to infer speciation modes based on the phylogeny and current distributions of the species of a lineage.

10. Rationale for deriving conservation strategies from empirical biological/ecological data.

16. Extensive evaluation of how DNA barcoding performs at different spatial and evolutionary scales.

34. First study to compare the relative importance of different drivers of range size in a phylogenetically controlled framework.

58. Oldest known Malagasay animal lineage.

76. Overview of beetle diversity and biology, with discussion on what constitutes a water beetle.

92. Key review of generic diversity of major aquatic family.

98. On of the few insect studies on the origin of a trait (tolerance to salinity) combining both phylogenetic and experimental physiological data.

110. A review on the habitat stability hypothesis applied to differences in lotic-lentic habitats.

131. Up-to-date review of water beetle taxonomic diversity.

RELATED RESOURCES

Beutel RG, Leschen RAB. 2016. *Coleoptera, Beetles Volume 1: Morphology and Systematics (Archostemata, Adephaga, Myxophaga, Polyphaga partim)*, 2nd edition Handbook of Zoology, Berlin: Walter de Gruyter GmbH. [chapters on all major water beetle families].

Hansen M. 1998. World Catalogue of Insects, Volume 1: Hydraenidae (Coleoptera).

Stenstrup: Apollo Books

Hansen M. 1999. World Catalogue of Insects, Volume 2: Hydrophiloidea

(Coleoptera). Stenstrup: Apollo Books

Nilsson AN. 2001. World Catalogue of Insects, Volume 3: Dytiscidae (Coleoptera).

Stenstrup: Apollo Books

Nilsson AN, van Vondel BJ. 2005. World Catalogue of Insects, Volume 7:

Amphizoidae, Aspidytidae, Haliplidae, Noteridae and Paelobiidae (Coleoptera,

Adephaga). Stenstrup: Apollo Books

Jäch MA, Kodada J, Čiampor Jr F. 2016. World Catalogue of Insects, Volume 14:

Coleoptera: Elmidae and Protelmidae. Leiden: Brill

Figure legends

Figure 1. Representative time-calibrated phylogeny of Coleoptera showing the relative phylogenetic position and antiquity of each major water beetle lineage. The tree topology is simplified from McKenna et al. (85), with the exception that Adephaga is drawn as a polytomy between Geoadephaga, Grynidae, Haliplidae, and Dytiscoidea. Divergence time estimates of crown lineages (red circles) are based on Toussaint et al. (148) (but see e.g. 85 or 161 for an alternative dating), and the oldest known fossil taxa (black circles) are taken from Pomonorenko & Prokin (105). Primarily terrestrial lineages that contain some aquatic taxa are marked with an asterisk. See Supplementary File 1 for additional explanations.

Figure 2. Sexually selected characters in water beetles: (a-c) *Acilius sulcatus* (Dytiscidae) (a) male habitus – note expanded fore and mid tarsi; (b) ventral surface of male fore-tarsus – note plunger-like articulo-setae which function as attachment devices; (c) female habitus – note furrowed elytra and hairs on pronotum and elytra, resistance traits making male attachment more difficult during pairing. (d-g) *Limnebius* species (Hydraenidae) (d) *Limnebius truncatellus* male (above) and female (below) habitus – note strong sexual dimorphism in body size and leg modifications; (e) relatively complex male genitalia of *Limnebius truncatellus*; (f) *Limnebius evanescens* male (above) and female (below) habitus – note limited sexual dimorphism in body size; (g) relatively simple male genitalia of *Limnebius evanescens*. (h-i) sperm of *Hygrotus sayi* (Dytiscidae), epifluorescence microscopy with only DNA-stained heads visible (h) isolated, dimorphic sperm, some with broad heads and basal spurs, others with filamentous heads; (i) sperm conjugation –

sperm with broad heads stack (like traffic cones), forming a scaffold which sperm with filamentous heads attach to. Scale bars as follows: a & $c = 5$ mm, b , d & $f = 1$ mm, e & $g = 500$ μm , h - $l = 20$ μm .