


Indonesian J. Pharm.
Volume 29 Issue 4 (2018)
October-December



ISSN : 2338-9427
Formerly ISSN : 0126-1037

Indonesian Journal of Pharmacy (*Indonesian J. Pharm.*)

Accredited by DGHE (DIKTI) No. 58/DIKTI/Kep/2013



STT NO. 1652/SK/DITJEN PPG/SST/1990

Faculty of Pharmacy
Universitas Gadjah Mada



Editorial Team

Editor in Chief

Prof. Sugiyanto Sugiyanto, Universitas Gadjah Mada, Department of Pharmacology and Clinical Pharmacy, Indonesia

Editorial Board

Prof. Dr. Abdul Rohman, Faculty of Pharmacy, Department of Pharmaceutical Chemistry, Universitas Gadjah Mada, Indonesia

Prof. Dr. Shufeng Zhou, Department of Pharmaceutical Sciences, University of South Florida Tampa, United States

Prof. Dr. Kazutaka Maeyama, Ehime University, Department of Pharmacology, Japan

Prof. Dr. Masashi Kawaichi, Nara Institute of Science and Technology, Division of Gene Function in Animals, Japan

Prof. Dr. Gunawan Indrayanto, Universitas Airlangga, Faculty of Pharmacy, Indonesia

Prof. Dr. Veeresh P. Veerapur, Sree Siddaganga College of Pharmacy, Pharmaceutical Chemistry Department, India

Prof. Dr. Agung Endro Nugroho, Universitas Gadjah Mada, Faculty of Pharmacy, Department of Pharmacology and Clinical Pharmacy, Indonesia

Prof. Dr. Lee E. Kirsch, University of Iowa, Division of Pharmaceutics and Translational Therapeutics, United States

Prof. Dr. Henk Timmerman, Vrije Universiteit Amsterdam, Division of Medicinal Chemistry, Netherlands

Prof. Dr. Jeroen Kool, Vrije Universiteit Amsterdam, Division of BioAnalytical Chemistry, Netherlands

Dr. Saikat Kumar Basu, University of Lethbridge, Department of Biological Sciences, Canada

Dr. Joseph David Francis Tucci, La Trobe University, School of Pharmacy and Applied Science, Australia

Dr. Chuda Chittasupho, Srinakharinwirot University, Department of Pharmaceutical Technology, Thailand

Dr. Rina Kuswahyuning, Universitas Gadjah Mada, Faculty of Pharmacy, Department of Pharmaceutics, Indonesia

Dr. Supang Khonde, University of Phayao, School of Pharmaceutical Sciences, Thailand

Dr. Pudjono Pudjono, Universitas Gadjah Mada, Faculty of Pharmacy, Department of Pharmacology and Clinical Pharmacy, Indonesia

Dr. Montarat Thavorncharoensap, Faculty of Pharmacy, Department of Pharmacy, Mahidol University, Thailand

Dr. Karuna Shanker, Central Institute of Medicinal and Aromatic Plants India, Department of Analytical Chemistry, India

Dr. Jun An, Sun Yat-Sen University, Department of Cardiothoracic Surgery, China

Dr. Mohammed Emamussalehin Choudhury, Department of Pharmacology, Bangladesh Agriculture University, Bangladesh













Dr. Abdul Wahab, Department of Pharmacy, Kohat University of Science and Technology (KUST), Pakistan

Dr. Tony Hadibarata, Curtin University Sarawak Malaysia, Department of Environmental Engineering, Malaysia

Dr. Shahin Gavanji, Department of Biotechnology, Faculty of Advanced Sciences and Technologies, University of Isfahan, Isfahan, Iran, Islamic Republic of

Table of Contents

Articles

Screening of Antibacterial and Anticancer Activity of Soft Corals from Togeang Islands, Indonesia	
 <i>Muhammad Sulaiman Zubair, Subehan Lallo, Rusmiyanti Rusmiyanti, Arsa Wahyu Nugrahani, Ibrahim Jantan</i>	173
DOI:10.14499/indonesianjpharm29iss4pp173 Abstract view:100 PDF download:29	
Cytotoxicity of Tetrahydropentagamavunon-0 (THPGV)-0 and Tetrahydropentagamavunon-1 (THPGV-1) in Several Cancer Cell Lines	
 <i>Muthi Ikawati, Heri Purwanto, Niar Nurul Imaniyati, Anis Afifah, Marrita Langgeng Sagiyo, Jasson Yohanes, Sismindari Sismindari, Ritmaleni Ritmaleni</i>	179
DOI:10.14499/indonesianjpharm29iss4pp179 Abstract view:92 PDF download:9	
Effects of Antioxidant, Anti-Collagenase, Anti-Elastase, Anti-Tyrosinase of The Extract and Fraction From <i>Turbinaria decurrens</i> Bory.	
 <i>Arief Nurrochmad, Wirasti Wirasti, Arifin Dirman, Endang Lukitaningsih, Adillah Rahmawati, Nanang Fakhruddin</i>	188
DOI:10.14499/indonesianjpharm29iss4pp188 Abstract view:98 PDF download:15	
Synthesis and Molecular Docking Studies of N'-benzoylsalicylhydrazide derivatives as antituberculosis through InHA enzym inhibition	
 <i>Harry Santosa, Galih Satrio Putra, Tegar Achsendo Yuniarta, Tutuk Budiati</i>	198
DOI:10.14499/indonesianjpharm29iss4pp198 Abstract view:86 PDF download:9	
α-Glucosidase Inhibition Activity of Stem Bark <i>Ceiba pentandra</i> Linn. In Vitro and In Silico	
 <i>Vicri Syihabudin</i>	206
DOI:10.14499/indonesianjpharm29iss4pp206 Abstract view:78 PDF download:14	
Efficacy of Thymol and Eugenol Against Polymicrobial Biofilm	
 <i>Hasyrul Hamzah, Sylvia Utami Tunjung Pratiwi, Triana Hertiani</i>	214
DOI:10.14499/indonesianjpharm29iss4pp214 Abstract view:113 PDF download:13	

Synthesis and Molecular Docking Studies of N'-benzoysalicylhydrazide Derivatives as Antituberculosis through InhA Enzyme Inhibition

Harry Santosa^{1*}, Galih Satrio Putra¹, Tegar Achsendo Yuniarta¹, Tutuk Budiati²

¹Faculty of Pharmacy,
University of Surabaya, Jalan
Raya Kali Rungkut Surabaya,
60293, Indonesia

²Faculty of Pharmacy,
Universitas Airlangga, Jalan
Dharmawangsa Dalam
Surabaya, 60286, Indonesia

Submitted: 03-10-2018

Revised: 23-11-2018

Accepted: 17-12-2018

*Corresponding author
Harry Santosa

Email:
harry_s@staff.ubya.ac.id

ABSTRACT

The specific aims of this study is to synthesize and to study the possible mechanism of N'-benzoysalicylhydrazide derivatives as an antituberculosis agent through InhA (Enoyl acyl carrier protein reductase) inhibition using *in silico* method. Five analogues of N'-benzoysalicylhydrazide were synthesized using microwave irradiation from methyl salicylate as starting material, which yielded 80-90% product on average. This indicates a considerable improvement in terms of effectivity and efficiency, compared to the more conventional method using reflux condition. Characterization of the compounds were subsequently carried out by UV, FTIR, ¹H-NMR, ¹³C-NMR spectroscopy, which confirmed that the compounds had been successfully synthesized. Ultimately, molecular docking was performed using Molegro Virtual Docker (MVD) on the active site of InhA enzyme to predict the activity of the compounds. The results showed that all compounds performed comparatively well against N-(4-Methylbenzoyl)-4-benzylpiperidine as the native ligand and also yielded lower docking score than isoniazide (INH). From this study it can be concluded that N'-benzoysalicylhydrazide derivatives could be synthesized using microwave irradiation with good product yield and all of the synthesized analogues are suggested to possess antituberculosis activity via InhA enzyme inhibition. *In vitro* activity will have to be determined in the future to validate whether N'-benzoysalicylhydrazide derivatives perform well as a potential antituberculosis agent.

Key words: Antituberculosis, Benzoysalicylhydrazide, Methyl Salicylate, InhA, Microwave synthesis, Molecular docking

INTRODUCTION

Lower respiratory tract infections were one of the top 10 deadliest diseases in 2010-2015. Several types of these diseases such as pneumonia, bronchitis, tuberculosis were ranked among the top 10 deadliest diseases in the world. According to WHO data, lower respiratory tract infections account for around 5.7% of deaths in the world or around 3.2 million people died in 2015 (Pietrangelo and Holland, 2018). One of the lower respiratory tract infections that has become a concern for medicals and researchers is tuberculosis infection. Tuberculosis infection accounts for half of the total mortality from lower respiratory tract infections. Therefore, this disease still ranks 9th out of the list of the top 10 deadliest diseases in the world. Based on

data from the WHO in 2015 around 1.3 million (2.4%) people died from this infection (Pietrangelo and Holland, 2018). Indonesia, as one of the most populated developing country in the world, is still facing this challenge. Based on the recent data in 2007-2013, the lower respiratory tract infections occurred frequently, with no significant change in the number of prevalence observed during this period (Indonesian Ministry of Health, 2103).

Mycobacterium tuberculosis had been found to be resistant to tuberculosis treatment with rifampin and isoniazid (INH) which is often referred to as multi-drugs resistance (MDR). Many cases of MDR has occurred during the treatment using rifampin and isoniazid as the main therapy (Dipiro *et al.*, 2008). Rifampin possesses bactericidal

mechanism against *Mycobacterium tuberculosis* as it binds strongly to the β -subunit of bacterial DNA-dependent RNA polymerase and thereby inhibits RNA synthesis (Katzung *et al.*, 2009). Resistance against rifampin can be caused by one of the several possible point mutations in *rpoB*, the gene for the β -subunit of RNA polymerase. These mutations prevent binding of rifampin to RNA polymerase (Katzung *et al.*, 2009). As for Isoniazid (INH), its mechanism as antituberculosis against *Mycobacterium tuberculosis* is via inhibition of mycolic acids synthesis, which plays an important role as major components of mycobacterial cell walls. Isoniazid is a prodrug that is activated by KatG, the mycobacterial catalase-peroxidase. The activated form of isoniazid exerts its lethal effect by forming a covalent complex with an acyl carrier protein (AcpM) and KasA, a β -ketoacyl carrier protein synthetase, which blocks mycolic acid synthesis (Katzung *et al.*, 2009). Resistance to isoniazid has been associated with mutations resulting in overexpression of *InhA*, which encodes an NADH-dependent acyl carrier protein reductase; mutation or deletion of KatG, promote mutations resulting in overexpression of *ahpC*. The existence of mutations in KatG expresses high levels of resistance in isoniazid (Katzung *et al.*, 2009).

To overcome MDR problems, it is necessary to develop new drugs to reduce the high cases of MDR in tuberculosis infections. Isoniazid has a simpler chemical structure compared to rifampin, therefore it is easier to develop this compound to overcome resistance in isoniazid. Some of hydrazide derivatives were reported to have antibacterial (Maheswari *et al.*, 2015; Kumar *et al.*, 2014; Thomas *et al.*, 2014) and antituberculosis activity (Maheswari *et al.*, 2015; Lourenço *et al.*, 2011). In this study, we synthesize five compounds of N^2 -benzoylsalicylhydrazide derivatives (Figure 1). Based on retrosynthetic analysis (Figure 2), we proposed to synthesize N^1 -benzoylsalicylhydrazide derivatives from methyl salicylate, hydrazine hydrate and benzoyl chloride as starting materials. Methyl salicylate would be reacted with hydrazine hydrate to produce an acylhydrazide compound, which subsequently reacted with benzoyl chloride to

yield the final products. Previous study in hydrazide derivative synthesis indicates stirring method at room temperature for 30min, while some others suggest the application of microwave irradiation (450-900watts) for 30-180s (Widiyana *et al.*, 2017). In this study, we applied microwave irradiation on the first stage of the reaction to obtain the optimal of time and product yield.




Figure 1. Isoniazid (a) and N^2 -benzoylsalicylhydrazide derivatives (b) chemical structure.




Figure 2. Retrosynthetic analysis of N^2 -benzoylsalicylhydrazide derivatives

MATERIAL AND METHODS

All reagents such as methyl salicylate, hydrazine, some of benzoyl chloride derivatives and all solvent such as THF were purchased from standard commercial suppliers. Reactions were monitored with TLC using pre-coated aluminum sheets with GF254 silica gel, 0.2mm layer thickness (E.Merck). Mobile phase used for TLC were chloroform: ethyl acetate (3:1) and the spots were visualized in UV chamber under UV light 254nm. Melting points of the synthesized compounds were measured with an Sybron Thermolyne MP 12615 melting point apparatus. UV spectra were obtained using Shimadzu UV-1800 spectrophotometer. IR spectra were obtained using a Perkin Elmer Spectrum spectrophotometer using KBr disks. $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectra were obtained on JEOL JNM-ECS400 ($^1\text{H-NMR}$: 400MHz, $^{13}\text{C-NMR}$: 100 MHz). Deuterated DMSO was used as solvent for these analysis.




Figure 3. Synthesis of *N'*-benzoylsalicylhydrazide derivatives

General synthesis

Synthesis process to obtain *N'*-benzoylsalicylhydrazide derivatives on this study (Figure 3). The first stage of reaction, methyl salicylate was reacted with hydrazine hydrate to obtain Salicylbenzohydrazide. The second stage of reaction, Salicylbenzohydrazide was reacted with some of benzoyl chloride derivatives to obtain *N'*-benzoylsalicylhydrazide derivatives. The detail of condition of reaction is explained in the subsequent part.

Synthesis of Compound 1 (Salicylbenzohydrazide)

Methyl salicylate (10mmol) was reacted with hydrazine hydrate (80%, 20mmol) and homogenized with magnetic stirrer. The mixture was heated in microwave irradiation at 360W for 3min until a solid mass was formed. The mixture was then cooled and distilled water (10mL) was added to the mixture. The separated solid was collected by filtration. The solid mass was crystallized by ethanol 96% to obtain Compound 1.

Synthesis of Compounds 2a-e (*N'*-benzoylsalicylhydrazide derivatives)

Compound 1 (8.5mmol) was added with 10 mL of THF as solvent and homogenized with magnetic stirrer. The mixture was then added by benzoyl chloride derivatives (9mmol) dropwise with constant stirring at 0°C for 30min, before the addition of 10% sodium bicarbonate solution. The latter was performed until the effervescence process ceased. The separated solid was allowed to settle down

before filtered off. Ultimately, the solid was crystallized by ethanol 96% to obtain compound 2a-e.

Molecular docking study

This study was performed in notebook equipped with processor AMD A9-9420 Radeon R5, RAM 4.00 GB and 64 bit operating system. Molegro Virtual Docker (MVD) ver 5.5 (CLC Bio) was utilized for molecular docking and CS ChemBioDraw Ultra ver 11.0 (Cambridge Soft) was used to prepare our compounds (2a-e) prior to docking process. The structure of InhA protein (PDB ID: 2NSD) was obtained from the Protein Data Bank (www.rcsb.org) (Xin *et al.*, 2007). Compounds 2a-e structures were built with ChemBioDraw Ultra 11.0 and their geometry optimization were performed using MMFF94 (Halgren, 1996). The validation of docking was carried out by redocking the native ligand of the enzyme (*N*-(4-Methylbenzoyl)-4-benzylpiperidine) into its active site. Criteria of acceptance is set with the value of Root Mean Square Deviation (RMSD) below 2.0 Å (Figure 4). After redocking process, the compounds 2a-e were docked into active site of InhA enzyme. The binding affinity between ligand and enzyme (docking score) was evaluated using Rerank Score, and we also compared the score of the native ligand with our compounds. Rerank Score is one of the evaluation methods of docking pose which is based on MolDock Score with some adjustments in the weight of each term of the score. In addition, sp²-sp² torsion and Lennard-Jones 12-6 potential were also introduced to further improve the accuracy




Figure 4. Comparison of its native ligand (green) with the docking result simulation (purple) by Molegro Virtual Docker (MVD) software Ver.5.5. The RMSD is 0.45Å




Figure 5. The mechanism reaction of the addition of nucleophilic hydrazine (-NH₂) to the electrophilic carbonyl group of the ester group




Figure 6. The mechanism reaction of the addition of nucleophilic (-NH₂) compound (1) to the electrophilic carbonyl group of the acyl carbonyl groups

of docking score (Thomsen and Christensen, 2006). Low score means smaller amount of energy required in forming drug-receptor interaction which leads to an assumption that the compound is more suitable to occupying active site of the receptor.

RESULT AND DISCUSSION

The reaction of methyl salicylate with hydrazine was performed by applying microwave irradiation at 360W for 3min. This procedure yielded (1) via nucleophilic addition of hydrazine (-NH₂) to the electrophilic carbonyl group of the ester group (methyl salicylate) (Figure 5). The reaction of compound (1) with some of benzoyl chloride derivatives (4-methoxy benzoyl chloride; 4-chloro benzoyl chloride; 2,4-dichloro benzoyl chloride; 3,4-dichloro benzoyl chloride) under water-free condition has successfully yielded several benzohydrazide derivatives (compounds 2a-e).

The mechanism of this reaction is the addition of nucleophilic (-NH₂) compound (1) to the electrophilic carbonyl group of the acyl carbonyl groups through S_NAcyl mechanism (Figure 6).








Compound 1 (Salicylbenzohydrazide)

Obtained in white powders (85 %), mp : UV λ_{max} (nm) : 233; FTIR (KBr) cm⁻¹ : 3320 (-OH); 3270 (-NH); 3055 (C-H sp²); 1647 (C=O); 1586 and 1485 (C=C aromatic); 1351 (C-N). ¹H-NMR (400 MHz, DMSO-d₆, δ, ppm): δ 12.48 (1H, s); δ 10.07 (2 H, s); δ 7.80 (1 H, d, 9, 1 Hz); δ 7.37 (1 H, t, 8.8 Hz); δ 6.89 (1 H, d, 8.5 Hz); δ 6.85 (1H; t, 8.1 Hz); δ 4.66 (1 H, s).

Compound 2a (N'-benzoyl-2-hydroxybenzohydrazide)

Obtained in white crystals (85 %), mp : 239-240 °C, UV λ_{max} (nm) : 302; FTIR (KBr) cm⁻¹ : 3302 (-OH); 3270 (-NH); 1655 (C=O); 1586 and 1485 (C=C aromatic); 1387(C-N).

Table I. Molecular docking result on InhA protein

Compound	Rerank score (kcal/mol)	Docked Pose	Hydrogen bond	Residues involved	Steric interaction	Residues involved
	-97.86	✓	1	Ile 194	2	Ala 157 Ile 202
4PI						
	-29.67	✓	2	Ile 194 Thr 196	3	Asp 148 Ala 191 Met 99
INH						
	-88.52	✓	1	Ile 194	-	-
(2a)						
	-94.56	✓	-	-	4	Phe 149 Tyr 158 Pro 193 Glu 219
(2b)						
	-80.84	✓	1	Ile 194	1	Ala 157
(2c)						
	-92.82	✓	-	-	3	Met 155 Met 199 Glu 219
(2d)						
	-89.21	✓	1	Ile 194	1	Leu 218
(2e)						

¹H-NMR (400MHz, DMSO-d₆, δ, ppm) : δ 11.91 (1H, s); δ 10.65 (2H, s); δ 7.89 (3 H, d, 10 Hz); δ 7.58 (1 H, t, 9.2 Hz); δ 7.51-7.41 (3 H, m); δ 6.95- 6.90 (2H, m); ¹³C-NMR (100 MHz, DMSO-d₆, δ, ppm) : δ 168.3; δ 1661.1; 159,8; δ 134,7; δ 132.8; δ 129.1 (2C); δ 128.8; δ 128.0 (2C); δ 119.6; δ 117.9; δ 115.0. All these spectral data are in agreement with the structure of compound *N'*-benzoyl-2-hydroxybenzohydrazide

Compound 2b (2-hydroxy-*N'*-(4-methoxybenzoyl)benzohydrazide)

Obtained in white powders (82 %), mp : 215-216°C ; UV λ_{max} (nm) : 301; FTIR (KBr) cm⁻¹ : 3309 (-OH); 3270 (-NH); 3054 (C-H sp²); 1606 (C=O); 1586 and 1483 (C=C

aromatic); 1284 (C-O); 1264 (C-N); 1174 (-CH sp³); ¹H-NMR (400 MHz, DMSO-d₆, δ, ppm) : δ 11.93 (1 H, s); δ 10,60 (1H, s); δ 10,49 (1H, s); 7,90-7,86 (3 H, m); δ 7.44-7.40 (1 H, m); δ 7.15 (1 H, d, 11,5 Hz); 6.94-6.92 (2H, m); δ 3.79 (3H, s); ¹³C-NMR (100 MHz, DMSO-d₆, δ, ppm) : δ 168.4; δ 155.6; δ 162.6; δ 134.7; δ 129.9 (2C); δ 128.7; δ 124.9; δ 119.6; δ 117.9; δ 115.0; δ 114.3 (2C); δ 55.9. All these spectral data are in agreement with the structure of compound 2-hydroxy-*N'*-(4-methoxybenzoyl)-benzohydrazide.

Compound 2c (*N'*-(4-chlorobenzoyl)-2-hydroxybenzohydrazide)

Obtained in white crystals (92%), mp : 233-234°C, UV λ_{max} (nm) : 302; FTIR (KBr)

cm⁻¹ : 3259 (-OH); 3270 (-NH); 1607 (C=O); 1565 and 1475 (C=C aromatic); 1175 (C-N); 737 (C-Cl); ¹H-NMR (400MHz, DMSO-d₆, δ, ppm) : δ 11.85 (1H, s); δ 10.90 (1H, s); δ 10.75 (1H, s); δ 7.89 (3 H, t, 10,2 Hz); δ 7.58 (2H, d, 11.0 Hz); δ 7.42 (1H, t, 8.7 Hz); δ 6.95-6.92 (2H, m); ¹³C-NMR (100 MHz, DMSO-d₆, δ, ppm) : δ 168.1; δ 165.1; δ 159.7; δ 137.4; δ 134.7; δ 131.5; δ 129.9 (2C); δ 129.2 (2C); δ 128.9; δ 119.6; δ 117.9; δ 115.1. All these spectral data are in agreement with the structure of compound N¹-(4-chlorobenzoyl)-2-hydroxy-benzohydrazide.

Compound 2d (2,4-dichloro-N¹-(2-hydroxybenzoyl)benzohydrazide)

Obtained in white crystals (81%), mp : 236-238 °C, UV λ_{max} (nm) : 294; FTIR (KBr) cm⁻¹ : 3321 (-OH); 3290 (-NH); 3065 (C-H sp²); 1635 (C=O); 1607 and 1475 (C=C aromatic); 1375 (C-N); 571 (C-Cl); ¹H-NMR (400MHz, DMSO-d₆, δ, ppm) : δ 11.87 (1H, s); δ 10.75 (1H, s); δ 10.71 (1H, s), δ 7.85 (1H, d, 10.0 Hz); δ 7.73 (1 H, s); δ 7.55-7.41 (3 H, m); δ 6.96-6.90 (2 H. m); ¹³C-NMR (100 MHz, DMSO-d₆, δ, ppm) : δ 167.7; δ 164.9; δ 159.6; δ 135.9; δ 134.8; δ 133.7; δ 132.3; δ 131.3; δ 130.1; δ 129.0; δ 128.0; δ 117.9; δ 115.0. All these spectral data are in agreement with the structure of compound 2,4-dichloro-N¹-(2-hydroxybenzoyl)benzohydrazide.

Compound 2e (3,4-dichloro-N¹-(2-hydroxybenzoyl)benzohydrazide)

Obtained in white crystals (93%), mp : 249-241 °C, UV λ_{max} (nm) : 300; FTIR (KBr) cm⁻¹ : 3190 (-OH); 3120 (-NH); 1606 (C=O); 1504 and 1470 (C=C aromatic); 1382 (C-N); 642 (C-Cl); ¹H-NMR (400MHz, DMSO-d₆, δ, ppm) : δ 11.80; δ 10.89; δ 10.70; δ 8.11 (1H, d, 3.0 Hz); δ 7.86-6.94 (6 H, m); ¹³C-NMR (100 MHz, DMSO-d₆, δ, ppm) : δ 167.9; δ 163.8; δ 159.5; δ 135.4; 134.8; δ 133.1; δ 132.1; δ 131.6; 129.9; δ 129.0; δ 128.3; δ 119.7; δ 117.9; δ 115.2 All these spectral data are in agreement with the structure of compound 3,4-dichloro-N¹-(2-hydroxybenzoyl) benzohydrazide

Molecular docking study

Compounds 2a-e were docked in to the active site of enoyl acyl carrier protein reductase (InhA) This enzyme was chosen,

since it catalyzes the reduction of long-chain trans-2-enoyl-ACP in the type II fatty acid biosynthesis pathway of Mycobacterium tuberculosis. Inhibition of InhA disrupts the biosynthesis of the mycolic acids that were main component of the mycobacterial cell wall. The mechanism of action of isoniazid is via inhibition of the enzyme. However, Mycobacterium tuberculosis has been found to be resistant to isoniazid. Compounds 2a-e are N¹-benzoyl-2-hydroxybenzohydrazide derivatives that were developed are found to be similar with isoniazid. Thus, compounds 2a-e were suggested to be potent to inhibit InhA.

Based on molecular docking result (Table I; Figure 7), it is shown that compounds 2a-e have lower docking score (-94.56 to -80.84kcal/mol) than isoniazid (-29.67kcal/mol) and some of the compounds possess comparable score to N-(4-Methylbenzoyl)-4-benzyl-piperidine (-97.86kcal/mol). Interaction between native ligand and isoniazid with InhA showed a hydrogen bond with Ile 194 residues and some steric interactions on residues such as Ala 157 and Ile 202. Compound 2a; 2c; 2e were predicted to form hydrogen bond interaction with Ile 194 and some steric interaction. Hydrogen bond on Ile 194 residue was predicted as one of the most important interaction for a ligand to inhibit InhA activity. Both N-(4-Methylbenzoyl)-4-benzylpiperidine and isoniazid interact with this residue in aforementioned manner. Based on the results of docking score and analysis of hydrogen bond and steric interactions with InhA,

compound 2a-e were suggested to have antituberculosis activity through the mechanism of inhibition of InhA enzyme. This prediction can be used as a foundation for further investigation in enzymatic and other in vitro assay.

CONCLUSION

The method to obtain N¹-benzoyl-2-hydroxybenzohydrazide derivatives by microwave irradiation is more time efficient with satisfactory product yield (80-90%). Compounds 2a-e are suggested to possess antituberculosis activity by inhibiting InhA enzyme based on molecular docking that have been conducted.




Figure 7. The Interaction between native ligand (4PI), INH, and compounds 2a-e into active site InhA protein

ACKNOWLEDGEMENT

We would like to thank Professor Siswandono from Airlangga University for license of docking program (Molegro Virtual Docker software Ver.5.5.)

REFERENCES

Indonesian Ministry of Health. 2013. Riset Kesehatan Dasar.
Joseph D., Robert L., Gary C., Gary R., Barbara G., Michael PL. 2008. *Pharmacotherapy: a pathophysiologic approach 7th*. New

York, McGraw-Hill Medical. pp 1839-2096.

Katzung BG., Masters SB., Trevor AJ. 2009. *Basic & Clinical Pharmacology Edition 11th*. New York: McGraw-Hill Companies. pp 1089-1093

Lourenço MCS., Souza MVN., Pinheiro A, C., Ferreira ML., Gonçalves RSB., Nogueira TC., Peralta MA. 2007. Evaluation of anti-tubercular activity of nicotinic and isoniazid analogues. *ARKIVOC* (xv) 181-191

- Maheswari R., Manjula J. 2015. Synthesis, Characterization and Biological Applications of Benzohydrazide derivatives. *Int J of App Res.* 1(10): 587-592
- Pietrangelo A., 2017. Holland K. The Top 10 Deadliest Diseases. Retrieved: September 5th, 2018, from <https://www.healthline.com/health/top-10-deadliest-diseases#cad>
- Pradeep K., Narasimhan B., Ramasamy K., Mani V., Mishra RK., Majeed ABA. 2014. Synthesis, antimicrobial, anticancer evaluation and QSAR studies of 2/3-bromo-N0-(substituted benzylidene/3-phenylallylidene)-benzohydrazides. *Arab J of Chemistry.* XXX.
- Thomas AB., Nanda RK., Kothapalli LP., Deshpande AV. 2011. Synthesis and Antimicrobial Activity of N-[2-(aryl/substituted aryl)-4-oxo-1,3-thiazolidin-3-yl]pyridine-4-carboxamide. *J of the Kor Chem Soc.* 55, 6, 960-968
- Thomas HA. 1996. Merck Molecular Force Field. I Basis, form, scope, parametrization, and performance of MMFF94. *J. Com. Chem.* 17(5-6): 490-519.
- Thomsen R., Christensen MH. 2006. MolDock: A New Technique for High-Accuracy Molecular Docking. *J. Med. Chem.* 49: 3315-3321.
- Widiyana AP., Putra GS., Sulistyowaty MI., Hardjono S., Budiati T. 2017. Method Optimization of 2,3-Disubstitutedquinazolin-4(3H)-One Derivatives. *J of Chem and Pharm Res.* 9(8):24-28
- Xin H., Alian, A., Montellano, PRO. 2007. Inhibition of the Mycobacterium tuberculosis enoyl acyl carrier protein reductase InhA by arylamides. *Bioorg Med Chem.* 15(21): 6649–6658