

Deep mantle H₂O recycling at 3.3Ga?

A.V. SOBOLEV^{1,2}, E.V. ASAFOV², A. A. GURENKO³, N. T. ARNDT², V.G. BATANOVA^{1,2}, M.V. PORTNYAGIN^{2,4}, D. GARBE-SCHÖNBERG⁵, S. P. KRASHENINNIKOV², A. H. WILSON⁶ AND G.R. BYERLY⁷ ¹²³

¹ UGA, France; alexander.sobolev@univ-grenoble-alpes.fr

² Vernadsky Institute, Russia; evasafov@gmail.com

³ CRPG, Nancy, France, agurenko@crpg.cnrs-nancy.fr

⁴ GEOMAR, Kiel, Germany; mportnyagin@geomar.de

⁵ CAU Kiel Univ., Germany; d.garbe-schoenberg@gmx.de

⁶ University of the Witwatersrand; Allan.Wilson@wits.ac.za

⁷ LSU, Baton Rouge, USA; glbyer@lsu.edu

Knowledge of water in the Earth's interior and its evolution through time is of great importance for understanding global geodynamics. MgO-rich ultramafic magmas – komatiites - result from high-degree mantle melting at high pressures and thus are excellent probes of H₂O contents in the deep mantle. Recently, a significant excess of H₂O over elements of similar behaviour in magmatic processes (e.g. Ce) was found in melt inclusions in Mg-rich olivine from 2.7 Ga komatiites (Sobolev et al, Nature 2016; Asafov et al, Chem. Geol. 2018). These data were interpreted as the evidence of a deep hydrated mantle reservoir, probably at the transition zone, at the Neoproterozoic time. In this paper we confirm the mantle origin of elevated H₂O in komatiites using H isotope compositions of the melt inclusions and present similar data for 3.3 Ga old Weltevreden komatiites (S. Africa). The reconstructed initial hydrogen isotope composition of komatiites ($\delta D < -110\text{‰}$) is significantly more depleted in deuterium than surface reservoirs and typical mantle and argues that dehydrated subducted lithosphere was the source of H₂O accumulated in the transition zone by ringwoodite or wadsleyite (Bercovici & Karato, Nature, 2003). This interpretation is supported by a significant Cl excess and the temporal trend of Pb/Ce in the mantle sources of komatiites. These results argue that subduction occurred on the Earth well before 3.3 Ga.