

Received: 21 June 2018


Revised: 10 October 2018

Accepted: 13 October 2018

DOI: 10.1111/jeb.13393

## SHORT COMMUNICATION

WILEY

JOURNAL OF Evolutionary Biology 

## Niche differentiation among clones in asexual grass thrips

Casper J. van der Kooi\*  | Karim Ghali\*  | David Amptmeijer | Tanja Schwander 

Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland

**Correspondence**

Casper J. van der Kooi, University of Groningen, Groningen Institute for Evolutionary Life Sciences, Nijenborgh 7, NL-9747 AG Groningen, The Netherlands.  
Email: C.J.van.der.Kooi@rug.nl

**Present address**

Casper J. van der Kooi, University of Groningen, Groningen Institute for Evolutionary Life Sciences, Groningen, The Netherlands

**Funding information**

This research was funded by Grant PP00P3\_139013 of the Swiss National Science Foundation (SNSF).

**Abstract**

Many asexual animal populations comprise a mixture of genetically different lineages, but to what degree this genetic diversity leads to ecological differences remains often unknown. Here, we test whether genetically different clonal lineages of *Aptinothrips* grass thrips differ in performance on a range of plants used as hosts in natural populations. We find a clear clone-by-plant species interactive effect on reproductive output, meaning that clonal lineages perform differently on different plant species and thus are characterized by disparate ecological niches. This implies that local clonal diversities can be driven and maintained by frequency-dependent selection and that resource heterogeneity can generate diverse clone assemblies.

**KEY WORDS**

clonal diversity, frozen niche variation, general-purpose genotype, parthenogenesis, Poaceae, reproduction

## 1 | INTRODUCTION


The continuous generation of genetic variation for fitness is regarded as one of the main explanations for the predominance of sex in eukaryotes (Bell, 1982). Theoretical studies have shown that under some conditions, sex can be favoured over asexuality because it creates novel and/or rare genotypes (Bell, 1982; Otto, 2009). Such novel and rare genotypes are favoured under negative frequency-dependent selection or in habitats with spatially heterogeneous or temporally fluctuating resources. For example, one component of the “parasite hypothesis for sex” implies negative frequency-dependent selection, as it predicts that parasites evolve to track common host genotypes (Dagan, Liljeroos, Jokela, & Ben-Ami, 2013; Decaestecker et al., 2007; Morran, Schmidt, Gelarden, Parrish, & Lively, 2011). Similar ideas are available for selection imposed by resources; that is rare genotypes are able to exploit resources where competition is low (reviewed by Scheu & Drossel, 2007).

Theories for the maintenance of sex generally assume genetically uniform populations of asexuals, whereas in nature, many asexual animal and plant populations comprise genetically distinct

clones (Bell, 1982; Ellstrand & Roose, 1987; Jokela, Dybdahl, & Lively, 1999; Parker, 1979). Indeed, high genetic diversity is the case in many asexual populations (Fontcuberta García-Cuenca, Dumas, & Schwander, 2016; Kearney et al., 2009; Moritz, Donnellan, Adams, & Baverstock, 1989; Neiman, Jokela, & Lively, 2005; Pannebakker, Zwaan, Beukeboom, & Van Alphen, 2004; Tully & Ferrière, 2008; Vorburger, Lancaster, & Sunnucks, 2003), but whether clonal diversity is generally associated with phenotypic and ecological diversity remains unknown. Genetic diversity in asexuals can be due to lineage diversification after the transition to asexuality (owing to mutation accumulation) and because asexuals have different sexual ancestors; that is they arose via independent transitions to asexuality (Janko, Drozd, Flegr, & Pannell, 2008). Such independent transitions are especially likely to generate ecological differentiation (Bell, 1982; Janko et al., 2008; King, Jokela, & Lively, 2011). More generally, it will be difficult to explain the maintenance of sex via its effect on genetic diversity if sexual species compete with genetically diverse rather than genetically uniform asexuals (Judson, 1997).

In this study, we test whether genetic differentiation in asexual *Aptinothrips* (Thysanoptera: Thripidae) grass thrips is neutral or

\*Shared first authorship.



**FIGURE 1** Number of offspring produced by different asexual *Aptinothrips* lineages on seven host plant species. Every panel depicts the asexual lineages of an independently evolved asexual clade (identified in van der Kooi and Schwander (2014)). For every lineage, the mean  $\pm$  95% confidence interval (assuming a normal distribution) is shown

ecologically relevant. *Aptinothrips* is a system where asexuality has evolved several times independently and is associated with a large geographic distribution and extreme genetic diversity (van der Kooi & Schwander, 2014). We test whether thirteen genetically different clonal lineages of *Aptinothrips rufus* and *Aptinothrips stylifer* occupy different ecological niches by examining their performance on seven common host plant species. In a breeding experiment and using the number of offspring produced as a proxy for fitness, we show that different clonal lineages indeed feature different ecological niches.

## 2 | MATERIALS AND METHODS

### 2.1 | Study species and breeding

*Aptinothrips* grass thrips are small (1–2 mm), wingless insects that feed on various species of Poaceae. For *A. rufus*, asexuality evolved at least twice independently (van der Kooi & Schwander, 2014) as inferred from phylogenies based on mitochondrial sequences. Asexual populations of this species can be found all over the world and exhibit very high levels of mitochondrial diversity, whereas sexual populations only occur in the Mediterranean region and exhibit a relatively low mitochondrial diversity (Fontcuberta García-Cuenca et al., 2016; van der Kooi & Schwander, 2014). The species *A. stylifer* evolved asexuality independently from *A. rufus* (van der Kooi & Schwander, 2014), and different populations also comprise a pool of genetically different individuals (see Results). Nine and four genetically different females of *A. rufus* and *A. stylifer* were isolated from three locations in Switzerland (Supporting Information). These individuals were used to establish thirteen iso-female lines (hereafter: clonal lineages); for each lineage, third-generation individuals were used in the experiment.

### 2.2 | Experimental design

We tested the performance of asexual females on seven grass species that are commonly found in natural habitats: meadow brome (*Bromus erectus*), alpine bluegrass (*Poa alpina*), common meadowgrass (*Poa pratensis*), red fescue (*Festuca rubra*), sheep fescue (*Festuca ovina*), Timothy grass (*Phleum pratense*) and orchard grass (*Dactylis glomerata*). For each clonal lineage, 7–14 replicate individuals were tested on each host plant species; this yielded a total of 735 replicates (see Supporting Information). The experiment was performed in July–August 2016; for logistic reasons, it was performed in two batches separated by 4 days.

### 2.3 | Data analysis

To test for ecological variation among species and clonal lineages of *Aptinothrips*, we used linear mixed models (LMMs) as implemented in the R package *lme4* (Bates, Maechler, Bolker, & Walker, 2014) in R 3.4.2. We tested whether thrips species, independent transitions to asexuality within *A. rufus*, clonal lineage, the grass species and their interactions affected the number of offspring. Batch was included as a random factor in the model. Significance was tested using a permutation ANOVA (Supporting Information).

## 3 | RESULTS

We observed clear differences in the number of offspring produced by asexual *Aptinothrips* (Figure 1). There was an effect of thrips species ( $df = 1$ ,  $p < 0.01$ ) and an effect of clade, that is, independent transitions to asexuality ( $df = 2$ ,  $p < 0.01$ ), on the number of offspring produced. In line with our hypothesis of ecological divergence,

**TABLE 1** Overview of species that show ecological diversification of clonal lineages

Asexual species investigated	Common name	Approach	Results	Reference
<i>Erigeron annuus</i>	Eastern daisy fleabane	Experimentally test the number and size of seeds, biomass allocation and seed germination for different lineages.	Variation in seed size, allocation to roots and seed germination for different lineages.	Stratton (1991)
<i>Taraxacum officinale</i>	Dandelion	Experimentally test the effect of drought and shade on biomass allocation for different lineages.	Lineages often perform differently under different conditions.	Oplaat and Verhoeven (2015)
<i>Rhopalosiphum padi</i>	Cherry oat aphid	Experimentally test fecundity and longevity of clones on different plant species at two temperatures.	Small differences between lineages; performance in laboratory experiment is correlated with abundance in natural populations.	Valenzuela, Ridland, Weeks, and Hoffmann (2010)
<i>Myzus persicae</i>	Peach potato aphid	Experimentally test reproductive success of lineages on different host plant species.	Lineages perform differently on different host plant species.	Vorburger, Sunnucks, and Ward (2003)
<i>Folsomia candida</i>	Springtail	Experimentally test reproductive investment and survival for different lineages.	Significant variation between lineages in number and size of offspring produced.	Tully and Ferrière (2008)
<i>Daphnia pulex</i>	Water flea	Experimentally test lineage performance in different locations with different salinities.	Lineages perform differently in different regions, consistent with local adaptation.	Weider and Hebert (1987)
<i>Potamopyrgus antipodarum</i>	Fresh water snail	Field sampling in different populations.	In different populations, different lineages feature specific dietary preferences.	Jokela et al. (1999)
<i>Heterontia binoei</i>	Gecko	Field sampling of lineages in natural populations with different climates.	Lineages with different origins occupied different geographic regions and niches.	Kearney, Moussalli, Strasburg, Lindenmayer, and Moritz (2003)
<i>Lepidodactylus lugubris</i>	Gecko	Field sampling of lineages in natural habitats.	Two common lineages feature different niches, though with some overlap.	Hanley, Bolger, and Case (1994)
<i>Aptinothrips rufus</i> & <i>A. stylifer</i>	Grass thrips	Experimentally test performance of different lineages on seven common host plant species.	Lineages perform differently on different host plant species	This study

different clonal lineages are characterized by different ecologies, as revealed by significant clone-by-plant species interaction effects on clonal lineage performances ( $df = 72$ ,  $p < 0.01$ ). Ecological differences among clonal lineages are not solely explained by independent transitions to asexuality as clonal lineages within one *A. rufus* clade (clade B in Figure 1) also differ from each other ( $df = 12$ ,  $p = 0.04$ ); the small sample sizes in *A. stylifer* preclude performing a similar analysis.

## 4 | DISCUSSION

Many asexual populations comprise a large number of different clonal lineages (Bell, 1982; Parker, 1979), but whether genetically different clones feature different ecologies is largely unknown. In this study, we showed that different clonal *Aptinothrips* lineages feature different ecological niches (Figure 1). Ecological divergence of clonal lineages is pronounced both within and between independently derived asexual clades. Although assessing the relative importance of independent transitions versus within clade divergence requires deeper sampling, the multiple origins of asexuality in *Aptinothrips* (van der Kooi & Schwander, 2014) have increased the genetic diversity in asexuals and likely contribute to their extreme ecological success.

The facts that asexual *Aptinothrips* populations exhibit a higher genetic diversity than sexuals (Fontcuberta García-Cuenca et al., 2016; van der Kooi & Schwander, 2014) and that this genetic diversity translates to ecological divergence (this study) may explain why asexuals outcompete sexuals in mesocosm experiments (Lavanchy et al., 2016). Our findings further suggest that locations with different grass species would select for different sets of clonal lineages. High grass species diversity is expected to promote and maintain clonal diversity in the thrips population, because more grass species reflect more niches available for different clonal lineages. An interesting open question is whether temporal fluctuations in plant species frequency lead to fluctuating clone frequencies, as was previously found in the aphid *Myzus persicae* by Vorburger (2006).

A review of literature showed that clonal diversity is associated with ecological diversity in a suite of at least nine additional plant and animal species (Table 1). This includes aquatic and terrestrial asexual systems with very diverse ecologies. Taken together, these studies convincingly show that populations of clonal animals comprise ecologically relevant genetic variation and that clonal diversity should therefore not be disregarded in theory on the maintenance of sex in nature.

Asexuals are frequently more ecologically successful than related sexuals, which is often believed to be due to the reproductive assurance conferred by asexuality (e.g., Bell, 1982), but this may not be the sole explanation. For example, a recent large-scale comparative study showed that asexuals have 2–3.5 times as many host species and occur in up to five times as many countries compared to their sexual counterparts, highlighting the (ecological) success of asexuality (van der Kooi, Matthey-Doret, & Schwander, 2017). Our

current study shows that in addition to the reproductive advantage of asexuality, high genetic diversity can contribute to the success of asexuals.

To summarize, genetic diversity in asexual *Aptinothrips* lineages is associated with ecological diversity. Under such ecological differentiation of clonal lineages, which appears to occur in a suite of asexual animal and plant systems (Table 1), natural selection may maintain clonal diversity rather than sex. Future theory on the maintenance of sex would benefit from incorporating clonal diversity, different clonal diversity-generating mechanisms and their ecological significance.

## ACKNOWLEDGMENTS

We acknowledge the members of the Schwander group for interesting discussions.

## CONFLICT OF INTEREST

The authors declare no competing interests.

## AUTHOR CONTRIBUTIONS

CJvdK, KG and TS designed the study. CJvdK, DA and KG carried out the experiment and analysed the data. CJvdK and KG wrote the first draft of the manuscript. All authors edited the manuscript and approved the final version.

## ORCID

Casper J. Kooi  <https://orcid.org/0000-0003-0613-7633>

Karim Ghali  <https://orcid.org/0000-0003-1509-7602>

Tanja Schwander  <https://orcid.org/0000-0003-1945-5374>

## REFERENCES

- Bates, D., Maechler, M., Bolker, B., & Walker, S. (2014). lme4: Linear mixed-effects models using Eigen and S4. *R package version, 1*, 1–23.
- Bell, G. (1982). *The masterpiece of nature: The evolution and genetics of sexuality*. Berkeley, CA: University of California Press.
- Dagan, Y., Liljeroos, K., Jokela, J., & Ben-Ami, F. (2013). Clonal diversity driven by parasitism in a freshwater snail. *Journal of Evolutionary Biology, 26*, 2509–2519. <https://doi.org/10.1111/jeb.12245>
- Decaestecker, E., Gaba, S., Raeymaekers, J. A., Stoks, R., Van Kerckhoven, L., Ebert, D., & De Meester, L. (2007). Host-parasite 'Red Queen' dynamics archived in pond sediment. *Nature, 450*, 870. <https://doi.org/10.1038/nature06291>
- Ellstrand, N. C., & Roose, M. L. (1987). Patterns of genotypic diversity in clonal plant species. *American Journal of Botany, 74*, 123–131. <https://doi.org/10.1002/j.1537-2197.1987.tb08586.x>
- Fontcuberta García-Cuenca, A., Dumas, Z., & Schwander, T. (2016). Extreme genetic diversity in asexual grass thrips populations. *Journal of Evolutionary Biology, 29*, 887–899. <https://doi.org/10.1111/jeb.12843>
- Hanley, K. A., Bolger, D. T., & Case, T. J. (1994). Comparative ecology of sexual and asexual gecko species (*Lepidodactylus*) in French

- Polynesia. *Evolutionary Ecology*, 8, 438–454. <https://doi.org/10.1007/BF01238194>
- Janko, K., Drozd, P., Flegel, J., & Pannell, J. R. (2008). Clonal turnover versus clonal decay: A null model for observed patterns of asexual longevity, diversity and distribution. *Evolution*, 62, 1264–1270. <https://doi.org/10.1111/j.1558-5646.2008.00359.x>
- Jokela, J., Dybdahl, M., & Lively, C. (1999). Habitat-specific variation in life-history traits, clonal population structure and parasitism in a freshwater snail (*Potamopyrgus antipodarum*). *Journal of Evolutionary Biology*, 12, 350–360. <https://doi.org/10.1046/j.1420-9101.1999.00035.x>
- Judson, O. P. (1997). A model of asexuality and clonal diversity: Cloning the Red Queen. *Journal of Theoretical Biology*, 186, 33–40. <https://doi.org/10.1006/jtbi.1996.0339>
- Kearney, M., Fujita, M., Ridenour, J., Schön, I., Martens, K., & van Dijk, P. (2009). Lost sex in the reptiles: Constraints and correlations. In I. Schön, P. van Dijk, & K. Martens (Eds.), *Lost sex: The evolutionary biology of parthenogenesis* (pp. 447–474). Berlin, Germany: Springer. <https://doi.org/10.1007/978-90-481-2770-2>
- Kearney, M., Moussalli, A., Strasburg, J., Lindenmayer, D., & Moritz, C. (2003). Geographic parthenogenesis in the Australian arid zone: I. A climatic analysis of the *Heteronotia binoei* complex (Gekkonidae). *Evolutionary Ecology Research*, 5, 953–976.
- King, K. C., Jokela, J., & Lively, C. M. (2011). Parasites, sex, and clonal diversity in natural snail populations. *Evolution*, 65, 1474–1481. <https://doi.org/10.1111/j.1558-5646.2010.01215.x>
- Lavanchy, G., Strehler, M., Llanos Roman, M. N., Lessard-Therrien, M., Humbert, J. Y., Dumas, Z., ... Schwander, T. (2016). Habitat heterogeneity favors asexual reproduction in natural populations of grasshoppers. *Evolution*, 70, 1780–1790. <https://doi.org/10.1111/evo.12990>
- Moritz, C., Donnellan, S., Adams, M., & Baverstock, P. (1989). The origin and evolution of parthenogenesis in *Heteronotia binoei* (Gekkonidae): Extensive genotypic diversity among parthenogens. *Evolution*, 43, 994–1003. <https://doi.org/10.1111/j.1558-5646.1989.tb02545.x>
- Morran, L. T., Schmidt, O. G., Gelarden, I. A., Parrish, R. C., & Lively, C. M. (2011). Running with the Red Queen: Host-parasite coevolution selects for biparental sex. *Science*, 333, 216–218. <https://doi.org/10.1126/science.1206360>
- Neiman, M., Jokela, J., & Lively, C. (2005). Variation in asexual lineage age in *Potamopyrgus antipodarum*, a New Zealand snail. *Evolution*, 59, 1945–1952. <https://doi.org/10.1111/j.0014-3820.2005.tb01064.x>
- Oplaat, C., & Verhoeven, K. J. (2015). Range expansion in asexual dandelions: Selection for general-purpose genotypes? *Journal of Ecology*, 103, 261–268. <https://doi.org/10.1111/1365-2745.12347>
- Otto, S. P. (2009). The evolutionary enigma of sex. *The American Naturalist*, 174, S1–S14. <https://doi.org/10.1086/599084>
- Pannebakker, B. A., Zwaan, B. J., Beukeboom, L. W., & Van Alphen, J. J. (2004). Genetic diversity and Wolbachia infection of the *Drosophila* parasitoid *Leptopilina clavipes* in western Europe. *Molecular Ecology*, 13, 1119–1128. <https://doi.org/10.1111/j.1365-294X.2004.02147.x>
- Parker, E. D. (1979). Ecological implications of clonal diversity in parthenogenetic morphospecies. *American Zoologist*, 19, 753–762. <https://doi.org/10.1093/icb/19.3.753>
- Scheu, S., & Drossel, B. (2007). Sexual reproduction prevails in a world of structured resources in short supply. *Proceedings of the Royal Society of London B*, 274, 1225–1231. <https://doi.org/10.1098/rspb.2007.0040>
- Stratton, D. A. (1991). Life history variation within populations of an asexual plant, *Erigeron annuus* (Asteraceae). *American Journal of Botany*, 78, 723–728. <https://doi.org/10.1002/j.1537-2197.1991.tb12596.x>
- Tully, T., & Ferrière, R. (2008). Reproductive flexibility: Genetic variation, genetic costs and long-term evolution in a Collembola. *PLoS ONE*, 3, e3207. <https://doi.org/10.1371/journal.pone.0003207>
- Valenzuela, I., Ridland, P. M., Weeks, A. R., & Hoffmann, A. A. (2010). Patterns of genetic variation and host adaptation in an invasive population of *Rhopalosiphum padi* (Hemiptera: Aphididae). *Annals of the Entomological Society of America*, 103, 886–897. <https://doi.org/10.1603/AN09113>
- van der Kooi, C. J., Matthey-Doret, C., & Schwander, T. (2017). Evolution and comparative ecology of parthenogenesis. *Evolution Letters*, 1, 304–316. <https://doi.org/10.1002/evl3.30>
- van der Kooi, C. J., & Schwander, T. (2014). Evolution of asexuality via different mechanisms in grass thrips (Thysanoptera: Aptinothrips). *Evolution*, 68, 1883–1893. <https://doi.org/10.1111/evo.12402>
- Vorburger, C. (2006). Temporal dynamics of genotypic diversity reveal strong clonal selection in the aphid *Myzus persicae*. *Journal of Evolutionary Biology*, 19, 97–107. <https://doi.org/10.1111/j.1420-9101.2005.00985.x>
- Vorburger, C., Lancaster, M., & Sunnucks, P. (2003). Environmentally related patterns of reproductive modes in the aphid *Myzus persicae* and the predominance of two 'superclones' in Victoria, Australia. *Molecular Ecology*, 12, 3493–3504. <https://doi.org/10.1046/j.1365-294X.2003.01998.x>
- Vorburger, C., Sunnucks, P., & Ward, S. A. (2003). Explaining the coexistence of asexuals with their sexual progenitors: No evidence for general-purpose genotypes in obligate parthenogens of the peach-potato aphid, *Myzus persicae*. *Ecology Letters*, 6, 1091–1098. <https://doi.org/10.1046/j.1461-0248.2003.00536.x>
- Weider, L. J., & Hebert, P. D. (1987). Ecological and physiological differentiation among low-arctic clones of *Daphnia pulex*. *Ecology*, 68, 188–198. <https://doi.org/10.2307/1938819>

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** van der Kooi CJ, Ghali K, Amptmeijer D, Schwander T. Niche differentiation among clones in asexual grass thrips. *J Evol Biol*. 2019;32:126–130. <https://doi.org/10.1111/jeb.13393>