



Open Archive TOULOUSE Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in : <http://oatao.univ-toulouse.fr/>
Eprints ID : 10390

To cite this version :

Mougel, Jerome and Fabre, David and Lacaze, Laurent *Instability of a potential swirling flow with a free surface*. (2012) In: 9th European Fluid Mechanics Conference - EFMC 9, 09 September 2012 - 13 September 2012 (Rome, Italy). (Unpublished)

Any correspondence concerning this service should be sent to the repository administrator: staff-oatao@listes-diff.inp-toulouse.fr

Instability of a potential swirling flow with a free surface

Jérôme Mougel ^{*†}, David Fabre^{*}, Laurent Lacaze^{*‡}

Rotating flows with free surface are commonly observed in everyday-life, the most popular example being the drain vortex observed in bathtubs and sinks. These flows are known to support spectacular patterns arising from breaking of the axial symmetry. A situation which has been particularly investigated through experiments is the case of a partially filled container with a rotating bottom. In this situation, an instability leads to the appearance of rotating polygons along the free surface ^{1 2}. The axisymmetric base flow preexisting the instability is rather complex, involving strong recirculations, but includes a region where the rotation is approximately potential, ³ suggesting that the potential vortex may be a relevant model to investigate the instability mechanism.

Consequently, we investigate here the linear inviscid stability of the potential swirling flow with a free surface; namely, the flow defined $V_\theta = \Gamma/(2\pi r)$, with a free surface given by $h_0(r) \sim \frac{1}{R_1^2} - \frac{1}{r^2}$, where R_1 corresponds to the radius of the dry area. For the study, we use both a global stability approach based on the finite element software Freefem++ and a shallow water approach. Our results show that unstable modes exist in narrow ranges of the Froude number (defined by $Fr = \Gamma/2\pi\sqrt{gR_0^3}$, R_0 being the radius of the cylinder), for all non-zero values of the azimuthal wavenumber m (see figure 1). The instability seems to be related to the "strato-rotational" instability mechanism existing in stratified rotating fluids⁴.

^{*}Institut de Mécanique des Fluides de Toulouse (IMFT), Université de Toulouse, 2 Allée du Professeur Camille Soula, 31400, Toulouse, France

[†]Direction Générale de l'Armement (DGA), France

[‡]CNRS, IFMT

¹Vatistas et al, *PRL* **100**, 174503 (2008).

²Jansson et al, *PRL* **96**, 174502 (2006).

³Tophøj, *Phd Thesis*, Danmark Technical University (2012).

⁴Le Dizés and Riedinger, *J. Fluid Mech.* **660**, 147 (2010).




Figure 1: (a) 3D free surface reconstruction of the most amplified mode for an azimuthal wave number $m = 5$ (b) Amplification rate as a fonction of the Froude number for $m = 5$ (the liquid volume in the container is kept constant as the Froude number is varied).