



Open Archive TOULOUSE Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in : <http://oatao.univ-toulouse.fr/>
Eprints ID : 10470

To cite this version : Abbas, Micheline and Loisel, Vincent and Masbernat, Olivier and Climent, Eric. *Laminar-turbulent transition of channel flows: the effect of neutrally buoyant finite-size particles*. (2013) In: 66th Annual APS-DFD Meeting, 24 November 2013 - 26 November 2013 (Pittsburg, United States).

Abstract Submitted
for the DFD13 Meeting of
The American Physical Society

Laminar-turbulent transition of channel flows: the effect of neutrally buoyant finite-size particles¹ MICHELINE ABBAS, VINCENT LOISEL, OLIVIER MASBERNAT, University of Toulouse - Laboratoire de Genie Chimique, ERIC CLIMENT, University of Toulouse - Institut de Mecanique des Fluides de Toulouse — Numerical simulations were performed on channel flows laden with resolved finite-size neutrally buoyant particles at moderate volumetric concentration. In the case of fluctuating flows close to laminar-turbulent transition, the particle volume fraction is homogeneously distributed in the channel except an accumulation layer in the near-wall region (particle migration is driven by inertia). Particles increase the level of perturbations close to the wall leading to significant enhancement of both the velocity fluctuations and the wall friction coefficient. Additionally, particles break down the large-scale flow structures into smaller, more numerous and sustained eddies. When the flow Reynolds number is decreased, flow relaminarization occurs at critical Reynolds number Re_{cS} (based on the effective suspension viscosity) significantly below the critical Reynolds number Re_c of single-phase flow transition. In the case of laminar flows, the suspension segregates into pure fluid and particle laden wall layers due to cross-stream migration. An instability is observed characterized by the formation of dune-like patterns at the separation between pure fluid and concentrated suspension. Increasing the Reynolds number yields transition to turbulence at a threshold above Re_{cS} .

¹This work was granted access to the HPC resources of CALMIP and GENCI under the allocations 2012-P1002 and x20132a6942 respectively.

Micheline Abbas
University of Toulouse - Laboratoire de Genie Chimique