

UNIVERSITY OF BIRMINGHAM

Research at Birmingham

On the integer max-linear programming problem

Butkovic, Peter; MacCaig, Marie

DOI:

[10.1016/j.dam.2013.08.007](https://doi.org/10.1016/j.dam.2013.08.007)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Butkovic, P & MacCaig, M 2014, 'On the integer max-linear programming problem', *Discrete Applied Mathematics*, vol. 162, pp. 128–141. <https://doi.org/10.1016/j.dam.2013.08.007>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Eligibility for repository : checked 02/04/2014

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



On the integer max-linear programming problem[☆]



Peter Butkovič^{*}, Marie MacCaig

School of Mathematics, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

ARTICLE INFO

Article history:

Received 7 December 2012

Received in revised form 19 July 2013

Accepted 13 August 2013

Available online 13 September 2013

Keywords:

Max-linear system

Integer vector

Max-linear program

ABSTRACT

Max-linear programs have been used to describe optimisation problems for multiprocessor interactive systems. In some instances the variables used in this model are required to be integer; however, no method seems to exist for finding integer solutions to max-linear programs.

For a generic class of matrices, we show that integer solutions to two-sided max-linear systems and programs can be found in polynomial time. For general matrices, we adapt the existing methods for finding real solutions to obtain algorithms for finding integer solutions.

© 2013 The Authors. Published by Elsevier B.V. All rights reserved.

1. Introduction

This paper deals with the task of finding integer solutions to two-sided max-linear systems and the integer max-linear programming problem.

In max-algebra, for $a, b \in \bar{\mathbb{R}} = \mathbb{R} \cup \{-\infty\}$, we define $a \oplus b = \max(a, b)$, $a \otimes b = a + b$ and extend the pair (\oplus, \otimes) to matrices and vectors in the same way as in linear algebra, that is (assuming compatibility of sizes),

$$(A \oplus B)_{ij} = a_{ij} \oplus b_{ij},$$

$$(A \otimes B)_{ij} = \bigoplus_k a_{ik} \otimes b_{kj} \quad \text{and}$$

$$(\alpha \otimes A)_{ij} = \alpha \otimes a_{ij}.$$

Throughout this paper we will use ε to denote the zero element $-\infty$ as well as any vector or matrix whose every entry is $-\infty$.

Except for complexity arguments, all multiplications in this paper are in max-algebra, and where appropriate we will omit the \otimes symbol.

A two-sided max-linear system (TSS) is of the form

$$Ax \oplus c = Bx \oplus d,$$

where $A, B \in \bar{\mathbb{R}}^{m \times n}$ and $c, d \in \bar{\mathbb{R}}^m$. If $c = d = \varepsilon$, then we say that the system is *homogeneous*; otherwise, it is called *nonhomogeneous*. Nonhomogeneous systems can be transformed to homogeneous systems [3]. If $B \in \bar{\mathbb{R}}^{m \times k}$, a system of the form

$$Ax = By$$

is called a *system with separated variables*.

[☆] This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

^{*} Corresponding author. Tel.: +44 1214146600; fax: +44 1214143389.

E-mail addresses: P.Butkovic@bham.ac.uk, peter.butkovic@virgin.net (P. Butkovič), mxm779@bham.ac.uk (M. MacCaig).

The problems of finding solutions to

$$Ax = By \tag{1.1}$$

and

$$Ax = Bx \tag{1.2}$$

have been studied previously, and one solution approach is to use the Alternating Method [3,9]. If A and B are integer matrices, then the solution found by the Alternating Method is integer; however, this cannot be guaranteed if A and B are real.

In Section 3, we show that we can adapt the Alternating Method in order to obtain algorithms which determine whether integer solutions to these problems exist for real matrices A and B , and find one if it exists. Note that various other methods for solving a TSS are known [1,5,12], but none of them has been proved polynomial, and there is no obvious way of adapting them to integrality constraints. In Section 4, we show that, for a certain class of matrices, which represents a generic case, the problem of finding an integer solution to both systems can be solved in strongly polynomial time, and give a method in this case.

If $f \in \mathbb{R}^n$, then the function $f(x) = f^T \otimes x$ is called *max-linear*. *Max-linear programming problems* (MLPs) seek to minimise or maximise a max-linear function subject to constraints given by max-linear equations. Note that unlike in linear programming there is no obvious way of converting maximisation of max-linear functions to minimisation of the same type of functions, and vice versa.

For $A, B \in \mathbb{R}^{m \times n}$, $c, d \in \mathbb{R}^m$, $f \in \mathbb{R}^n$, the *integer max-linear program* (IMLP) is given by

$$\begin{aligned} f^T \otimes x &\rightarrow \min \text{ or } \max \\ \text{s.t. } Ax \oplus c &= Bx \oplus d \\ x &\in \mathbb{Z}^n. \end{aligned}$$

Solution methods to solve the MLP are known; for example, in [3,4], a bisection method is applied to obtain an algorithm that finds an approximate solution to the MLP. Also, a Newton-type algorithm has been designed [10] to solve a more general max-linear fractional programming problem by a reduction to a sequence of mean payoff games. Again, an integer solution is found for any instances of the MLP with integer entries, but the problem with integrality constraints is very different if the entries are real. In Section 5, we develop an algorithm based on the bisection method which will find an optimal solution to the IMLP, or determine that no solution exists. In Section 6, we show that, in the generic case, the IMLP can be solved in polynomial time.

It should be noted that the solution set of a max-linear system with all variables on one side can be described in terms of set coverings and can easily be solved [2,3]. Consequently, the corresponding systems with integrality constraints can also be easily solved, as can max-linear programs and integer max-linear programs with constraints of this type.

As an example of an application of a TSS and the IMLP, consider the following *multiprocessor interactive system* (MPIS) as described in [4]; see also [8].

Products P_1, \dots, P_m are made up of a number of components which are prepared using n processors. Each processor contributes to the final product P_i by producing one of its components. We assume that each processor can work on a component for every product simultaneously, and that work begins on all products as soon as the processor is switched on.

Let a_{ij} be the time taken for the j th processor to complete its component for P_i ($i = 1, \dots, m$; $j = 1, \dots, n$). Denote the starting time of the j th processor by x_j ($j = 1, \dots, n$). Then, for each product P_i , all of its components will be completed at time $\max(x_1 + a_{i1}, \dots, x_n + a_{in})$.

Now suppose that, independently, k other processors prepare components for products Q_1, \dots, Q_m and that the duration and starting times of these are b_{ij} and y_j , respectively. Then the *synchronisation problem* is to find starting times of all $n + k$ processors so that each pair (P_i, Q_i) ($i = 1, \dots, m$) is completed at the same time. This task is equivalent to solving the system of equations

$$\max(x_1 + a_{i1}, \dots, x_n + a_{in}) = \max(y_1 + b_{i1}, \dots, y_k + b_{ik}) \quad (i = 1, \dots, m).$$

If, additionally, we require that P_i and Q_i are not completed before a particular time, c_i and d_i , say, then the equations are

$$\max(x_1 + a_{i1}, \dots, x_n + a_{in}, c_i) = \max(y_1 + b_{i1}, \dots, y_k + b_{ik}, d_i) \quad (i = 1, \dots, m),$$

which is equivalent to the two-sided system $Ax \oplus c = By \oplus d$.

When solving the MPIS it may be required that the starting times are restricted to discrete values, in which case we would want to look for integer solutions to the TSS.

In applications it may also be required that the starting times of the MPIS are optimised with respect to a given criterion. In this paper, we consider the case when the objective function is also max-linear and it has to be either minimised or maximised. As an example, it may be required that all processors in an MPIS begin as soon [late] as possible; that is, the latest starting time of a processor is as small [big] as possible. In this case, we would set $f_j = 0$ and seek to minimise [maximise] $f(x) = \max(x_1, \dots, x_n)$.

With this extra requirement we obtain the following MLP:

$$f^T \otimes x \rightarrow \min \text{ or } \max$$

$$\text{s.t. } Ax \oplus c = Bx \oplus d.$$

If the starting times are restricted to discrete values, then the MLP is transformed to an IMLP.

2. Preliminaries

We will use the following standard notation and terminology. The symbol α^{-1} stands for $-\alpha$, and for a vector γ we use γ^{-1} to mean the vector with entries γ_i^{-1} . A vector [matrix] whose every entry belongs to \mathbb{R} is called *finite* as is any scalar from \mathbb{R} . If a matrix has no ε rows [columns], then it is called *row [column] \mathbb{R} -astic*, and it is called *doubly \mathbb{R} -astic* if it is both row and column \mathbb{R} -astic [3,8].

For $a \in \mathbb{R}$, the *fractional part* of a is $\text{fr}(a) := a - \lfloor a \rfloor$. For a matrix $A \in \overline{\mathbb{R}}^{m \times n}$, we use $\lfloor A \rfloor$ ($\lceil A \rceil$) to denote the matrix with (i, j) entry equal to $\lfloor a_{ij} \rfloor$ ($\lceil a_{ij} \rceil$), and similarly for vectors. We define $\lfloor \varepsilon \rfloor = \varepsilon = \lceil \varepsilon \rceil$.

For positive integers m, n, k , we denote $M = \{1, \dots, m\}$, $N = \{1, \dots, n\}$ and $K = \{1, \dots, k\}$. If $A = (a_{ij}) \in \overline{\mathbb{R}}^{n \times n}$, then $\lambda(A)$ denotes the *maximum cycle mean*; that is,

$$\lambda(A) = \max \left\{ \frac{a_{i_1 i_2} + \dots + a_{i_l i_1}}{l} : (i_1, \dots, i_l) \text{ is a cycle, } l = 1, \dots, n \right\},$$

where $\max(\emptyset) = \varepsilon$ by definition. Note that this definition is independent of whether we allow cycles to contain repeated nodes [3]. The maximum cycle mean can be calculated in $\mathcal{O}(n^3)$ time [3,11].

An $n \times n$ matrix is called *diagonal*, written $\text{diag}(d_1, \dots, d_n) = \text{diag}(d)$, if its diagonal entries are $d_1, \dots, d_n \in \mathbb{R}$ and off-diagonal entries are ε . We use I to denote the identity matrix, $I = \text{diag}(0, \dots, 0)$, of appropriate size.

If $a, b \in \overline{\mathbb{R}} = \mathbb{R} \cup \{+\infty\}$, then we define $a \oplus' b = \min(a, b)$ and $a \otimes' b = a + b$ if at least one of a, b is finite, $(-\infty) \otimes (+\infty) = (+\infty) \otimes (-\infty) = -\infty$ and $(-\infty) \otimes' (+\infty) = (+\infty) \otimes' (-\infty) = +\infty$. The pair of operations (\oplus', \otimes') is extended to matrices and vectors similarly as (\oplus, \otimes) . For $A \in \overline{\mathbb{R}}^{m \times n}$, we define $A^\# = -A^T \in \overline{\mathbb{R}}^{n \times m}$. It can be shown [3,8] that $(A \otimes B)^\# = B^\# \otimes' A^\#$.

Next, we give an overview of some basic properties.

Lemma 2.1. *Let $A \in \overline{\mathbb{R}}^{m \times n}$, $x \in \mathbb{R}^n$.*

- (a) *If A is row \mathbb{R} -astic, then $A \otimes x$ is finite.*
- (b) *If A is column \mathbb{R} -astic, then $A^\# \otimes' x$ is finite.*

Proof. Straightforward from the definitions. \square

Lemma 2.2 ([3,8]). *If $A \in \overline{\mathbb{R}}^{m \times n}$ and $x, y \in \overline{\mathbb{R}}^n$, then*

$$x \leq y \Rightarrow A \otimes x \leq A \otimes y \quad \text{and} \quad A \otimes' x \leq A \otimes' y.$$

Corollary 2.3 ([3,8]). *If $A, B \in \overline{\mathbb{R}}^{m \times n}$ and $x \leq y$, then*

$$B^\# \otimes' (A \otimes x) \leq B^\# \otimes' (A \otimes y).$$

Lemma 2.4 ([3]). *Let $A, B \in \overline{\mathbb{R}}^{m \times n}$, $c, d \in \overline{\mathbb{R}}^m$. Then there exists $x \in \mathbb{R}^n$ satisfying $Ax \oplus c = Bx \oplus d$ if and only if there exists $z \in \mathbb{R}^{n+1}$ satisfying $(A|c)z = (B|d)z$.*

The cancellation law in max-algebra is as follows.

Lemma 2.5 ([3]). *Let $v, w, a, b \in \mathbb{R}$, $a > b$. Then, for any real number x , we have*

$$v \oplus a \otimes x = w \oplus b \otimes x \Leftrightarrow v \oplus a \otimes x = w.$$

For any matrices of compatible sizes [3,8],

$$X \otimes (X^\# \otimes' Y) \leq Y, \tag{2.1}$$

$$X \otimes (X^\# \otimes' (X \otimes Z)) = X \otimes Z. \tag{2.2}$$

If $A \in \overline{\mathbb{R}}^{m \times n}$ and $b \in \mathbb{R}^m$, then, for all $j \in N$, define

$$M_j(A, b) = \left\{ t \in M : a_{ij} \otimes b_t^{-1} = \max_i a_{ij} \otimes b_t^{-1} \right\}.$$

Proposition 2.6 ([7]). Let $A \in \overline{\mathbb{R}}^{m \times n}$, $b \in \mathbb{R}^m$, and $\bar{x} = A^\# \otimes' b$.

(a) An integer solution to $Ax \leq b$ exists if and only if \bar{x} is finite. If an integer solution exists, then all integer solutions can be described as the integer vectors x satisfying $x \leq \bar{x}$.

(b) An integer solution to $Ax = b$ exists if and only if

$$\bigcup_{j:\bar{x}_j \in \mathbb{Z}} M_j(A, b) = M.$$

If an integer solution exists, then all integer solutions can be described as the integer vectors x satisfying $x \leq \bar{x}$ with

$$\bigcup_{j:x_j=\bar{x}_j} M_j(A, b) = M.$$

We define $\hat{x} = \lfloor A^\# \otimes' b \rfloor$. Then, from Proposition 2.6 and (2.2), we conclude the following.

Corollary 2.7. Let $A \in \overline{\mathbb{R}}^{m \times n}$, $b \in \overline{\mathbb{R}}^m$, $c \in \mathbb{Z}^n$. Then the following hold.

- (a) \hat{x} is the greatest integer solution to $Ax \leq b$ (provided \hat{x} is finite).
- (b) $Ax = b$ has an integer solution if and only if \hat{x} is an integer solution.
- (c) $A \otimes \lfloor A^\# \otimes' (A \otimes c) \rfloor = A \otimes c$.

Consider the matrix inequality $AX \leq B$, where $A \in \overline{\mathbb{R}}^{m \times n}$, $B \in \overline{\mathbb{R}}^{m \times k}$, $X \in \overline{\mathbb{R}}^{n \times k}$, and let $\hat{X} = \lfloor A^\# \otimes' B \rfloor$. This system can be written as a set of inequalities of the form $Ax \leq b$ in the following way, using the notation X_r, B_r to denote the r th column of X and B , respectively:

$$AX_r \leq B_r, \quad r = 1, \dots, k.$$

This allows us to state the following result.

Corollary 2.8. Let $A \in \overline{\mathbb{R}}^{m \times n}$, $B \in \overline{\mathbb{R}}^{m \times k}$, $C \in \mathbb{Z}^{n \times k}$. Then the following hold.

- (a) \hat{X} is the greatest integer solution to $AX \leq B$ (provided \hat{X} is finite); that is, $A \otimes \lfloor A^\# \otimes' B \rfloor \leq B$.
- (b) $AX = B$ has an integer solution if and only if \hat{X} is an integer solution.
- (c) $A \otimes \lfloor A^\# \otimes' (A \otimes C) \rfloor = A \otimes C$.

A vector $x \in \overline{\mathbb{R}}^n - \{\varepsilon\}$ [$x \in \mathbb{Z}^n$] satisfying $Ax \leq \lambda x$ is called a [integer] subeigenvector of A with respect to subeigenvalue λ . In this paper, we only need results on finite subeigenvectors. The set of all [integer] subeigenvectors with respect to subeigenvalue λ is denoted

$$V^*(A, \lambda) = \{x \in \mathbb{R}^n : Ax \leq \lambda x\}$$

$$[IV^*(A, \lambda) = \{x \in \mathbb{Z}^n : Ax \leq \lambda x\}].$$

The existence of [integer] subeigenvectors can be determined, and the whole set can be described, using the following result.

Theorem 2.9 ([3,7]). Let $A \in \overline{\mathbb{R}}^{n \times n}$, $\lambda \in \mathbb{R}$.

- (i) $V^*(A, \lambda) \neq \emptyset$ if and only if $\lambda(\lambda^{-1}A) \leq 0$.
- (ii) If $V^*(A, \lambda) \neq \emptyset$, then $V^*(A, \lambda) = \{\lambda^{-1}A^*u : u \in \mathbb{R}^n\}$.
- (iii) $IV^*(A, \lambda) \neq \emptyset$ if and only if $\lambda(\lceil \lambda^{-1}A \rceil) \leq 0$.
- (iv) If $IV^*(A, \lambda) \neq \emptyset$, then $IV^*(A, \lambda) = \{\lceil \lambda^{-1}A \rceil^*z : z \in \mathbb{Z}^n\}$.

Corollary 2.10. For $A \in \overline{\mathbb{R}}^{n \times n}$, it is possible to decide whether $IV^*(A, \lambda) \neq \emptyset$ in $\mathcal{O}(n^3)$ time.

3. Alternating Method for integer solutions

In this section, we show that the Alternating Method [3,9] can be easily adapted to design algorithms that determine whether integer solutions to (1.1) or (1.2) exist, and if so find one. This method will be crucial for the solution of the IMLP. We first detail an algorithm to solve systems with separated variables, and then a second algorithm to solve general systems. Since the justification behind the construction of the two algorithms in this section is similar to the arguments in [3,9], we only outline the key results here; full details can be found in [6].

If the i th row of either A or B is ε , then we have $(Ax)_i = \varepsilon = (By)_i$, which, since x and y are finite, means that the i th row of the other matrix is also ε . Thus we may remove the redundant i th equation from the equality. If, instead, either of A or B has an ε column, then this column may be removed without affecting the solution. Hence we assume without loss of generality that A, B are doubly \mathbb{R} -astic.

For any matrix $Y \in \mathbb{R}^{m \times n}$, let

$$K(Y) = \left[\max\{|y_{ij}| : i \in M, j \in N\} \right]. \quad (3.1)$$

We propose the following algorithm to find integer solutions to the system with separated variables (1.1).

Algorithm: SEP-INT-TSS

Input: $A \in \bar{\mathbb{R}}^{m \times n}$, $B \in \bar{\mathbb{R}}^{m \times k}$ doubly \mathbb{R} -astic, any starting vector $x(0) \in \mathbb{Z}^n$.

Output: An integer solution (x, y) to $Ax = By$ or indication that no such solution exists.

1. $r := 0$.
2. $y(r) := \lfloor B^\# \otimes' (A \otimes x(r)) \rfloor$.
3. $x(r+1) := \lfloor A^\# \otimes' (B \otimes y(r)) \rfloor$.
4. If $x_i(r+1) < x_i(0)$ for all $i \in N$, then STOP (no solution).
5. If $A \otimes x(r+1) = B \otimes y(r)$, then STOP (solution found).
6. Go to 2.

Theorem 3.1 ([6]). *Algorithm SEP-INT-TSS is correct and terminates after*

$$\mathcal{O}(mn(n+k)K(A))$$

operations, if applied to instances where the matrix A is finite.

Lemma 2.4 allows us to write any general two-sided system as a homogeneous system, (1.2), so to find integer solutions to general systems it is sufficient to develop a method to solve homogeneous systems. The following statement is obvious.

Proposition 3.2. *Let $A, B \in \bar{\mathbb{R}}^{m \times n}$. The problem of finding $x \in \mathbb{Z}^n$ satisfying $Ax = Bx$ is equivalent to finding $x \in \mathbb{Z}^n, y \in \mathbb{R}^m$ such that*

$$\begin{pmatrix} A \\ B \end{pmatrix} x = \begin{pmatrix} I \\ I \end{pmatrix} y,$$

where $I \in \bar{\mathbb{R}}^{m \times m}$.

We propose the following algorithm to find integer solutions to (1.2).

Algorithm: GEN-INT-TSS

Input: $A', B' \in \bar{\mathbb{R}}^{m \times n}$ doubly \mathbb{R} -astic, $I \in \bar{\mathbb{R}}^{m \times m}$, any starting vector $x(0) \in \mathbb{Z}^n$.

Output: A solution $x \in \mathbb{Z}^n$ to $A'x = B'x$ or indication that no such vectors exist.

1. $r := 0, A := \begin{pmatrix} A' \\ B' \end{pmatrix}, B := \begin{pmatrix} I \\ I \end{pmatrix}$.
2. $y(r) := B^\# \otimes' (A \otimes x(r))$.
3. $x(r+1) := \lfloor A^\# \otimes' (B \otimes y(r)) \rfloor$.
4. If $x_i(r+1) < x_i(0)$ for all $i \in N$, then STOP (no solution).
5. If $A \otimes x(r+1) = B \otimes y(r)$, then STOP (solution found).
6. Go to (2).

Theorem 3.3 ([6]). *Algorithm GEN-INT-TSS is correct and terminates after*

$$\mathcal{O}(K(A'|B')(mn(m+n)))$$

operations, if applied to instances where both of the matrices A', B' are finite.

4. A polynomially solvable case

In this section, we give a generic condition on the matrices A, B , which, if satisfied, means that we can determine in polynomial time whether a solution to (1.1) or (1.2) exists, and if so find one. We then show that the method for these matrices can be extended to find integer solutions to any two-sided systems, but at a cost to efficiency.

4.1. Description of the generic case: Property OneFP

A key observation is that, if $Ax = z = By$, where x and y are integer vectors, then, for each i , there exist j, t such that

$$\text{fr}(z_i) = \text{fr}(a_{ij}) = \text{fr}(b_{it}).$$

Recall that we assume without loss of generality that A, B are doubly \mathbb{R} -astic.

We define the following properties on a pair of matrices (A, B) .

- (a) We say that the pair (A, B) satisfies Property *ZeroFP* if there exists $i \in M$ such that there is no pair (a_{ij}, b_{it}) with the same fractional part.
- (b) We say that the pair (A, B) satisfies Property *One⁺FP* if, for each $i \in M$, there is at least one pair (a_{ij}, b_{it}) with the same fractional part.
- (c) We say that the pair (A, B) satisfies Property *OneFP* if, for each $i \in M$, there is exactly one pair (a_{ij}, b_{it}) with the same fractional part and $a_{ij}, b_{it} \in \mathbb{Z}$ (this can be assumed without loss of generality).

Further, for either a TSS or the IMLP, we will say that the system satisfies Property *OneFP* if the pair of input matrices (A, B) satisfies Property *OneFP*.

Proposition 4.1. Let $A \in \overline{\mathbb{R}}^{m \times n}, B \in \overline{\mathbb{R}}^{m \times k}$. If an integer solution to $Ax = By$, or (if $n = k$) $Ax = Bx$, exists, then

$$(\forall i \in M)(\exists j \in N, t \in K) \quad \text{fr}(a_{ij}) = \text{fr}(b_{it}).$$

Proof. Assume that $x \in \mathbb{Z}^n, y \in \mathbb{Z}^k$ satisfy $Ax = By$. Then

$$(\forall i \in M) \quad \max_j(a_{ij} + x_j) = \max_j(b_{ij} + y_j).$$

Therefore, for each i , there exist $r(i), r'(i) \in N$ such that

$$\text{fr}(a_{i,r(i)} + x_{r(i)}) = \text{fr}(b_{i,r'(i)} + y_{r'(i)}).$$

But $\text{fr}(a_{i,r(i)} + x_{r(i)}) = \text{fr}(a_{i,r(i)})$ and $\text{fr}(b_{i,r'(i)} + y_{r'(i)}) = \text{fr}(b_{i,r'(i)})$. Hence

$$(\forall i)(\exists j \in N, t \in K) \quad \text{fr}(a_{ij}) = \text{fr}(b_{it}). \quad \square$$

So a necessary condition for a two-sided system to have an integer solution is that (A, B) satisfies Property *One⁺FP*, and therefore any TSS where the pair (A, B) satisfies Property *ZeroFP* has no integer solutions. The aim of this section is to show that, when the pair (A, B) satisfies Property *OneFP*, the problem of finding integer solutions can be solved in polynomial time.

For such a pair of matrices, we define the pair $(r(i), r'(i))$ to be the indices such that $\text{fr}(a_{i,r(i)}) = \text{fr}(b_{i,r'(i)})$, $i = 1, \dots, m$.

Without loss of generality we may assume that the entries $(a_{i,r(i)}, b_{i,r'(i)})$ are integer, and that no other entries in the equation for either matrix are integer (this is since we may subtract a constant from each row of the system without affecting the answer to the question).

If we randomly generated two real matrices $A \in \overline{\mathbb{R}}^{m \times n}, B \in \overline{\mathbb{R}}^{m \times k}$, then we would expect there to be very few pairs of entries $(a_{ir(i)}, b_{ir'(i)})$ which share the same fractional part. So, when given a random two-sided system, the most likely outcome is that there is no integer solution (that the matrices satisfy Property *ZeroFP*). Among matrices which satisfy Property *One⁺FP*, the next most likely outcome is that the pair (A, B) satisfies Property *OneFP*. While this discussion is not mathematically rigorous, it does allow us to conclude that matrices (A, B) satisfying either Property *ZeroFP* or Property *OneFP* represent a generic case.

We say that an element a_{ij} of A is *active* with respect to x if $a_{ij} + x_j = (Ax)_i$, and inactive otherwise. A consequence of Proposition 4.1 is the following.

Corollary 4.2. Let $A \in \overline{\mathbb{R}}^{m \times n}, B \in \overline{\mathbb{R}}^{m \times k}$ satisfy Property *OneFP*. Then the entries $a_{i,r(i)} [b_{i,r'(i)}]$ are the only possible active entries in the matrix $A [B]$ with respect to any integer solution $x [y]$ satisfying $Ax = By$.

4.2. Systems with separated variables

Let $A \in \overline{\mathbb{R}}^{m \times n}, B \in \overline{\mathbb{R}}^{m \times k}$. First, we consider the question of whether there exist $x \in \mathbb{Z}^n, y \in \mathbb{Z}^k$ such that $Ax = By$ when (A, B) satisfies Property *OneFP*.

Observe that

$$Ax = z^{-1} = By \Leftrightarrow \text{diag}(z)Ax = 0 = \text{diag}(z)By.$$

Thus we conclude the following.

Proposition 4.3. Let $A \in \overline{\mathbb{R}}^{m \times n}, B \in \overline{\mathbb{R}}^{m \times k}$ satisfy Property *OneFP*. Then (x, y) is an integer solution to (1.1) if and only if there exists $z \in \mathbb{Z}^m$ such that (x, y) satisfy

$$\text{diag}(z) \otimes A \otimes x = 0 \tag{4.1}$$

and

$$\text{diag}(z) \otimes B \otimes y = 0. \tag{4.2}$$

Proposition 4.4. Let $A \in \overline{\mathbb{R}}^{m \times n}$, $B \in \overline{\mathbb{R}}^{m \times k}$ satisfy Property OneFP. Suppose that $z \in \mathbb{Z}^m$ satisfies (4.1) and (4.2) for some integer vectors x, y . If there exists a column of $\text{diag}(z)A$ containing more than one integer entry, then these entries are equal. Similarly for $\text{diag}(z)B$.

Proof. Assume that z satisfies (4.1). Then, for each i , the entry $a_{i,r(i)}$ is the only active entry of A in the i th row (equivalently, $z_i + a_{i,r(i)}$ is the only active entry in the i th row of $\text{diag}(z)A$). This implies that, if there exist $i, t \in M$ such that $r(i) = r(t)$, then

$$z_i + a_{i,r(i)} + x_{r(i)} = 0 = z_t + a_{t,r(t)} + x_{r(t)} \Rightarrow z_i + a_{i,r(i)} = z_t + a_{t,r(t)}. \quad \square$$

Proposition 4.5. Simultaneously solving (4.1) and (4.2) is equivalent to the problem of finding $z \in \mathbb{Z}^m$ satisfying

$$(\forall i, t \in M) \quad z_i - z_t \geq \lceil a_{t,r(i)} \rceil - a_{i,r(i)}$$

and

$$(\forall i, t \in M) \quad z_i - z_t \geq \lceil b_{t,r'(i)} \rceil - b_{i,r'(i)}.$$

Proof. Consider (4.1). We have that, for each i , the integer entry $a_{i,r(i)} + z_i$ is the only possible active entry of $\text{diag}(z)A$ with respect to an integer vector x . From Proposition 2.6 we have that an integer solution to (4.1) exists exactly when the integer column maxima of $\text{diag}(z)A$ cover all rows. A similar argument holds for (4.2). Hence we require that

$$a_{i,r(i)} + z_i > a_{t,r(i)} + z_t \quad \text{for } t \neq i \text{ and } a_{t,r(i)} \notin \mathbb{Z}, \tag{4.3}$$

$$a_{i,r(i)} + z_i = a_{t,r(i)} + z_t \quad \text{for } a_{t,r(i)} \in \mathbb{Z}, \tag{4.4}$$

$$b_{i,r'(i)} + z_i > b_{t,r'(i)} + z_t \quad \text{for } t \neq i \text{ and } b_{t,r'(i)} \notin \mathbb{Z} \quad \text{and}$$

$$b_{i,r'(i)} + z_i = b_{t,r'(i)} + z_t \quad \text{for } b_{t,r'(i)} \in \mathbb{Z}.$$

For any other column (those not containing integer entries) we do not get any additional constraints, since we may set the corresponding entry of x or y to be small enough so that the column has no effect on the product Ax or By .

This set of inequalities is equivalent to

$$\forall i, t \in M \quad a_{i,r(i)} + z_i \geq \lceil a_{t,r(i)} \rceil + z_t \tag{4.5}$$

and

$$\forall i, t \in M \quad b_{i,r'(i)} + z_i \geq \lceil b_{t,r'(i)} \rceil + z_t. \tag{4.6}$$

To see this, note that (4.3) and (4.4) imply (4.5). For the other direction, assume that (4.5) holds. If $a_{t,r(i)} \notin \mathbb{Z}$, then we have $a_{i,r(i)} + z_i > a_{t,r(i)} + z_t$, as required. If, instead, $a_{t,r(i)} \in \mathbb{Z}$, then $r(t) = r(i)$, and from (4.5) we have

$$a_{i,r(i)} + z_i \geq \lceil a_{t,r(i)} \rceil + z_t \quad \text{and} \quad a_{t,r(t)} + z_t \geq \lceil a_{i,r(t)} \rceil + z_i,$$

which together imply equality. Similar arguments hold for the inequalities with entries from B .

The result is obtained by rearranging inequalities (4.5) and (4.6). \square

Let $W = (w_{ij}) \in \overline{\mathbb{Z}}^{m \times m}$, where

$$w_{ij} = \max(\lceil a_{j,r(i)} \rceil - a_{i,r(i)}, \lceil b_{j,r'(i)} \rceil - b_{i,r'(i)}).$$

Then, by Proposition 4.5, for solving (1.1), we need to determine whether there exists $z \in \mathbb{Z}^m$ satisfying

$$(\forall i, j \in M) z_i - z_j \geq w_{ij}$$

$$\Leftrightarrow (\forall i) \max_j (w_{ij} + z_j) \leq z_i$$

$$\Leftrightarrow W \otimes z \leq z.$$

This is exactly the condition for $z \in IV^*(W, 0)$, which can be checked using Theorem 2.9. We conclude the following.

Theorem 4.6. Let $A \in \overline{\mathbb{R}}^{m \times n}$, $B \in \overline{\mathbb{R}}^{m \times k}$ satisfy Property OneFP. For all $i, j \in M$, let

$$w_{ij} = \max(\lceil a_{j,r(i)} \rceil - a_{i,r(i)}, \lceil b_{j,r'(i)} \rceil - b_{i,r'(i)}).$$

Then an integer solution to $Ax = By$ exists if and only if $\lambda(W) \leq 0$. If this is the case, then $Ax = By = z^{-1}$, where $z \in IV^*(W, 0)$, and x and y can be found using Proposition 2.6.

Note that, for all $i \in M$, $w_{ii} = 0$, and therefore W is doubly \mathbb{R} -astic and $\lambda(W) \geq 0$. Therefore the condition for existence of an integer solution is equivalent to $\lambda(W) = 0$.

From Theorem 4.6 and Corollary 2.10, we obtain the following.

Corollary 4.7. For $A \in \overline{\mathbb{R}}^{m \times n}$, $B \in \overline{\mathbb{R}}^{m \times k}$, it is possible to decide whether an integer solution to $Ax = By$ exists in $\mathcal{O}(m^3 + n + k)$ time.

4.3. General two-sided systems

We now consider finding integer solutions to (1.2) under the condition that (A, B) satisfy Property OneFP. The following statement is obvious.

Proposition 4.8. *Let $A, B \in \overline{\mathbb{R}}^{m \times n}$ satisfy Property OneFP. The problem of finding $x \in \mathbb{Z}^n$ such that $Ax = Bx$ is equivalent to finding $x \in \mathbb{Z}^n, y \in \mathbb{Z}^n$ such that*

$$\begin{pmatrix} A \\ I \end{pmatrix} x = \begin{pmatrix} B \\ I \end{pmatrix} y.$$

Observe that, if (A, B) satisfies Property OneFP, then so does (\hat{A}, \hat{B}) , where

$$\hat{A} = \begin{pmatrix} A \\ I \end{pmatrix}, \quad \hat{B} = \begin{pmatrix} B \\ I \end{pmatrix}.$$

Thus to solve a general two-sided system satisfying Property OneFP we may convert it into a system with separated variables and solve using Theorem 4.6. By Corollary 4.7, we have the following.

Corollary 4.9. *For $A, B \in \overline{\mathbb{R}}^{m \times n}$ satisfying Property OneFP, we can decide whether an integer solution to $Ax = Bx$ exists in $\mathcal{O}((m+n)^3)$ time.*

Remark 1. The transformation described in Proposition 3.2 is not suitable here, since it has $y \in \mathbb{R}$, whereas to use Theorem 4.6 we want to be able to look for integer solutions. Conversely, the transformation described in Proposition 4.8 is not suitable to use when discussing the Alternating Method, since we need at least one of the matrices to be finite for our complexity arguments to hold.

4.4. Some special cases

We now present a couple of cases where we can give simpler conditions, both for systems with separated variables.

The first case occurs when we have $r(1) = \dots = r(m)$ or $r'(1) = \dots = r'(m)$. We assume without loss of generality that the former occurs.

Proposition 4.10. *Assume that $A \in \overline{\mathbb{R}}^{m \times n}, B \in \overline{\mathbb{R}}^{m \times k}$ satisfy Property OneFP. Suppose further that $r(1) = \dots = r(m) = p$. Let A' and B' be the matrices obtained from A and B by subtracting a_{ip} from the i th row. Then an integer solution to $Ax = By$ exists if and only if*

$$B' \otimes \lfloor (B')^\# \otimes' 0 \rfloor = 0.$$

Proof. Observe that an integer solution to $Ax = By$ exists if and only if an integer solution to $A'x = B'y$ exists.

Assume first that x, y are integer vectors satisfying $A'x = B'y$. Now, from Corollary 4.2, we know that the active entries in A' with respect to x are the zero entries in column p . Thus $A' \otimes x = (x_p, x_p, \dots, x_p)^T$.

Therefore $B'y = (x_p, x_p, \dots, x_p)^T$, which implies that $B(x_p^{-1} \otimes y) = 0$, and hence, using Corollary 2.7, we know that $B' \otimes \lfloor (B')^\# \otimes' 0 \rfloor = 0$.

For the other direction, assume that $B' \otimes \lfloor (B')^\# \otimes' 0 \rfloor = 0$. Choosing $x \in \mathbb{Z}^n$ such that $x = (x_1, \dots, x_{p-1}, 0, x_{p+1}, \dots, x_n)^T$ with x_j small for $j \neq p$ gives us that $A' \otimes x = 0$, and thus setting $y = \lfloor (B')^\# \otimes' 0 \rfloor$ gives $A' \otimes x = B' \otimes y$, as required. \square

Remark 2. If $r(1) = \dots = r(m) = p$ and $r'(1) = \dots = r'(m) = q$, then the only candidates for active entries are found in columns A_p and B_q . So, if $Ax = By$, then $x_p A_p = y_q B_q$, and the other components of x and y are small enough not to affect the outcome. Thus a solution to (1.1) exists if and only if A_p is a max-multiple of B_q .

The second case occurs when A, B are square, satisfy Property OneFP, and for one matrix the active entries are spread over all columns. Without loss of generality, assume that it is matrix A , so $\{r(1), \dots, r(m)\} = M$.

Proposition 4.11. *Assume that $A, B \in \overline{\mathbb{R}}^{m \times m}$ satisfy Property OneFP. Suppose further that $r(i) \neq r(t)$ for all $i, t \in M$ with $i \neq t$. Let A' be obtained from A by subtracting $a_{i,r(i)}$ from row i for each i and permuting the columns so that the zero entries appear on the leading diagonal. If*

$$\lambda(\lceil A' \rceil) \neq 0,$$

then no integer solution to $A \otimes x = B \otimes y$ exists.

Proof. Let B' be obtained from B by subtracting $a_{i,r(i)}$ from row i for each i . Assume that an integer solution to $Ax = By$ exists. Then an integer solution to $A'x = B'y$ exists, and the active entries in A' are the zeros on the diagonal by Corollary 4.2. Thus $(A'x)_i = a_{ii} + x_i = x_i$, and hence $x \in IV^*(A', 0)$. By Theorem 2.9 we have $\lambda(\lceil A' \rceil) = 0$. \square

4.5. Adapting the method to solve any TSS

We end this section by giving a brief description of how the solution method for systems satisfying Property OneFP could be adapted to find integer solutions to any TSS, but that in doing so we may lose efficiency. Since we can convert any general two-sided system into a system with separated variables, we discuss systems with separated variables only.

Let $A \in \mathbb{R}^{m \times n}$, $B \in \mathbb{R}^{m \times k}$, and suppose that (A, B) satisfies Property One⁺FP. In this case, for each row i of $Ax = By$, we will have a number of pairs

$$(a_{ir(i,s)}, b_{ir'(i,s)})$$

for some integer $s \leq nk$, such that $\text{fr}(a_{ir(i,s)}) = \text{fr}(b_{ir'(i,s)})$. Observe that, for any $x \in \mathbb{Z}^n$, $y \in \mathbb{Z}^k$ satisfying $Ax = By$, we can identify a single pair of active entries for each row of the equation, and hence the pair (x, y) is also an integer solution to the system $Ax = B^-y$, where (A, B^-) satisfies Property OneFP and B^- is obtained from B by subtracting a small real number $0 < \delta \ll 1$ from every entry in B except the single active entry per row in B with respect to y .

In general, $Ax = By$ if and only if there exists an m -tuple (k_1, \dots, k_m) , $k_i \in K$, and a matrix $B^- = (b_{ij}^-) \in \mathbb{R}^{m \times k}$ with

$$b_{ij}^- = \begin{cases} b_{ij} & \text{if } j = k_i \\ b_{ij} - \delta & \text{otherwise,} \end{cases}$$

such that $Ax = B^-y$.

Hence, given a pair (A, B) satisfying Property One⁺FP, we can generate a number of pairs $(A, B^{(g)})$, $g \in \mathbb{N}$ such that x, y is an integer solution to $Ax = By$ if and only if there exists g such that $Ax = B^{(g)}y$. Note that each $B^{(g)}$ is obtained from B by subtracting δ from all but one element, $b_{ir'(i,s)}$, per row, and that the pairs $(A, B^{(g)})$ satisfy Property OneFP. We can therefore determine whether an integer solution to $Ax = B^{(g)}y$ exists in strongly polynomial time.

Unfortunately, in the worst case, there could be as many as nk pairs per row, and thus $(nk)^m$ matrices to check, so the complexity of this method is $\mathcal{O}(m^3 n^m k^m)$. However we can say that, for fixed m , a polynomial method for finding integer solutions to a TSS exists.

5. Integer max-linear programs

In this section, we develop algorithms to solve the IMLP for problems with finite entries. We will write IMLP^{\max} to mean the integer max-linear program which maximises $f^T x$, and IMLP^{\min} to denote the program minimising $f^T x$, where $f \in \mathbb{R}^n$. For $A, B \in \mathbb{R}^{m \times n}$, $c, d \in \mathbb{R}^m$, we denote

$$\begin{aligned} S &= \{x \in \mathbb{R}^n : Ax \oplus c = Bx \oplus d\}, \\ IS &= S \cap \mathbb{Z}^n, \\ IS^{\min} &= \{x \in IS : f(x) \leq f(z) \forall z \in IS\}, \\ IS^{\max} &= \{x \in IS : f(x) \geq f(z) \forall z \in IS\}. \end{aligned}$$

The method described here is based on the bisection method used to find real optimal solutions to max-linear programs described in [3].

The justification for using a bisection method for problems with integrality requirements is given by the following.

Proposition 5.1. *Let $x, y \in IS$, with $f(x) = \alpha < \beta = f(y)$. Then, for all $\gamma \in (\alpha, \beta)$ with $\text{fr}(\gamma) = \text{fr}(\beta)$, there exists $z \in IS$ such that $f(z) = \gamma$.*

Proof. Take $\lambda = 0$ and $\mu = \gamma \otimes \beta^{-1} \in \mathbb{Z}$. Let $z = \lambda \otimes x \oplus \mu \otimes y$. Now S is a max-convex set [3], and so, since $\lambda \oplus \mu = 0$, we have $z \in S \cap \mathbb{Z}^n = IS$. Finally,

$$f(z) = \lambda \otimes f(x) \oplus \mu \otimes f(y) = \alpha \oplus \gamma = \gamma. \quad \square$$

We also need the following results. The first follows from the cancellation rule, Lemma 2.5.

Lemma 5.2 ([3]). *Let $\alpha, \alpha' \in \mathbb{R}$, $\alpha' < \alpha$ and $f(x) = f^T \otimes x, f'(x) = f'^T \otimes x$, where $f'_j < f_j$ for every $j \in N$. Then the following holds for every $x \in \mathbb{R} : f(x) = \alpha$ if and only if $f(x) \oplus \alpha' = f'(x) \oplus \alpha$.*

Since the result holds for real vectors x , it clearly also holds for integer x . Using this and the cancellation law, we can check whether $f(x)$ attains some value α by checking whether a two-sided system has an integer solution.

Proposition 5.3. *$f(x) = \alpha$ for some $x \in IS$ if and only if the following integer max-linear system has a solution:*

$$\begin{aligned} A \otimes x \oplus c &= B \otimes x \oplus d, \\ f(x) \oplus \alpha' &= f'(x) \oplus \alpha, \\ x &\in \mathbb{Z}^n, \end{aligned}$$

where $\alpha' < \alpha$ and $f'_j < f_j$ for every $j \in N$.

We know from Section 3 that we can decide whether a two-sided system has an integer solution by applying Algorithm GEN-INT-TSS.

5.1. When the objective function is unbounded

We now consider the question of when optimal solutions exist. We denote the minimum and maximum of f by f^{\min} and f^{\max} , respectively. Without loss of generality, we assume that $c \geq d$, and we denote $M^> = \{i \in M : c_i > d_i\}$. Let

$$L_r = \min_{j \in N} (f_j \otimes c_r \otimes b_{rj}^{-1})$$

and

$$L = \left[\max_{r \in M^>} L_r \right]. \tag{5.1}$$

Note that $L = -\infty$ if $M^> = \emptyset$.

In the rest of this section, we follow similar arguments as appear in [3] regarding the max-linear programming problem. For any result stated without a proof, the result either follows immediately from a theorem in [3], or can be obtained by a simple adaptation of the proof found there.

Lemma 5.4. *If $c \geq d$, then $f(x) \geq L$ for every $x \in IS$.*

Theorem 5.5. *$f^{\min} = -\infty$ if and only if $c = d$.*

For the upper bound, we need the following results.

Lemma 5.6. *Let $c \geq d$. If $x \in IS$ and $(Ax)_i > c_i$ for all $i \in M$, then $x' = \alpha x \in IS$ and $(Ax')_i \leq c_i \otimes 1$ for some $i \in M$, where*

$$\alpha = \left[\max_{i \in M} \left(c_i (Ax)_i^{-1} \right) \right]. \tag{5.2}$$

Proof. Assume that $x \in IS$. Then $Ax = Bx$, since $Ax > c \geq d$. From the choice of α , we get that

$$(A(\alpha x))_i = \alpha (Ax)_i \geq c_i$$

for all $i \in M$. Further, since $A(\beta x) = B(\beta x)$ for any $\beta \in \mathbb{Z}$, we have $x' \in IS$.

Finally, let $j \in M$ be an index at which the value of α is attained. Then

$$(Ax')_j = \lceil c_j (Ax)_j^{-1} \rceil (Ax)_j \leq c_j \otimes (Ax)_j^{-1} \otimes 1 \otimes (Ax)_j \leq c_j \otimes 1. \quad \square$$

Let

$$U = \left[\max_{r \in M} \max_{j \in N} (f_j \otimes a_{rj}^{-1} \otimes c_r \otimes 1) \right]. \tag{5.3}$$

Lemma 5.7. *If $c \geq d$, then the following hold.*

- (a) *If $x \in IS$ and $(Ax)_r \leq c_r \otimes 1$ for some $r \in M$, then $f(x) \leq U$.*
- (b) *If $Ax = Bx$ has no integer solution, then $f(x) \leq U$ for every $x \in IS$.*

Theorem 5.8. *$f^{\max} = +\infty$ if and only if $Ax = Bx$ has an integer solution.*

So far, we have shown that we can determine immediately when f^{\min} is unbounded and can check whether f^{\max} is unbounded, for example by applying Algorithm GEN-INT-TSS. We now need to argue that, when the objective function value is bounded, there exist integer vectors for which f^{\max} and f^{\min} are attained.

5.2. Attainment of optimal values

For all $j \in N$, let

$$\begin{aligned} h_j &= \left[\min \left(\min_{r \in M} (a_{rj}^{-1} \otimes c_j), \min_{r \in M} (b_{rj}^{-1} \otimes d_j), f_j^{-1} \otimes L \right) \right], \\ h'_j &= \left[\min \left(\min_{r \in M} (a_{rj}^{-1} \otimes c_j), \min_{r \in M} (b_{rj}^{-1} \otimes d_j) \right) \right], \end{aligned} \tag{5.4}$$

$h = (h_1, \dots, h_n)^T$, and $h' = (h'_1, \dots, h'_n)^T$. Observe that h' is finite and that h is finite if and only if $f^{\min} > -\infty$.

Proposition 5.9. Let h and h' be as defined above.

- (i) For any $x \in IS$, the vector $x' = x \oplus h \in IS$ satisfies $x' \geq h$ and $f(x) = f(x')$.
- (ii) For any $x \in IS$, the vector $x' = x \oplus h' \in IS$ satisfies $x' \geq h'$ and $f(x) \leq f(x')$.

Corollary 5.10. Let h and h' be as defined above.

- (i) If $f^{\min} > -\infty$ and $IS \neq \emptyset$, then

$$f^{\min} = \min_{x \in IS} f(x),$$

where $\overline{IS} = IS \cap \{x \in \mathbb{Z}^n : h_j \leq x_j \leq f_j^{-1} \otimes f(\bar{x}), j \in N\}$.

- (ii) If $f^{\max} < +\infty$, then

$$f^{\max} = \max_{x \in \overline{IS}'} f(x),$$

where $\overline{IS}' = IS \cap \{x \in \mathbb{Z}^n : h'_j \leq x_j \leq f_j^{-1} \otimes U, j \in N\}$.

Proof. Similar to the proof of Corollaries 10.2.12 and 10.2.17 in [3]. \square

We summarise these results as follows.

Corollary 5.11. If $IS \neq \emptyset$ and $f^{\min} > -\infty$ [$f^{\max} < +\infty$], then $IS^{\min} \neq \emptyset$ [$IS^{\max} \neq \emptyset$].

Finally, we need a finite lower bound on f^{\max} , since L will not work in the case when $c = d$.

Corollary 5.12. Let $L' = \lfloor f(h') \rfloor$. If $x \in IS$, then $x' = x \oplus h'$ is such that $f(x') \geq L'$, and hence $f^{\max} \geq L'$.

5.3. The algorithms

5.3.1. Algorithm when minimising the objective function

For minimisation, we know that an optimal solution exists provided that $c \neq d$. We first check whether $f^{\min} = L$ using **Proposition 5.3**: if so, we are done: if not, then we find any feasible x^0 using Algorithm GEN-INT-TSS and, if necessary, scale it using (5.2) so that $f(x^0) \leq U$. Then we know that $f^{\min} \in (L, f(x^0)]$.

Once we know that, for any $x \in IS$, we have that $f(x)$ satisfies

$$L \leq f(x) \leq f(x^0) = U,$$

we can set

$$\Theta = \{\theta : \theta \in (L, U] \text{ and } \text{fr}(\theta) = \text{fr}(U)\},$$

and apply the bisection method on the set Θ as follows.

1. Order $\theta_i \in \Theta$ from smallest to largest, and test whether the middle value, θ , is attained for any $x \in IS$.
2. If it is, then we have a new upper bound: $f(x) \leq \theta$.
3. If it is not, then we have a new lower bound: $f(x) > \theta$.

In 3, we use the fact that, if θ is unattainable, then no value in $(L, \theta]$ is attainable. To see this, note that, if there exists $\alpha \in (L, \theta]$ attainable, then by **Proposition 5.1** all values in (α, U) with fractional part equal to that of U work, but θ satisfied this condition, and was not attainable. In each case, we have halved the number of values that we need to test.

Using this idea, we obtain the following algorithm for solving IMLP^{min}.

Algorithm: INT-MAXLINMIN

Input: $A, B \in \mathbb{R}^{m \times n}$, $c, d \in \mathbb{R}^m$, $c \geq d$, $c \neq d$, $f \in \mathbb{R}^n$.

Output: $x \in IS$ such that $f(x) = f^{\min}$.

1. Calculate L from (5.1). If $L = f(x)$ for some $x \in IS$, then STOP ($f^{\min} = L$).
2. Find an $x^0 \in IS$. If $(A \otimes x^0)_i > c_i$ for all $i \in M$, then scale x^0 by α defined in (5.2).
3. $L(0) := L$, $U(0) := f(x^0)$, $r := 0$.
4. $\Theta := \{\theta : \theta \in (L(r), U(r)] \text{ and } \text{fr}(\theta) = \text{fr}(U(r))\}$, $\eta := |\Theta|$. If $\eta = 1$, then STOP ($f^{\min} = U(r)$).
5. Take $\theta \in \Theta \cap \left[\frac{L(r)+U(r)}{2} - \frac{1}{2}, \frac{L(r)+U(r)}{2} + \frac{1}{2} \right]$.
6. Check whether $f(x) = \theta$ for some $x \in IS$, and if so find one.
If yes, then $U(r+1) = \theta$, $L(r+1) = L(r)$.
If no, then $U(r+1) := U(r)$, $L(r+1) = \theta$.
7. $r := r + 1$, go to 4.

Theorem 5.13. Algorithm INT-MAXLINMIN is correct and terminates after at most $\mathcal{O}(\log(\lceil U - L \rceil))$ iterations.

Let

$$\bar{K} = \left\lceil \max\{|a_{ij}|, |b_{ij}|, |c_i|, |d_i|, |f_j| : i \in M, j \in N\} \right\rceil.$$

Observe that $L, L', U \in [-3\bar{K}, 3\bar{K}]$.

Corollary 5.14. If Algorithm GEN-INT-TSS is used to perform the checks in steps 1 and 6, then Algorithm INT-MAXLINMIN has complexity $\mathcal{O}(mn(m+n)\bar{K} \log \bar{K})$.

Proof. The number of iterations is $\mathcal{O}(\log(\lceil U - L \rceil)) \leq \mathcal{O}(\log 6\bar{K}) = \mathcal{O}(\log \bar{K})$. Each iteration uses Algorithm GEN-INT-TSS which, from Theorem 3.3, requires $\mathcal{O}(K(X|Y)(m'n'(m'+n')))$ operations, where $K(X|Y)$ is defined in (3.1), $m' = m + 1$, $n' = n + 1$, and

$$X = \begin{pmatrix} A & c \\ f^T & \alpha' \end{pmatrix}, \quad Y = \begin{pmatrix} B & d \\ f^T & \alpha \end{pmatrix}.$$

We can choose α' and f' so that $\alpha - 1 \leq \alpha' \leq \alpha, f_j - 1 \leq f'_j \leq f_j$, and hence $K(X|Y) \leq \bar{K} + 1$. Therefore the number of operations in a single iteration is $\mathcal{O}(\bar{K}mn(m+n))$. \square

5.3.2. Algorithm when maximising the objective function

For maximisation, we cannot assume that $c \neq d$, since this is not the criterion for f^{\max} to be unbounded. We must first check that $f^{\max} < +\infty$ by verifying that $Ax = Bx$ has no integer solution, which can be done using Algorithm GEN-INT-TSS. We then check whether $f^{\max} = U$ (where U is defined in (5.3)) using Proposition 5.3. If not, then we find any feasible solution x^0 , and set $x^0 := x^0 \oplus h'$ so that $f(x^0) \geq L$.

Further, when maximising, it is no longer enough to only check values in the interval with a single fractional part, because the upper bound is not attained, and so we can no longer guarantee that the optimal value shares its fractional part with U . However, we do know that there are only a finite number of possible fractional parts that could be attained; these are $\text{fr}(f_i)$ for all i , because $f^T \otimes x$ for $x \in \mathbb{Z}^n$ can only take its fractional part from the elements of f . Once we know, for all $x \in IS$, that $L = f(x^0) \leq f(x) \leq U$, we proceed as follows.

1. Let $[J, J + 1)$ be an interval contained halfway between L and U .
2. Test each of the (at most n) values in this interval that share the same fractional part as a component of f to see whether they are attained by some $x \in IS$.
3. If one exists, then the largest becomes a new lower bound.
4. If none in the interval are attained, then Proposition 5.1 guarantees that no value higher than J can be attained, and thus we have a new upper bound.

Continue in this way, each time approximately halving the length of the interval until $U - L \leq 2$. In this case, the interval $[J, J + 1)$ may not be contained entirely in (L, U) , and so testing points in this smaller interval is no longer efficient, since we will check unnecessary points, or find L again. So, instead, check the remaining $\leq 2n$ possible points, and choose the one with smallest value.

We obtain the following algorithm for IMLP^{max}.

Algorithm: INT-MAXLINMAX

Input: $A, B \in \mathbb{R}^{m \times n}, c, d \in \mathbb{R}^m, c \geq d, f \in \mathbb{R}^n$.

Output: $x \in IS$ such that $f(x) = f^{\max}$.

1. Calculate U from (5.3). If $U = f(x)$ for some $x \in IS$, then STOP ($f^{\max} = U$).
2. Check whether $Ax = Bx$ has an integer solution. If yes, STOP ($f^{\max} = +\infty$).
3. Find an $x^0 \in IS$. Set $x^0 := x^0 \oplus h'$ as defined in (5.4).
4. $L(0) := f(x^0), U(0) := U, r := 0$.
5. If $U - L \leq 2$, go to 8. Else let $J := \frac{1}{2}(U(r) + L(r))$.
6. Using a bivalent search, find the biggest $\sigma \in [J, J + 1)$ such that $(\exists j)\text{fr}(f_j) = \text{fr}(\sigma)$ and $f(x) = \sigma$ for some $x \in IS$.
If none exist, then $U(r + 1) := J, L(r + 1) = L(r)$.
Otherwise, $U(r + 1) = U(r), L(r + 1) = \sigma$.
7. $r := r + 1$, go to 5.
8. Using a bivalent search, find the biggest $\gamma \in (L(r), U(r))$ such that $(\exists j)\text{fr}(f_j) = \text{fr}(\gamma)$ and $f(x) = \gamma$ for some $x \in IS$.
If none exist, then STOP ($f^{\max} = L(r)$).
Otherwise STOP, $f^{\max} = \gamma$.

Theorem 5.15. Algorithm INT-MAXLINMAX is correct and terminates after at most $\mathcal{O}(\log(U - L'))$ iterations, where $L' = \lfloor f(h') \rfloor$.

Corollary 5.16. If Algorithm GEN-INT-TSS is used to perform the checks in steps 1, 6, and 8, then Algorithm INT-MAXLINMAX has complexity $\mathcal{O}(mn(m+n) \log(n)\bar{K} \log \bar{K})$.

Proof. The same as the proof of Corollary 5.14, with L replaced by L' , but here we have that each iteration uses Algorithm GEN-INT-TSS at most $\log(2n)$ times. \square

6. IMLP and systems with Property OneFP

In Section 4, we showed that, for systems satisfying Property OneFP, we could find integer solutions to a TSS in strongly polynomial time; here, we show that, if Property OneFP is satisfied, then our solution methods for the IMLP can also be made more efficient.

6.1. The adapted Alternating Method is pseudopolynomial for systems with Property OneFP

Let $A \in \overline{\mathbb{R}}^{m \times n}$, $B \in \overline{\mathbb{R}}^{m \times k}$ be doubly \mathbb{R} -astic such that the pair (A, B) satisfies Property OneFP. Let $\tilde{A} = (\tilde{a}_{ij})$ and $\tilde{B} = (\tilde{b}_{it})$, where

$$\tilde{a}_{ij} = \begin{cases} a_{ij} & \text{if } a_{ij} \in \mathbb{Z} \\ \lfloor a_{ij} \rfloor + s_1 & \text{otherwise} \end{cases} \quad \text{and} \quad \tilde{b}_{it} = \begin{cases} b_{it} & \text{if } b_{it} \in \mathbb{Z} \\ \lfloor b_{it} \rfloor + s_2 & \text{otherwise} \end{cases}$$

for some real numbers $0 < s_1, s_2 < 1$, $s_1 \neq s_2$. Note that the pair (\tilde{A}, \tilde{B}) also satisfies Property OneFP.

Proposition 6.1. *Let $A, B, \tilde{A}, \tilde{B}$ be as described above. The pair (x, y) [vector x] is an integer solution to $Ax = By$ [$Ax = Bx$] if and only if (x, y) [x] is an integer solution to $\tilde{A}x = \tilde{B}y$ [$\tilde{A}x = \tilde{B}x$].*

Proof. Assume that $Ax = By$. We know by Corollary 4.2 that the only active entries in A and B are the integer entries, $a_{ir(i)}$ and $b_{ir'(i)}$. We have

$$\begin{aligned} & (\forall i \in M)(\forall j \in N, j \neq r(i)) a_{ir(i)} + x_{r(i)} > a_{ij} + x_j \\ \therefore & (\forall i \in M)(\forall j \in N, j \neq r(i)) a_{ir(i)} + x_{r(i)} \geq \lceil a_{ij} \rceil + x_j \\ \therefore & (\forall i \in M)(\forall j \in N, j \neq r(i)) a_{ir(i)} + x_{r(i)} > \lfloor a_{ij} \rfloor + s_1 + x_j \\ \therefore & (\forall i \in M)(\forall j \in N, j \neq r(i)) \tilde{a}_{ir(i)} + x_{r(i)} > \tilde{a}_{ij} + x_j. \end{aligned}$$

Hence $Ax = \tilde{A}x$, and it can also be shown that $By = \tilde{B}y$. A similar argument holds for the other direction. The case $Ax = Bx$ is proved analogously. \square

Remark 3. Another way to prove this would be to calculate W from (A, B) as described in Theorem 4.6 and \tilde{W} from (\tilde{A}, \tilde{B}) similarly. Then note that $W = \tilde{W}$, and therefore for any integer solutions x, \tilde{x} we have $Ax = Bx = z = \tilde{A}\tilde{x} = \tilde{B}\tilde{x}$ for any $z \in IV^*(W, 0)$. Finally, observe that this implies that $x = \tilde{x}$ using Corollary 4.2.

Proposition 6.1 allows us to transform any system satisfying Property OneFP into a system with the same solution set, also satisfying Property OneFP, with the additional property that all non-integer entries of each matrix have the same fractional part (but different from the fractional part in the other matrix). Without loss of generality, we can set, for instance, $s_1 = 0.1$ and $s_2 = 0.2$ so that no entry in the system has more than one decimal place. Hence the maximum number of symbols in an entry is

$$\log \kappa + 1,$$

where $\kappa = K(\tilde{A}|\tilde{B})$ is as defined in (3.1).

For systems with Property OneFP where A, B are finite, we can input this transformed system into Algorithms SEP-INT-TSS and GEN-INT-TSS, which we know to have complexity

$$\mathcal{O}(mn(n+k)K(\tilde{A})) \quad \text{and} \quad \mathcal{O}(\kappa(mn(m+n))),$$

respectively, by Theorems 3.1 and 3.3. Therefore both Algorithm SEP-INT-TSS and Algorithm GEN-INT-TSS are pseudopolynomial when applied to instances with finite matrices satisfying Property OneFP.

Finally, note that the computationally expensive step in Algorithms INT-MAXLINMIN and INT-MAXLINMAX is the use of Algorithm GEN-INT-TSS in each iteration. Therefore, for systems satisfying Property OneFP, these algorithms are pseudopolynomial.

6.2. Polynomial algorithm for IMLP for systems with Property OneFP

In Section 6.1, we argued that for pairs (A, B) satisfying Property OneFP the algorithms for solving the IMLP are pseudopolynomial. Here we show that, if Property OneFP is satisfied, we can perform the checks in each iteration by the condition from Theorem 4.6 instead of Algorithm GEN-INT-TSS, and obtain a polynomial algorithm for solving IMLP.

In each iteration, Algorithm GEN-INT-TSS is used to determine whether there exists some $x \in IS$ satisfying $f(x) = \alpha$ by finding if an integer solution to the following two-sided system exists:

$$\begin{pmatrix} A & c \\ f^T & \alpha' \end{pmatrix} \otimes x = \begin{pmatrix} B & d \\ f'^T & \alpha \end{pmatrix} \otimes x,$$

where $f' < f, \alpha' < \alpha$.

Now note that we always test an α that shares its fractional part with some element of f , and we can always choose f' , α' so as not to add any fractional parts which already appear in the $(m + 1)$ st row. As before, we can transform this system into one in which every entry has at most one decimal place, and therefore the maximum number of symbols in an entry is

$$v = \log \kappa' + 1,$$

where

$$\kappa' = K(X|Y), \quad X = \begin{pmatrix} A & c \\ f^T & \alpha' \end{pmatrix}, \quad Y = \begin{pmatrix} B & d \\ f'^T & \alpha \end{pmatrix}.$$

So we obtain the following proposition.

Proposition 6.2. Let $A' = (A|c)$, $B' = (B|d)$. Under the assumptions that

- (i) the pair (A', B') satisfies Property OneFP, and
- (ii) $(\forall j, t \in N) \text{fr}(f_t) \neq \text{fr}(f_j)$

Algorithm INT-MAXLINMIN has complexity $\mathcal{O}((m + n)^3 \log \kappa')$ and Algorithm INT-MAXLINMAX has complexity $\mathcal{O}((m + n)^3 \log n \log \kappa')$. Hence the algorithms are polynomial in v .

Proof. The same as the proofs of Corollaries 5.14 and 5.16, except for the complexity of checking whether $f(x) = \alpha$. This requires us to determine whether

$$\left(\begin{array}{c|c} A & c \\ \hline f^T & \alpha' \\ \hline I & \end{array} \right) x = \left(\begin{array}{c|c} B & d \\ \hline f'^T & \alpha \\ \hline I & \end{array} \right) y$$

has an integer solution. This can be done in $\mathcal{O}((m + n + 2)^3) = \mathcal{O}((m + n)^3)$ time by Corollary 4.9. \square

Acknowledgement

The first author was supported by EPSRC grant EP/J00829X/1.

References

- [1] X. Allamigeon, S. Gaubert, É. Goubault, The tropical double description method, in: J.-Y. Marion, Th. Schwentick (Eds.), Proceedings of the 27th International Symposium on Theoretical Aspects of Computer Science, STACS 2010, in: Leibniz International Proceedings in Informatics (LIPIcs), vol. 5, Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, Dagstuhl, Germany, 2010, pp. 47–58.
- [2] P. Butkovič, Max-algebra: the linear algebra of combinatorics? Linear Algebra and its Applications 367 (2003) 313–335.
- [3] P. Butkovič, Max-Linear Systems: Theory and Algorithms, Springer-Verlag, London, 2010.
- [4] P. Butkovič, A. Aminu, Max-linear programming, IMA Journal of Management Mathematics 20 (3) (2009) 233–249.
- [5] P. Butkovič, G. Hegedüs, An elimination method for finding all solutions of the system of linear equations over an extremal algebra, Ekonomicko-Matematický Obzor 20 (1984) 203–215.
- [6] P. Butkovič, M. MacCaig, The alternating method for finding integer solutions to two-sided systems, University of Birmingham, School of Mathematics, 2012/08.
- [7] P. Butkovič, M. MacCaig, On integer eigenvectors and subeigenvectors in the max-plus algebra, Linear Algebra and its Applications 438 (2013) 3408–3424.
- [8] R.A. Cuninghame-Green, Minimax Algebra, in: Lecture Notes in Economics and Math. Systems, vol. 166, Springer, Berlin, 1979.
- [9] R.A. Cuninghame-Green, P. Butkovič, The equation $Ax = By$ over $(\max, +)$, Theoretical Computer Science 293 (2003) 3–12.
- [10] S. Gaubert, R.D. Katz, S. Sergeev, Tropical linear-fractional programming and parametric mean-payoff games, Journal of Symbolic Computation 47 (2012) 1447–1478.
- [11] R.M. Karp, A characterization of the minimum cycle mean in a digraph, Discrete Mathematics 23 (1978) 309–311.
- [12] E.A. Walkup, G. Boriello, a general linear max-plus solution technique, in: J. Gunawardena (Ed.), Idempotency, Cambridge, 1988, pp. 406–415.