

# UNIVERSITY OF BIRMINGHAM

## Research at Birmingham

### New Binary and Exoplanet Candidates from STEREO Light Curves

Whittaker, Gemma; Sangaralingam, Vino; Stevens, Ian

DOI:

[10.1017/S1743921311027232](https://doi.org/10.1017/S1743921311027232)

License:

None: All rights reserved

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Whittaker, G, Sangaralingam, V & Stevens, I 2011, 'New Binary and Exoplanet Candidates from STEREO Light Curves', *Proceedings of the International Astronomical Union*, vol. 7, no. S282, pp. 143-144.  
<https://doi.org/10.1017/S1743921311027232>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

© Cambridge University Press

Eligibility for repository checked July 2014

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# New Binary and Exoplanet Candidates from STEREO Light Curves

Gemma Whittaker, Vino Sangaralingam and Ian Stevens

Astrophysics and Space Research, The University of Birmingham, Edgbaston,  
Birmingham, B15 2TT, UK  
email: [gemma@star.sr.bham.ac.uk](mailto:gemma@star.sr.bham.ac.uk)

**Abstract.** The Heliospheric Imagers (HI) onboard the STEREO satellites are observing an abundance of background stars as they follow their respective Sun-centered orbits. These are wide-angled CCD cameras with a  $20^\circ \times 20^\circ$  field of view, directed  $\sim 14^\circ$  from the solar disk. These imagers monitor 20% of the sky over one year, providing light curves for over 500,000 stars down to 12th magnitude and brighter than 7th. We are currently analysing the photometric data from the HI-1 cameras, obtained since March 2007. Following a standard data reduction of the raw photometric images, the resultant light curves underwent a sequence of detrending procedures to minimize systematics in the data, which can contribute to red noise. A transit search was performed using the BLS algorithm, which is sensitive to the box-like shape associated with planetary transits. The resulting candidates were subjected to a number of false-alarm tests to determine the most promising candidates and these were investigated further, visually and using available catalogue data. Possible new exoplanet and binary candidates will now be submitted for follow-up photometric and spectroscopic observations to confirm their nature.

**Keywords.** techniques: image processing, techniques: photometric, binaries: eclipsing, planetary systems, catalogs, methods: data analysis

---

## 1. STEREO Observations

STEREO (Solar TERrestrial RELations Observatory) was launched in March 2006 to study Coronal Mass Ejections (CMEs) in unprecedented detail by providing a stereoscopic view of the Sun. The twin observatories follow a heliocentric orbit, with one moving ahead of the Earth (STEREO-A) and the second lagging behind the Earth (STEREO-B). Among the four, identical suites of instruments onboard the two spacecraft is the SECCHI package (Sun Earth Connection Coronal and Heliospheric Investigation). SECCHI is comprised of an extreme UV imager, two optical coronagraphs and two heliospheric imagers (HI-1 and HI-2), designed (primarily) to study the 3-D evolution of CMEs along the Sun-Earth line.

The HI-1 instruments are optical CCD cameras with a  $20^\circ \times 20^\circ$  field of view. Within this field an estimated 29,000 stars between 7th and 12th magnitude may be visible in the background at any one time. These cameras have a 40 minute observing cadence, which provides 36 consecutive frames per day for 20 days after which a star will pass out of the field of view. STEREO has now been fully operational for over 4 years, hence most stars have been observed at least 4 times by each camera. This has provided a wealth of precision, photometric data for many stars which are not well studied at present.


## 2. Data Processing and Detrending

A standard data reduction is carried out for the raw photometric data, which are downloaded directly from the STEREO webpage†. This includes corrections for shutterless readout and saturated columns. Background subtraction is then performed to minimise contamination from the F-corona, using a 1-day running, minimum background. The NOMAD catalogue is used to track individual stars over consecutive frames and finally the light curves for these stars are extracted using aperture photometry. The HI-1 cameras have a moderate PSF of 3.7 arcmin (Cf. CoRoTs PSF which is 10.5 arcmin), meaning that the extracted light curves are susceptible to contamination from nearby stars.

## 3. Transit Analysis and Results

A transit search was performed, within a period range of 0.5 - 10 days, using the BLS transit detection algorithm (Kovács *et al.*, 2002). From an initial study including 82,863 light curves from HI-1A, 952 potential candidates were detected with a Signal Detection Efficiency (SDE) of greater than 9. A series of false-alarm tests were then used to determine the most promising candidates. These included spectral-type information from SIMBAD and the Exoplanet Diagnostic Tool,  $\eta^*$ , (Tingley & Sackett, 2005), which calculates the likelihood of a transit being from a planet as opposed to it being from a star or as a result of noise. The candidates passing all false-alarm tests were visually analysed to enable them to be classified into potential exoplanet or binary candidates.

Some binary targets have recently been submitted for preliminary spectroscopic follow-up using the WIRO telescope in Wyoming, USA. Present and future work is now focussed establishing a reliable set of targets for high resolution follow-up observations. This work will involve refining our search parameters and detrending procedures and the addition of the HI-1B data, which will increase the detection efficiency of BLS.



**Figure 1.** Example exoplanet and binary candidates. Figure (a) shows an F5 dwarf, which was detected by BLS with  $SDE = 15$ ,  $Period = 0.73$  days and  $\eta^* = 0.12$ , which lies well below the cutoff value of 1.0 (indicative of a planet). Figure (b) shows an A0 star, which was detected with  $SDE = 40$ ,  $Period = 2.1$  days and  $\eta^* = 0.96$ .

## References

- Kovács, G., Zucker, S. & Mazek, T. 2002 *A&A*, 391, 369  
 Tingley, B. & Sackett, P. 2005 *ApJ*, 627, 1011  
 Sangaralingam, V. & Stevens, I. 2011 *submitted*  
 Whittaker, G., Sangaralingam, V. & Stevens, I. 2011 *in preparation*

† [www.ukssdc.rl.ac.uk/solar/stereo/data.html](http://www.ukssdc.rl.ac.uk/solar/stereo/data.html)