

# UNIVERSITY OF BIRMINGHAM

## Research at Birmingham

### The thymus medulla fosters generation of natural Treg cells, invariant T Cells and invariant NKT cells : what we learn from intrathymic migration

Cowan, Jennifer; Jenkinson, William; Anderson, Graham

DOI:

[10.1002/eji.201445108](https://doi.org/10.1002/eji.201445108)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Cowan, JE, Jenkinson, WE & Anderson, G 2015, 'The thymus medulla fosters generation of natural Treg cells, invariant T Cells and invariant NKT cells : what we learn from intrathymic migration', *European Journal of Immunology*, vol. 45, no. 3, pp. 652-660. <https://doi.org/10.1002/eji.201445108>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

Checked for repository 18/03/2015

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Thymus medulla fosters generation of natural Treg cells, invariant $\gamma\delta$ T cells, and invariant NKT cells: What we learn from intrathymic migration

Jennifer E. Cowan, William E. Jenkinson and Graham Anderson

MRC Centre for Immune Regulation, Institute for Biomedical Research, Medical School, University of Birmingham, Birmingham, UK

The organization of the thymus into distinct cortical and medullary regions enables it to control the step-wise migration and development of immature T-cell precursors. Such a process provides access to specialized cortical and medullary thymic epithelial cells at defined stages of maturation, ensuring the generation of self-tolerant and MHC-restricted conventional CD4<sup>+</sup> and CD8<sup>+</sup>  $\alpha\beta$  T cells. The migratory cues and stromal cell requirements that regulate the development of conventional  $\alpha\beta$  T cells have been well studied. However, the thymus also fosters the generation of several immunoregulatory T-cell populations that form key components of both innate and adaptive immune responses. These include Foxp3<sup>+</sup> natural regulatory T cells, invariant  $\gamma\delta$  T cells, and CD1d-restricted invariant natural killer T cells (iNKT cells). While less is known about the intrathymic requirements of these nonconventional T cells, recent studies have highlighted the importance of the thymus medulla in their development. Here, we review recent findings on the mechanisms controlling the intrathymic migration of distinct T-cell subsets, and relate this to knowledge of the microenvironmental requirements of these cells.

**Keywords:** Chemokine · iNKT cells ·  $\gamma\delta$  T cells · Natural Treg cell · Thymic medulla · Thymus

## Introduction

The immune system consists of a wide range of specialized cells within tissues that play key roles in the control of pathogen recognition, cellular stress, and tumor surveillance. Typically, descriptions of the immune system involve its separation into innate and adaptive components. For example, initial pathogen-mediated activation of innate cell types such as DCs can drive effector responses by lymphocyte components of the adaptive immune system. Appropriate stimulation of effector function in  $\alpha\beta$ TCR bearing T cells can take several days, and eventually leads to the formation of antigen-specific memory T-cell populations. However, as shown recently for  $\gamma\delta$  T cells [1], this separation of innate and adaptive immune cells is not as clearly defined as previously thought, and individual T-cell subsets can be involved in both innate and

adaptive arms of the immune response. From a developmental point of view, this is important as the generation of several of the cellular mediators of innate and adaptive immunity takes place at a common site, the thymus. While the role of the thymus in the generation of adaptive CD4<sup>+</sup> and CD8<sup>+</sup>  $\alpha\beta$  T cells is well known (reviewed in [2, 3]), the thymus also provides support for additional innate and adaptive T-cell subsets including Foxp3<sup>+</sup> natural regulatory T (Foxp3<sup>+</sup> nTreg) cells, invariant natural killer T cells (iNKT cells), and various  $\gamma\delta$  T-cell subsets in fetal and postnatal life [4–6].


Thymic microenvironments demonstrate a distinct anatomical organization that is directly linked to their function [7–9]. The medulla represents a site where single positive (SP) thymocytes accumulate prior to their exit into the periphery. Here, various subsets of medullary thymic epithelial cells (mTECs) and DCs are involved in multiple aspects of T-cell development, including negative selection and thymic emigration [10–12]. The organization and stromal cell heterogeneity of the thymus medulla is becoming increasingly well-defined [11], including further

**Correspondence:** Prof. Graham Anderson  
e-mail: g.anderson@bham.ac.uk

subdivision of mTECs using markers that include the CCR7 ligand CCL21 [13] and the atypical chemokine receptor CCRL1 [14]. These descriptions of medullary heterogeneity are helping to provide a clearer understanding of the mechanisms controlling conventional  $\alpha\beta$  T-cell development. Increasing evidence suggests that the thymic medulla is also important in the development of other T-cell populations that play important roles in both the innate and adaptive immune systems. In this review, we compare the thymic requirements of these distinct T-cell populations, drawing attention to our current understanding of the multiple roles played by the thymus medulla in the intrathymic migration and development of these cells.

## Conventional CD4<sup>+</sup> and CD8<sup>+</sup> $\alpha\beta$ T cells

Following positive selection in the cortex, the medullary dwell time of SP thymocytes has been suggested to be as short as 4–5 days (reviewed in [15]), to as long as 2 weeks [16]. During this time, medullary resident thymocytes undergo a series of phenotypically and functionally distinct maturation stages (Fig. 1), including downregulation of CD24, and upregulation of Qa2 and CD62L [17]. Thus, while newly generated CD4<sup>+</sup> and CD8<sup>+</sup> SP thymocytes in mice are characterized by a CD69<sup>+</sup>HSA<sup>high</sup>Qa2<sup>-</sup>CD62L<sup>-</sup> surface phenotype [15, 18, 19], with increasing maturity thymocytes downregulate HSA and CD69 whilst upregulating Qa2 and CD62L to become CD69<sup>-</sup>HSA<sup>low</sup>Qa2<sup>+</sup>CD62L<sup>+</sup> (Fig. 1) [18, 20, 21]. The differentiation program of CD4 SP thymocytes has been further subdivided with the aid of additional markers such as 6C10 [22] into four distinct maturation stages, each with unique molecular signatures [23]. The possible functional relevance of the majority of these surface markers, however, remains relatively unknown. More recently, the different stages in CD4 SP thymocyte development has been analyzed in relation to the expression of chemokine receptors linked to intrathymic migration (Fig. 1). For instance, newly selected CD69<sup>+</sup> CD4 SP murine thymocytes have been shown to express high levels of CCR9 and CCR4, while mature CD69<sup>-</sup> CD4 SP cells are CCR4<sup>-</sup>CCR9<sup>-</sup> [24]. Relevant to the process of medullary entry, visualization of thymocyte mobility within whole thymic lobes via two-photon laser scanning microscopy showed that the positive selection of cells moving randomly within cortical regions triggers their rapid and directed migration toward the medulla [25]. Moreover, treatment of mice with pertussis toxin (an inhibitor of G-protein coupled receptors including chemokine receptors) was shown to prevent SP cells from crossing the cortico-medullary junction, resulting in their retention within cortical regions of the thymus [26, 27]. Of the large family of chemokine receptors, CCR7 has been identified as the prominent receptor mediating this vital relocation step in intrathymic T-cell development, a finding that maps well with the CCR7 expression pattern in SP thymocytes (Fig. 1) [28, 29]. TCR engagement during cortical positive selection induces CCR7 surface expression on developing thymocytes [30], which then migrate into medullary thymic microenvironments containing CCL19- and CCL21-producing stromal cells [31, 32]. Disruption of CCR7-mediated migration results



**Figure 1.** Conventional and Foxp3<sup>+</sup> regulatory T-cell development in thymic microenvironments. Both conventional and Foxp3<sup>+</sup> regulatory CD4<sup>+</sup> T cells are generated from CD4<sup>+</sup>CD8<sup>+</sup> double positive (DP) thymocytes following interactions with self-peptide/MHC complexes expressed by thymic stroma. CCR7 expression during positive selection facilitates the migration of positively selected cells from the cortex to the medulla. Further TCR/MHC interactions with mTECs are involved in the generation of Foxp3<sup>+</sup> nTreg cells from two distinct precursor populations, CD25<sup>+</sup>Foxp3<sup>-</sup> and CD25<sup>-</sup>Foxp3<sup>+</sup> cells. In contrast, the generation of phenotypically mature conventional CD4<sup>+</sup> T cells from recently selected CCR7<sup>lo</sup>CCR9<sup>+</sup>CCR4<sup>+</sup> thymocytes can occur independently of mTECs.

in defective thymocyte migration, cortical accumulation of SP cells [28, 29] and disrupted negative selection [30, 33], leading to a breakdown of central tolerance [34]. Two-photon microscopy monitoring the live migration of CCR7-deficient CD4 SP thymocytes added to thymic slices, suggests that CCR7 may also play a role in the medullary accumulation of thymocytes [27]. In addition, further studies suggest that TCR-mediated signaling is impaired in the absence of CCR7, which may contribute to the altered negative selection of thymocytes observed in *Ccr7*<sup>-/-</sup> mice [30].

Further to the induction of CCR7, positive selection of conventional thymocytes alters the expression of other chemokine receptors that may be linked to cortex to medulla migration. For example, CCR9 has been shown to be involved in multiple stages in intrathymic T-cell development in mice, including CD4 SP generation [35]. While newly selected CD4 SP thymocytes have been shown to retain CCR9 expression (Fig. 1), their expression of PlexinD1 is thought to suppress CCR9-CCL25 signaling, thereby

preventing retention in the cortex and enabling entry into the medulla [36]. Along with CCR7 and PlexinD1 orchestrating thymocyte trafficking, other G-protein coupled receptors may also control medullary access, as CCR7 deficiency appears to have a lesser effect on cortex-to-medulla migration compared with the total block observed with pertussis toxin treatment [26–28]. CCR4 is upregulated on the thymocyte cell surface after the initiation of positive selection (Fig. 1) [37, 38] and its ligands CCL17 and CCL22 are expressed by CD80<sup>high</sup>Aire<sup>+</sup> mTECs [37, 39]. However, *Ccr4*<sup>-/-</sup> mice show no obvious disruption in CD4 and CD8 SP T-cell development or their accumulation within medullary regions [38]. In addition, thymic stromal cells express CCRL1, an atypical chemokine receptor for CCL19, CCL21, and CCL25 [14, 40]. Interestingly however, and in contrast to earlier reports [41], *Ccr11*<sup>-/-</sup> mice show grossly normal thymus architecture and T-cell development [42].

Such findings may reflect redundancy within the extensive chemokine–chemokine receptor system that influences the intrathymic development and location of immature thymocytes [38]. Although the importance of thymocyte entry to the medulla in tolerance induction through negative selection is well established, the role of mTECs in the multistage developmental program of differentiating thymocytes is relatively unclear. To explore this issue, SP thymocyte development has been examined in mice with deficiencies known to disrupt mTEC development, such as *Relb*<sup>-/-</sup> and *Aire*<sup>-/-</sup> mice [17]. Analysis of the latter is of interest and potentially relevant to intrathymic thymocyte migration, as Aire has been linked to the expression of multiple chemokines in the thymic medulla including CCL17, CCL19, CCL21, CCL22, and XCL1 [39, 43, 44]. In both *Relb*<sup>-/-</sup> and *Aire*<sup>-/-</sup> mice, a reduction in mature Qa2<sup>+</sup>CD69<sup>-</sup> CD4 SP thymocytes has been reported [17], suggesting that the transition from immature to mature stages in conventional CD4 SP thymocyte development is dependent upon the presence of mTEC via a mechanism linked to their expression of Aire. However, the consequences of *Relb* deficiency are not exclusive to mTECs development/function, and mice deficient in *Relb* display a complex phenotype including a reduction in thymic DCs, failed lymph node organogenesis and fatal multiorgan autoimmunity [45–47]. To specifically address the role of *Relb*-dependent mTECs in the presence of an otherwise normal immune system, we previously transplanted *Relb*<sup>-/-</sup> fetal thymic stroma into WT mice [24]. Analysis showed the presence of mature Qa2<sup>+</sup> CD69<sup>-</sup> CD4 SP cells within *Relb*<sup>-/-</sup> mTEC-deficient grafts [24]. Moreover, a single cohort of intravenously transferred immature CD4 SP thymocytes was found to undergo late-stage differentiation extrathymically [24]. Although these findings suggest that conventional SP thymocyte development can occur independently of interaction with mTECs (Fig. 1), CD11c<sup>+</sup> DCs present within *Relb*-dependent mTEC-deficient grafts may influence late-stage thymocyte differentiation [38]. More extensive investigations are required to examine the role of thymic DC subsets in fostering conventional CD4 SP thymocyte development, in addition to the division of labor between mTECs and DCs populations in the generation of conventional  $\alpha\beta$  T cells.

## Foxp3<sup>+</sup> natural regulatory T cells


While negative selection plays an important role in shaping the developing  $\alpha\beta$  TCR repertoire, some potentially auto-reactive cells escape intrathymic deletion. Control of unwanted autoimmune responses mediated by these cells requires the thymus to generate a subset of natural CD4<sup>+</sup> regulatory T (nTreg) cells that possesses potent immunosuppressive properties [48]. Commitment to this nTreg lineage occurs in the thymus and depends upon the transcription factor Foxp3, which may be induced as a consequence of increasing affinity of  $\alpha\beta$  TCR/peptide-MHC interactions (reviewed in [49]). Importantly, expression of Foxp3 during nTreg-cell development renders these thymocytes prone to apoptosis unless rescued by signaling through  $\gamma\text{c}$ -dependent cytokines [50]. These findings support a model in which Foxp3<sup>+</sup> nTreg-cell development in the thymus is a multistage process, involving a sequential requirement for TCR and costimulation followed by cytokine receptor signaling. While the precise timing of the branch point of nTreg lineage commitment from that of conventional  $\alpha\beta$  T-cell development is uncertain, two distinct nTreg precursor subsets, CD25<sup>+</sup>Foxp3<sup>-</sup> [51] and CD25<sup>-</sup>Foxp3<sup>+</sup> thymocytes [50], have been identified within CD4 SP  $\alpha\beta$  TCR<sup>+</sup> cells in mice (Fig. 1). While identification of the latter is perhaps consistent with studies reporting Foxp3 expression in some CD4<sup>+</sup>CD8<sup>+</sup> thymocytes [52], the appearance of CD25<sup>+</sup>Foxp3<sup>-</sup> nTreg precursors in CD4SP thymocytes argues against expression of Foxp3 prior to the SP stage [53]. Whatever the developmental sequence may be in Foxp3<sup>+</sup> nTreg-cell generation, access of thymocytes to the medulla has been shown to play an important role in this process (Fig. 1). For example, both CD25<sup>+</sup>Foxp3<sup>-</sup> nTreg precursors and CD25<sup>+</sup>Foxp3<sup>+</sup> nTreg cells are dramatically reduced in the absence of *Relb*-dependent mTECs [24], and it has been suggested that the importance of mTECs during early stages of nTreg-cell development correlates with their provision of appropriate TCR and costimulatory ligands [54–56]. That the availability of mTECs plays a key role in controlling nTreg-cell development is supported by several recent studies. For example, increased mTEC numbers in mice, caused by either loss of TGF- $\beta$ -mediated or OPG-mediated negative control of medullary growth, was found to correlate with increased numbers of thymic Foxp3<sup>+</sup> nTreg cells [57, 58], while diminished mTEC numbers caused by a NIK mutation in *aly/aly* mice was shown to lead to reduced nTreg-cell development [59]. Interestingly, a partial reduction in mTECs observed in mice lacking the adapter molecule Sin did not alter numbers of Foxp3<sup>+</sup> T cells in lymph nodes [60]. Whether this reflects normal Foxp3<sup>+</sup> T-cell development in the presence of diminished mTEC numbers, or postthymic expansion of nTreg in peripheral tissues is not clear. In addition to the role played by mTECs, several studies have also suggested the involvement of DCs in nTreg-cell development in the thymus. While their role in this process has been unclear, particularly in relation to their known involvement in negative selection, recent evidence demonstrates the importance of the combined presence of both mTECs and DCs, in that *Batf3*-dependent CD8 $\alpha$ <sup>+</sup> DCs have been shown to support Foxp3<sup>+</sup> T-cell development via presentation of Aire-dependent antigens provided by mTECs [12].



Although the studies described above highlight the importance of the thymus medulla in nTreg-cell generation, the mechanisms that guide the migration of these cells and their lineage-restricted precursors to this region are not fully understood. For example, while the majority of  $\text{Foxp3}^+\text{CD4}^+$  thymocytes expresses CCR7 (Fig. 1) and demonstrate heterogeneity with regard to expression of CXCR4 and CCR8 [61], it is not clear how this heterogeneity relates to the stages in nTreg-cell development described above. More recent analysis shows that  $\text{CD25}^+\text{Foxp3}^-$  precursors in mice express CCR4 and CCR7, while their  $\text{CD25}^+\text{Foxp3}^+$  nTreg progeny are uniformly  $\text{CCR4}^-\text{CCR7}^+$  [24, 38]. Despite the transient nature of CCR4 expression,  $\text{Foxp3}^+$  nTreg-cell development appears normal in  $\text{Ccr4}^{-/-}$  mice [38]. In contrast,  $\text{Ccr7}^{-/-}$  mice do show alterations in the development and intrathymic positioning of nTreg cells [38, 52, 62], consistent with the idea that conventional  $\alpha\beta$  T cells and developing nTreg cells share a requirement for CCR7 to enter the medulla. This scenario is also consistent with the diminished number of  $\text{Foxp3}^+$  nTreg cells in mice lacking podoplanin, a molecule that controls the intrathymic localization of CCL21, a ligand for CCR7 that is involved in thymocyte entry to the medulla [63]. Interestingly however, nTreg-cell numbers are not decreased in  $\text{Ccr7}^{-/-}$  mice [38]. In fact, while the frequency of  $\text{CD25}^+\text{Foxp3}^-$  nTreg precursors is unaltered,  $\text{CD25}^+\text{Foxp3}^+$  nTreg cells are actually increased in  $\text{Ccr7}^{-/-}$  mice [38, 62]. Of further relevance to the involvement of CCR7 during nTreg-cell development is the recent identification and characterization of the mTEC subset expressing CCL21. Lkhagvasuren et al. [13] showed that the development of  $\text{CCL21}^+$  mTECs is controlled by  $\text{LT}\beta\text{R}$  signaling, with a decrease in this mTEC subset evident in  $\text{Ltbr}^{-/-}$  mice. Moreover,  $\text{CCL21}^+$  mTECs are largely distinct from the  $\text{Aire}^+$  mTECs subset. However, while  $\text{Aire}^{-/-}$  mice have reported defects in  $\text{Foxp3}^+$  T-cell development [43, 64],  $\text{Ltbr}^{-/-}$  mice do not [65, 66]. Taken together, these findings suggest that in the absence of CCR7 expression by nTreg precursors, or  $\text{LT}\beta\text{R}$ -mediated CCL21 production by mTECs, nTreg precursors can still develop. This suggests that other chemokines distinct from those triggered by  $\text{LT}\beta\text{R}$  signaling are able to control the access to mTECs that is required for nTreg-cell generation. The reasons behind the increase in  $\text{CD25}^+\text{Foxp3}^+$  nTreg cells in  $\text{Ccr7}^{-/-}$  mice remain unexplained. It is unclear whether this occurs as a result of defective emigration leading to increased thymic retention of nTreg cells, or is caused by the reentry of peripheral  $\text{Foxp3}^+$  Treg cells back to the thymus. Furthermore, whether these possibilities also account for the large frequency of  $\text{Rag2p-GFP}^- \text{Foxp3}^+$  nTreg cells observed in  $\text{RAG2p-GFP}$  reporter mice [15, 67, 68] requires additional investigation.

## CD1d-restricted iNKT cells

Of the other  $\alpha\beta$  T-cell subsets that are generated intrathymically, arguably the most well described are Type 1 invariant natural killer T cells (iNKT cells), which recognize glycolipid antigens bound to CD1d molecules (reviewed in [69–71]). Unlike conventional  $\alpha\beta$  T cells, iNKT cells express an invariant  $\text{V}\alpha 14\text{-J}\alpha 18$  TCR $\alpha$  chain that pairs with a limited number of TCR $\beta$  chains. Moreover,



**Figure 2.** CD1d-restricted iNKT-cell development in thymic microenvironments. Initiation of iNKT-cell development in the thymus occurs in the cortex following recognition of glycolipid/CD1d complexes on  $\text{CD4}^+\text{CD8}^+$  DP thymocytes. The migration of developing iNKT-cells from the cortex to the medulla involves upregulation of CCR7, and so may involve  $\text{CCL21}^+$  mTECs (Stage 0, St. 0). In the medulla, later stages of iNKT-cell development involve IL-15 transpresentation from a subset of  $\text{mTEC}^{\text{low}}$  cells (Stage 2, St. 2). Any relationship between the  $\text{mTEC}^{\text{low}}$  cells which provide CCL21 and those which trans-present IL-15 is not known. In addition to the importance of the thymus medulla for iNKT-cell generation, developing iNKT cells express RANKL to drive  $\text{Aire}^+$  mTEC differentiation. Stages in iNKT-cell development [102]: St0:  $\text{CD24}^+\text{CD44}^-\text{NK1.1}^-$ , St1:  $\text{CD24}^-\text{CD44}^-\text{NK1.1}^-$ , St2:  $\text{CD44}^+\text{NK1.1}^-$ , St3:  $\text{CD44}^+\text{NK1.1}^+$ .


the positive selection of iNKT cells (Fig. 2) depends upon the recognition of CD1d/glycolipid complexes expressed by cortex resident  $\text{CD4}^+\text{CD8}^+$  thymocytes and not TEC [5, 72]. Although cell surface markers (CD24, CD44, NK1.1), cytokine production (IL-4, IFN- $\gamma$ , IL-17) and transcription factor expression (PLZF, T-bet, GATA3) identify heterogeneity among iNKT cells during their intrathymic development [73–75], relatively little is known about their positioning in the thymus. This is likely due, at least in part, to the technical difficulty of using CD1d tetramers to detect iNKT cells in tissue sections. However, that positive selection of iNKT cells occurs in the cortex draws parallels with conventional  $\alpha\beta$  T cells, and suggests that iNKT cells also undergo cortex-to-medulla migration during their intrathymic development (Fig. 2). Indeed, PCR analysis of thymic iNKT-cell subsets separated on the basis of CD4 and IL17RB expression shows their differential expression of mRNAs encoding the chemokine

receptors CCR4, CCR6, and CCR7, the ligands for which are expressed in the medulla [76].

While the relationship between expression of these chemokine receptors and the intrathymic location of particular iNKT-cell subsets is unclear, expression of the chemokine receptor CXCR6 is induced during iNKT-cell positive selection, a process that occurs in the cortex [77]. However, while expression of a CXCR6 ligand (CXCL16) has been detectable in thymus medulla [78] no defects were reported during intrathymic iNKT-cell development in CXCR6-deficient mice [77]. By performing flow cytometric analysis to compare chemokine receptor expression during conventional  $\alpha\beta$  T-cell and iNKT-cell development [38], we found that CCR7 expression was most abundant in CD24<sup>+</sup>CD44<sup>-</sup>NK1.1<sup>-</sup> (stage 0) iNKT cells, suggesting CCR7 induction takes place following their positive selection to mediate trafficking from the cortex to the medulla. Consistent with this, mice lacking CCR7 were shown to have fewer PBS57<sup>+</sup> iNKT cells, notably the CD44<sup>-</sup>NK1.1<sup>-</sup> (stage 1), CD44<sup>+</sup>NK1.1<sup>-</sup> (stage 2) and CD44<sup>+</sup>NK1.1<sup>+</sup> (stage 3) subsets (Fig. 2). The involvement of the thymus medulla in iNKT-cell development fits well with the requirement for costimulation via B7 family members, which are predominantly expressed in the medulla [79, 80]. It also correlates with the reduction in iNKT cells in thymus grafts devoid of Relb-dependent mTECs [81–83] that act as a source of IL-15/IL-15R $\alpha$  trans-presentation [81]. While further studies are necessary to fully understand the impact of mTECs on iNKT-cell development, expression of Aire by mTEC has been reported to be dispensable [84]. This suggests that although the absence of Aire expression may alter the intrathymic expression of several chemokines [43, 44], these changes do not necessarily impact on iNKT-cell development. Finally, the link between medullary microenvironments and intrathymic iNKT-cell development has been strongly highlighted in studies exploring “long-term resident” NK1.1<sup>+</sup>CD44<sup>+</sup> iNKT cells, which have been shown to remain in the thymus for extended periods [85]. While the functional significance of these cells is not yet known, their thymic residency requires the chemokine receptor CXCR3, the ligands for which are expressed by mTECs [86]. Collectively, the studies outlined above are important as they support the idea that, as with conventional  $\alpha\beta$  T cells and Foxp3<sup>+</sup> Treg cells (Fig. 1), iNKT cell populations in the thymus (Fig. 2) depend upon both the cortical and medullary microenvironments. Furthermore, that iNKT cells undergo stepwise migration through these sites and can influence mTECs via their expression of RANKL and CD40L [81] are additional features shared with conventional  $\alpha\beta$  T cells and Foxp3<sup>+</sup> Treg cells.

## $\gamma\delta$ T cells

In addition to supporting the development of the  $\alpha\beta$  T-cell subsets described above, the thymus fosters the development of multiple  $\gamma\delta$  T-cell subsets. It has long been known that the appearance of heterogeneous  $\gamma\delta$  T-cell populations follows a temporally ordered program of development, characterized by defined TCR- $\gamma$  variable gene usage, which is in turn accompanied by biased tissue



**Figure 3.** Fetal and adult phases of  $\gamma\delta$  T-cell development in thymic microenvironments. The thymus supports the generation of distinct subsets of  $\gamma\delta$  T cells in both fetal and adult life. In the fetal thymus, the first  $\gamma\delta$  T cells to emerge express the V $\gamma$ 5<sup>+</sup> T-cell receptor and represent progenitors of dendritic epidermal T cells. As with the intrathymic development of iNKT cells and conventional T cells, fetal  $\gamma\delta$  T cells (right) also express RANKL and influence the development of Aire<sup>+</sup> mTECs in the medulla. Note that although V $\gamma$ 5<sup>+</sup> thymocytes express CCR4 and CCR7, they are not required for their intrathymic development. In contrast, CCR6 expression by fetal  $\gamma\delta$  T cells has been linked with their thymic exit. Interestingly, in contrast to fetal thymus (right), CCR7 has been shown to play a role on the medullary accumulation of  $\gamma\delta$  T cells in the adult thymus (left).

tropism [77]. In this regard, during murine fetal development,  $\gamma\delta$  T cells utilizing restricted V $\gamma$ 5 gene segments (Fig. 3) have been shown to arise first, (colonizing epidermal sites and therefore being termed dendritic epidermal T cells DETCs), followed by a wave of V $\gamma$ 6-bearing  $\gamma\delta$  T cells (lung, reproductive tract, and tongue), and subsequently V $\gamma$ 1 and V $\gamma$ 4 cells (dermis, secondary lymphoid tissues) [87, 88]. Moreover, such defined  $\gamma\delta$  T-cell subsets, in a similar fashion to recently described iNKT-cell subsets and novel natural Th17 cells [75, 89], appear to preferentially express particular cytokines (IFN- $\gamma$ , IL-17, and IL-4) and associated transcription factors (Tbet, ROR $\gamma$ t, and PLZF) [90]. In contrast to development of  $\alpha\beta$  T cells, the intrathymic developmental requirements of  $\gamma\delta$  T cells remain comparatively poorly defined, including their dependency upon positive and negative selection. However, evidence has implicated a role for instructive signaling in the direction of  $\gamma\delta$  T-cell maturation, including conditioning by conventional CD4<sup>+</sup>8<sup>+</sup>  $\alpha\beta$  T cells via the lymphotoxin pathway [91], and importantly, interaction of fetal V $\gamma$ 5<sup>+</sup> DETCs

with RANK-dependent mTECs expressing the immunoglobulin-like molecule Skint-1 (Selection and upKeep of INtraepithelial T cells, Fig. 3) [92, 93]. Critically, conditioning of  $V\gamma 5^+$  DETCs by Skint-1<sup>+</sup> mTECs appears to bias DETCs toward IFN- $\gamma$  production, with Skint-1 mutant mice conversely producing IL-17<sup>+</sup> DETCs [94]. Consistent with a potential role for the medulla in the conditioning of fetal  $\gamma\delta$  T cells, initial experiments analyzing the developmental window during which  $\gamma\delta$  T cells appeared within the embryonic thymus revealed an accumulation of  $\gamma\delta$  TCR<sup>+</sup> cells from approximately E15 of gestation, coinciding with early phases of thymic medullary islet formation [6]. Furthermore, histological analysis of the intrathymic anatomical positioning of  $\gamma\delta$  T cells at later stages of fetal thymus development revealed a preferential accumulation of  $\gamma\delta$  T cells within medullary foci [95, 96]. Interestingly however, the positioning of adult  $\gamma\delta$  T cells has been reported to subsequently demonstrate a reduced bias to medullary regions, with the majority of  $\gamma\delta$  T cells found to localize to cortical, including subcapsular, and cortico-medullary regions [96, 97]. Significantly, the functional importance of the medullary accumulation of fetal  $\gamma\delta$  T cells appears not only important for the conditioning of IFN- $\gamma^+$   $V\gamma 5^+$  DETCs, but in a similar fashion to both conventional and iNKT  $\alpha\beta$  T cells, fetal  $\gamma\delta$  T cells were reported to play a role in the reciprocal conditioning of mTECs (Fig. 3), via provision of RANKL to drive the development of the earliest cohorts of Aire<sup>+</sup> mTECs [92].

Although the requirements for a strictly ordered progression of conventional  $\alpha\beta$  T-cell sequential migration through cortical and medullary microenvironments has been relatively well characterized, the dependency of  $\gamma\delta$  T cells upon ordered intrathymic migration, and the mechanisms that control this potential process, both remain much less well-defined. Significantly however, in vitro treatment of E15 fetal thymus organ cultures with pertussis toxin was found to prevent the accumulation  $\gamma\delta$  T cells within medullary regions, suggesting that in parallel with  $\alpha\beta$  T cells, the medullary localization of fetal  $\gamma\delta$  T cells is highly dependent upon G-protein coupled receptor signaling [26, 95]. Interestingly, initial studies analyzing the development of fetal thymic DETCs revealed that the putative selection of  $V\gamma 5^+$  thymocytes resulted in the down-regulation of CCR4 and CCR6 with concomitant upregulation of CCR10, indicating changes in chemokine receptor expression may relate to migration of developing  $\gamma\delta$  T cells through the thymus [98]. Despite the expression of the relevant cognate ligands for CCR4 and CCR10 by mTECs [39], the development of  $V\gamma 5^+$  thymocytes occurs normally in both *Ccr4*- and *Ccr10*-deficient fetal thymuses [95, 99]. In contrast, analysis of an intrathymic role for CCR6 expression by  $V\gamma 5^+$  DETCs revealed that the selection-associated loss of CCR6 plays a critical role in releasing mature  $V\gamma 5^+$   $\gamma\delta$  T cells from CCL20-producing medullary microenvironments [100], thereby facilitating thymic egress [101]. Consistent with  $\alpha\beta$  T-cell populations,  $\gamma\delta$  T cells additionally express CCR7 at both fetal and adult stages [95, 99]. Although CCR7-deficient fetal  $\gamma\delta$  T cells demonstrated no reduction in medullary localization [95], recent studies analyzing the intrathymic positioning of adult  $\gamma\delta$  T cells revealed a significant reduction in medulla-associated  $\gamma\delta$  T cells in *Ccr7*<sup>-/-</sup> mice [97]. However, beyond the develop-

mental requirement of fetal  $V\gamma 5^+$  DETC progenitors for reciprocal signaling interactions with mTECs, the significance of medullary localization and interactions for additional fetal and adult  $\gamma\delta$ T-cell subsets remains unclear and warrants further investigation.

## Conclusions

The medulla has long been recognized as a site that contains thymocyte subsets that represent late stages in T-cell development. Through the process of negative selection, it plays an essential role in conventional  $\alpha\beta$  T-cell development that ensures the generation of self-tolerant T cells. In addition, increasing evidence suggests that the medulla also influences the development of multiple T-cell lineages that are generated during both fetal and adult life. Recent studies are also now highlighting functional complexity within the mTEC compartment, which may help to explain the contributions made by these cells in supporting the development of diverse T-cell populations. As the cellular and molecular complexity of medullary microenvironments becomes clearer, we anticipate that this will lead to a better understanding of the processes that govern intrathymic T-cell development.

**Acknowledgment:** This work was supported by an MRC Programme Grant to G.A.

**Conflict of interest:** The authors declare no financial or commercial conflict of interest.

## References

- 1 Wencker, M., Turchinovich, G., Di Marco Barros, R., Deban, L., Jandke, A., Cope, A., Hayday, A. C. et al., Innate-like T cells straddle innate and adaptive immunity by altering antigen-receptor responsiveness. *Nat. Immunol.* 2014. 15: 80–87.
- 2 Takahama, Y., Journey through the thymus: stromal guides for T-cell development and selection. *Nat. Rev. Immunol.* 2006. 6: 127–135.
- 3 Boehm, T. and Swann, J. B., Thymus involution and regeneration: two sides of the same coin? *Nat. Rev. Immunol.* 2013. 13: 831–838.
- 4 Fontenot, J. D., Dooley, J. L., Farr, A. G. and Rudensky, A. Y., Developmental regulation of Foxp3 expression during ontogeny. *J. Exp. Med.* 2005. 202: 901–906.
- 5 Bendelac, A., Positive selection of mouse NK1+ T cells by CD1-expressing cortical thymocytes. *J. Exp. Med.* 1995. 182: 2091–2096.
- 6 Havran, W. L. and Allison, J. P., Developmentally ordered appearance of thymocytes expressing different T-cell antigen receptors. *Nature* 1988. 335: 443–445.
- 7 Griffith, A. V., Fallahi, M., Nakase, H., Gosink, M., Young, B. and Petrie, H. T., Spatial mapping of thymic stromal microenvironments reveals

- unique features influencing T lymphoid differentiation. *Immunity* 2009. 31: 999–1009.
- 8 Anderson, G., Jenkinson, E. J. and Rodewald, H. R., A roadmap for thymic epithelial cell development. *Eur. J. Immunol.* 2009. 39: 1694–1699.
  - 9 Alves, N. L., Takahama, Y., Ohigashi, I., Ribeiro, A. R., Baik, S., Anderson, G., Jenkinson, W. E. et al., Serial progression of cortical and medullary thymic epithelial microenvironments. *Eur. J. Immunol.* 2014. 44: 16–22.
  - 10 Breart, B., Ramos-Perez, W. D., Mendoza, A., Salous, A. K., Gobert, M., Huang, Y., Adams, R. H. et al., Lipid phosphate phosphatase 3 enables efficient thymic egress. *J. Exp. Med.* 2011. 208: 1267–1278.
  - 11 Anderson, G. and Takahama, Y., Thymic epithelial cells: working class heroes for T cell development and repertoire selection. *Trends Immunol.* 2012. 33: 256–263.
  - 12 Perry, J. S., Lio, C. W., Kau, A. L., Nutsch, K., Yang, Z., Gordon, J. I., Murphy, K. M. and Hsieh, C. S., Distinct contributions of aire and antigen-presenting-cell subsets to the generation of self-tolerance in the thymus. *Immunity* 2014. 41: 414–426.
  - 13 Lkhagvasuren, E., Sakata, M., Ohigashi, I. and Takahama, Y., Lymphotoxin beta receptor regulates the development of CCL21-expressing subset of postnatal medullary thymic epithelial cells. *J. Immunol.* 2013. 190: 5110–5117.
  - 14 Ribeiro, A. R., Meireles, C., Rodrigues, P. M. and Alves, N. L., Intermediate expression of CCRL1 reveals novel subpopulations of medullary thymic epithelial cells that emerge in the postnatal thymus. *Eur. J. Immunol.* 2014. 44: 2918–2924.
  - 15 McCaughy, T. M., Wilken, M. S. and Hogquist, K. A., Thymic emigration revisited. *J. Exp. Med.* 2007. 204: 2513–2520.
  - 16 Scollay, R. and Godfrey, D. I., Thymic emigration: conveyor belts or lucky dips? *Immunol. Today* 1995. 16: 268–273; discussion 273–264.
  - 17 Li, J., Li, Y., Yao, J. Y., Jin, R., Zhu, M. Z., Qian, X. P., Zhang, et al., Developmental pathway of CD4+CD8- medullary thymocytes during mouse ontogeny and its defect in Aire-/- mice. *Proc. Natl. Acad. Sci. USA* 2007. 104: 18175–18180.
  - 18 Lucas, B., Vasseur, F. and Penit, C., Production, selection, and maturation of thymocytes with high surface density of TCR. *J. Immunol.* 1994. 153: 53–62.
  - 19 Kelly, K. A. and Scollay, R., Analysis of recent thymic emigrants with subset- and maturity-related markers. *Int. Immunol.* 1990. 2: 419–425.
  - 20 Ge, Q. and Chen, W. F., Phenotypic identification of the subgroups of murine T-cell receptor alpha beta+ CD4+ CD8- thymocytes and its implication in the late stage of thymocyte development. *Immunology* 1999. 97: 665–671.
  - 21 Tian, T., Zhang, J., Gao, L., Qian, X. P. and Chen, W. F., Heterogeneity within medullary-type TCRalpha beta(+)/CD3(+)/CD4(-)/CD8(+) thymocytes in normal mouse thymus. *Int. Immunol.* 2001. 13: 313–320.
  - 22 Jin, R., Wang, W., Yao, J. Y., Zhou, Y. B., Qian, X. P., Zhang, J., Zhang, Y. and Chen, W. F., Characterization of the in vivo dynamics of medullary CD4+CD8- thymocyte development. *J. Immunol.* 2008. 180: 2256–2263.
  - 23 Teng, F., Zhou, Y., Jin, R., Chen, Y., Pei, X., Liu, Y., Dong, J. et al., The molecular signature underlying the thymic migration and maturation of TCRalpha beta+ CD4+ CD8 thymocytes. *PLoS One* 2011. 6: e25567.
  - 24 Cowan, J. E., Parnell, S. M., Nakamura, K., Caamano, J. H., Lane, P. J., Jenkinson, E. J., Jenkinson, W. E. and Anderson, G., The thymic medulla is required for Foxp3+ regulatory but not conventional CD4+ thymocyte development. *J. Exp. Med.* 2013. 210: 675–681.
  - 25 Witt, C. M., Raychaudhuri, S., Schaefer, B., Chakraborty, A. K. and Robey, E. A., Directed migration of positively selected thymocytes visualized in real time. *PLoS Biol* 2005. 3: e160.
  - 26 Suzuki, G., Sawa, H., Kobayashi, Y., Nakata, Y., Nakagawa, K., Uzawa, A., Sakiyama, H. et al., Pertussis toxin-sensitive signal controls the trafficking of thymocytes across the corticomedullary junction in the thymus. *J. Immunol.* 1999. 162: 5981–5985.
  - 27 Ehrlich, L. I., Oh, D. Y., Weissman, I. L. and Lewis, R. S., Differential contribution of chemotaxis and substrate restriction to segregation of immature and mature thymocytes. *Immunity* 2009. 31: 986–998.
  - 28 Ueno, T., Saito, F., Gray, D. H., Kuse, S., Hieshima, K., Nakano, H., Kakuchi, T. et al., CCR7 signals are essential for cortex-medulla migration of developing thymocytes. *J. Exp. Med.* 2004. 200: 493–505.
  - 29 Kwan, J. and Killeen, N., CCR7 directs the migration of thymocytes into the thymic medulla. *J. Immunol.* 2004. 172: 3999–4007.
  - 30 Davalos-Misslitz, A. C., Worbs, T., Willenzon, S., Bernhardt, G. and Forster, R., Impaired responsiveness to T-cell receptor stimulation and defective negative selection of thymocytes in CCR7-deficient mice. *Blood* 2007. 110: 4351–4359.
  - 31 Ueno, T., Hara, K., Willis, M. S., Malin, M. A., Hopken, U. E., Gray, D. H., Matsushima, K. et al., Role for CCR7 ligands in the emigration of newly generated T lymphocytes from the neonatal thymus. *Immunity* 2002. 16: 205–218.
  - 32 Halkias, J., Melichar, H. J., Taylor, K. T., Ross, J. O., Yen, B., Cooper, S. B., Winoto, A. and Robey, E. A., Opposing chemokine gradients control human thymocyte migration in situ. *J. Clin. Invest.* 2013. 123: 2131–2142.
  - 33 Nitta, T., Nitta, S., Lei, Y., Lipp, M. and Takahama, Y., CCR7-mediated migration of developing thymocytes to the medulla is essential for negative selection to tissue-restricted antigens. *Proc. Natl. Acad. Sci. USA* 2009. 106: 17129–17133.
  - 34 Kurobe, H., Liu, C., Ueno, T., Saito, F., Ohigashi, I., Seach, N., Arakaki, R. et al., CCR7-dependent cortex-to-medulla migration of positively selected thymocytes is essential for establishing central tolerance. *Immunity* 2006. 24: 165–177.
  - 35 Svensson, M., Marsal, J., Uronen-Hansson, H., Cheng, M., Jenkinson, W., Cilio, C., Jacobsen, S. E. et al., Involvement of CCR9 at multiple stages of adult T lymphopoiesis. *J. Leukoc. Biol.* 2008. 83: 156–164.
  - 36 Choi, Y. I., Duke-Cohan, J. S., Ahmed, W. B., Handley, M. A., Mann, F., Epstein, J. A., Clayton, L. K. and Reinherz, E. L., PlexinD1 glycoprotein controls migration of positively selected thymocytes into the medulla. *Immunity* 2008. 29: 888–898.
  - 37 Campbell, J. J., Pan, J. and Butcher, E. C., Cutting edge: developmental switches in chemokine responses during T cell maturation. *J. Immunol.* 1999. 163: 2353–2357.
  - 38 Cowan, J. E., McCarthy, N. I., Parnell, S. M., White, A. J., Bacon, A., Serge, A., Irla, M. et al., Differential requirement for CCR4 and CCR7 during the development of innate and adaptive alpha beta T cells in the adult thymus. *J. Immunol.* 2014. 193: 1204–1212.
  - 39 Laan, M., Kisand, K., Kont, V., Moll, K., Tserel, L., Scott, H. S. and Peterson, P., Autoimmune regulator deficiency results in decreased expression of CCR4 and CCR7 ligands and in delayed migration of CD4+ thymocytes. *J. Immunol.* 2009. 183: 7682–7691.
  - 40 Heinzl, K., Benz, C. and Bleul, C. C., A silent chemokine receptor regulates steady-state leukocyte homing in vivo. *Proc. Natl. Acad. Sci. USA* 2007. 104: 8421–8426.
  - 41 Comerford, I., Nibbs, R. J., Litchfield, W., Bunting, M., Harata-Lee, Y., Haylock-Jacobs, S., Forrow, S. et al., The atypical chemokine receptor CCX-CKR scavenges homeostatic chemokines in circulation and suppresses Th17 responses. *Blood* 2010. 116: 4130–4140.



- 42 Lucas, B., White, A. J., Ulvmar, M. H., Nibbs, R. J., Sitnik, K. M., Agace, W. W., Jenkinson, W. E. et al., CCRL1/ACKR4 is expressed in key thymic microenvironments but is dispensable for T lymphopoiesis at steady state in adult mice. *Eur. J. Immunol.* 2015. 45: 574–583.
- 43 Lei, Y., Ripen, A. M., Ishimaru, N., Ohigashi, I., Nagasawa, T., Jeker, L. T., Bosl, M. R. et al., Aire-dependent production of XCL1 mediates medullary accumulation of thymic dendritic cells and contributes to regulatory T cell development. *J. Exp. Med.* 2011. 208: 383–394.
- 44 Hubert, F. X., Kinkel, S. A., Davey, G. M., Phipson, B., Mueller, S. N., Liston, A., Proietto, A. I. et al., Aire regulates the transfer of antigen from mTECs to dendritic cells for induction of thymic tolerance. *Blood* 2011. 118: 2462–2472.
- 45 Weih, F., Carrasco, D., Durham, S. K., Barton, D. S., Rizzo, C. A., Ryseck, R. P., Lira, S. A. and Bravo, R., Multiorgan inflammation and hematopoietic abnormalities in mice with a targeted disruption of RelB, a member of the NF-kappa B/Rel family. *Cell* 1995. 80: 331–340.
- 46 Burkly, L., Hession, C., Ogata, L., Reilly, C., Marconi, L. A., Olson, D., Tizard, R. et al., Expression of relB is required for the development of thymic medulla and dendritic cells. *Nature* 1995. 373: 531–536.
- 47 Wu, L., D'Amico, A., Winkel, K. D., Suter, M., Lo, D. and Shortman, K., RelB is essential for the development of myeloid-related CD8alpha-dendritic cells but not of lymphoid-related CD8alpha+ dendritic cells. *Immunity* 1998. 9: 839–847.
- 48 Lio, C. W. and Hsieh, C. S., Becoming self-aware: the thymic education of regulatory T cells. *Curr. Opin. Immunol.* 2011. 23: 213–219.
- 49 Klein, L., Kyewski, B., Allen, P. M. and Hogquist, K. A., Positive and negative selection of the T cell repertoire: what thymocytes see (and don't see). *Nat. Rev. Immunol.* 2014. 14: 377–391.
- 50 Tai, X., Erman, B., Alag, A., Mu, J., Kimura, M., Katz, G., Guinter, T. et al., Foxp3 transcription factor is proapoptotic and lethal to developing regulatory T cells unless counterbalanced by cytokine survival signals. *Immunity* 2013. 38: 1116–1128.
- 51 Lio, C. W. and Hsieh, C. S., A two-step process for thymic regulatory T cell development. *Immunity* 2008. 28: 100–111.
- 52 Liston, A., Nutsch, K. M., Farr, A. G., Lund, J. M., Rasmussen, J. P., Koni, P. A. and Rudensky, A. Y., Differentiation of regulatory Foxp3(+) T cells in the thymic cortex. *Proc. Natl. Acad. Sci. USA* 2008. 105: 11903–11908.
- 53 Lee, H. M. and Hsieh, C. S., Rare development of Foxp3+ thymocytes in the CD4+CD8+ subset. *J. Immunol.* 2009. 183: 2261–2266.
- 54 Mahmud, S. A., Manlove, L. S., Schmitz, H. M., Xing, Y., Wang, Y., Owen, D. L., Schenkel, J. M. et al., Costimulation via the tumor-necrosis factor receptor superfamily couples TCR signal strength to the thymic differentiation of regulatory T cells. *Nat. Immunol.* 2014. 15: 473–481.
- 55 Hinterberger, M., Wirnsberger, G. and Klein, L., B7/CD28 in central tolerance: costimulation promotes maturation of regulatory T cell precursors and prevents their clonal deletion. *Front Immunol.* 2011. 2: 30.
- 56 Verhagen, J., Gabrysova, L., Minaee, S., Sabatos, C. A., Anderson, G., Sharpe, A. H. and Wraith, D. C., Enhanced selection of Foxp3+ T-regulatory cells protects CTLA-4-deficient mice from CNS autoimmune disease. *Proc. Natl. Acad. Sci. USA* 2009. 106: 3306–3311.
- 57 Hauri-Hohl, M., Zuklys, S., Hollander, G. A. and Ziegler, S. F., A regulatory role for TGF-beta signaling in the establishment and function of the thymic medulla. *Nat. Immunol.* 2014. 15: 554–561.
- 58 Akiyama, N., Shinzawa, M., Miyauchi, M., Yanai, H., Tateishi, R., Shimo, Y. et al., Limitation of immune tolerance-inducing thymic epithelial cell development by Spi-B-mediated negative feedback regulation. *J. Exp. Med.* 2014. 211: 2425–2438.
- 59 Mouri, Y., Nishijima, H., Kawano, H., Hirota, F., Sakaguchi, N., Morimoto, J. and Matsumoto, M., NF-kappaB-inducing kinase in thymic stroma establishes central tolerance by orchestrating cross-talk with not only thymocytes but also dendritic cells. *J. Immunol.* 2014. 193: 4356–4367.
- 60 Danzl, N. M., Donlin, L. T. and Alexandropoulos, K., Regulation of medullary thymic epithelial cell differentiation and function by the signaling protein Sin. *J. Exp. Med.* 2010. 207: 999–1013.
- 61 Lee, J. H., Kang, S. G. and Kim, C. H., FoxP3+ T cells undergo conventional first switch to lymphoid tissue homing receptors in thymus but accelerated second switch to nonlymphoid tissue homing receptors in secondary lymphoid tissues. *J. Immunol.* 2007. 178: 301–311.
- 62 Schneider, M. A., Meingassner, J. G., Lipp, M., Moore, H. D. and Rot, A., CCR7 is required for the in vivo function of CD4+ CD25+ regulatory T cells. *J. Exp. Med.* 2007. 204: 735–745.
- 63 Fuertbauer, E., Zaujec, J., Uhrin, P., Raab, I., Weber, M., Schachner, H., Bauer, M. et al., Thymic medullary conduits-associated podoplanin promotes natural regulatory T cells. *Immunol. Lett.* 2013. 154: 31–41.
- 64 Malchow, S., Leventhal, D. S., Nishi, S., Fischer, B. I., Shen, L., Paner, G. P., Amit, A. S. et al., Aire-dependent thymic development of tumor-associated regulatory T cells. *Science* 2013. 339: 1219–1224.
- 65 Martins, V. C., Boehm, T. and Bleul, C. C., Ltbeta signaling does not regulate Aire-dependent transcripts in medullary thymic epithelial cells. *J. Immunol.* 2008. 181: 400–407.
- 66 Zhu, M., Chin, R. K., Tumanov, A. V., Liu, X. and Fu, Y. X., Lymphotoxin beta receptor is required for the migration and selection of autoreactive T cells in thymic medulla. *J. Immunol.* 2007. 179: 8069–8075.
- 67 Cuss, S. M. and Green, E. A., Abrogation of CD40-CD154 signaling impedes the homeostasis of thymic resident regulatory T cells by altering the levels of IL-2, but does not affect regulatory T cell development. *J. Immunol.* 2012. 189: 1717–1725.
- 68 Yang, E., Zou, T., Lechner, T. M., Zhang, S. L. and Kambayashi, T., Both retention and recirculation contribute to long-lived regulatory T-cell accumulation in the thymus. *Eur. J. Immunol.* 2014. 44: 2712–2720.
- 69 Godfrey, D. I., Stankovic, S. and Baxter, A. G., Raising the NKT cell family. *Nat. Immunol.* 2010. 11: 197–206.
- 70 Engel, I. and Kronenberg, M., Transcriptional control of the development and function of valpha14i NKT cells. *Curr. Top Microbiol. Immunol.* 2014. 381: 51–81.
- 71 Constantinides, M. G. and Bendelac, A., Transcriptional regulation of the NKT cell lineage. *Curr. Opin. Immunol.* 2013. 25: 161–167.
- 72 Wei, D. G., Lee, H., Park, S. H., Beaudoin, L., Teyton, L., Lehuen, A. and Bendelac, A., Expansion and long-range differentiation of the NKT cell lineage in mice expressing CD1d exclusively on cortical thymocytes. *J. Exp. Med.* 2005. 202: 239–248.
- 73 Benlagha, K., Wei, D. G., Veiga, J., Teyton, L. and Bendelac, A., Characterization of the early stages of thymic NKT cell development. *J. Exp. Med.* 2005. 202: 485–492.
- 74 Coquet, J. M., Chakravarti, S., Kyparissoudis, K., McNab, F. W., Pitt, L. A., McKenzie, B. S., Berzins, S. P. et al., Diverse cytokine production by NKT cell subsets and identification of an IL-17-producing CD4-NK1.1-NKT cell population. *Proc. Natl. Acad. Sci. USA* 2008. 105: 11287–11292.
- 75 Lee, Y. J., Holzappel, K. L., Zhu, J., Jameson, S. C. and Hogquist, K. A., Steady-state production of IL-4 modulates immunity in mouse strains and is determined by lineage diversity of iNKT cells. *Nat. Immunol.* 2013. 14: 1146–1154.
- 76 Watarai, H., Sekine-Kondo, E., Shigeura, T., Motomura, Y., Yasuda, T., Satoh, R., Yoshida, H. et al., Development and function of invariant natural killer T cells producing T(h)2- and T(h)17-cytokines. *PLoS Biol.* 2012. 10: e1001255.

- 77 Germanov, E., Veinotte, L., Cullen, R., Chamberlain, E., Butcher, E. C. and Johnston, B., Critical role for the chemokine receptor CXCR6 in homeostasis and activation of CD1d-restricted NKT cells. *J. Immunol.* 2008. **181**: 81–91.
- 78 Matloubian, M., David, A., Engel, S., Ryan, J. E. and Cyster, J. G., A transmembrane CXC chemokine is a ligand for HIV-coreceptor Bonzo. *Nat. Immunol.* 2000. **1**: 298–304.
- 79 Williams, J. A., Lumsden, J. M., Yu, X., Feigenbamm, L., Zhang, J. J., Steinberg, S. M. and Hodes, R. J., Regulation of thymic NKT cell development by the B7-CD28 costimulatory pathway. *J. Immunol.* 2008. **181**: 907–917.
- 80 Chung, Y., Nurieva, R., Esashi, E., Wang, Y. H., Zhou, D. P., Gapin, L. and Dong, C., A critical role of costimulation during intrathymic development of invariant NKT cells. *Faseb. J.* 2008. **180**: 2276–2283.
- 81 White, A. J., Jenkinson, W. E., Cowan, J. E., Parnell, S. M., Bacon, A., Jones, N. D., Jenkinson, E. J. and Anderson, G., An essential role for medullary thymic epithelial cells during the intrathymic development of invariant NKT cells. *J. Immunol.* 2014. **192**: 2659–2666.
- 82 Elewaut, D., Shaikh, R. B., Hammond, K. J., DeWinter, H., Leishman, A. J., Sidobre, S., Turovskaya, O. et al., NIK-dependent RelB activation defines a unique signaling pathway for the development of V alpha 14i NKT cells. *J. Exp. Med.* 2003. **197**: 1623–1633.
- 83 Sivakumar, V., Hammond, K. J., Howells, N., Pfeffer, K. and Weih, F., Differential requirement for Rel/nuclear factor kappa B family members in natural killer T cell development. *J. Exp. Med.* 2003. **197**: 1613–1621.
- 84 Pitt, L. A., Hubert, F. X., Scott, H. S., Godfrey, D. I. and Berzins, S. P., NKT cell development in the absence of the autoimmune regulator gene (Aire). *Eur. J. Immunol.* 2008. **38**: 2689–2696.
- 85 Berzins, S. P., McNab, F. W., Jones, C. M., Smyth, M. J. and Godfrey, D. I., Long-term retention of mature NK1.1+ NKT cells in the thymus. *J. Immunol.* 2006. **176**: 4059–4065.
- 86 Drennan, M. B., Franki, A. S., Dewint, P., VanBeneden, K., Seeuws, S., vande Pavert, S. A., Reilly, E. C. et al., Cutting edge: the chemokine receptor CXCR3 retains invariant NK T cells in the thymus. *J. Immunol.* 2009. **183**: 2213–2216.
- 87 Xiong, N. and Raulet, D. H., Development and selection of gammadelta T cells. *Immunol. Rev.* 2007. **215**: 15–31.
- 88 Ito, K., Bonneville, M., Takagaki, Y., Nakanishi, N., Kanagawa, O., Krecko, E. G. and Tonegawa, S., Different gamma delta T-cell receptors are expressed on thymocytes at different stages of development. *Proc. Natl. Acad. Sci. USA* 1989. **86**: 631–635.
- 89 Marks, B. R., Nowyhed, H. N., Choi, J. Y., Poholek, A. C., Odegard, J. M., Flavell, R. A. and Craft, J., Thymic self-reactivity selects natural interleukin 17-producing T cells that can regulate peripheral inflammation. *Nat. Immunol.* 2009. **10**: 1125–1132.
- 90 Prinz, I., Silva-Santos, B. and Pennington, D. J., Functional development of gammadelta T cells. *Eur. J. Immunol.* 2013. **43**: 1988–1994.
- 91 Silva-Santos, B., Pennington, D. J. and Hayday, A. C., Lymphotoxin-mediated regulation of gammadelta cell differentiation by alphabeta T cell progenitors. *Science* 2005. **307**: 925–928.
- 92 Roberts, N. A., White, A. J., Jenkinson, W. E., Turchinovich, G., Nakamura, K., Withers, D. R., McConnell, F. M. et al., Rank signaling links the development of invariant gammadelta T cell progenitors and Aire(+) medullary epithelium. *Immunity* 2012. **36**: 427–437.
- 93 Lewis, J. M., Girardi, M., Roberts, S. J., Barbee, S. D., Hayday, A. C. and Tigelaar, R. E., Selection of the cutaneous intraepithelial gammadelta+ T cell repertoire by a thymic stromal determinant. *Nat. Immunol.* 2006. **7**: 843–850.
- 94 Turchinovich, G. and Hayday, A. C., Skint-1 identifies a common molecular mechanism for the development of interferon-gamma-secreting versus interleukin-17-secreting gammadelta T cells. *Immunity* 2011. **35**: 59–68.
- 95 Nakamura, K., White, A. J., Parnell, S. M., Lane, P. J., Jenkinson, E. J., Jenkinson, W. E. and Anderson, G., Differential requirement for CCR4 in the maintenance but not establishment of the invariant Vgamma5(+) dendritic epidermal T-cell pool. *PLoS One* 2013. **8**: e74019.
- 96 Farr, A., Hosier, S., Nelson, A., Itoharu, S. and Tonegawa, S., Distribution of thymocytes expressing gamma delta receptors in the murine thymus during development. *J. Immunol.* 1990. **144**: 492–498.
- 97 Reinhardt, A., Ravens, S., Fleige, H., Haas, J. D., Oberdorfer, L., Lyszkiewicz, M., Forster, R. and Prinz, I., CCR7-mediated migration in the thymus controls gammadelta T-cell development. *Eur. J. Immunol.* 2014. **44**: 1320–1329.
- 98 Xiong, N., Kang, C. and Raulet, D. H., Positive selection of dendritic epidermal gammadelta T cell precursors in the fetal thymus determines expression of skin-homing receptors. *Immunity* 2004. **21**: 121–131.
- 99 Jin, Y., Xia, M., Sun, A., Saylor, C. M. and Xiong, N., CCR10 is important for the development of skin-specific gammadelta T cells by regulating their migration and location. *J. Immunol.* 2010. **185**: 5723–5731.
- 100 Bunting, M. D., Comerford, I., Kara, E. E., Korner, H. and McColl, S. R., CCR6 supports migration and differentiation of a subset of DN1 early thymocyte progenitors but is not required for thymic nTreg development. *Immunol. Cell Biol.* 2014. **92**: 489–498.
- 101 Hu, S. and Xiong, N., Programmed downregulation of CCR6 is important for establishment of epidermal gammadelta T cells by regulating their thymic egress and epidermal location. *J. Immunol.* 2013. **190**: 3267–3275.
- 102 Stritesky, G. L., Jameson, S. C. and Hogquist, K. A., Selection of self-reactive T cells in the thymus. *Annu. Rev. Immunol.* 2012. **30**: 95–114.

**Abbreviations:** mTEC: medullary thymic epithelial cell · SP: single positive

**Full correspondence:** Prof. Graham Anderson, MRC Centre for Immune Regulation, Institute for Biomedical Research, Medical School, Vincent Drive, Edgbaston, Birmingham B15 2TT, UK  
 Fax: +44-121-414-6815  
 e-mail: g.anderson@bham.ac.uk

Received: 31/10/2014

Revised: 16/1/2015

Accepted: 19/1/2015

Accepted article online: 23/1/2015