

# UNIVERSITY OF BIRMINGHAM

## Research at Birmingham

### Secondary structure and $^1\text{H}$ , $^{13}\text{C}$ , $^{15}\text{N}$ resonance assignments of the endosomal sorting protein sorting nexin 3

Overduin, Michael; Rajesh, Sandya; Gruenberg, Jean; Lenoir, Marc

DOI:

[10.1007/s12104-015-9609-z](https://doi.org/10.1007/s12104-015-9609-z)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Overduin, M, Rajesh, S, Gruenberg, J & Lenoir, M 2015, 'Secondary structure and  $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^{15}\text{N}$  resonance assignments of the endosomal sorting protein sorting nexin 3', *Biomolecular NMR Assignments*.  
<https://doi.org/10.1007/s12104-015-9609-z>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

Eligibility for repository : checked 23/06/2015

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Secondary structure and $^1\text{H}$ , $^{13}\text{C}$ , $^{15}\text{N}$ resonance assignments of the endosomal sorting protein sorting nexin 3

Michael Overduin<sup>1</sup> · Sandya Rajesh<sup>1</sup> · Jean Gruenberg<sup>2</sup> · Marc Lenoir<sup>1</sup>

Received: 6 January 2015 / Accepted: 11 April 2015

© The Author(s) 2015. This article is published with open access at Springerlink.com

**Abstract** Sorting nexin 3 (SNX3) belongs to a sub-family of sorting nexins that primarily contain a single Phox homology domain capable of binding phosphoinositides and membranes. We report the complete  $^1\text{H}$ ,  $^{13}\text{C}$  and  $^{15}\text{N}$  resonance assignments of the full-length human SNX3 protein and identification of its secondary structure elements, revealing a canonical fold and unstructured termini.

**Keywords** SNX3 · PX domain · Phosphoinositide recognition · Phosphatidylinositol 3-phosphate · Endosome · NMR

## Biological context

The SNX3 protein is localized within the early endosome through its interaction with phosphatidylinositol 3-phosphate (PI3P) (Xu et al. 2001). It associates with the cargo-selective retromer complex VPS26-VPS29-VPS35 in a trafficking pathway, thus forming a retromer capable of transporting Wntless from the endosome to the trans-Golgi network (Harterink et al. 2011). SNX3 has also been implicated in the formation of the intra-endosomal vesicles that are found in multivesicular bodies (Pons et al. 2008) and in the maturation of the Salmonella-induced vacuole in infected cells (Braun et al. 2010). Together with the SNX10, SNX11, SNX12,

SNX22, SNX23 and SNX24 proteins, SNX3 belong to a sub-family of sorting nexins that contain a single PX domain and apparently unstructured termini (Cullen and Korswagen 2012). PX domain-containing proteins are well known for their phosphatidylinositide binding activities, and are typically 120 amino acids in length. Their structural domains typically contain three antiparallel  $\beta$  strands followed by an unstructured loop with a conserved PXXP motif followed by three  $\alpha$  helices. The structures, dynamics and interactions of full-length sorting nexins are less well understood.

Here, we have assigned the resonances of the wild-type human SNX3 protein to identify its secondary structure and disordered regions, revealing a classical topology with an unstructured loop at the vicinity of the PI3P binding site. The assignment of the resonances provides a basis for the structural and dynamic analysis of this full length protein and its ligand and membrane binding mechanisms.

## Methods and experimental


### Expression and purification of SNX3

A pET45b (Merck) vector including the entire *SNX3* sequence encoding residues 2–162 was expressed in *E.coli* strain BL21(DE3). Cells were grown in M9 media supplemented by  $^{15}\text{NH}_4\text{Cl}$  and  $^{13}\text{C}$ -glucose. Expression was induced by addition of 1 mM IPTG until the  $\text{OD}_{600}$  reached 0.6. The cells were harvested by centrifugation at 6000g for 20 min and resuspended in 20 mM TrisHCl buffer pH 7.5, 100 mM NaCl, 1 mM DTT, 20 mM imidazole and Complete EDTA-free protease inhibitors (Roche). The cells were lysed with an Emulsiflex (17,000 psi) and the lysate was centrifuged at 75,000g for 45 min. The protein was bound to a HisTrap FF column (GE Healthcare) and eluted by an imidazole gradient

✉ Marc Lenoir  
m.lenoir@bham.ac.uk; lenoirM@bham.ac.uk


<sup>1</sup> Henry Wellcome Building for Biomolecular NMR Spectroscopy, School of Cancer Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

<sup>2</sup> Department of Biochemistry Sciences II, University of Geneva, 30, Quai Ernest-Ansermet, 1211 Geneva 4, Switzerland



**Fig. 1**  $^1\text{H}$ - $^{15}\text{N}$  HSQC of SNX3 (700  $\mu\text{M}$ ) in 20 mM sodium phosphate pH 6.5, 100 mM NaCl, 1 mM  $\text{NaN}_3$  and 10 % (v/v)  $\text{D}_2\text{O}$  collected at 298 K on a Varian INOVA 800 MHz spectrometer.

Residue numbers are indicated for cross peaks corresponding to backbone amides. Aliased residue peaks are indicated in blue and alternate conformer peaks in red



**Fig. 2** Secondary structure of SNX3 predicted by CCPNMR analysis. The arrows and bars indicate the positions of  $\beta$  strands and  $\alpha$  helices, respectively

(20–500 mM). Fractions containing SNX3 were further purified on a Superdex S75 HiLoad column connected to an ÄKTA Purifier (G.E. Healthcare) in a 20 mM sodium phosphate buffer containing 100 mM NaCl, DTT, and 1 mM  $\text{NaN}_3$ . NMR samples were prepared with 10 %  $\text{D}_2\text{O}$  (v/v).

### NMR spectroscopy

NMR spectra were acquired at 298 K on a Varian Inova 800 spectrometer equipped with a triple resonance  $^1\text{H}/^{13}\text{C}/^{15}\text{N}$  cryogenic probe and z-axis pulsed field gradients with a uniformly  $^{13}\text{C}/^{15}\text{N}$  labeled sample of 800  $\mu\text{M}$  SNX3 protein. The backbone assignment was obtained using Biopack pulse sequences (Varian) to collect HNC(O), HN(CA)CO, HN(CO)CA, HNCA, CBCA(CO)NH, HNCACB, H(C)CH-TOCSY, and (H)CCH-TOCSY spectra. The  $^{15}\text{N}$ -edited NOESY-HSQC and  $^{13}\text{C}$ -edited NOESY-HSQC experiment were acquired with mixing times of 100 ms. Spectra were processed using Nmrpipe (Delaglio et al. 1995) and analyzed with the Ccpnmr suite (Vranken et al. 2005). Proton chemical shifts were directly referenced against external 4,4-dimethyl-4-silapentane-1-sulfonic acid (DSS) standard while the  $^{15}\text{N}$  and  $^{13}\text{C}$  chemical shifts were referenced indirectly from the gyromagnetic ratios (Wishart et al. 1995).

### Assignments and data deposition

The  $^1\text{H}$ - $^{15}\text{N}$  HSQC spectrum of the full length SNX3 shows the assignment of essentially all the resonances (Fig. 1). The backbone assignment was complete except for the

N-terminal  $\text{His}_6$  tag. Beyond the affinity tag, most of the backbone HN (99 %), non-proline N (99 %),  $\text{C}\alpha$  (100 %),  $\text{C}\beta$  (100 %),  $\text{C}'$  (96 %) and  $\text{H}\alpha$  (100 %) resonances were assigned.

A second conformer was likely due to the isomerization of Pro102 and represented c.a. 30 % of the population based on the relative intensities of the peaks. The second set of resonances was unambiguously assigned for residues Leu101, Phe103, Gly105, Asp107, Ile109, Asp111, Phe114 and Arg118 situated within a predicted loop between  $\alpha 1$  and  $\alpha 2$  helices and at the start of the  $\alpha 2$  helix while the weaker resonances of the Arg104, Asp106, Gly108, Phe110, Asp112 and Asn113 residues were overlapped or obscured.

The secondary structure of SNX3 was predicted from  $\text{H}\alpha$ ,  $\text{C}\alpha$ ,  $\text{C}\beta$ ,  $\text{C}'$  chemical shifts using Ccpnmr analysis suite and the corresponding chemical shift index (Fig. 2). Three  $\beta$ -strands (29–39, 46–55, 64–69) followed by three  $\alpha$ -helices (71–84, 112–130, 133–145) were predicted, consistent with the structural characteristics of PX domains and SNX12-PX in particular (pdb 2CSK). The third helix may contain an irregularity midway based on an interrupted CSI pattern, while both termini exhibit random coil resonances and are intrinsically disordered.

The assignments of the apo state have been deposited to the BioMagResBank with the accession number 25402.

**Acknowledgments** We thank Cancer Research UK and EU-PRISM for funding (M.O.) and gratefully acknowledge the HWB-NMR staff for spectrometer access and discussions, and the Wellcom Trust for support of the national NMR facility at the University of Birmingham.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Braun V, Wong A, Landekic M, Hong WJ, Grinstein S, Brumell JH (2010) Sorting nexin 3 (SNX3) is a component of a tubular endosomal network induced by Salmonella and involved in maturation of the Salmonella-containing vacuole. *Cell Microbiol* 12:1352–1367
- Cullen PJ, Korswagen HC (2012) Sorting nexins provide diversity for retromer-dependent trafficking events. *Nat Cell Biol* 14:29–37
- Delaglio F, Grzesiek S, Vuister GW, Zhu G, Pfeifer J, Bax A (1995) NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *J Biomol NMR* 6:277–293
- Harterink M, Port F, Lorenowicz MJ, McGough IJ, Silhankova M, Betist MC, van Weering JR, van Heesbeen RG, Middelkoop TC, Basler K, Cullen PJ, Korswagen HC (2011) A SNX3-dependent retromer pathway mediates retrograde transport of the Wnt sorting receptor Wntless and is required for Wnt secretion. *Nat Cell Biol* 13:914–923
- Pons V, Luyet PP, Morel E, Abrami L, van der Goot FG, Parton RG, Gruenberg J (2008) Hrs and SNX3 functions in sorting and membrane invagination within multivesicular bodies. *PLoS Biol* 6:e214
- Vranken WF, Boucher W, Stevens TJ, Fogh RH, Pajon A, Llinas M, Ulrich EL, Markley JL, Ionides J, Laue ED (2005) The CCPN data model for NMR spectroscopy: development of a software pipeline. *Proteins* 59:687–696
- Wishart DS, Bigam CG, Yao J, Abildgaard F, Dyson HJ, Oldfield E, Markley JL, Sykes BD (1995) <sup>1</sup>H, <sup>13</sup>C and <sup>15</sup>N chemical shift referencing in biomolecular NMR. *J Biomol NMR* 6:135–140
- Xu Y, Hortsman H, Seet L, Wong SH, Hong W (2001) SNX3 regulates endosomal function through its PX-domain-mediated interaction with PtdIns(3)P. *Nat Cell Biol* 3:658–666