

UNIVERSITY OF BIRMINGHAM

Research at Birmingham

Realistic activity propagation for mean field models of human cortex

Bojak, Ingo; Liley, David T J

DOI:

[10.1186/1471-2202-10-S1-P290](https://doi.org/10.1186/1471-2202-10-S1-P290)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Bojak, I & Liley, DTJ 2009, Realistic activity propagation for mean field models of human cortex. in BMC Neuroscience: Eighteenth Annual Computational Neuroscience Meeting: CNS*2009. vol. 10 (Suppl 1), P290, BMC Neuroscience , no. Suppl 1, vol. 10, Eighteenth Annual Computational Neuroscience Meeting, Berlin, Germany, 18/07/09. <https://doi.org/10.1186/1471-2202-10-S1-P290>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Poster presentation

Open Access

Realistic activity propagation for mean field models of human cortex

Ingo Bojak*¹ and David TJ Liley²

Address: ¹Donders Institute for Brain, Cognition and Behaviour, Centre for Neuroscience, Radboud University Nijmegen Medical Centre, P.O. Box 9101//126, 6500 HB Nijmegen, The Netherlands and ²Brain Sciences Institute (BSI), Swinburne University of Technology, P.O. Box 218, Victoria 3122, Australia

Email: Ingo Bojak* - i.bojak@donders.ru.nl

* Corresponding author

from Eighteenth Annual Computational Neuroscience Meeting: CNS*2009
Berlin, Germany. 18–23 July 2009

Published: 13 July 2009

BMC Neuroscience 2009, 10(Suppl 1):P290 doi:10.1186/1471-2202-10-S1-P290

This abstract is available from: <http://www.biomedcentral.com/1471-2202/10/S1/P290>

© 2009 Bojak and Liley; licensee BioMed Central Ltd.

Inspired by the great density of neurons in cortex (about 10^6 per macrocolumn), continuum mean field models (CMFMs) treat the cortex as one continuous neural medium. Interactions between neurons thus become flows of activity spreading in this medium. The most popular propagation model [1] is derived by a two-fold ansatz: a pulse of activity will on one hand spread isotropically with a conduction velocity c , and on the other hand its amplitude will decay exponentially with a distance scale σ . This ansatz reflects action potentials traveling with constant speed through axons and that the number of synapses axons form falls roughly exponentially with distance. However, a pulse in a CMFM means a mass of 10^5 to 10^6 neurons pulsing *together*. It is well known that axons can vary greatly in conduction speed, depending on their myelination and diameter. Hence it is natural to expect that such a mass has a broad conduction velocity distribution $f(v)$, rather than a singular one $f(v) = \delta(v-c)$. Furthermore, one still has to expand for large wavelengths in order to derive a manageable partial differential equation (PDE) – a damped wave equation as it turns out. If one calculates the velocity distribution of this approximation, one sees that the Dirac delta peak has softened only towards lower velocities, leaving an unnatural distribution with a hard velocity cut-off. We hence propose a new propagation PDE:

$$\left[\frac{\partial}{\partial t} + \frac{c}{2\sigma} (1 - \sigma^2 \bar{v}^2) \right]^n \Phi(\bar{x}, t) = \frac{N^\alpha c^n}{2^n \sigma^n} S(\bar{x}, t),$$

where Φ is the activity being propagated, S is a local source (e.g., a firing rate function), N^α the total number of connections, c is the conduction velocity and σ the decay parameter, and $n > 0$ will usually be chosen as an integer.

This new PDE has the following advantages:

For $n > 1$ the connectivity implied by the PDE remains finite for small distances.

The PDE has a smooth velocity distribution, the shape of which depends only on n and c .

The velocity distribution for $n = 3$ and $c = 14.35$ m/s fits human data [2] well. This value of c implies a mean conduction velocity of 7.87 m/s. The distribution for the damped wave equation fits this data very badly; see Figure 1.

There is no switch between finite wave number Hopf-Turing and Hopf bifurcations as is the case for the damped wave equation. Self-sustained oscillations can emerge as spatial patterns with arbitrarily small wavenumber.




Figure 1
Number of fibres faster than a velocity threshold.
 Aboitiz et al. data [2] compared to a fit with the new PDE (blue), and with the damped wave equation matching the new PDE's median speed (green) or fit to the first three data points (cyan, a fit to all points is here not feasible due to severe functional mismatch).

We have investigated the bifurcations of this propagator both analytically and numerically on large grids. We have also compared to more detailed data from the rat [3], but find that they do not scale to human in a simple manner. Our new propagation PDE provides now for CMFMs the best match to activity conduction in humans, and can be easily adjusted as more human data become available later.

References

1. Jirsa VK, Haken H: **Field theory of electromagnetic brain activity.** *Phys Rev Lett* 1996, **77**:960-963.
2. Aboitiz F, et al.: **Fiber composition of the human corpus callosum.** *Brain Res* 1992, **598**:143-153.
3. Partadiredja G, et al.: **The number, size, and type of axons in rat subcortical white matter on left and right sides: A stereological, ultrastructural study.** *J Neurocytol* 2003, **32**:1165-1179.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."
 Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp