

# UNIVERSITY OF BIRMINGHAM

## Research at Birmingham

### Remember, remember the 5th of November

Singh, Ajit; Bloss, William; Pope, Francis

DOI:

[10.1002/wea.2587](https://doi.org/10.1002/wea.2587)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Singh, A, Bloss, WJ & Pope, FD 2015, 'Remember, remember the 5th of November: gunpowder, particles and smog', *Weather*, vol. 70, no. 11, pp. 320-324. <https://doi.org/10.1002/wea.2587>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

Eligibility for repository: Checked on 18/12/2015

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

1 **Remember, remember the 5<sup>th</sup> of November; gunpowder, particles and smog**

2 Ajit Singh, William J. Bloss and Francis D. Pope\*

3 School of Geography, Earth and Environmental Sciences, University of Birmingham,  
4 Birmingham, B15 2TT, UK

5  
6 \*Corresponding author: [f.pope@bham.ac.uk](mailto:f.pope@bham.ac.uk)

7  
8  
9 **Abstract**

10 In the UK on 5 November, Guy Fawkes Night is celebrated with bonfires and  
11 fireworks. An undesirable consequence of these activities is a statistically significant  
12 reduction (~25%) in atmospheric visibility nationwide. This reduction is caused by  
13 increased loading of atmospheric particulate matter generated by bonfires and  
14 fireworks. The effect of this increased loading on visibility is investigated in greater  
15 detail for the city of Nottingham where larger visibility decreases compared to the  
16 national average are observed. Visibility reduction is more significant when the  
17 background particulate matter loading and/or the atmospheric relative humidity are  
18 high.

19  
20 **Introduction**

21 Fireworks are often used to celebrate festivals and special days, for example,  
22 Bastille Day, Independence Day, New Year's Eve, Diwali, Chinese New Year and  
23 large sporting events. Within the UK, the biggest fireworks event is Guy Fawkes  
24 Night which is celebrated around the 5 November every year. It commemorates the  
25 events of that day in the year 1605 when the Gunpowder Plot, involving the  
26 eponymous Mr Fawkes, failed to blow up the Houses of Parliament. Typically the  
27 celebrations involve both bonfires and fireworks (ground and air detonating). For  
28 public safety, the UK government has legislation surrounding the purchase and use

1 of fireworks during the celebration period which limits the dates on which fireworks  
2 displays can occur.

3 While fireworks and bonfires have tremendous and enduring appeal, some of their  
4 effects are not always beneficial. Incorrect handling and use of fireworks can lead to  
5 injury (Vernon, 1988) and allergy (Becker *et al.*, 2000). They can be distracting  
6 through various human senses (sight, sound and smell). They can generate dense  
7 clouds of smoke in the lower atmosphere (Drewnick *et al.*, 2006) and are sources of  
8 pollution including gas phase species (e.g. sulphur dioxide, ozone, etc.) and  
9 particulate phase species including black carbon and metals (Ravindra *et al.*, 2003;  
10 Seidel and Birnbaum, 2015), some of which have negative associated health effects  
11 (Ravindra *et al.*, 2001). Within the UK, several studies have linked the use of  
12 fireworks with changes in air quality (Knox *et al.*, 2008; Godri *et al.*, 2010). Of  
13 particular interest to this study is the effect of the smoke on visibility. Reductions in  
14 visibility can deleteriously affect public safety, transportation, and tourism (Singh and  
15 Day, 2012). Over the past few years several studies have connected fireworks to  
16 visibility issues (for example, Wang *et al.*, 2007; Vecchi *et al.*, 2008; Saha *et al.*,  
17 2014; Kong *et al.*, 2015). Local scale and short term pollution episodes can cause  
18 large decreases in visibility potentially with devastating consequences. For example,  
19 in 2011 a tragic incident occurred on the M5 motorway near Taunton, Somerset,  
20 where seven people died and 57 were injured due to a car crash. The resulting  
21 investigation found that a local fireworks display near the road might have  
22 contributed to the poor visibility on an already foggy night albeit no blame was  
23 assigned to the organizer of the fireworks event (Rose, 2014).

24 Particulate matter (PM) can scatter and absorb light, to an extent dependent upon its  
25 size and composition, and hence its introduction into the atmosphere (via fireworks,  
26 bonfires or any other source) can lead to reductions in visibility (Appel *et al.*, 1985).  
27 The ability of PM to scatter and absorb light is dependent upon the size and  
28 composition (refractive index) of the PM. Absorption increases with increasing PM  
29 mass. The amount of scattering for a given PM ensemble is much more finely tuned  
30 to the given particle size and composition distribution. In general, the closer the PM  
31 size matches with the wavelength of the light, the greater the scattering (Mie, 1908).

1 Nearly all atmospheric PM are hygroscopic to some degree, that is to say, their  
2 water content is dependent upon the local relative humidity. As relative humidity  
3 increases, so does the water content of the PM. This effect changes both the  
4 particle size and the average composition. For a given PM starting composition, its  
5 size will increase with increasing relative humidity, and its refractive index will be  
6 lowered because the refractive index of water is typically lower than the other  
7 common PM components (e.g. minerals, organics, sulphates, nitrates) (Harrison *et*  
8 *at.*, 2004).

9 Mathematically, visibility ( $V$ ) can be represented as a function of the extinction  
10 coefficient ( $\beta_{ext}$ ) via equation 1, where the constant  $k$  is equal to 3.912 which  
11 assumes a visibility contrast threshold of 2 % (Koschmeider, 1924).

12

$$13 \quad V = k/\beta_{ext} \quad (1)$$

14

15 The extinction coefficient is dependent upon both the scattering (*sca*) and absorption  
16 (*abs*) due to PM and gases ( $\beta_{ext} = \beta_{gas,sca} + \beta_{gas,abs} + \beta_{PM,sca} + \beta_{PM,abs}$ ). Typically the  
17 contribution of PM to the extinction coefficient far outweighs the contribution of gases  
18 with the one possible exception being nitrogen dioxide (NO<sub>2</sub>) which has a significant  
19 visible absorption coefficient (Groblicki *et al.*, 1981) and may be present with  
20 appreciable abundance. For PM with a given size distribution and composition the  
21 extinction coefficient is linearly dependent upon the PM number concentration,  
22 hence, visibility can be used as a metric of atmospheric pollutant levels.

23 In this paper we provide evidence for a significant reduction in visibility in the UK on  
24 the 5 November which is largely caused by the PM generated from the combination  
25 of fireworks and bonfires.

26

## 27 **Data Collection and Methodology**

28 Two distinct data series were generated for this paper: 1. UK average visibility maps  
29 for the dates of the 2 - 8 November inclusive, and 2. Average visibility time series for

1 Nottingham in the months of October and November. Due to limited availability of  
2 visibility stations which co-located with pollution sensors and limited data-sets,  
3 Nottingham has been selected. The visibility maps allow for the detection of regional  
4 changes in visibility associated with Guy Fawkes Night, while the longer time series  
5 of the Nottingham data allows for the attribution of the change in visibility on Guy  
6 Fawkes Night to the change in atmospheric composition.

7 The hourly values of horizontal visibility (maximum visible distance along a horizontal  
8 line at the earth's surface) and relative humidity data were obtained from the British  
9 Atmospheric Data Centre (BADC), ([www.badc.nerc.ac.uk](http://www.badc.nerc.ac.uk)), for 34 stations situated  
10 throughout the United Kingdom. For the locations of the stations see Figure 1.

11 Daily mean pollutant data were obtained for a city centre location in Nottingham. In  
12 particular, data for the gas phase species nitrogen dioxide (NO<sub>2</sub>) and sulphur dioxide  
13 (SO<sub>2</sub>) were obtained in addition to three PM metrics: PM with diameter less than 10  
14 µm (PM<sub>10</sub>), PM with diameter less than 2.5 µm (PM<sub>2.5</sub>), and black carbon (BC) PM.  
15 These data were obtained from the DEFRA Automatic Urban and Rural Monitoring  
16 Network (AURN) (<http://uk-air.defra.gov.uk/>).

17 Both the meteorological and pollutant data sets were obtained for a 13 year period  
18 encompassing the years 2000-2012. The length of the time period investigated was  
19 constrained by the availability of the AURN data. The PM<sub>2.5</sub> data are only available  
20 for the period 2009-2012. As official fireworks displays typically start after 1900 h,  
21 we selected visibility and humidity data-set for 2100 h for study. Whilst many  
22 fireworks displays occur on the 5 November not all do; often official displays will be  
23 performed on the weekend preceding or subsequent to the 5. Within our 13 year  
24 data set, the number of days that November the 5 fell on a given day of the week  
25 was: Monday (3), Tuesday (1), Wednesday (2), Thursday (1), Friday (2), Saturday  
26 (2) and Sunday (2).

27 The available visibility data network is not sufficiently dense to allow for the visibility  
28 mapping of the whole of the UK. The maps were produced by assuming that the  
29 measurement at each station was reliable until the distance exceeded 32 km, which  
30 represents a typical upper limit to visibility for the UK measured in November (albeit  
31 not for Guy Fawkes Night). Where two measurements overlap, the visibility is  
32 approximated by inverse distance weighting interpolation of the multiple

1 measurements. This approach necessarily assumes that visibility is homogenous  
2 within the area defined by the visibility measurement; in reality, visibility is affected  
3 by local orography and other effects and can be highly heterogeneous. Data  
4 manipulation was carried out using R software, where ggplot tool has been used for  
5 spatial mapping. The meteorology and pollution data were analysed using Matlab  
6 (2014a) software.

7

## 8 **Results and Discussion**

9 The change in the UK visibility over Guy Fawkes Night and the surrounding days is  
10 shown in Figure 1. The spatial trend analysis is presented using 34 stations with  
11  $0.29^\circ \times 0.29^\circ$  distance resolution. This is a latitude equivalent to 32 km and a  
12 longitude equivalent to 20.3 km. 32 km represents a typical upper limit value for  
13 visibility in the UK in November. In Figure 1, a dramatic dip in visibility can clearly be  
14 seen for November 5 compared with preceding and subsequent days. The average  
15 visibilities observed at the 34 stations on the 2, 3, 4, 5, 6, 7, and 8 November are  
16  $20.0 \pm 4.6$ ,  $17.8 \pm 4.0$ ,  $17.6 \pm 5.1$ ,  $14.8 \pm 5.2$ ,  $16.9 \pm 4.4$ ,  $18.3 \pm 3.8$  and  $20.5 \pm 4.1$  km  
17 respectively, with a mean reduction of ~25% on 5 November compared with other  
18 days (2, 3, 4, 6, 7, and 8 November). It can be seen that the 6 November has the  
19 next lowest visibility after the 5 which may be due to higher atmospheric PM loading  
20 persisting following Guy Fawkes on the preceding night. The effect of the fireworks  
21 on visibility is observed at all sites. Typically, urban locations show the greatest  
22 visibility loss. 82% of the UK population lives within urban locations  
23 (<http://data.worldbank.org>) and hence poor visibility in these regions has greater  
24 potential for disruption.

25 To better understand the change in visibility caused by Guy Fawkes Night, we  
26 investigate the 13 year average 2100 h visibility values for a single site (Nottingham-  
27 Watnall, SYNOP code number = 3354, station source ID = 556) for the months of  
28 October and November. Again there is a clear dip in visibility occurring on the 5  
29 November showing the effect of Guy Fawkes Night (a reduction of more than 55%  
30 compares to the 2, 3, 4, 6, 7, and 8 November). Investigation of the 24 hourly data  
31 over Nottingham also indicates that visibility starts decreasing from 1900 h on the 5

1 November and is stable after 0100 h on the 6 November, due to changes in the  
2 loading of firework-derived PM.

3 This Guy Fawkes driven dip in visibility becomes even more pronounced when the  
4 effect of relative humidity is taken into account. The daily relative humidity  
5 (measured at 2100 h) shows the expected seasonal increase as the date progresses  
6 from the start of October to the end of November, see Figure 2 panel (a). The  
7 relative humidity increases from ~85 to 90 % over the course of the two months, with  
8 significant variability. Whilst this represents only a modest increase in relative  
9 humidity, such a change of 5 % in this region of a PM hygroscopicity growth curve  
10 can be significant (e.g. Pope *et al.*, 2010). As discussed in the introduction, as  
11 relative humidity increases so does particle water content with the corresponding  
12 changes in size, composition, refractive index and hence extinction coefficient.  
13 Since the composition and size distribution of the PM is unknown, the effect of  
14 relative humidity cannot be explicitly calculated but a simple correlation curve  
15 between visibility and relative humidity can be generated. The relationship between  
16 visibility and relative humidity (RH) can be reasonably approximated using a linear  
17 relationship,  $V = m \times RH + c$ , where the constants  $m$  and  $c$  are best described by -  
18 1.19 % and 119.42 km, respectively, with a  $R^2$  value of 0.60. This approach  
19 assumes that the non-water component of the PM loading, composition and size  
20 stays constant over the time period analysed, clearly this necessitates the exclusion  
21 of the data from Guy Fawkes Night.

22 Using the relationship between visibility and relative humidity, the visibility  
23 anticipated for the 5 November in the absence of Guy Fawkes Night can be  
24 predicted, see Figure 2 panel (b). It is discernible that the predicted visibility shows  
25 reasonable agreement with the observed visibility except on the 5 Nov. Panel (c) in  
26 Figure 2 shows the difference between observed visibility and predicted visibility and  
27 the effect of Guy Fawkes Night on visibility is even more apparent. This analysis  
28 suggests that Guy Fawkes Night in Nottingham reduces visibility by approximately  
29 10 km which corresponds to a 64% reduction in visibility. This reduction in visibility  
30 is highly significant as it lies below four standard deviations ( $\pm 8.88$  km) of the  
31 average visibility, see figure 2 panel (c).

1 The change in visibility due to Guy Fawkes Night is compared with atmospheric  
2 pollutant concentrations in Figure 3. There are clear spikes in the PM<sub>2.5</sub>, PM<sub>10</sub> and  
3 black carbon PM matter on Guy Fawkes Night which suggest that these pollutants  
4 are, at least in part, responsible for the reduction in visibility for this night. PM<sub>2.5</sub>, PM<sub>10</sub>  
5 and black carbon PM concentrations were ~253%, ~62% and ~201% greater,  
6 respectively. It is emphasized again that the PM<sub>2.5</sub> data set only contains 4 years of  
7 data whereas the PM<sub>10</sub>, black carbon and visibility data sets encompass 13 years.  
8 The corresponding reduction in visibility for the time period of the PM<sub>2.5</sub>  
9 measurements is 63%. The PM<sub>2.5</sub> data are noisier compared with the other data  
10 streams. This is due to shorter measurement period which introduces more  
11 statistical noise in addition to a greater 'day of the week' effect on the average data.  
12 Over longer averaging periods, the likelihood of the 5 November falling preferentially  
13 on specific days of the week becomes less likely. The data suggest that the average  
14 concentration of PM<sub>2.5</sub> was higher on both 5 (31.25 µg m<sup>-3</sup>) and 6 November (38.75  
15 µg m<sup>-3</sup>) compared with rest of the investigated dates, see Figure 3, panel d. In 2010  
16 there was an exceptionally large signal on the 6 November (Saturday) which  
17 explains the Guy Fawkes Night PM<sub>2.5</sub> peak occurring on both the 5 and 6 November.

18 The influence of NO<sub>2</sub> on visibility was also investigated. The NO<sub>2</sub> concentration did  
19 not show any significant change over the Guy Fawkes Night period. Hence none of  
20 the visibility loss can be ascribed to gas phase absorption of light. Likewise gas  
21 phase scattering will not be affected by Guy Fawkes Night. Therefore the change in  
22 the extinction coefficient over the celebration period must be due to changes in PM  
23 absorption and scattering. This is expected because the contribution of gases to the  
24 extinction coefficient is typically negligible except in pristine conditions or conditions  
25 with exceptionally high NO<sub>2</sub>.

26 The most significant component of PM that absorbs light is black carbon. The  
27 incremental loading of black carbon on Guy Fawkes night can be directly converted  
28 into an PM absorption coefficient ( $\beta_{PM,abs}$ ) through use of the recommended value of  
29 the mass normalized absorption coefficient ( $\sigma$ ), 7.5 m<sup>2</sup> g<sup>-1</sup> (Bond and Bergstrom,  
30 2006) and equation 2, where BC is the black carbon mass concentration (g m<sup>-3</sup>).

31

32

$$\beta_{PM,abs} = \sigma \times BC \quad (2)$$



1

2 The average observed visibility for Guy Fawkes Night (2000-2012) is 5.1 km and the  
3 predicted visibility in the absence of celebrations is 14.8 km which correspond to  
4 extinction coefficients of 0.77 and 0.26 km<sup>-1</sup>, respectively. The average black carbon  
5 concentration on Guy Fawkes Night was 6.04µgm<sup>-3</sup>. The average non Guy Fawkes  
6 Night background black carbon concentration was 2.00µgm<sup>-3</sup>. The corresponding  
7 average PM absorption coefficients for Guy Fawkes Night and non-Guy Fawkes  
8 Night period are 0.045 and 0.015 km<sup>-1</sup>, respectively. Hence the PM absorption  
9 coefficient only accounts for 5.9 and 5.8 % of the total extinction coefficient for the  
10 two different time periods. Even though the absolute loadings are very different on  
11 Guy Fawkes Night compared with the normal background conditions, the percentage  
12 contribution of black carbon to total visibility reduction is largely independent of Guy  
13 Fawkes Night which is surprising.

14

15 In contrast to the PM absorption coefficient, it is more difficult to estimate the  
16 enhancement in the PM scattering coefficient ( $\beta_{PM,sca}$ ) due to Guy Fawkes Night,  
17 because key parameters are unknown, including the distribution in the particle size  
18 and composition and how these differ from the background PM. Nevertheless the  
19 increment in PM<sub>2.5</sub> loading over the Guy Fawkes Night period can be used to  
20 estimate the effect if several assumptions are made: 1) the distributions of aerosol  
21 size and composition are the same as the background PM; 2) the only difference  
22 between the PM distributions is the number concentration of PM, i.e. Guy Fawkes  
23 Night introduces an additional source of PM which has identical properties to the  
24 background aerosol; 3) PM<sub>2.5</sub> is a good proxy for light scattering PM which are  
25 largely sub-micron in size; 4) the component of the extinction coefficient due to gas  
26 phase species is negligible. With these assumptions the scattering coefficient can  
27 be calculated through knowledge of the extinction and absorption coefficients using  
28 equation 3.

29

$$\beta_{PM,sca} = \beta_{ext} - \beta_{PM,abs} \quad (3)$$

31

1 The validity of these assumptions can be tested by comparing the scaling constant  
2 (k) required to convert the PM<sub>2.5</sub> concentration into a scattering coefficient, i.e.  
3  $\beta_{PM,sca} = \delta \times PM_{2.5}$ . The calculated scattering coefficients for Guy Fawkes Night and  
4 non-Guy Fawkes Night period are 0.67 and 0.25 km<sup>-1</sup>. The corresponding values for  
5  $\delta$  are 0.021 and 0.020 m<sup>3</sup>  $\mu\text{g}^{-1}$  km<sup>-1</sup>. Given the assumptions used in deriving the  
6 scattering coefficients, the two derived values of  $\delta$  are in remarkably good  
7 agreement. The contribution of the scattering coefficient to the total extinction  
8 coefficient for Guy Fawkes Night and non-Guy Fawkes Night is 94.4 and 96.1 %.

9 In addition to visibility altering pollutants, the gas phase concentration of SO<sub>2</sub> was  
10 also investigated. No significant changes in SO<sub>2</sub> were observed over the celebration  
11 period. Therefore for the site investigated, SO<sub>2</sub> is not a good chemical marker for  
12 Guy Fawkes Night. This is surprising since enhanced SO<sub>2</sub> has previously been  
13 associated with fireworks (Ravindra *et al.*, 2003).

14

## 15 **Conclusions**

16 In the present study, the role and influence of firework and bonfire emissions on  
17 visibility and short term air quality has been quantified. Although Guy Fawkes Night  
18 only occurs once a year, the associated celebrations occur over a wider time range  
19 (on the order of a week). Using data from 34 meteorological stations, sharp  
20 reductions in nationwide visibility were clearly observed on Guy Fawkes Night.  
21 However, the effect is short-lived, with average visibility returning to normal within  
22 two days due to the dispersal of the additional PM loading. More detailed analysis on  
23 the data from a single urban area (Nottingham) showed that the effect of Guy  
24 Fawkes Night is even more pronounced when the effect of relative humidity, through  
25 its influence on the extinction coefficient of PM, is taken into account.

26 The reduction in visibility on Guy Fawkes Night is caused by increases in  
27 atmospheric PM loading which is generated through bonfires and fireworks. In  
28 particular there is clear anti-correlation between the PM<sub>2.5</sub>, PM<sub>10</sub> and black carbon  
29 PM with the observed visibility. It is interesting to note that as the UK becomes ever  
30 more multicultural, we might expect the frequency and magnitude of

1 festivals/celebrations that use fireworks to increase with corresponding visibility  
2 reduction hotspots.

3 The data and analysis shown above, indicates that the public should be made aware  
4 of the possibility of low visibility on Guy Fawkes Night which can be very localized.  
5 In particular, care should be taken when relative humidity is predicted to be high and  
6 the atmosphere already has a high PM loadings. Since fireworks displays are  
7 planned months in advance, weather and pollution forecasts of sufficient skill are  
8 unavailable to help in their planning. However, if forecasts subsequently suggest  
9 that the planned display will coincide with conditions likely to exacerbate poor  
10 visibility then the organizers and local authorities should be prepared to issue poor  
11 visibility warnings in advance. This precautionary step can help to increase  
12 enjoyment, or at least prevent unnecessary accidents, during Guy Fawkes Night  
13 which is a much loved and highly anticipated celebration.

14

15

## 16 **Acknowledgements**

17 A.S. Thanks for the University of Birmingham for funding through the Elite Scholar  
18 scheme. Nick Davidson is thanked for providing proof reading. We are also very  
19 thankful to both reviewers for their valuable comments and suggestions.

20

## 21 **References**

22 **Appel BR, Tokiwa Y, Hsu J, Kothny EL, Hahn E.** 1985. Visibility as related to  
23 atmospheric aerosol constituents. *Atmos. Environ.* (1967), **19(9)**: 1525-1534.

24 **Becker JM, Iskandrian S, Conkling J.** 2000. Fatal and near-fatal asthma in children  
25 exposed to fireworks. *Ann. All. Asth. Immu.*, **85(6)**: 512-513.

26 **Bond TC, Bergstrom RW,** 2006. Light absorption by carbonaceous particles: An  
27 investigative review. *Aero. Sci. tech.*, **40(1)**: 27-67.

- 1 **Drewnick F, Hings SS, Curtius J, Eerdekens G, Williams J.** 2006. Measurement  
2 of fine particulate and gas-phase species during the New Year's fireworks 2005 in  
3 Mainz, Germany. *Atmos. Environ.*, **40(23)**: 4316-4327.
- 4 **Godri KJ et al.** 2010. Particulate oxidative burden associated with firework  
5 activity. *Environ. Sci. & Tech.*, **44(21)**: 8295-8301.
- 6 **Groblicki PJ, Wolff GT, Countess RJ.** 1981. Visibility-reducing species in the  
7 Denver "brown cloud"—I. Relationships between extinction and chemical  
8 composition. *Atmos. Environ.*, **15(12)**: 2473-2484.
- 9 **Harrison RM, Jones AM, Lawrence RG.** 2004. Major component composition of  
10 PM 10 and PM 2.5 from roadside and urban background sites. *Atmos.*  
11 *Environ.*, **38(27)**: 4531-4538.
- 12 **Knox FA et al.** 2008. A British Ophthalmological Surveillance Unit study on serious  
13 ocular injuries from fireworks in the UK. *Eye.*, **22(7)**: 944-947.
- 14 **Koschmieder H.** 1924. Theorie der horizontalen Sichtweite. *Beitr. z. Phys. Freien.*  
15 *Atm.*, **12**: 33-53
- 16 **Mie G.** 1908. Beiträge zur Optik trüber Medien, speziell kolloidaler  
17 Metallösungen. *Ann. Der. Phys.*, **330(3)**: 377-445.
- 18 **Kong SF et al.** 2015. The impacts of firework burning at the Chinese Spring Festival  
19 on air quality: insights of tracers, source evolution and aging processes. *Atmos.*  
20 *Chem. and Phys.*, **15(4)**: 2167-2184.
- 21 **Pope FD, Dennis-Smith BJ, Griffiths PT, Clegg SL, Cox RA.** 2010. Studies of  
22 single aerosol particles containing malonic acid, glutaric acid, and their mixtures with  
23 sodium chloride. I. Hygroscopic growth. *J. Phys. Chem. A*, **114(16)**: 5335-5341.
- 24 **Ravindra K, Mittal, AK, Van GR.** 2001. Health risk assessment of urban suspended  
25 particulate matter with special reference to polycyclic aromatic hydrocarbons: a  
26 review. *Rev. Environ. Healt.*, **16(3)**: 169-189.
- 27 **Ravindra K, Mor S, Kaushik CP.** 2003. Short-term variation in air quality associated  
28 with firework events: A case study. *J. Environ, Monit.*, **5(2)**: 260-264.

1 **Rose M.** 2014. M5(Seven). *Report on Prevent future deaths (1) in category of Road*  
2 *(Highways Safety) related deaths, Ref: 2014-0654*

3 **Saha U, Talukdar S, Jana S, Maitra A.** 2014. Effects of air pollution on  
4 meteorological parameters during Deepawali festival over an Indian urban  
5 metropolis. *Atmos. Environ.*, **98**: 530-539.

6 **Seidel DJ, Birnbaum AN.** 2015. Effects of Independence Day Fireworks on  
7 Atmospheric Concentrations of Fine Particulate Matter in the United States. *Atmos.*  
8 *Enviro.*, **115**: 192-198.

9 **Singh A, Dey S.** 2012. Influence of aerosol composition on visibility in megacity  
10 Delhi. *Atmos. Environ.*, **62**: 367-373.

11 **Vecchi R et al.** 2008. The impact of fireworks on airborne particles. *Atmos.*  
12 *Environ.*, **42(6)**: 1121-1132.

13 **Vernon SA.** 1988. Fireworks and the eye. *J. R. Soc. Medic.*, **81(10)**: 569-571.

14 **Wang Y, Zhuang G, Xu C, An Z.** 2007. The air pollution caused by the burning of  
15 fireworks during the lantern festival in Beijing. *Atmos. Environ.*, **41(2)**: 417-431.

16

17

18

19

20

21

22

23

24

25

26


1

2

3


1

## Figures




2

3 **Figure 1** Visibility mapping of the UK before (2, 3, and 4 Nov), during (5 Nov), and  
4 after (6, 7 and 8 Nov) the Guy Fawkes Night using 13 years (2000-2012) data. The  
5 filled circles in the UK map represent the available meteorological stations (=34)  
6 measuring visibility. The green coloured circle represents the Watnall station  
7 (Nottingham). The meteorological stations (=20) used for the spatial mapping are  
8 indicated within the black box. The scale bar provides the visibility range.





1 **Figure 2** Dependency of relative humidity on visibility with 13 years (2000-1012)  
2 data-set, where panel **a)** Average relative humidity **b)** Comparison of observed  
3 ( $VIS_O$ ) and predicted visibility ( $VIS_P$ ) **and c)** Difference between observed and  
4 predicted visibility, where grey shading strata represent standard deviations ( $\sigma$ ) from  
5 the mean. Lightest grey =  $1\sigma$  and darkest grey =  $4\sigma$ .



1 **Figure 3** Comparison of pollutant concentrations with visibility measurement  
2 (predicted – observed). The visibility, PM<sub>10</sub> and black carbon data sets represent 13  
3 year averages (2000-2012), whereas the PM<sub>2.5</sub> data set represents a 4 year average  
4 (2009-2012).

5

6

7

1 **Definitions of technical terms**

2

3 **Absorption coefficient** – Parameter expressing the ability of PM to absorb  
4 radiation.

5 **Contrast threshold-** Smallest difference that the eye can discern between two  
6 different stimuli.

7 **Extinction coefficient** – Sum of the light scattering and absorption coefficient.

8 **Inverse distance weighting interpolation** - A mathematical technique used to  
9 estimate the visibility at locations laying between two different measurement  
10 locations.

11 **Mass normalized absorption coefficient** – Ratio of absorption coefficient to mass  
12 concentration of the absorbing material.

13 **Scattering coefficient** – Parameter expressing the ability of PM to scatter radiation.

14