

UNIVERSITY OF BIRMINGHAM

Research at Birmingham

The Papaver rhoeas S determinants confer self-incompatibility to Arabidopsis thaliana in planta

Lin, Zongcheng; Eaves, Deborah J; Sanchez-Moran, Eugenio; Franklin, Chris ; Franklin-Tong, Veronica

DOI:

[10.1126/science.aad2983](https://doi.org/10.1126/science.aad2983)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Lin, Z, Eaves, DJ, Sanchez-Moran, E, Franklin, FCH & Franklin-Tong, VE 2015, 'The Papaver rhoeas S determinants confer self-incompatibility to Arabidopsis thaliana in planta', Science, vol. 350, no. 6261, pp. 684-687. <https://doi.org/10.1126/science.aad2983>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Title: The *Papaver rhoeas* S-determinants confer self-incompatibility to *Arabidopsis thaliana* in planta

Authors: Zongcheng Lin[†], Deborah J. Eaves, Eugenio Sanchez-Moran, F. Christopher H. Franklin and Veronica E. Franklin-Tong*

Affiliations:

School of Biosciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, U.K.

[†] current address: Department of Plant Systems Biology, VIB, 9052 Belgium.

* To whom correspondence should be addressed.

email: V.E.Franklin-Tong@bham.ac.uk; fax. +44 121 414 5925

Abstract:

Self-incompatibility (SI) is a major genetically controlled system used to prevent inbreeding in higher plants. S-determinants regulate allele-specific rejection of “self” pollen by the pistil. SI is an important model system for cell-cell recognition/signaling and could be potentially useful for F₁ hybrid breeding. To date, transfer of S-determinants has utilized complementation of orthologs to “restore” SI in close relatives. We expressed the *Papaver rhoeas* S-determinants, *PrsS* and *PrpS*, in *Arabidopsis thaliana*. This enabled pistils to reject pollen expressing cognate *PrpS*. Moreover, plants co-expressing cognate *PrpS* and *PrsS* exhibit robust SI. This demonstrates that *PrsS* and *PrpS* are sufficient for a functional synthetic S-locus *in vivo*. This represents the first transfer of novel S-determinants into a highly divergent species (>140 *m.y.* apart) with no orthologs.

One Sentence Summary:

The *Papaver rhoeas* S-determinants *PrsS* and *PrpS* confer self-incompatibility to *Arabidopsis thaliana* in planta

Main Text:

Many plants are hermaphrodites, with male and female organs in close proximity. As this risks self-fertilization and undesirable inbreeding depression, many plants utilize self-incompatibility (SI) as a mechanism to prevent selfing. SI is controlled by a S-locus allowing self/non-self

recognition between pistil and pollen (1, 2). SI in *Papaver rhoeas* is gametophytically controlled and specified by a pistil *S*-determinant, *PrsS* (*P. rhoeas stigma S* (3)) and a pollen *S*-determinant, *PrpS* (*P. rhoeas pollen S* (4)). *PrsS* and *PrpS* interact to trigger a signaling network in incompatible pollen, resulting in Programmed Cell Death (PCD) (5-8). *Arabidopsis thaliana* is a self-fertile member of the Brassicaceae. Self-compatibility in *Arabidopsis* originated recently (<0.5 mya) (9), through loss/inactivation of their *S*-determinants, *SRK* (10) and *SCR* (11). We recently demonstrated that pollen from *A. thaliana* expressing *PrpS-GFP* was inhibited by cognate recombinant *PrsS* proteins and displayed hallmark features of *Papaver* SI (8). Here we have expressed *PrsS* in *A. thaliana* pistils and show that they reject cognate pollen. Moreover, *A. thaliana* pistils co-expressing *PrsS* and *PrpS* set no self-seed. This demonstrates that *PrsS* and *PrpS* function as *S*-determinants and are the sole additional requirement to elicit SI. Intergeneric transfer of *S*-determinants has only been achieved between closely related species with orthologs of the *Brassica* *S*-determinants, effectively involving complementation using *SRK* and *SCR* pairs to “restore” SI (12-14). Because the Papaveraceae diverged from the Brassicaceae ~140 mya (8, 15) and *Arabidopsis* lacks *PrsS* and *PrpS* orthologs, finding that they function *in planta* in *A. thaliana* is a milestone.

PrsS encodes a small secreted protein, specifically and developmentally expressed in *P. rhoeas* stigmas (3). The promoter of *S Locus-Related 1* (*SLR1*) gene from *Brassica oleracea* directs stigma-specific, developmentally-regulated gene expression (16, 17), and exhibits maximal expression at flower maturity (16). Here we show that expression of *SLR1p* in *Arabidopsis* (**Fig 1A, Fig S1**) is spatiotemporally indistinguishable from *SRK* in *B. oleracea* (16). Therefore, we used *SLR1p* to drive expression of *PrsS₁* in *Arabidopsis* by transforming a *SLR1p::PrsS₁*

construct into *Col-0* (*At-PrsS₁* lines). RT-PCR analysis of pistil mRNA from 10 independent lines revealed differing *PrsS₁* transcript levels (**Fig 1B**). Western analysis of pistil extracts confirmed the presence of PrsS₁ (**Fig 1C**). We also transformed *Col-0* with a *SLR1p::PrsS₃* construct to make *At-PrsS₃* lines. RT-PCR analysis revealed similar transgene expression to the highest-expressing *At-PrsS₁* line (**Fig 1D**).

To test the functionality of the *At-PrsS₁* lines we performed *semi-vivo* pollination assays (**Fig S2**) on excised pistils from the *At-PrsS₁* lines, using pollen from an *A. thaliana* line expressing *PrpS₁-GFP* (8), referred to as *At-PrpS₁* hereafter, examining the ability of *At-PrsS₁* stigmas to inhibit *At-PrpS₁* (“incompatible”) pollen tube growth.

At-PrsS₁ line 9 pistils inhibited *At-PrpS₁* pollen tubes more strongly than line 4, while *Col-0* pollen was not inhibited (**Fig 2A**). Quantitation revealed that in *At-PrsS₁* pistils, *At-PrpS₁* pollen tubes were significantly shorter than *Col-0* pollen tubes ($P < 0.001$, *t-test*, $n = 40$; **Fig 2B**). After 70 min on *At-PrsS₁* pistils, *At-PrpS₁* pollen tubes from eight out of ten lines were $< 300 \mu\text{m}$; *Col-0* controls were $> 300 \mu\text{m}$ ($n = 40$; **Fig 2B**). Thus, *At-PrsS₁* pistils support pollen tube growth, but reject *At-PrpS₁* pollen. At 110 min, *At-PrpS₁* pollen tubes in *At-PrsS₁* pistils remained shorter than controls (**Fig S3**). The level of inhibition of *At-PrpS₁* pollen tubes in *At-PrsS₁* pistils correlated with *PrsS₁* expression levels (**Fig S4**). This provides strong evidence that *PrsS₁* functions in *A. thaliana* pistils to inhibit *At-PrpS₁* pollen.

A key feature of SI is *S*-allele specific inhibition of pollen. To test this, we pollinated excised *At-PrsS₁* or *At-PrsS₃* pistils with *At-PrpS₁* pollen or pollen from a line expressing *PrpS₃-GFP* (8),

referred to as *At-PrpS*₃ hereafter (**Fig 2C**). Strong pollen tube inhibition was observed only with cognate combinations of *At-PrsS* with *At-PrpS* (**Fig 2C, i, v**). Pollinations using non-cognate combinations of *At-PrsS* with *At-PrpS* resulted in normal pollen tube growth (**Fig 2C, ii, iv**). Controls (**Fig 2C, iii, vi, vii, viii, ix**) had long pollen tubes. *In vivo* pollinations of *At-PrsS* pistils also revealed differential inhibition of pollen tubes after 18 h (**Fig S5**). This demonstrates *S*-specific pollen tube inhibition by *A. thaliana* expressing PrsS.

With SI, pollination between cognate pollen and pistil *S*-alleles results in no seed production. *In planta* pollinations on *At-PrsS*₁ and *At-PrsS*₃ stigmas using cognate (incompatible) *At-PrpS*₁ and *At-PrpS*₃ pollen gave dramatically reduced silique lengths (6.2 ± 1.4 and 6.3 ± 1.7 mm respectively) compared to *Col-0* controls ($p < 0.001$ ***, *t*-test, $n=10$; **Fig 2D-E, Fig S6, Table 1**). In contrast, *At-PrsS*₁ and *At-PrsS*₃ pistils pollinated with non-cognate (compatible) pollen resulted in normal silique lengths, like *At-PrsS*₁ and *At-PrsS*₃ pistils pollinated with *Col-0* pollen ($p=0.397$, ANOVA, $n=10$), and pollination of *Col-0* stigmas with *At-PrpS*₁ or *At-PrpS*₃ pollen ($p=0.871$, ANOVA, $n=10$; **Table 1**) resulted in normal siliques, similar to selfed *Col-0* siliques, demonstrating *At-PrsS* stigma and *At-PrpS* pollen are functional.

Analyzing self-seed set, many siliques were completely empty (7/10 for *At-PrsS*₁ pollinated with *At-PrpS*₁ and 6/10 for *At-PrsS*₃ pollinated with *At-PrpS*₃); seed-set for cognate pollinations was between 0.5 ± 1.0 and 1.2 ± 1.8 seeds/silique ($n=20$; **Table 1**). Pollinations between non-cognate combinations resulted in normal seed-set (50.6 ± 5 , 50.0 ± 3.9 , $n=10$), significantly different from those with cognate combinations ($p < 0.001$, ***, *t*-test, $n=10$). Pollinations between *Col-0* pistils and *At-PrpS*₁ or *At-PrpS*₃ pollen gave normal seed-set (49.9 ± 3.7 and 47.6 ± 3.7 , $n=10$), so

transgenic stigmas and pollen are fully functional. Thus, the *Papaver* *S*-determinants function *in vivo* in an *S*-specific manner, resulting in failure of fertilization with cognate, but not non-cognate, pollen expressing *PrpS*.

We generated *Col-0* lines co-expressing *PrsS₁* and *PrpS₁* (*SI₁*-lines) by transforming homozygous *At-PrpS₁*-GFP plants with *SLR1p::PrsS₁*. Lines co-expressing *PrsS₃* and *PrpS₁* (*SC*-lines) were also generated. Expression of *PrsS₁* and *PrpS₁* was examined in three *SI₁*-lines (**Fig 3A**). Fluorescence microscopy of pollen from these *SI₁*-lines confirmed the expression of *PrpS₁*-GFP (**Fig 3B**). The *SI₁*- and *SC*-lines had a similar vegetative phenotype to *Col-0*, *At-PrpS₁* and *At-PrsS₁* plants (**Fig S7**). However, when left to set self-seed naturally, the *SI₁*-line plants had small siliques (**Fig 3C, Fig S7F**), between 3 ± 0.5 and 7 ± 1.4 mm long (n=470; **Fig S8A**), significantly shorter than siliques of control plants (**Fig 3C, Fig S7F**) *Col-0* (15.5 ± 0.6 mm), *At-PrpS₁* (16.3 ± 1.0), *At-PrsS₁* (15.9 ± 0.5) and *SC* plants (15.3 ± 0.5 mm; p<0.001 ***, *t*-test; n=10 per plant). Twelve of the *SI₁*-lines set no seed; the remaining 35 plants had between 0.1 ± 0.3 and 7.0 ± 1.4 seeds/silique (n=350; **Fig S8B**). This was significantly less (p<0.001 ***) than the 58 ± 1.6 seeds/silique in *Col-0* plants, *At-PrpS₁* plants (57.7 ± 2.8), *At-PrsS₁* plants (58.3 ± 1.6) and *SC*-lines (57.1 ± 1.7 ; n=10). Total self seed-set from these *SI₁*-lines gave between 0 and 680 seeds; ~60% had <100 seeds per plant (**Fig S8C**). Self seed-set of control plants was >8,500 seeds/plant (n=12). This SI response is stronger than previously obtained using the *S*-determinants from *A. lyrata* (12, 18) and similar to that achieved by (19). Lines co-expressing *PrsS₃* and *PrpS₃* (*SI₃*-lines) had a similar vegetative phenotype to *Col-0* plants except for short siliques (**Fig S9**). Self-seed-set analysis revealed small siliques (**Fig S10A, B**) and no/very low seed-set (**Fig S10B, C**), which were similar to those for the *SI₁*-lines. Analysis of naturally self-

pollinated pistils from *SI*-lines revealed that pollen tubes were inhibited in the upper pistil, while comparable self-pollinated *Col-0* pistils had pollen tubes extending through the pistil (**Fig 3E, F, Fig S11, Fig S12**). Together, these data provide compelling evidence that the *SI*-lines are self-incompatible.

To confirm that *SI*-lines were fully functional, pistils from representative *SI_I*-lines (*SI_I-9*, *SI_I-18* and *SI_I-32*) were pollinated with *At-PrpS₃* or *Col-0* pollen (**Fig 3D**, n=9). Siliques obtained were not significantly different from those pollinated using *Col-0* stigmas (p=0.246, p=0.703, ANOVA; n=3). Pollen from *SI_I*-lines was also pollinated to *At-PrpS₁* stigmas. They produced siliques and seed set not significantly different from *At-PrpS₁* stigmas pollinated with *Col-0* pollen (p=0.931, p=0.803, ANOVA; n=3; **Fig S13A, B**). As pollen and pistils from these *SI*-lines are functional, the reason why these *SI*-lines set no self-seed is not because they have a fertility defect, but because they are self-incompatible.

In summary, our data provide compelling evidence that the *Papaver S*-determinants co-expressed in *A. thaliana* make plants self-incompatible and are the sole additional requirement to establish SI in this highly diverged self-compatible species. This is a milestone, as successful transfer of *S*-determinants to date has been between close relatives sharing an ancestral SI system, using complementation to “restore” SI (12, 13). Because the *Papaveraceae* and Brassicaceae are evolutionarily separated by ~140 million years (8, 15), our finding that they function *in planta* in *Col-0* to display a robust SI rejection response is of considerable interest. We are not “restoring” a SI system as SI in Brassica/Arabidopsis has genetically and functionally distinct *S*-determinants. As we previously showed that recombinant PrsS can trigger SI-PCD in

Arabidopsis pollen expressing PrpS (8) and there is no evidence that Brassica/Arabidopsis SI involves PCD, the most economical explanation is that the *Papaver* *S*-determinants can interface with, and activate, a network of common signaling components that mediate PCD to induce a “*Papaver*-like” SI response in Arabidopsis pollen. *Papaver* SI uses Ca^{2+} , reactive oxygen species and pH (7, 20), which have all been described in Arabidopsis signaling networks achieving various physiological responses, including PCD (21). We hypothesize that these common signaling components are co-opted downstream of PrsS-PrpS interaction to mediate SI. Our findings reinforce proposals that SI may recruit pre-existing signaling networks from other biological processes (8, 22). This raises questions about how SI systems evolved, as well as about recruitment and functional diversification of pre-existing components (23).

Wide transgenera functionality of the *Papaver* SI system opens up the possibility that transfer of these *S*-determinants may, in the longer-term, provide a tractable SI system for crop plants. Use of the *SLRI* promoter from Brassica (16, 17) allows *PrsS* to be expressed in mature *Col-0* pistils unlike older *Col-0* pistils expressing *SCRb-SRKb* (12, 18). The production of F₁ hybrid plants in normally self-compatible species typically utilizes laborious, expensive manual emasculation to prevent self-fertilization. Transferral of a SI system into self-compatible species as an alternative method for the production of F₁ hybrids has been a long-term goal of SI research, with implications for solving Food Security issues.

REFERENCES & NOTES:

1. V. E. Franklin-Tong, Ed., *Self-Incompatibility in Flowering Plants: Evolution, Diversity, and Mechanisms*, (Springer-Verlag, Berlin, Heidelberg, 2008).
2. T. Dresselhaus, V. E. Franklin-Tong, Male–Female Crosstalk during Pollen Germination, Tube Growth and Guidance, and Double Fertilization. *Molecular Plant* **6**, 1018-1036 (2013).
3. H. C. C. Foote, et al., Cloning and expression of a distinctive class of self-incompatibility (*S*) gene from *Papaver rhoeas* L. *Proceedings of the National Academy of Sciences (USA)* **91**, 2265-2269 (1994).
4. M. J. Wheeler et al., Identification of the pollen self-incompatibility determinant in *Papaver rhoeas*. *Nature* **459**, 992-995 (2009).
5. S. G. Thomas, V. E. Franklin-Tong, Self-incompatibility triggers programmed cell death in *Papaver* pollen. *Nature* **429**, 305-309 (2004).
6. M. Bosch, V. E. Franklin-Tong, Temporal and spatial activation of caspase-like enzymes induced by self-incompatibility in *Papaver* pollen. *Proceedings of the National Academy of Sciences (USA)* **104**, 18327-18332 (2007).
7. K. A. Wilkins, N. S. Poulter, V. E. Franklin-Tong, Taking one for the team: self-recognition and cell suicide in pollen. *Journal of Experimental Botany* **65**, 1331-1342 (2014).
8. B. H. de Graaf et al., The *Papaver* Self-Incompatibility pollen *S*-determinant, PrpS, functions in *Arabidopsis thaliana*. *Current Biology* **22**, 154-159 (2012).
9. J. S. Bechsgaard, V. Castric, D. Charlesworth, X. Vekemans, M. H. Schierup, The transition to Self-Compatibility in *Arabidopsis thaliana* and evolution within *S*-Haplotypes over 10 Myr. *Molecular Biology and Evolution* **23**, 1741-1750 (2006).
10. J. C. Stein, B. Howlett, D. C. Boyes, M. E. Nasrallah, J. B. Nasrallah, Molecular cloning of a putative receptor protein kinase gene encoded at the self-incompatibility locus of *Brassica oleracea*. *Proc. Natl Acad. Sci. (USA)*. **88**, 8816-8820 (1991).
11. C. R. Schopfer, M. E. Nasrallah, J. B. Nasrallah, The male determinant of self-incompatibility in Brassica. *Science* **286**, 1697-1700 (1999).
12. M. E. Nasrallah, P. Liu, J. B. Nasrallah, Generation of Self-Incompatible *Arabidopsis thaliana* by Transfer of Two *S* Locus Genes from *A. lyrata*. *Science* **297**, 247-249 (2002).
13. N. A. Boggs, K. G. Dwyer, P. Shah, A. A. McCulloch, J. Bechsgaard, M. H. Schierup, M. E. Nasrallah, J. B. Nasrallah, Expression of distinct Self-Incompatibility specificities in *Arabidopsis thaliana*. *Genetics* **182**, 1313-1321 (2009).
14. M. Yamamoto, T. Nishio, Commonalities and differences between Brassica and Arabidopsis self-incompatibility. *Horticulture Research* **1**, 14054 (2014).
15. C. D. Bell, D. E. Soltis, P. S. Soltis, The age and diversification of the angiosperms re-revisited. *American Journal of Botany* **97**, 1296-1303 (2010).
16. B. A. Lalonde et al., A highly conserved *Brassica* gene with homology to the *S*-locus-specific glycoprotein structural gene. *The Plant Cell* **1**, 249–258 (1989).
17. R. M. Hackett, M. J. Lawrence, F. C. H. Franklin, A Brassica *S*-locus related gene promoter directs expression in both pollen and pistil of tobacco. *The Plant Journal* **2**, 613-617 (1992).
18. N. A. Boggs, J. B. Nasrallah, M. E. Nasrallah, Independent *S*-Locus Mutations Caused Self-Fertility in *Arabidopsis thaliana*. *PLoS Genet* **5**, e1000426 (2009).
19. D. R. Goring, E. Indriolo, M. A. Samuel, The ARC1 E3 Ligase promotes a strong and stable Self-Incompatibility response in Arabidopsis species: Response to the Nasrallah and Nasrallah Commentary. *The Plant Cell* **26**, 3842-3846 (2014).
20. K. A. Wilkins et al., Self-Incompatibility-Induced Programmed Cell Death in Field Poppy Pollen Involves Dramatic Acidification of the Incompatible Pollen Tube Cytosol. *Plant Physiology* **167**, 766-779 (2015).
21. T. van Hautegeem, A. J. Waters, M. K. Nowack, Only dying in life: programmed cell death during plant development. *Trends in Plant Sciences* **20**, 1360-1385 (2015).
22. T. Tantikanjana, M. E. Nasrallah, J. B. Nasrallah, Complex networks of self-incompatibility signaling in the Brassicaceae. *Current Opinion in Plant Biology* **13**, 520-526 (2010).
23. F. M. Ausubel, Are innate immune signaling pathways in plants and animals conserved? *Nature Immunology* **6**, 973 - 979 (2005).

Acknowledgments: We thank Daphne Goring for the Binary Ti vector pORE O3 containing *SLR1* promoter. Thanks to Steve Price for technical assistance. Z.L. held a PhD studentship from the China Scholarship Council (C.S.C.). D.J.E. was funded by the Biotechnology and Biological Sciences Research Council (B.B.S.R.C.).

V.E.F-T. and F.C.H.F. are co-inventors on a patent application (2691/KOLNP/2011) filed by University of Birmingham relating to PrsS and PrpS. Materials will be freely available upon request for research purposes.

The authors declare that they do not have other competing financial interests.

The data reported here are available in Supplementary Materials.

Fig. 1. Expression of the *SLRI* promoter is developmental-, tissue-specific and drives expression of *PrsS₁* in *A. thaliana*.

- (A) RT-PCR (*top*) shows *SLRI_p::GFP* developmentally expressed in pistils and not expressed in stamen, petal, or leaf tissue. *GAPC* shows equal loading.
- (B) RT-PCR of pistils from *At-PrsS₁* lines shows expression of *PrsS₁* (*top*) and quantitation of *PrsS₁* expression relative to *GAPC* (n=3, ± S.D., *below*).
- (C) Western blot (α -PrsS₁ antisera) shows expression of PrsS₁ in *At-PrsS₁*.
- (D) RT-PCR: expression of *PrsS₃* in *At-PrsS₃* line 8 is comparable to *At-PrsS₁* line 9.

Fig. 2. *At-PrpS* pollen is inhibited on cognate *At-PrsS* pistils, demonstrating S-specificity.

- (A) Aniline blue staining of representative *semi-in-vivo* pollinations of *At-PrsS₁* pistils with *At-PrpS₁* or *Col-0* pollen.
- (B) Quantitation of pollen tube lengths on *At-PrsS₁* pistils using *At-PrpS₁* pollen (*left*) or *Col-0* pollen (*right*); n= 4 stigmas/*At-PrsS₁* line.
- (C) *At-PrsS₁* and *At-PrsS₃* pistils pollinated *semi-in-vivo* with *At-PrpS₃* or *At-PrpS₁* pollen. *At-PrpS* pollen tubes were inhibited on cognate *At-PrsS* pistils (i,v), while controls did not.
- (D) Representative *in-vivo* pollination of an *At-PrsS₁* stigma with *At-PrpS₁* pollen resulted in a small, empty silique.
- (E) *Col-0* pollinated with *Col-0* pollen had normal length silique and many seeds.

Fig 3. *A. thaliana* co-expressing *PrsS₁* and *PrpS₁* are self-incompatible and set no seed.

- (A) RT-PCR of 3 *A. thaliana* *SI₁*-lines co-expressing *PrsS₁* and *PrpS₁*.
- (B) Pollen from *SI₁*-lines exhibits GFP fluorescence (*top*); *Col-0* pollen has weak autofluorescence.
- (C) Self-seed set: *SI₁*-lines formed short siliques; controls, including a *SC*-line co-expressing *PrsS₃* and *PrpS₁-GFP*, set normal siliques
- (D) A selfed *SI₁*-plant gave small siliques; pollinations with *Col-0* or *At-PrpS₃* pollen gave normal siliques.
- (E) Aniline blue staining of a self-pollinated *SI₁*-line pistil; pollen tubes are inhibited in the stigma/style.
- (F) Self-pollinated *Col-0* pistil had long pollen tubes.

Table 1. *In vivo* pollination of *At-PrsS* stigmas with cognate *At-PrpS* pollen resulted in shorter siliques and no seed set.

Pollination of emasculated *At-PrsS₁* stigmas with *At-PrpS₁* pollen resulted in short siliques and reduced seed number, as did pollination of *At-PrsS₃* with *At-PrpS₃* pollen. Other control pollinations: non-cognate pollination of *At-PrsS₁* stigmas with *At-PrpS₃* pollen, *At-PrsS₃* stigmas with *At-PrpS₁* or *Col-0* pollen, *Col-0* stigmas with *Col-0*, *At-PrpS₁* or *At-PrpS₃* pollen gave normal silique length and seed number (mean \pm S.D., n=10).

♀ \ ♂		<i>At-PrpS₁</i>	<i>At-PrpS₃</i>	<i>Col-0</i>
<i>At-PrsS₁</i> (line 9)	Silique lengths (mm)	6.2 \pm 1.4	16.1 \pm 0.8	16.4 \pm 0.7
	Seeds per silique	0.5 \pm 1.0	50.6 \pm 5.1	49.3 \pm 5.3
<i>At-PrsS₃</i> (line 8)	Silique lengths (mm)	16.4 \pm 0.8	6.3 \pm 1.7	16.5 \pm 0.5
	Seeds per silique	50.0 \pm 3.9	1.2 \pm 1.8	50.0 \pm 3.2
<i>Col-0</i>	Silique lengths (mm)	16.6 \pm 1.0	16.6 \pm 0.8	16.4 \pm 0.7
	Seeds per silique	49.9 \pm 3.7	47.6 \pm 3.7	47.7 \pm 3.6




Fig 1.




Fig 2.




Fig 3.