

# UNIVERSITY OF BIRMINGHAM

## Research at Birmingham

### Biliary epithelium and liver B cells exposed to bacteria activate intrahepatic MAIT cells through MR1

Jeffery, Hannah; van Wilgenburg, Bonnie; Kurioka, Ayako; Parekh, Krishan; Stirling, Kathryn; Roberts, Sheree; Dutton, Emma; Hunter, Stuart; Geh, Daniel; Kaur, Manjit; Rajanayagam, Jeremy; Iqbal, Tariq; Pinkney, Thomas; Brown, Rachel; Withers, David; Adams, David; Klenerman, Paul; Oo, Ye Htun

DOI:

[10.1016/j.jhep.2015.12.017](https://doi.org/10.1016/j.jhep.2015.12.017)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Jeffery, HC, van Wilgenburg, B, Kurioka, A, Parekh, K, Stirling, K, Roberts, S, Dutton, EE, Hunter, S, Geh, D, Braitch, MK, Rajanayagam, J, Iqbal, T, Pinkney, T, Brown, R, Withers, DR, Adams, DH, Klenerman, P & Oo, Y 2016, 'Biliary epithelium and liver B cells exposed to bacteria activate intrahepatic MAIT cells through MR1', *Journal of Hepatology*, vol. 64, no. 5, pp. 1118–1127. <https://doi.org/10.1016/j.jhep.2015.12.017>

[Link to publication on Research at Birmingham portal](#)

#### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## Biliary epithelium and liver B cells exposed to bacteria activate intrahepatic MAIT cells through MR1

Hannah C. Jeffery<sup>1,†</sup>, Bonnie van Wilgenburg<sup>2,†</sup>, Ayako Kurioka<sup>2</sup>, Krishan Parekh<sup>1</sup>, Kathryn Stirling<sup>1</sup>, Sheree Roberts<sup>1</sup>, Emma E. Dutton<sup>3</sup>, Stuart Hunter<sup>1</sup>, Daniel Geh<sup>1</sup>, Manjit K. Braitch<sup>1</sup>, Jeremy Rajanayagam<sup>1</sup>, Tariq Iqbal<sup>4</sup>, Thomas Pinkney<sup>4</sup>, Rachel Brown<sup>4</sup>, David R. Withers<sup>3</sup>, David H. Adams<sup>1,4</sup>, Paul Klenerman<sup>2,‡</sup>, Ye H. Oo<sup>1,4,\*</sup>

<sup>1</sup>Centre for Liver Research and NIHR Biomedical Research Unit in Liver Disease, Institute of Immunology & Immunotherapy, University of Birmingham, United Kingdom; <sup>2</sup>Peter Medawar Building for Pathogen Research, University of Oxford, United Kingdom; <sup>3</sup>Institute of Immunology and Immunotherapy, University of Birmingham, United Kingdom; <sup>4</sup>University Hospital of Birmingham NHS Foundation Trust, United Kingdom

See Editorial, pages 1008–1010

**Background & Aims:** Mucosal-Associated Invariant T (MAIT) cells are innate-like T cells characterised by the invariant TCR-chain, V $\alpha$ 7.2-J $\alpha$ 33, and are restricted by MR1, which presents bacterial vitamin B metabolites. They are important for antibacterial immunity at mucosal sites; however, detailed characteristics of liver-infiltrating MAIT (LI-MAIT) and their role in biliary immune surveillance remain unexplored.

**Methods:** The phenotype and intrahepatic localisation of human LI-MAIT cells was examined in diseased and normal livers. MAIT cell activation in response to *E. coli*-exposed macrophages, biliary epithelial cells (BEC) and liver B cells was assessed with/without anti-MR1.

**Results:** Intrahepatic MAIT cells predominantly localised to bile ducts in the portal tracts. Consistent with this distribution, they expressed biliary tropic chemokine receptors CCR6, CXCR6, and integrin  $\alpha$ E $\beta$ 7. LI-MAIT cells were also present in the hepatic sinusoids and possessed tissue-homing chemokine receptor CXCR3 and integrins LFA-1 and VLA-4, suggesting their recruitment via hepatic sinusoids. LI-MAIT cells were enriched in the parenchyma of acute liver failure livers compared to chronic diseased livers. LI-MAIT cells had an activated, effector memory phenotype, expressed  $\alpha$ 4 $\beta$ 7 and receptors for IL-12, IL-18, and IL-23. Importantly, in response to *E. coli*-exposed macrophages, liver B cells and BEC, MAIT cells upregulated IFN- $\gamma$  and CD40 Ligand and degranulated in an MR1-dependent, cytokine-independent

manner. In addition, diseased liver MAIT cells expressed T-bet and ROR $\gamma$ t and the cytokines IFN- $\gamma$ , TNF- $\alpha$ , and IL-17.

**Conclusions:** Our findings provide the first evidence of an immune surveillance effector response for MAIT cells towards BEC in human liver; thus they could be manipulated for treatment of biliary disease in the future.

© 2016 European Association for the Study of the Liver. Published by Elsevier B.V. All rights reserved.

### Introduction

Mucosal-associated invariant T (MAIT) cells are a recently identified subset of T cells with an evolutionarily conserved invariant T cell antigen receptor (TCR)  $\alpha$ -chain, composed of the invariant  $\alpha$ -chain V $\alpha$ 7.2-J $\alpha$ 33/J $\alpha$ 20/J $\alpha$ 12 in humans and V $\alpha$ 19-J $\alpha$ 33 in mice [1,2]. They are restricted to the CD161<sup>+</sup> population and are abundant in human blood, the intestinal mucosa and mesenteric lymph nodes [3–5]. MAIT cells respond to antigen presented on the highly phylogenetically conserved major histocompatibility complex (MHC) class I-related molecule, MR1, which possesses a unique antigen-binding cleft for vitamin B metabolites from pathogenic and/or commensal bacteria, and distinguishes MAIT cells from peptide- or lipid-recognizing  $\alpha$  $\beta$  T cells [1,6,7]. MAIT cells can be activated by a wide variety of bacterial strains *in vitro*, and importantly they are crucial in mucosal immune defence in bacterial infection [8–10]. They respond in an MR1-dependent manner to antigen presenting cells (APC) cultured with bacteria and can also be activated via IL-12 and IL-18 in a TCR-independent manner [11,12]. MAIT cell frequencies have been reported to be lower in bacterially-infected patients' blood [10,13].

Both hepatic sinusoids and biliary epithelial cells (BEC) are crucial in first-line defence towards pathogens in both the steady and disease state as the human liver is continuously exposed to intestinally-derived antigens from portal venous blood and biliary reflux [14]. A recent study suggested that immune cells in the

**Keywords:** Human liver; Mucosal-associated invariant T cells; Biliary epithelium; *E. coli*; Immune response; Biliary firewall.

Received 21 May 2015; received in revised form 17 December 2015; accepted 20 December 2015; available online 29 December 2015

\* DOI of original article: <http://dx.doi.org/10.1016/j.jhep.2016.02.003>.

\* Corresponding author. Address: Centre for Liver Research and National Institute of Health Research Birmingham Biomedical Research Unit, University of Birmingham, Vincent Drive, Edgbaston, Birmingham B15 2TT, UK. Tel.: +44 121 415 8700; fax: +44 121 415 8701.

E-mail address: [y.h.oo@bham.ac.uk](mailto:y.h.oo@bham.ac.uk) (Y.H. Oo).

<sup>†</sup> These contributed equally as joint first authors.

<sup>‡</sup> These authors share senior authorship.



ELSEVIER

hepatic sinusoids function as a firewall to prevent the systemic spread of gut-derived pathogens that evade the mesenteric immune system [15]. The presence of MAIT cells has been reported in healthy human liver sinusoidal fluids [16], however, their role in mucosa defence at the bile ducts, which are continuous with the gut lumen and its microbes, and form the first-line protection against biliary pathogens, is still unexplored [17,18]. BEC are known to express antigen presenting molecules and can activate lymphocytes [19]. A recent report indicated that MAIT cells could efficiently lyse epithelial cells of the HeLa cell line that are infected with bacteria [20]. Taken together, these findings indicate that MAIT cells are likely to be important contributors to the maintenance of steady state immunity and the pathogenesis of inflammatory and biliary liver diseases, especially in response to bacterial exposure. Thus, in the current study, we used primary human liver tissues, obtained from both normal and diseased explanted human livers, to investigate the localisation and phenotype of intrahepatic/liver-infiltrating MAIT (LI-MAIT) cells, as well as their functional response to bacterially-exposed biliary epithelial surfaces in inflammatory biliary liver diseases.

## Materials and methods

*Isolation of liver-infiltrating lymphocytes (LIL), peripheral blood lymphocytes (PBL), and BEC*

Venous blood, collected in EDTA, was obtained from healthy donors, and patients with inflammatory and autoimmune liver diseases (primary sclerosing cholangitis (PSC), primary biliary cirrhosis (PBC) and alcoholic liver disease (ALD)). Explanted diseased liver tissue was obtained from patients who underwent liver transplantation for end-stage liver diseases including PSC, PBC, ALD, and non-alcoholic steatohepatitis (NASH) or for acute liver failure from seronegative (NonA NonB (NANB)) hepatitis. Non-diseased liver tissues were obtained from unused donor liver tissues. All samples were collected with appropriate patient consent and local research ethics committee approval (LREC ref. CA/5192, 06/Q2708/11). Human LIL, PBL, and BEC cells were isolated from fresh liver tissue and peripheral blood as described previously [21].

*Localisation of TCR V $\alpha$ 7.2-expressing cells*

Human liver tissues were stained with purified anti-TCR V $\alpha$ 7.2 (50  $\mu$ g/ml, 3C10, BioLegend) or IgG1 isotype control to detect the localisation of V $\alpha$ 7.2<sup>+</sup> cells. See [Supplementary material](#).

*Phenotyping of intrahepatic and peripheral blood MAIT cells*

Liver-infiltrating and blood MAIT cells were phenotyped directly *ex vivo* for the expression of surface markers, transcription factors and intracellular cytokines. See [Supplementary material](#).

*MAIT cell response to E. coli-exposed antigen presenting cells*

APCs: blood monocyte-derived macrophages, THP1, liver B cells or BEC were incubated overnight with paraformaldehyde-fixed *Escherichia coli* (*E. coli*) (DH5 $\alpha$ , Invitrogen) at 25, 20, 1000, or 1000 bacteria per cell respectively. CD8<sup>+</sup> T cells isolated from blood using CD8 Microbeads (Miltenyi Biotec) or CD3<sup>+</sup> T cells isolated from liver by flow sorting were cultured with the *E. coli*-exposed APCs, in the presence of anti-CD107a (Pe or PeCy5) and blocking antibodies against IL-12p40/70 (5  $\mu$ g/ml, C8.6, eBioscience), IL-18 (5  $\mu$ g/ml, 125-2H, MBL International, USA) and MR1 (10  $\mu$ g/ml) [22] as indicated. In some assays, anti-CD40 Ligand (CD40L)-PeCy7 was added. MAIT cell intracellular and surface markers were stained and data were acquired on a MACSQuant (Miltenyi Biotec) or CyAN (Dako) flow cytometer and analyzed using FlowJo (Tree Star Inc.). Autologous liver-infiltrating B cells and T cells were cell sorted by a MoFlo Astrios cell sorter (Beckman Coulter). Cells were labeled with anti-CD3-PeCy7 and anti-CD19-APCv770 to identify CD3<sup>+</sup> T cells and CD19<sup>+</sup> B cells respectively. Blood

macrophages were generated by culturing CD14<sup>+</sup> monocytes, isolated from blood using CD14 microbeads (Miltenyi Biotec), with 100 ng/ml M-CSF (R and D Systems) for 7–9 days.

*Statistical analysis*

GraphPad Prism 5.0 software (GraphPad software, San Diego, CA, USA) was used for statistical analysis. Comparisons of two populations were by the Mann-Whitney *U* test or *t* test. Comparisons of more than one population were by Friedman's test with Dunn's Multiple Comparison post-hoc test or by one-way ANOVA with Bonferroni's multiple comparison post hoc test as indicated in the figure legend. Statistical significance was defined as *p* value <0.05. Error bars on graphs are presented as median  $\pm$  interquartile range or mean  $\pm$  SEM. Values in text are given as median and overall range (in brackets).


## Results

*Intrahepatic MAIT cells preferentially reside in peri-biliary areas of portal tracts*

We examined the localisation of LI-MAIT cells in normal and diseased human livers by immunohistochemistry staining for TCR V $\alpha$ 7.2. Most V $\alpha$ 7.2<sup>+</sup> cells resided around bile ducts in portal tracts with few detected in the parenchyma (Fig. 1A, B; [Supplementary Fig. 2](#)). The distribution was similar in normal, autoimmune, and non-autoimmune diseased livers (Fig. 1C; [Supplementary Fig. 2](#)) similar to other immune subsets ([Supplementary Fig. 1](#)). Interestingly, in acute, seronegative liver failure, increased infiltration of V $\alpha$ 7.2<sup>+</sup> cells to the parenchyma was noted (Fig. 1A iii, vi, 1C; [Supplementary Fig. 3](#)) when compared to normal livers or any of the chronic liver diseases studied (Fig. 1A i, iv). The overall frequency of V $\alpha$ 7.2<sup>+</sup> cells appeared increased in PSC compared to the other liver diseases (Fig. 1C). By flow cytometry, we showed that the majority of V $\alpha$ 7.2<sup>+</sup> lymphocytes in normal livers (63.6% (24.4–93.2%)) and over one-third in diseased (40.5% (11.6–75.2%)) were CD3<sup>+</sup>CD161<sup>++</sup> MAIT cells ([Supplementary Fig. 4](#)). We confirmed the predominant localisation of CD3<sup>+</sup> CD161<sup>+</sup> V $\alpha$ 7.2<sup>+</sup> MAIT cells in peri-biliary regions of portal tracts by both immunohistochemistry (Fig. 1Aii, v; 1C) and confocal microscopy (Fig. 2).

*Frequencies of MAIT cells are reduced in liver diseases, with an increase in the CD4<sup>+</sup> MAIT cells*

Next, using flow cytometry we compared frequencies of CD3<sup>+</sup> CD161<sup>++</sup> V $\alpha$ 7.2<sup>+</sup> MAIT cells in intrahepatic liver infiltrates and in blood from normal and diseased tissues. Increased frequency of MAIT cells in liver compared to blood was observed in both normal and diseased states (Fig. 3A, B). The frequency of liver and blood MAIT cells in total CD3<sup>+</sup> T cells was decreased in chronic liver diseases (Fig. 3A, B). In liver as in blood, CD8<sup>+</sup> cells represented the major MAIT cell subset (Fig. 3C, D). However, in disease, the proportion of CD4<sup>+</sup> MAIT cells was significantly increased in both the blood and liver, which in liver, was compensated for by a significant reduction in the CD8<sup>+</sup> MAIT cell frequency (Fig. 3C, D). MAIT cells were unique among the T cell subsets that we examined in showing a reduced frequency with disease (Fig. 3E). We observed a negative correlation between total MAIT cells and total CD4<sup>+</sup> T cells in normal livers but found no sign of this correlation in disease. Conversely there was a trend towards a positive correlation of MAIT cells with CD8<sup>+</sup> T cells in normal livers. In non-autoimmune livers we noticed a positive correlation with CD161<sup>+</sup> T cells. No relationships were




**Fig. 1. Peri-biliary localisation of Va7.2+ cells in chronic liver diseases.** (A) Representative staining for Va7.2 on frozen liver sections viewed at 10× (i and iii) or 40× (ii, iv, v, and vi) magnification. Distribution of Va7.2+ cells in the parenchyma (i and iv) and portal tract (i, ii, and v) in PSC and in the parenchyma (iii and vi) in seronegative acute liver failure. (B) Densities of Va7.2+ cells in parenchyma and portal tracts of normal and chronically diseased livers (\*\**p* < 0.01; \*\*\**p* < 0.0001 by Mann-Whitney *U* test). (C) Va7.2+ cell density data according to diseases. Data are median ± interquartile range.

found between MAIT cells and CD4<sup>-</sup>CD8<sup>-</sup> double negative (DN) T cells in either normal or diseased livers (Fig. 3F). In disease, the proportion of CD4<sup>+</sup> cells within the MAIT cell population was approximately 2-fold greater than that occurring for total CD4<sup>+</sup> cells within the total T cell population; however the frequency of CD4<sup>+</sup> MAIT cells among total T cells did not alter with disease, rather the CD8<sup>+</sup> and DN MAIT cell frequencies among total T cells decreased significantly in disease, accounting for the rise in the proportion of CD4<sup>+</sup> cells within the MAIT cell population (Fig. 3E).

*Tissue-homing chemokine receptor, integrin and cytokine receptor expressions of intrahepatic MAIT cells*


Chemokine receptors, CXCR6 and CCR6 and integrin αEβ7 have been implicated in lymphocyte recruitment to biliary epithelium [21,23,24]. All three were expressed by LI-MAIT cells from both diseased and normal livers [CXCR6: (normal: 29% (14–33%);



**Fig. 2. CD3<sup>+</sup>CD161<sup>+</sup>Va7.2<sup>+</sup> cells reside close to bile ducts in portal tracts.** Representative confocal immunofluorescence staining for CD3, CD161, and Va7.2 on frozen sections from explanted human livers diagnosed with Alcoholic liver disease (A) and Primary Biliary Cirrhosis (B). DAPI nuclear stain reveals liver architecture indicating sites of bile ducts. Images are representative of staining of four different diseased livers, scale bar shows 100 μm.

diseased: 22% (4–52%)), CCR6 (normal: 36% (12–72%); diseased: 53% (7–81%)), αEβ7 (normal: 4% (4–12%); diseased: 16% (2–37%))] (Fig. 4A).


Sinusoidal recruitment to inflamed tissue involves the chemokine receptor CXCR3, which responds to interferon-dependent ligands, CXCL9/10/11 in inflamed tissues, and the integrins LFA-1 and VLA-4 [25]. Almost all LI-MAIT cells expressed LFA-1 (normal: 96% (92–98%); diseased: 91% (81–93%)) and most expressed VLA-4 (normal: 57% (13–75%); diseased: 58% (19–71%)) (Fig. 4B). LI-MAIT cells from diseased livers showed significant upregulation of CXCR3 compared to LI-MAIT cells from normal tissue (normal: 21% (17–31%); diseased: 81% (42–89%), *p* < 0.01) (Fig. 4B). Increased expression of CX3CR1 and CCR5 was also noticed in disease (Fig. 4C). Since MAIT cells are believed to expand in the gut in response to bacterial antigens, we screened for expression of the gut-homing chemokine receptor CCR9 and integrin α4β7 [26] but detected little expression on LI-MAIT cells from either normal or diseased tissue (Fig. 4C). Given the change in the balance of CD4<sup>+</sup>/CD8<sup>+</sup>/DN<sup>+</sup> MAIT cells in diseased livers we



**Fig. 3.** Flow cytometric analysis of MAIT cell frequencies in chronic liver disease and their correlations with other immune subsets. Representative FACS plots (A) and summary frequency data for total CD3<sup>+</sup> MAIT (B) and CD4<sup>+</sup>, CD8<sup>+</sup> and CD4<sup>+</sup>CD8<sup>+</sup> (DN) MAIT subsets in normal and diseased blood (C) and liver (D). (E) Frequencies of intrahepatic MAIT and other immune cells. (F) Correlation of CD3<sup>+</sup> MAIT cell frequencies with total CD4<sup>+</sup>, CD8<sup>+</sup>, DN, and CD161<sup>+</sup> T cell frequencies in normal and diseased livers. Data are median ± interquartile range. \**p* < 0.05; \*\**p* < 0.01; \*\*\**p* < 0.001 by Mann-Whitney *U* test (B–E) or Spearman's rank correlation (F).

assessed whether there were any subset-specific differences in homing receptor expression profile. We observed little difference in percentage and intensity of expression across subsets in their expression of CCR6, CCR5 and CX3CR1. Interestingly, however, higher frequencies of CD4<sup>+</sup> MAIT expressed CXCR3

compared to CD8<sup>+</sup> or DN MAIT in normal livers, although the intensity of expression did not differ across subsets. In disease, frequencies of CD4<sup>+</sup> and CD8<sup>+</sup> CXCR3-expressing cells did not differ but the intensity of expression on CD4<sup>+</sup> cells was significantly greater than on CD8<sup>+</sup> or DN cells (Supplementary Fig. 5).



**Fig. 4. Expressions of tissue-homing chemokine, integrin and cytokine receptors on intrahepatic MAIT cells in normal and chronic liver diseases.** Chemokine receptor and integrin expression profiles (A, B, and C) and cytokine receptor profiles (D) were determined by flow cytometry gating on the total CD3<sup>+</sup> MAIT cell population. Representative overlays for marker (line) and isotype control (grey shading) and total summary data are shown. Summary data are median ± interquartile range. \**p* < 0.05; \*\**p* < 0.01; \*\*\**p* < 0.001 by Mann-Whitney *U* test.

We also evaluated the expression of cytokine receptors whose cognate cytokines are known to be secreted by liver-resident cells and to mediate MAIT cell activation *in vitro* [12]. The pattern of cytokine receptor expression was similar in diseased and non-diseased liver tissue. IL-18R was highly expressed (normal: 97% (93–99%); diseased: 83% (65–96%)). There was low-to-moderate expression of IL-12R (normal: 7.5%; diseased: 15.7%) and IL-23R (normal: 23%; diseased: 16%). IL-6R was not detected on LI-MAIT cells (Fig. 4D).


*Phenotypic characterisation of liver-infiltrating MAIT cells*

There was no overlap of the LI-MAIT cell population with  $\gamma\delta$  T cells (Supplementary Fig. 6A). Although LI-MAIT cells had moderate expression of the Natural Killer (NK) cell marker CD56, they had low expression of other NK cell phenotypic markers such as NKG2D and NKp46 (Supplementary Fig. 6B). LI-MAIT cells were mostly CD45RA<sup>+</sup>CCR7<sup>-</sup> effector memory (Supplementary Fig. 7A) and expressed the activation marker CD69 but lacked CD40L (Supplementary Fig. 7B). LI-MAIT cells showed ubiquitous surface expression of CD95 (93%) but

lacked CD95L (Supplementary Fig. 7C). They were also found to have constitutive expression of CD26 and to express the adenosine-pathway receptors CD39, and CD73 (Supplementary Fig. 8A).

*Diseased liver-infiltrating MAIT cells produce IFN- $\gamma$ , TNF- $\alpha$ , IL-17, and granzyme B*

To determine possible effector functions of LI-MAIT cells, we examined the cytokines and cytotoxic granzyme produced by MAIT cells *ex vivo*. LI-MAIT cells showed high frequencies of IFN- $\gamma$  (55%) and TNF- $\alpha$  expression (89%) and low frequencies of IL-17 production (3.5%) but IL-22 and Th2-cytokines including IL-4, IL-5, and IL-13 were barely detected (Fig. 5A, B). We noticed that approximately 50% of IL-17-producing cells had a dual Th1/Th17 phenotype, secreting IFN- $\gamma$ . Consistent with their ability to produce both Th1 and Th17-type cytokines, LI-MAIT cells expressed the transcription factors T-bet and ROR $\gamma$ , (Supplementary Fig. 8B). Examining *ex vivo* stores of cytotoxicity factor, we found moderate frequencies of granzyme B-expressing LI-MAIT cells (10% (5–25%)) (Fig. 5F).



**Fig. 5. Cytokine and cytolytic factor expression by liver-infiltrating MAIT cells.** Intrahepatic MAIT cell production of Th1 (IFN- $\gamma$ , TNF- $\alpha$ ), Th2 (IL-4, IL-5, IL-13) and Th17 (IL-17, IL-22) cytokines was examined by flow cytometry. Representative dot-plots for each cytokine vs. IFN- $\gamma$  expression (A) and summary data by MAIT cells (B) and MAIT cell subsets (C and D) are shown. Summary data for *ex vivo* granzyme B expression in total MAIT and MAIT cell subsets (E). Data are median  $\pm$  interquartile range. \* $p < 0.05$ ; \*\* $p < 0.01$ ; by Dunn's post hoc test following Friedman's test.

*Activation of blood and liver-infiltrating MAIT cells by antigen presenting cells in an MR1 dependent manner*

MAIT cells can be activated in response to bacterial metabolites in an MR1- and/or IL-12/IL-18-dependent manner by professional and non-professional APC, such as HeLa cells, B cells and THP1 cells [20,27]. As a model to study activation by macrophages, we exposed blood MAIT cells to monocyte-derived macrophages that had been treated with or without *E. coli* and examined the expression of the degranulation marker CD107a, IFN- $\gamma$ , and CD40L. MAIT cells expressed CD107a and IFN- $\gamma$  in an MR1-dependent manner. They also showed a tendency for CD40L expression (Fig. 6A). These responses were MR1-dependent but independent of IL-12 or IL-18. Both THP1 cells (Fig. 6B) and liver-infiltrating B cells (Fig. 6C), pre-treated with *E. coli*, activated LI-MAIT cells from diseased livers by inducing the expression of CD107a, IFN- $\gamma$  and TNF- $\alpha$  in an MR1-dependent manner.


*Bacterially-exposed primary human biliary epithelial cells activate MAIT cells in an MR1-dependent, cytokine-independent manner*

Since we observed LI-MAIT cells around bile ducts in the portal tracts we proceeded to examine whether MAIT cells may respond to bacterial infection associated with the biliary epithelium. We

therefore co-cultured blood derived MAIT cells with primary human BEC with or without *E. coli* and observed selective activation by the MAIT population in the presence of BEC presenting *E. coli*. The V $\alpha$ 7.2<sup>+</sup>CD161<sup>-</sup> cells within the same culture did not respond (Fig. 7A). Increased expression of CD107a and IFN- $\gamma$  was MR1-dependent, however, blocking the cytokines IL-12 and IL-18 did not have any effect (Fig. 7B). CD40L upregulation was significantly inhibited by blocking either MR1 or the cytokines IL-12 and IL18 (Fig. 7B). We performed the same assay with T cells isolated from diseased livers. LI-MAIT responded to BEC presenting *E. coli*, upregulating CD107a, IFN- $\gamma$  and TNF- $\alpha$  in an MR1-dependent manner (Fig. 7C).

**Discussion**

Although MAIT cells have recently been reported in normal human liver perfusate, detailed characteristics of liver-infiltrating MAIT cells in normal and diseased state, and their role in biliary epithelial mucosa protection remained unexplored [3,16,28,29]. The biliary epithelium is in continuity with the intestinal gut flora and thus provides another potential portal of pathogen entry from the gut to the liver. As such, resident immune cells including MAIT cells that localised around bile




**Fig. 6.** MR1-dependent activation of blood and liver MAIT cells by professional and non-professional antigen presenting cells (APC) exposed to bacteria. APC exposed to *E. coli* were co-cultured with MAIT cells and their activation examined by flow cytometry for CD107a, IFN- $\gamma$ , TNF- $\alpha$ , CD40L. Dependency on MR1 or IL-12 and IL-18 was assessed by antibody blocking. Activation of blood MAIT by blood macrophages (A) and activation of liver MAIT by THP1 cells (B) and autologous intrahepatic B cells (C) were analyzed. Data are mean  $\pm$  SEM (A) and median  $\pm$  interquartile range (B and C). In overlays: untreated (shaded); *E. coli* (solid line); *E. coli* +  $\alpha$ MR1 (dotted line). \* $p$  < 0.05; \*\*\* $p$  < 0.001 by paired  $t$  test. UT = untreated.

ducts in the portal tracts play an important role in protection against invading bacteria. MAIT cells respond to microbial antigens to provide protection at epithelial and mucosal surfaces. The liver regulates tolerance to food antigens and at the same time acts as a firewall to prevent intestinal microbes entering the systemic circulation [15]. We observed the overall MAIT cell frequency to be significantly higher in the normal liver compared to diseased liver, which differed from conventional CD4<sup>+</sup>, CD8<sup>+</sup>, or DN T cells. This suggests that MAIT cells are a major class of T cell recruited to the normal liver in order to act as a firewall and protect the biliary epithelium, therefore playing an important role in immune surveillance and homeostasis at the biliary mucosal barrier.

Immunohistochemical and confocal fluorescence staining of human liver tissue demonstrated that intrahepatic MAIT cells are present in the sinusoids. During an immune response to invading infection (e.g. bacterial infection) or during an inflammatory reaction, lymphocytes are recruited to liver tissue in response to combinations of locally expressed chemokines [14]. Hepatic inflammation leads to upregulation of IFN-inducible chemokines CXCL9/10/11, ligands for CXCR3, along with increased expression of the adhesion molecules ICAM-1 and VCAM-1 on inflamed human liver sinusoids [25,30,31]. We detected a significantly higher level of the chemokine receptor, CXCR3, on intrahepatic MAIT cells in the diseased state, as well as the presence of integrins LFA-1 and VLA-4. VLA-4 is essential





**Fig. 7. Bacterially-exposed primary human biliary epithelial cells activate both blood and liver MAIT cells in an MR1 dependent manner.** Biliary epithelial cells exposed to *E. coli* were co-cultured with sorted blood CD8<sup>+</sup> (A and B) or liver CD3<sup>+</sup> (C) T cells and activation of CD161<sup>+</sup> Va7.2<sup>+</sup> MAIT cells examined by flow cytometry for CD107a, IFN $\gamma$ , TNF $\alpha$ , CD40L. Dependency on MR1 or cytokines IL-12 and IL-18 was assessed using function-blocking antibodies. Data are mean  $\pm$  SEM (A and B) and median  $\pm$  interquartile range (C). In overlays: untreated (shaded); *E. coli* (solid line); *E. coli* +  $\alpha$ MR1 (dotted line). \**p* < 0.05; \*\*\**p* < 0.001 by t test. UT = untreated.

in CD8<sup>+</sup> MAIT cell infiltration to central nervous system in multiple sclerosis [32]. Thus, these molecules are involved in MAIT cell recruitment from blood. Intrahepatic V $\alpha$ 7.2<sup>+</sup> CD161<sup>+</sup> MAIT cells were highly enriched for IL-18R $\alpha$  expression in the livers, which would allow the cells to respond to high local levels of IL-18 in the inflamed hepatic microenvironment [12,33] and to interact with IL-18 secreted by Kupffer cells, which we observed to reside in the hepatic sinusoid [15,34].

We observed enrichment of MAIT cells in the liver compared to in blood by flow cytometry and also demonstrated by immunohistochemistry and confocal microscopy that they were concentrated preferentially in portal tracts, where the majority of CD3<sup>+</sup> CD161<sup>+</sup> Va7.2<sup>+</sup> cells localised in the peri-biliary regions, often in close contact with bile ducts. BEC in their normal state secrete the chemokine CCL20 and express the cell adhesion molecule E-cadherin, both of which are enhanced in the diseased state [21]. Intrahepatic MAIT cells in either normal or diseased state expressed the chemokine receptors CCR6 and CXCR6, and the

E-cadherin receptor, integrin  $\alpha$ E $\beta$ 7, which would allow them to migrate to the peri-biliary region in response to CCL20 and CXCL16 secreted by BEC [21,23,35]. We propose that CCR6<sup>+</sup>, CXCR6<sup>+</sup> and  $\alpha$ E $\beta$ 7-expressing intrahepatic MAIT cells are retained close to the bile ducts in steady state to provide protection against ascending bacterial infection from the gut [23,24,36,37]. VLA-4 on intrahepatic MAIT cells would also interact with VCAM-1 on the bile ducts, an interaction known to provide survival signals for lymphocytes [38]. Thus, activated, effector memory MAIT cells in the human liver are ready to protect the biliary mucosa in both the steady and diseased state [39,40].

A higher prevalence of intrahepatic MAIT cells around bile ducts was apparent by immunohistochemistry in PSC compared to other chronic liver diseases, including other biliary diseases such as PBC. This is an interesting observation as PSC is a biliary disease driven by mucosal T cells and associated with recurrent ascending infections [41]. Surprisingly, although MAIT cells are found in the gut and associated with inflammatory bowel disease

## Research Article

and colonic cancer [29,40], we detected very few cells that expressed the gut-homing integrin  $\alpha 4\beta 7$  or chemokine receptor CCR9, suggesting that hepatic MAIT cells are most likely not derived from the gut.

Increased frequency of MAIT cells in the hepatic parenchyma of patients with severe seronegative acute liver failure was also noted with immunohistochemistry. Seronegative hepatitis is characterised by a progressive, marked hepatocyte-necrosis leading to acute liver failure and is associated with bacterial translocation [42]. Therefore, in this scenario, the effector functions of MAIT cells might contribute to acute liver injury. Intrahepatic cells not only expressed CD26 constitutively [39,43], but also expressed CD39 and/or CD73, two ectoenzymes involved in immune regulation via generation of immunosuppressive adenosine [44]. Thus, hepatic MAIT cells include cells with both effector and regulatory functions which may confer either proinflammatory or immune-regulatory properties depending on the context and timing of the hepatic inflammation.

There needs to be local protection against intestinal pathogens at the biliary epithelium due to its continuity with the gut flora where both commensal and pathogenic bacteria reside [45]. Our data suggest that MAIT cells could play an important role in this. The recognition of antigen by MAIT cells is mediated via the MR1 molecule, which can present microbial vitamin B-derived compounds in its antigen-binding cleft [7]. MR1 is expressed on APC such as B cells [27]. We noted close localisation of B cells around bile ducts, and they mediated the activation of diseased liver-infiltrating MAIT cells in the presence of *E. coli*. Human BEC can also act as non-professional APC by expressing MHC and co-stimulatory molecules under some circumstances [19,46]. We found that indeed, both LI-MAIT and blood MAIT cells co-cultured with BEC exposed to *E. coli* degranulated and secreted IFN- $\gamma$ . Importantly, this mechanism was MR1-dependent but independent of IL-12 and IL-18 cytokines, despite the presence of these receptors on intrahepatic MAIT cells. This finding suggests that MAIT cells will only be fully activated in the presence of bacteria that have breached the epithelial barrier. Intrahepatic MAIT cells also secreted IL-17 and therefore likely possess not only an antibacterial function but also mucosa-regeneration properties similar to other Th17 cells [21,47].

CD40L upregulation was also observed on MAIT cells in response to bacterial presentation by BEC. CD40L on immune cells such as lymphocytes and macrophages can induce BEC apoptosis via epithelial CD40 [33,48,49]. CD40L upregulation was mediated both by *E. coli* presented by MR1 as well as IL-12 and IL-18, providing a mechanism through which MAIT cells could drive bile duct damage in inflammatory liver disease in the absence of infection, in a non-specific manner. Intrahepatic CD4<sup>+</sup>, CD8<sup>+</sup>, and DN MAIT cells also secreted TNF- $\alpha$  and IFN- $\gamma$ , which would be expected to contribute to their proinflammatory functional activity, and expressed granzyme B, which is crucial for cytolytic activity [11,39,50,51].

Taken together, our findings provide the first evidence that intrahepatic MAIT cells in the human liver can respond to bacterial antigens presented by the biliary epithelium, B cells or macrophages by expressing IFN- $\gamma$ , TNF- $\alpha$ , CD40L, and degranulating, and they have the capacity to secrete IL-17 upon activation. This suggests that intrahepatic MAIT cells play an important part in the biliary firewall that prevents bacteria from the gut entering the normal liver and then the systemic circulation via the bile ducts. We therefore propose that intrahepatic MAIT cells act as

guardians in biliary mucosa protection at steady normal state. Whether they play a role in the pathogenesis of inflammatory liver disease requires further study.

### Financial support

Dr Ye Htun Oo was funded by Clinician Scientist Award from the Medical Research Council (G1002552), United Kingdom; Queen Elizabeth Hospital Charity, United Kingdom; NIHR Biomedical Research Unit in liver disease Birmingham, United Kingdom and Bowel Disease Research Foundation Grant, United Kingdom.

Professor Paul Klenerman was funded by the Wellcome Trust (WT 091663MA), United Kingdom, NIHR Biomedical Research Centre Oxford, NIHR Senior Investigator award, Oxford Martin School, United Kingdom. MRC STOP-HCV consortium, United Kingdom and NIH NIAID 1U19AI082630-02, USA.

### Conflict of interest

The authors who have taken part in this study declared that they do not have anything to disclose regarding funding or conflict of interest with respect to this manuscript.

### Authors' contributions

HJ, YO, BW, AK, and PK designed the study. HJ, BW, AK, KP, KS, SR, ED, SH, DG and MB collected and analysed the data. RB and DW advised on histological staining and analysis. JR, TP and TI contributed samples collection and intellectual input. HJ and YO drafted and wrote the manuscript. HJ, BW, AK, DA, PK and YO revised the manuscript critically for intellectual content. All authors gave intellectual input to the study and approved the final version of the manuscript.

### Acknowledgements

We would like to thank Gillian Muirhead and Chantelle Endeley for their assistance with cell isolation and Dr Matthew Graeme MacKenzie for his help with cell sorting. We would also like to thank medical, surgical and anesthetic staff from UHB NHS Foundation Trust for their help with tissue acquirement.

### Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.jhep.2015.12.017>.

### References

- [1] Treiner E, Duban L, Bahram S, Radosavljevic M, Wanner V, Tilloy F, et al. Selection of evolutionarily conserved mucosal-associated invariant T cells by MR1. *Nature* 2003;422:164–169.
- [2] Reantragoon R, Corbett AJ, Sakala IG, Gherardin NA, Furness JB, Chen Z, et al. Antigen-loaded MR1 tetramers define T cell receptor heterogeneity in mucosal-associated invariant T cells. *J Exp Med* 2013;210:2305–2320.
- [3] Billerbeck E, Kang YH, Walker L, Lockstone H, Grafmueller S, Fleming V, et al. Analysis of CD161 expression on human CD8<sup>+</sup> T cells defines a distinct functional subset with tissue-homing properties. *Proc Natl Acad Sci U S A* 2010;107:3006–3011.

- [4] Kang YH, Seigel B, Bengsch B, Fleming VM, Billerbeck E, Simmons R, et al. CD161(+)CD4(+) T cells are enriched in the liver during chronic hepatitis and associated with co-secretion of IL-22 and IFN-gamma. *Front Immunol* 2012;3:346.
- [5] Martin E, Treiner E, Duban L, Guerri L, Laude H, Toly C, et al. Stepwise development of MAIT cells in mouse and human. *PLoS Biol* 2009;7:e54.
- [6] Huang S, Gilfillan S, Cella M, Miley MJ, Lantz O, Lybarger L, et al. Evidence for MR1 antigen presentation to mucosal-associated invariant T cells. *J Biol Chem* 2005;280:21183–21193.
- [7] Kjer-Nielsen L, Patel O, Corbett AJ, Le Nours J, Meehan B, Liu L, et al. MR1 presents microbial vitamin B metabolites to MAIT cells. *Nature* 2012;491:717–723.
- [8] Ussher JE, Klenerman P, Willberg CB. Mucosal-associated invariant T-cells: new players in anti-bacterial immunity. *Front Immunol* 2014;5:450.
- [9] Meierovics A, Yankelevich WJ, Cowley SC. MAIT cells are critical for optimal mucosal immune responses during in vivo pulmonary bacterial infection. *Proc Natl Acad Sci U S A* 2013;110:E3119–E3128.
- [10] Booth JS, Salerno-Goncalves R, Blanchard TG, Patil SA, Kader HA, Safta AM, et al. Mucosal-associated invariant T cells in the human gastric mucosa and blood: role in *Helicobacter pylori* infection. *Front Immunol* 2015;6:466.
- [11] Kurioka A, Ussher JE, Cosgrove C, Clough C, Fergusson JR, Smith K, et al. MAIT cells are licensed through granzyme exchange to kill bacterially sensitized targets. *Mucosal Immunol* 2014.
- [12] Ussher JE, Bilton M, Attwood E, Shadwell J, Richardson R, de Lara C, et al. CD161++ CD8+ T cells, including the MAIT cell subset, are specifically activated by IL-12+IL-18 in a TCR-independent manner. *Eur J Immunol* 2014;44:195–203.
- [13] Le Bourhis L, Martin E, Peguillet I, Guihot A, Froux N, Core M, et al. Antimicrobial activity of mucosal-associated invariant T cells. *Nat Immunol* 2010;11:701–708.
- [14] Oo YH, Adams DH. The role of chemokines in the recruitment of lymphocytes to the liver. *J Autoimmun* 2010;34:45–54.
- [15] Balmer ML, Slack E, de Gottardi A, Lawson MA, Hapfelmeier S, Miele L, et al. The liver may act as a firewall mediating mutualism between the host and its gut commensal microbiota. *Sci Transl Med* 2014;6: 237ra266.
- [16] Tang XZ, Jo J, Tan AT, Sandalova E, Chia A, Tan KC, et al. IL-7 licenses activation of human liver intrasinusoidal mucosal-associated invariant T cells. *J Immunol* 2013;190:3142–3152.
- [17] Hubscher SG. Histological grading and staging in chronic hepatitis: clinical applications and problems. *J Hepatol* 1998;29:1015–1022.
- [18] Racanelli V, Rehermann B. The liver as an immunological organ. *Hepatology* 2006;43:S54–S62.
- [19] Schruppf E, Tan C, Karlsen TH, Sponheim J, Bjorkstrom NK, Sundnes O, et al. The biliary epithelium presents antigens to and activates natural killer T cells. *Hepatology* 2015.
- [20] Le Bourhis L, Dusseaux M, Bohineust A, Bessoles S, Martin E, Premel V, et al. MAIT cells detect and efficiently lyse bacterially-infected epithelial cells. *PLoS Pathog* 2013;9:e1003681.
- [21] Oo YH, Banz V, Kavanagh D, Liaskou E, Withers DR, Humphreys E, et al. CXCR3-dependent recruitment and CCR6-mediated positioning of Th-17 cells in the inflamed liver. *J Hepatol* 2012;57:1044–1051.
- [22] Miley MJ, Truscott SM, Yu YY, Gilfillan S, Fremont DH, Hansen TH, et al. Biochemical features of the MHC-related protein 1 consistent with an immunological function. *J Immunol* 2003;170:6090–6098.
- [23] Heydtmann M, Lalor PF, Eksteen JA, Hubscher SG, Briskin M, Adams DH. CXC chemokine ligand 16 promotes integrin-mediated adhesion of liver-infiltrating lymphocytes to cholangiocytes and hepatocytes within the inflamed human liver. *J Immunol* 2005;174:1055–1062.
- [24] Grant AJ, Lalor PF, Salmi M, Jalkanen S, Adams DH. Homing of mucosal lymphocytes to the liver in the pathogenesis of hepatic complications of inflammatory bowel disease. *Lancet* 2002;359:150–157.
- [25] Curbishley SM, Eksteen B, Gladue RP, Lalor P, Adams DH. CXCR3 activation promotes lymphocyte transendothelial migration across human hepatic endothelium under fluid flow. *Am J Pathol* 2005;167:887–899.
- [26] Eksteen B, Grant AJ, Miles A, Curbishley SM, Lalor PF, Hubscher SG, et al. Hepatic endothelial CCL25 mediates the recruitment of CCR9+ gut-homing lymphocytes to the liver in primary sclerosing cholangitis. *J Exp Med* 2004;200:1511–1517.
- [27] Salerno-Goncalves R, Rezwan T, Szein MB. B cells modulate mucosal associated invariant T cell immune responses. *Front Immunol* 2014;4:511.
- [28] Le Bourhis L, Mburu YK, Lantz O. MAIT cells, surveyors of a new class of antigen: development and functions. *Curr Opin Immunol* 2013;25:174–180.
- [29] Serriari NE, Eoche M, Lamotte L, Lion J, Fumery M, Marcelo P, et al. Innate mucosal-associated invariant T (MAIT) cells are activated in inflammatory bowel diseases. *Clin Exp Immunol* 2014;176:266–274.
- [30] Lalor PF, Adams DH. Adhesion of lymphocytes to hepatic endothelium. *Mol Pathol* 1999;52:214–219.
- [31] Lalor PF, Clements JM, Pigott R, Humphries MJ, Spragg JH, Nash GB. Association between receptor density, cellular activation, and transformation of adhesive behavior of flowing lymphocytes binding to VCAM-1. *Eur J Immunol* 1997;27:1422–1426.
- [32] Willing A, Leach OA, Ufer F, Attfield KE, Steinbach K, Kursawe N, et al. CD8(+) MAIT cells infiltrate into the CNS and alterations in their blood frequencies correlate with IL-18 serum levels in multiple sclerosis. *Eur J Immunol* 2014;44:3119–3128.
- [33] Alabraba EB, Lai V, Boon L, Wigmore SJ, Adams DH, Afford SC. Coculture of human liver macrophages and cholangiocytes leads to CD40-dependent apoptosis and cytokine secretion. *Hepatology* 2007.
- [34] Ikeda A, Aoki N, Kido M, Iwamoto S, Nishiura H, Maruoka R, et al. Progression of autoimmune hepatitis is mediated by IL-18-producing dendritic cells and hepatic CXCL9 expression in mice. *Hepatology* 2014;60:224–236.
- [35] Grant AJ, Lalor PF, Hubscher SG, Briskin M, Adams DH. MAdCAM-1 expressed in chronic inflammatory liver disease supports mucosal lymphocyte adhesion to hepatic endothelium (MAdCAM-1 in chronic inflammatory liver disease). *Hepatology* 2001;33:1065–1072.
- [36] Eksteen B, Miles A, Curbishley SM, Tselepis C, Grant AJ, Walker LS, et al. Epithelial inflammation is associated with CCL28 production and the recruitment of regulatory T cells expressing CCR10. *J Immunol* 2006;177:593–603.
- [37] Shields PL, Morland CM, Salmon M, Qin S, Hubscher SG, Adams DH. Chemokine and chemokine receptor interactions provide a mechanism for selective T cell recruitment to specific liver compartments within hepatitis C-infected liver. *J Immunol* 1999;163:6236–6243.
- [38] Afford SC, Humphreys EH, Reid DT, Russell CL, Banz VM, Oo Y, et al. Vascular cell adhesion molecule 1 expression by biliary epithelium promotes persistence of inflammation by inhibiting effector T-cell apoptosis. *Hepatology* 2014;59:1932–1943.
- [39] Dusseaux M, Martin E, Serriari N, Peguillet I, Premel V, Louis D, et al. Human MAIT cells are xenobiotic-resistant, tissue-targeted, CD161hi IL-17-secreting T cells. *Blood* 2011;117:1250–1259.
- [40] Sundstrom P, Ahlmanner F, Akeus P, Sundquist M, Alsen S, Yrliid U, et al. Human mucosa-associated invariant T cells accumulate in colon adenocarcinomas but produce reduced amounts of IFN-gamma. *J Immunol* 2015;195:3472–3481.
- [41] Hirschfield GM, Karlsen TH, Lindor KD, Adams DH. Primary sclerosing cholangitis. *Lancet* 2013;382:1587–1599.
- [42] Wigg AJ, Gunson BK, Mutimer DJ. Outcomes following liver transplantation for seronegative acute liver failure: experience during a 12-year period with more than 100 patients. *Liver Transpl* 2005;11:27–34.
- [43] Sharma PK, Wong EB, Napier RJ, Bishai WR, Ndung'u T, Kasprovicz VO, et al. High expression of CD26 accurately identifies human bacteria-reactive MR1-restricted MAIT cells. *Immunology* 2015;145:443–453.
- [44] Deaglio S, Dwyer KM, Gao W, Friedman D, Usheva A, Erat A, et al. Adenosine generation catalyzed by CD39 and CD73 expressed on regulatory T cells mediates immune suppression. *J Exp Med* 2007;204:1257–1265.
- [45] Ma HD, Wang YH, Chang C, Gershwin ME, Lian ZX. The intestinal microbiota and microenvironment in liver. *Autoimmun Rev* 2015;14:183–191.
- [46] Rong GH, Yang GX, Ando Y, Zhang W, He XS, Leung PS, et al. Human intrahepatic biliary epithelial cells engulf blebs from their apoptotic peers. *Clin Exp Immunol* 2013;172:95–103.
- [47] Harada K, Shimoda S, Sato Y, Isse K, Ikeda H, Nakanuma Y. Periductal interleukin-17 production in association with biliary innate immunity contributes to the pathogenesis of cholangiopathy in primary biliary cirrhosis. *Clin Exp Immunol* 2009;157:261–270.
- [48] Afford SC, Rhandawa S, Eliopoulos AG, Hubscher SG, Young LS, Adams DH. CD40 activation induces apoptosis in cultured human hepatocytes via induction of cell surface FasL expression and amplifies Fas mediated hepatocyte death during allograft rejection. *J Exp Med* 1999;189:441–446.
- [49] Humphreys EH, Williams KT, Adams DH, Afford SC. Primary and malignant cholangiocytes undergo CD40 mediated Fas dependent apoptosis, but are insensitive to direct activation with exogenous Fas ligand. *PLoS One* 2010;5: e14037.
- [50] Walker LJ, Kang YH, Smith MO, Tharmalingham H, Ramamurthy N, Fleming VM, et al. Human MAIT and CD8alpha cells develop from a pool of type-17 precommitted CD8+ T cells. *Blood* 2012;119:422–433.
- [51] Fergusson JR, Smith KE, Fleming VM, Rajorija N, Newell EW, Simmons R, et al. CD161 defines a transcriptional and functional phenotype across distinct human T cell lineages. *Cell Rep* 2014;9:1075–1088.