

# Decomposition of bounded degree graphs into $C_4$ -free subgraphs

Kang, Ross; Perarnau, Guillem

DOI:

[10.1016/j.ejc.2014.09.009](https://doi.org/10.1016/j.ejc.2014.09.009)

License:

Other (please provide link to licence statement)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Kang, R & Perarnau, G 2015, 'Decomposition of bounded degree graphs into  $C_4$ -free subgraphs', *European Journal of Combinatorics*, vol. 44, no. A, pp. 99-105. <https://doi.org/10.1016/j.ejc.2014.09.009>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



Contents lists available at ScienceDirect

European Journal of Combinatorics

journal homepage: [www.elsevier.com/locate/ejc](http://www.elsevier.com/locate/ejc)

# Decomposition of bounded degree graphs into $C_4$ -free subgraphs<sup>☆</sup>



Ross J. Kang<sup>a</sup>, Guillem Perarnau<sup>b</sup>

<sup>a</sup> Applied Stochastics, IMAPP, Radboud University Nijmegen, Netherlands

<sup>b</sup> School of Computer Science, McGill University, Montreal, Canada

## ARTICLE INFO

### Article history:

Received 8 August 2014

Accepted 23 September 2014

Available online 15 October 2014

## ABSTRACT

We prove that every graph with maximum degree  $\Delta$  admits a partition of its edges into  $O(\sqrt{\Delta})$  parts (as  $\Delta \rightarrow \infty$ ) none of which contains  $C_4$  as a subgraph. This bound is sharp up to a constant factor. Our proof uses an iterated random colouring procedure.

© 2014 Elsevier Ltd. All rights reserved.

## 1. Introduction

In this paper we consider the following question.

*Given a graph  $G = (V, E)$  with maximum degree  $\Delta$ , into how few parts can we partition  $E$  so that no part has a  $C_4$  subgraph?*

More generally, for any graph  $H$  with at least two edges, given  $G = (V, E)$  and a map  $f : E \rightarrow [m]$  for some positive integer  $m$ , we call  $f$  an  $H$ -free (edge-)colouring of  $G$  with  $m$  colours if there is no  $i \in [m]$  such that the graph  $(V, f^{-1}(i))$  contains  $H$  as a subgraph. (Note that this is not necessarily a proper colouring unless  $H$  is a two-edge path.) Let  $\phi_H(G)$  be the least  $m$  such that  $G$  admits an  $H$ -free colouring with  $m$  colours.

Using this notation, the above asks specifically about  $\phi_{C_4}$ , and in answer we show the following.

**Theorem 1.** *For every graph  $G$  with maximum degree  $\Delta$ ,  $\phi_{C_4}(G) = O(\sqrt{\Delta})$  as  $\Delta \rightarrow \infty$ .*

In words, every graph with maximum degree  $\Delta$  admits a partition of its edges (also called a decomposition) into  $O(\sqrt{\Delta})$   $C_4$ -free subgraphs.

<sup>☆</sup> This work began in April 2014 during the Workshop on Structural Graph Theory at McGill's Bellairs Institute. We warmly thank the organisers for the collaborative opportunity.

E-mail addresses: [ross.kang@gmail.com](mailto:ross.kang@gmail.com) (R.J. Kang), [guillem.perarnaullobet@mcgill.ca](mailto:guillem.perarnaullobet@mcgill.ca) (G. Perarnau).

Let  $K_n$  be the complete graph on  $n$  vertices. By an upper bound on the size of every colour class in an  $H$ -free colouring of  $K_{\Delta+1}$ , we have that

$$\phi_H(K_{\Delta+1}) \geq \frac{\binom{\Delta+1}{2}}{\text{ex}(\Delta + 1, H)}, \tag{1}$$

where  $\text{ex}(n, H)$  as usual denotes the maximum number of edges in an  $H$ -free graph on  $n$  vertices. Then it follows from an old result of Erdős [3] on the extremal number of  $C_4$  (see [13] for context and more detailed results) that  $\phi_{C_4}(K_{\Delta+1}) = \Omega(\sqrt{\Delta})$ . This not only shows [Theorem 1](#) to be best possible up to a constant factor, but also foreshadows a central role of the complete graph.

For a broader context, [Theorem 1](#) may be understood in terms of the degree Ramsey numbers as first considered in the 1970s by Burr, Erdős and Lovász [1]—they studied these numbers for complete graphs and stars. The more general setting for other graphs was recently revisited in [7]. The question we posed at the beginning is equivalent to finding the multicolour degree Ramsey number of  $C_4$ . In [6] it was shown that  $\phi_{C_4}(G) = O(\Delta^{9/14})$  for graphs of maximum degree  $\Delta$ , and the authors asked for the right order of growth. [Theorem 1](#) settles this.

We prove [Theorem 1](#) in [Section 3](#) by using the probabilistic method. In particular, we use an iterated random colouring procedure. At each step of the procedure we identify a collection of large  $C_4$ -free colour classes, the removal of which significantly reduces the maximum degree of the graph (see [Corollary 7](#)). In the proof, we deliberately make little effort to optimise constants, but we note here that it is possible to obtain a factor less than 45 in [Theorem 1](#) by being more careful at a few points.

Recently, together with Bruce Reed [12], the second author proved that every  $\Delta$ -regular graph  $G$  contains a spanning  $C_4$ -free graph with minimum degree  $\Omega(\sqrt{\Delta})$ . This result has some similarity to our [Corollary 7](#), where instead of looking at the minimum degree of the resulting subgraph, they look at the minimum degree of a given colour class. In a way that is analogous to their work, we essentially reduce our considered problem to the determination of  $\text{ex}(\Delta + 1, C_4)$ . (For us, this is reminiscent of the relationship between independence number and chromatic number found in other extremal colouring problems.)

More generally, we ask the following.

*For any graph  $H$  with at least two edges, is it true that  $\phi_H(G) = O(\phi_H(K_{\Delta+1}))$  for every graph  $G$  with maximum degree  $\Delta$ ?*

Otherwise stated, we ask if the complete graph on  $\Delta + 1$  vertices is essentially the hardest graph to  $H$ -free colour among all the graphs with maximum degree  $\Delta$ .

Trivially, this holds for  $H$  a two-edge path. [Theorem 1](#) shows this to be true for  $H = C_4$ . Using the methods in the proof of [Theorem 1](#), it is possible to confirm this for other bipartite graphs  $H$  such as cycles of order twice a prime, or complete bipartite graphs. Moreover, for every  $g \geq 4$ , we can also edge-colour graphs of maximum degree  $\Delta$ , each colour class having girth at least  $g$ , with an asymptotically tight number of colours. We encourage the reader to consult [12] to see a concrete discussion of how [Theorem 4](#) can be used to upper bound  $\phi_H(G)$  for other bipartite graphs  $H$ .

Another problem strongly related to our result (via the above displayed question) is to determine  $\phi_H(K_n)$ . Inequality (1) provides a lower bound on  $\phi_H(K_n)$  in terms of  $\text{ex}(n, H)$ . This prompts us to ask for which graphs  $H$  we have  $\phi_H(K_n) = O(n^2/\text{ex}(n, H))$  (as  $n \rightarrow \infty$ ).

This last statement does not hold if  $H$  is not bipartite. On the one hand, Turán’s theorem implies that  $\text{ex}(n, H) = \Omega(n^2)$ . On the other hand, it can be shown in this case that  $\phi_H(K_n) = \Omega(\log \log n)$ . First observe that  $\phi_H(K_n) \geq \phi_{H'}(K_n)$  for any  $H \subseteq H'$ . Write  $|V(H)| = k$  for some fixed  $k \geq 3$ . The Erdős–Szekeres bound on two-colour Ramsey numbers gives that  $R(k, \ell) \leq \binom{k+\ell-2}{k-1} = O(\ell^{k-1})$ , so every  $K_k$ -free graph of order  $n$  has an independent set of size  $\Omega(n^{1/(k-1)})$ . Let  $m = \phi_{K_k}(K_n)$  and let  $G_1, \dots, G_m$  denote the colour classes of a  $K_k$ -free colouring of  $K_n$  with  $m$  colours. Beginning with  $V_0 = V$ , define  $V_i$  to be a maximum independent set of  $G_i[V_{i-1}]$  for every  $0 < i \leq m$ . Then  $|V_i| = \Omega(n^{(k-1)^{-i}})$ , which implies  $m = \Omega(\log \log n)$ , as claimed.

Nevertheless,  $\phi_H(K_n) = O(n^2/\text{ex}(n, H))$  for some bipartite graphs  $H$  such as  $C_4$  [2,5],  $C_6$  and  $C_{10}$  [8].

Bounding or determining the Turán number of bipartite graphs is a central problem in extremal graph theory (see again [13] or, more generally, [4]), so determining for bipartite  $H$  the right order of  $\phi_H(G)$  in terms of  $\Delta(G)$  might be difficult in general.

## 2. Some probabilistic tools

For our proof we need the following lemmas, the uses of which are covered extensively in [9].

**Lemma 2** (Simple Concentration Bound). *Let  $X$  be a random variable determined by  $n$  trials  $T_1, \dots, T_n$  such that for each  $i$ , and any two possible sequences of outcomes  $t_1, \dots, t_i, \dots, t_n$  and  $t_1, \dots, t'_i, \dots, t_n$ ,*

$$|X(t_1, \dots, t_i, \dots, t_n) - X(t_1, \dots, t'_i, \dots, t_n)| \leq c.$$

Then

$$\Pr(|X - \mathbb{E}(X)| > t) \leq 2e^{-t^2/(2c^2n)}.$$

**Lemma 3** (Lovász Local Lemma). *Consider a set  $\mathcal{E}$  of events such that for each  $E \in \mathcal{E}$*

- $\Pr(E) \leq p < 1$ , and
- $E$  is mutually independent from the set of all but at most  $D$  of other events.

If  $4pD \leq 1$ , then with positive probability none of the events in  $\mathcal{E}$  occur.

## 3. Proof of Theorem 1

Before proceeding with the main proof, let us first consider the complete graph  $K_{\Delta+1}$ . It was shown in the 1970s independently by Chung and Graham [2] and by Irving [5] that, if  $\Delta = p^2 + p + 1$  for some prime power  $p$ , then  $\phi_{C_4}(K_{\Delta+1}) \leq p + 1$ .

By the density of the primes, it follows easily that

$$\phi_{C_4}(K_{\Delta+1}) \leq \lceil 2\sqrt{\Delta} \rceil, \tag{2}$$

for all large enough  $\Delta$ . We later use this in the proof of Theorem 1.

Given a graph  $G = (V, E)$ , we say that a map  $f : V \rightarrow [m]$  is 1-frugal if it holds for all  $i \in [m]$  and  $v \in V$  that  $|f^{-1}(i) \cap N(v)| \leq 1$ . We may alternatively view a 1-frugal map as a vertex colouring such that every neighbourhood is rainbow. The engine in our proof of Theorem 1 is the following result.

**Theorem 4.** *Let  $G = (V, E)$  be a graph with maximum degree  $\Delta$  and minimum degree  $\delta \geq \log^2 \Delta$  with  $\Delta$  sufficiently large. For every  $\alpha > 16$ , there exist  $\beta = \beta(\alpha) > 0$ , a spanning subgraph  $H$  and a (vertex)  $(2\lceil \alpha \Delta \rceil)$ -colouring  $\chi$  such that*

- $d_H(v) \geq \beta d_G(v)$  for every  $v \in V$  and
- $\chi$  is 1-frugal and proper in  $H$ .

**Proof.** First observe that there exists a spanning bipartite subgraph  $H_0$  such that  $d_{H_0}(v) \geq d_G(v)/2$  for every vertex  $v \in V$ . (Consider  $H_0$  to be a subgraph induced by a maximum edge-cut. This subgraph is clearly bipartite, so let  $V = A \cup B$  denote its bipartition. Suppose that  $d_{H_0}(v) < d_G(v)/2$  for some  $v \in V$ . We can assume that  $v \in A$ . Then the number of edges between  $A \setminus \{v\}$  and  $B \cup \{v\}$  is strictly larger than the number of edges between  $A$  and  $B$ , contradicting the maximum edge-cut assumption.) While colouring  $V$ , we also construct  $H$  as a subgraph of  $H_0$ , by sequentially removing edges. The colouring has two consecutive rounds, the first of which colours the vertices of  $A$ , the second colours  $B$ .

We begin by describing the first round colouring  $A$ ; this itself has two phases, a probabilistic one followed by a deterministic one.

- Phase I. Colour each vertex  $a \in A$  with a colour  $\chi_0(a)$  chosen uniformly at random from  $[[\alpha \Delta]]$ . From  $\chi_0$  we obtain a partial colouring  $\chi_1$  of  $A$  as follows. We uncolour a vertex  $a \in A$  if

$$|\{b \in N_{H_0}(a) : \exists a' \in N_{H_0}(b) \setminus \{a\}, \chi_0(a') = \chi_0(a)\}| \geq \frac{d_{H_0}(a)}{\sqrt{\alpha}}; \tag{3}$$

that is, if  $a$  certifies that too many of its neighbours have another neighbour in  $A$  with colour  $\chi_0(a)$ . Otherwise, let  $\chi_1(a) = \chi_0(a)$  and remove all edges from  $a$  to  $b \in N_{H_0}(a)$  where  $b$  is incident to  $a'$  with  $a \neq a'$  and  $\chi_0(a') = \chi_0(a)$ . Let  $H_1$  be the subgraph obtained after removing all these edges. We have ensured that, for any  $\chi_1$ -coloured  $a \in A$  and any  $b \in N_{H_1}(a)$ ,  $a$  is the only neighbour of  $b$  coloured  $\chi_1(a)$ .

We stress that condition (3) is always checked on the initial colouring  $\chi_0$  and that all the vertices that are uncoloured lose their colour simultaneously.

- Phase II. Order the uncoloured vertices  $a_1, \dots, a_{s-1}$ . For  $i = 1, 2, \dots, s - 1$ , let  $c \in [\lceil \alpha \Delta \rceil]$  be the colour minimising

$$|\{b \in N_{H_i}(a_i) : \exists a' \in N_{H_i}(b) \setminus \{a_i\}, \chi_i(a') = c\}|.$$

Delete from  $H_i$  all edges  $a_i b$  such that there exists  $a' \in N_{H_i}(b) \setminus \{a_i\}$  with  $\chi_i(a') = c$  and call the resulting subgraph  $H_{i+1}$ . Let  $\chi_{i+1}$  be the partial colouring obtained from  $\chi_i$  by also assigning  $a_i$  the colour  $c$ .

First we show that  $d_{H_s}(a)$  is large for every  $a \in A$ .

**Claim 5.** For every  $a \in A$

$$d_{H_s}(a) \geq \left(1 - \frac{1}{\sqrt{\alpha}}\right) d_{H_0}(a).$$

**Proof.** Note that we only delete edges incident to  $a$  at a step in the procedure when  $a$  retains its colour. If  $a \in A$  retained its colour in the probabilistic phase, we can conclude  $d_{H_s}(a) = d_{H_1}(a) \geq (1 - 1/\sqrt{\alpha})d_{H_0}(a)$ , since by (3), conditioned on retaining the colour  $\chi_0(a)$ , we delete at most  $d_{H_0}(a)/\sqrt{\alpha}$  edges incident to  $a$ . Otherwise,  $a = a_i$  for some  $i \in [s - 1]$ , coloured in the deterministic phase, and since there are at most  $d_{H_0}(a_i)\Delta$  edges incident to  $N_{H_i}(a_i)$ , there exists a colour  $c \in [\lceil \alpha \Delta \rceil]$  such that

$$|\{b \in N_{H_i}(a_i) : \exists a' \in N_{H_i}(b) \setminus \{a_i\}, \chi_0(a') = c\}| \leq \frac{d_{H_0}(a_i)\Delta}{\lceil \alpha \Delta \rceil} \leq \frac{d_{H_0}(a_i)}{\alpha}.$$

Thus  $d_{H_s}(a_i) = d_{H_{i+1}}(a_i) \geq (1 - 1/\alpha)d_{H_0}(a_i) \geq (1 - 1/\sqrt{\alpha})d_{H_0}(a_i)$ .  $\square$

**Claim 6.** There exist a spanning subgraph  $H'$  and a  $\lceil \alpha \Delta \rceil$ -colouring  $\chi'$  of  $A$  such that for every  $a \in A$

$$d_{H'}(a) \geq \left(1 - \frac{1}{\sqrt{\alpha}}\right) d_{H_0}(a),$$

and for every  $b \in B$

$$d_{H'}(b) \geq \left(1 - \frac{4}{\sqrt{\alpha}}\right) d_{H_0}(b),$$

and  $N_{H'}(b)$  is rainbow in  $\chi'$ .

**Proof.** Note that the subgraph  $H_s$  and colouring  $\chi_s$  we have constructed are random objects, so it suffices to show that they satisfy the required properties with positive probability (when  $\Delta$  is large enough). Note that two of the properties are guaranteed by the construction of  $H_s$  and  $\chi_s$  (partly using Claim 5). It only remains to check the degree condition from  $B$ .

Let  $b \in B$ . Observe that the number of coloured neighbours of  $b$  under the colouring  $\chi_s$  is at least the number of coloured neighbours of  $b$  under  $\chi_{s-1}$  (and so on), since in the deterministic phase an edge  $ab$  can only be deleted in a step when  $a$  is coloured. Thus we can show that  $d_{H_s}(b)$  is large by showing that the degree of  $b$  in  $H_1$  to the set of vertices coloured by  $\chi_1$  is large.

For a given  $a \in N_{H_0}(b)$ , let  $E_1$  be the event that there exists  $a' \in N_{H_0}(b) \setminus \{a\}$  such that  $\chi_0(a') = \chi_0(a)$  and let  $E_2$  be the event that  $a$  becomes uncoloured (as governed by the condition in (3)). Let  $Y_b$  be the random variable that counts the number of vertices  $a \in N_{H_0}(b)$  for which  $E_1$  holds. Let  $Z_b$  be the random variable that counts the number of vertices  $a \in N_{H_0}(b)$  for which  $E_2$  holds but  $E_1$  does not.

Notice that these random variables count disjoint sets of vertices. By the observation of the previous paragraph,

$$d_{H_b}(b) \geq d_{H_0}(b) - Y_b - Z_b.$$

We estimate  $Z_b$  by studying another random variable. We say that the colour  $c$  is *dangerous* for  $a$  if

$$|\{b' \in N_{H_0}(a) \setminus \{b\} : \exists a' \in N_{H_0}(b') \setminus \{a\}, \chi_0(a') = c\}| \geq \frac{d_{H_0}(a)}{\sqrt{\alpha}} - 1.$$

For a given  $a \in N_{H_0}(b)$ , let  $E_3$  be the event that  $a$  receives a dangerous colour. Let  $Z'_b$  be the random variable that counts the number of vertices  $a \in N_{H_0}(b)$  for which  $E_3$  holds but  $E_1$  does not.

The following observation is important: if  $a$  is counted by  $Z_b$  it means that  $a$  becomes uncoloured and  $\chi_0(a)$  is a unique colour within  $N_{H_0}(b)$ . Then  $a$  must have been assigned a dangerous colour since for every vertex  $a' \in N_{H_0}(b) \setminus \{a\}$ ,  $\chi_0(a') \neq \chi_0(a)$ , and thus  $a'$  does not change the number of  $b' \in N_{H_0}(a) \setminus \{b\}$  that have colour  $\chi_0(a)$  in  $N_{H_0}(b') \setminus \{a\}$ . Hence  $Z_b \leq Z'_b$  and it is enough to verify that not too many vertices receive dangerous colours.

We are going to show that  $X_b = Y_b + Z'_b$  is concentrated given any fixed colouring in  $A \setminus N_{H_0}(b)$ . This, together with an upper bound on the conditional expectation of  $X_b$ , suffices to establish an upper bound on  $X_b$  that holds unconditionally. During the rest of the proof, we will assume that all the random variables are conditioned to the colouring in  $A \setminus N_{H_0}(b)$ .

First we deal with the expected value of  $Y_b$ . Consider  $a \in N_{H_0}(b)$ . Observe that at most  $d_{H_0}(b) - 1 \leq \Delta$  colours appear in  $N_{H_0}(b) \setminus \{a\}$  under the random colouring  $\chi_0$ . Then the probability that  $a$  does not have a unique colour in  $N_{H_0}(b)$  is at most  $(d_{H_0}(b) - 1) / \lceil \alpha \Delta \rceil \leq 1/\alpha$ , and so  $\mathbb{E}(Y_b) \leq d_{H_0}(b)/\alpha$ .

Second we compute the expected value of  $Z'_b$ . Since the maximum degree of  $H_0$  is  $\Delta$  and a colour is considered dangerous if at least  $d_{H_0}(a)/\sqrt{\alpha} - 1$  many vertices  $b' \in N_{H_0}(a) \setminus \{b\}$  already have it in  $N_{H_0}(b') \setminus \{a\}$ , there are at most  $d_{H_0}(a)\Delta / (\Delta/\sqrt{\alpha} - 1) \leq 2\sqrt{\alpha}\Delta$  dangerous colours for  $a$ . Thus  $a$  receives a dangerous colour with probability at most  $2\sqrt{\alpha}\Delta / \lceil \alpha \Delta \rceil \leq 2/\sqrt{\alpha}$ . So  $\mathbb{E}(Z'_b) \leq 2d_{H_0}(b)/\sqrt{\alpha}$ .

Then

$$\mathbb{E}(X_b) = \mathbb{E}(Y_b) + \mathbb{E}(Z'_b) \leq \left( \frac{1}{\alpha} + \frac{2}{\sqrt{\alpha}} \right) d_{H_0}(b) \leq \frac{3d_{H_0}(b)}{\sqrt{\alpha}}.$$

We can now apply the Simple Concentration Bound to show that  $X_b$  is concentrated with polynomially small probability. Note that changing the colour of  $a \in N_{H_0}(b)$  can change by at most two the value of  $X_b$ :

- it can change by at most two the number of vertices that are unique in their colour class (including  $a$  itself), and
- it can change by at most one the number of vertices that receive a dangerous colour and do not satisfy  $E_1$ , since the colour classes are prescribed by the colouring given to  $A \setminus N_{H_0}(b)$ .

Moreover,  $X_b$  conditioned on the colouring of  $A \setminus N_{H_0}(b)$  is determined by at most  $d_{H_0}(b)$  many different trials. By the Simple Concentration Bound with the choices  $c = 2$  and  $n = d_{H_0}(b)$ , we have that  $X_b$  conditioned to any colouring in  $A \setminus N_{H_0}(b)$  is unlikely to be large:

$$\begin{aligned} \Pr \left( X_b \geq \frac{4d_{H_0}(b)}{\sqrt{\alpha}} \right) &\leq \Pr \left( X_b - \mathbb{E}(X_b) \geq \frac{d_{H_0}(b)}{\sqrt{\alpha}} \right) \\ &\leq 2 \exp \left( - \frac{d_{H_0}^2(b)}{8\alpha \cdot d_{H_0}(b)} \right) = e^{-\Omega(d_{H_0}(b))} = o(\Delta^{-6}). \end{aligned}$$

In the last equality we used that  $d_{H_0}(b) = \Omega(\log^2 \Delta)$ . Thus the previous inequality also holds for the unconditioned random variable  $X_b$ .

Observe that  $X_b$  depends on the vertices at a distance of at most 3 from  $b$ ; the fact that  $a \in N_{H_0}(b)$  retains its colour depends only on the colours assigned to vertices at a distance of 2 from  $a$ . Thus every event corresponding to  $X_b$  is mutually independent from the set of events corresponding to  $X_{b'}$  with

$b'$  at a distance of more than 6 from  $b$ , the Lovász Local Lemma yields that with positive probability  $X_b \leq 4d_{H_0}(b)/\sqrt{\alpha}$  for every  $b \in B$ . This completes the proof of the claim.  $\square$

In the second round, we can apply the same argument to colour the vertices of  $B$  using the subgraph  $H'$ . By Claim 6 and recalling that  $\alpha > 16$ , this graph has minimum degree at least  $(1 - 4/\sqrt{\alpha})\delta(H_0) = \Omega(\log^2 \Delta)$  and maximum degree at most  $\Delta$ . So we can apply the same procedure (and claims) to colour  $B$  with a new set of  $\lceil \alpha \Delta \rceil$  colours. Combined with the colouring  $\chi'$  of  $A$ , in this way we obtain a subgraph  $H \subseteq H'$  and a  $(2\lceil \alpha \Delta \rceil)$ -colouring  $\chi$  of  $V$  such that

- for every  $v \in V$

$$d_H(v) \geq \left(1 - \frac{4}{\sqrt{\alpha}}\right)^2 d_{H_0}(v) \geq \left(1 - \frac{4}{\sqrt{\alpha}}\right)^2 \frac{d_G(v)}{2}, \quad \text{and}$$

- $\chi$  is a 1-frugal proper colouring of  $H$ .

This proves the theorem with the choice  $\beta = \frac{1}{2} \left(1 - 4/\sqrt{\alpha}\right)^2$ .  $\square$

**Corollary 7.** *Let  $G$  be a graph with maximum degree  $\Delta$  and minimum degree  $\delta \geq \log^2 \Delta$  with  $\Delta$  sufficiently large. For every  $\alpha > 16$ , there exist  $\beta = \beta(\alpha) > 0$  and  $\ell \leq \lceil 2\sqrt{2\lceil \alpha \Delta \rceil} \rceil$  many  $C_4$ -free disjoint spanning subgraphs  $G_1, \dots, G_\ell$  such that for all  $v \in V$*

$$\sum_{i=1}^{\ell} d_{G_i}(v) \geq \beta d_G(v).$$

**Proof.** We use the subgraph  $H$  and the colouring  $\chi$  guaranteed by Theorem 4 to find many  $C_4$ -free spanning subgraphs. By (2), for any sufficiently large  $t$  there exists a decomposition of  $K_t$  into  $C_4$ -free subgraphs  $\mathcal{G}_1, \dots, \mathcal{G}_{\lceil 2\sqrt{t} \rceil}$ . Consider  $t = 2\lceil \alpha \Delta \rceil$  and for any  $i \in [\lceil 2\sqrt{t} \rceil]$  construct  $G_i$  as follows:

- $V(G_i) = V(G)$  and
- $uv \in E(G_i)$  if and only if  $uv \in E(H)$  and  $\chi(u)\chi(v) \in E(\mathcal{G}_i)$ .

These subgraphs  $G_i$  are disjoint and, since  $H$  contains no monochromatic edge, each edge of  $H$  appears in exactly one subgraph  $G_i$ . So the minimum degree condition for  $H$  implies the minimum degree sum condition demanded here. Moreover, each  $G_i$  is  $C_4$ -free: by  $\chi$  being 1-frugal and proper, all 4-cycles in  $H$  are rainbow; and if  $G_i$  contains such a 4-cycle  $C$ , then the colours  $\chi(C)$  form a 4-cycle in  $\mathcal{G}_i$ .  $\square$

Besides the above, we need the following bound on arboricity by degeneracy (which follows, for instance, from the folkloric Proposition 3.1 of [11] combined with an old result of Nash-Williams [10]).

**Lemma 8.** *Let  $G = (V, E)$  be a graph with an ordering  $(v_1, \dots, v_n)$  of  $V$  which satisfies that  $|N(v_i) \cap \{v_{i+1}, \dots, v_n\}| \leq k$  for all  $i \in [n]$ . Then  $E$  can be partitioned into  $k$  parts such that no part contains a cycle of  $G$ .*

**Proof of Theorem 1.** Let  $G = (V, E)$  be a graph with maximum degree  $\Delta$  and fix  $\alpha > 16$ . We perform the following procedure.

1. Let  $\tilde{G}^0 = G$  and  $G' = (V, \emptyset)$ .
2. Start with  $i = 0$  and repeat the following until  $i = \tau$ , where  $\tau$  is the smallest such that  $\Delta(\tilde{G}^\tau) \leq \log^2 \Delta$ :
  - (a) obtain  $G^i$  from  $\tilde{G}^i$  by successively removing all vertices of degree less than  $\log^2 \Delta$ , and adding all of their incident edges to  $G'$ ;
  - (b) apply Corollary 7 to  $G^i$  to obtain the disjoint  $C_4$ -free subgraphs  $G_1^i, G_2^i, \dots, G_{\lceil 2\sqrt{2\lceil \alpha \Delta(G^i)} \rceil}^i$ ;
  - (c) set  $\tilde{G}^{i+1} = (V(G^i), E(G^i) \setminus \bigcup_j E(G_j^i))$  and then increment  $i$ .
3. Add all edges of  $\tilde{G}^\tau$  to  $G'$ .

We can always apply [Corollary 7](#) at each iteration since in Step 2(a) we forced the minimum degree of  $G^i$  to be at least  $\log^2 \Delta \geq \log^2 \Delta(G^i)$ .

Let us see that the maximum degree  $\Delta(G^{i+1})$  is significantly smaller than  $\Delta(G^i)$ . By [Corollary 7](#), the removal of  $C_4$ -free subgraphs at iteration  $i$  removes at least  $\beta d_{G^i}(v)$  edges incident to  $v \in V$ . Thus

$$\Delta(G^{i+1}) \leq \Delta(\tilde{G}^{i+1}) \leq (1 - \beta)\Delta(G^i) \leq (1 - \beta)^i \Delta. \quad (4)$$

This implies that the procedure is guaranteed to stop after  $\tau = O(\log \Delta)$  iterations.

Step 2(b) of each iteration generates a number of disjoint spanning  $C_4$ -free subgraphs, each of which we give a new colour. During the  $i$ th iteration we produce  $\lceil 2\sqrt{2\lceil \alpha \Delta(G^i) \rceil} \rceil < 2\sqrt{2\alpha \Delta(G^i)} + 4$  such subgraphs, so by (4) and the bound on the number  $\tau$  of iterations we produce at most

$$\begin{aligned} & O(\log \Delta) + 2\sqrt{2\alpha \Delta} + 2\sqrt{2\alpha(1 - \beta)\Delta} + 2\sqrt{2\alpha(1 - \beta)^2 \Delta} + \dots \\ &= \frac{2\sqrt{2\alpha}}{1 - \sqrt{1 - \beta}} \cdot \sqrt{\Delta} + O(\log \Delta) \end{aligned} \quad (5)$$

$C_4$ -free subgraphs throughout all iterations.

It only remains to upper bound the number of colours needed in the remainder graph  $G'$ . By construction,  $G'$  admits a degeneracy ordering satisfying the hypothesis of [Lemma 8](#) for  $k = \log^2 \Delta$ . Thus we can partition its edges into at most  $\log^2 \Delta$  acyclic (and thus  $C_4$ -free) subgraphs. By (5) we obtain a partition of  $E$  into  $O(\sqrt{\Delta})$   $C_4$ -free subgraphs in total. This completes the proof of the theorem.  $\square$

## Acknowledgements

We thank the referees for their helpful comments and suggestions.

The first author was supported by a NWO Veni grant (project number 639.031.138).

## References

- [1] S.A. Burr, P. Erdős, L. Lovász, On graphs of Ramsey type, *Ars Combin.* 1 (1) (1976) 167–190.
- [2] F.R.K. Chung, R.L. Graham, On multicolor Ramsey numbers for complete bipartite graphs, *J. Combin. Theory Ser. B* 18 (1975) 164–169.
- [3] P. Erdős, On sequences of integers no one of which divides the product of two others and on some related problems, *Inst. Math. Mech. Univ. Tomsk* 2 (1938) 74–82.
- [4] Z. Füredi, M. Simonovits, The history of degenerate (bipartite extremal graph problems), in: *Erdős Centennial*, in: *Bolyai Soc. Math. Stud.*, vol. 25, János Bolyai Math. Soc., Budapest, 2013, pp. 169–264.
- [5] R.W. Irving, Generalised Ramsey numbers for small graphs, *Discrete Math.* 9 (1974) 251–264.
- [6] T. Jiang, K.G. Milans, D.B. West, Degree Ramsey numbers for cycles and blowups of trees, *European J. Combin.* 34 (2) (2013) 414–423.
- [7] W.B. Kinnersley, K.G. Milans, D.B. West, Degree Ramsey numbers of graphs, *Combin. Probab. Comput.* 21 (1–2) (2012) 229–253.
- [8] Y. Li, K.W. Lih, Multi-color Ramsey numbers of even cycles, *European J. Combin.* 30 (1) (2009) 114–118.
- [9] M. Molloy, B. Reed, Graph Colouring and the Probabilistic Method, in: *Algorithms and Combinatorics*, vol. 23, Springer-Verlag, Berlin, 2002.
- [10] C.St.J.A. Nash-Williams, Decomposition of finite graphs into forests, *J. Lond. Math. Soc.* 39 (1964) 12.
- [11] J. Nešetřil, P. Ossona de Mendez, Sparsity: Graphs, Structures, and Algorithms, in: *Algorithms and Combinatorics*, vol. 28, Springer, Heidelberg, 2012.
- [12] G. Perarnau, B. Reed, Existence of spanning  $\mathcal{F}$ -free subgraphs of regular graphs with large minimum degree, 2014. arXiv:1404.7764.
- [13] O. Pikhurko, A note on the Turán function of even cycles, *Proc. Amer. Math. Soc.* 140 (11) (2012) 3687–3692.