

UNIVERSITY OF BIRMINGHAM

Research at Birmingham

Low dose interleukin-2 promotes STAT5 phosphorylation, Treg survival and CTLA-4 dependent function in autoimmune liver disease

Jeffery, Hannah; Jeffery, Louisa; Lutz, Philipp; Corrigan, Margaret; Webb, Gwilym; Hirschfield, Gideon; Adams, David; Oo, Ye Htun

DOI:

[10.1111/cei.12940](https://doi.org/10.1111/cei.12940)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Jeffery, H, Jeffery, L, Lutz, P, Corrigan, M, Webb, G, Hirschfield, G, Adams, D & Oo, YH 2017, 'Low dose interleukin-2 promotes STAT5 phosphorylation, Treg survival and CTLA-4 dependent function in autoimmune liver disease: Low dose IL2 promotes regulatory T cell function in autoimmune liver diseases', *Clinical & Experimental Immunology*, vol. 188, no. 3, pp. 394–411. <https://doi.org/10.1111/cei.12940>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Low-dose interleukin-2 promotes STAT-5 phosphorylation, T_{reg} survival and CTLA-4-dependent function in autoimmune liver diseases

H. C. Jeffery,* L. E. Jeffery,[†] P. Lutz,*
M. Corrigan,*[‡] G. J. Webb,*[‡]
G. M. Hirschfield,*[‡] D. H. Adams*[‡]
and Y. H. Oo*[‡]

*Centre for Liver Research and National
Institute for Health Research Liver Biomedical
Research Unit, Institute of Immunology and
Immunotherapy, University of Birmingham,
UK, [†]Institute of Metabolism and Systems
Research, University of Birmingham, UK, and
[‡]Liver Transplant and Hepatobiliary Unit,
University Hospital of Birmingham NHS
Foundation Trust, Birmingham, UK

Accepted for publication 12 January 2017
Correspondence: Ye Htun Oo, Centre for
Liver Research and NIHR Birmingham
Biomedical Research Unit, University of
Birmingham, Vincent Drive, Edgbaston,
Birmingham B15 2TT, UK.
E-mail: y.h.oo@bham.ac.uk

Introduction

Naturally occurring CD4⁺CD25^{high}CD127^{low}forkhead box protein 3 (FoxP3⁺) regulatory T cells (T_{reg}) constitute 5–10% of peripheral CD4⁺ T cells and maintain peripheral self-tolerance in rodents and humans [1,2]. Functional impairment or quantitative deficiency of T_{regs} has been described in autoimmune liver diseases (AILD) [3,4], including those targeted at bile ducts [primary biliary cholangitis (PBC) [5,6] or primary sclerosing cholangitis (PSC)] and hepatocytes [autoimmune hepatitis (AIH)] [3,7–9]. Current therapies for AILD are non-curative, provide unsatisfactory control of hepatic inflammation and require long-term immunosuppressive medications that carry unfavourable side effects. Thus, autologous T_{reg} therapy is an attractive option for the treatment of AILD that

Summary

CD4⁺CD25^{high}CD127^{low}forkhead box protein 3 (FoxP3⁺) regulatory T cells (T_{reg}) are essential for the maintenance of peripheral tolerance. Impaired T_{reg} function and an imbalance between effector and T_{regs} contribute to the pathogenesis of autoimmune diseases. We reported recently that the hepatic microenvironment is deficient in interleukin (IL)–2, a cytokine essential for T_{reg} survival and function. Consequently, few liver-infiltrating T_{reg} demonstrate signal transducer and activator of transcription-5 (STAT-5) phosphorylation. To establish the potential of IL-2 to enhance T_{reg} therapy, we investigated the effects of very low dose Proleukin (VLDP) on the phosphorylation of STAT-5 and the subsequent survival and function of T_{reg} and T effector cells from the blood and livers of patients with autoimmune liver diseases. VLDP, at less than 5 IU/ml, resulted in selective phosphorylation of STAT-5 in T_{reg} but not effector T cells or natural killer cells and associated with increased expression of cytotoxic T lymphocyte antigen-4 (CTLA-4), FoxP3 and CD25 and the anti-apoptotic protein Bcl-2 in T_{reg} with the greatest enhancement of regulatory phenotype in the effector memory T_{reg} population. VLDP also maintained expression of the liver-homing chemokine receptor CXCR3. VLDP enhanced T_{reg} function in a CTLA-4-dependent manner. These findings open new avenues for future VLDP cytokine therapy alone or in combination with clinical grade T_{reg} in autoimmune liver diseases, as VLDP could not only enhance regulatory phenotype and functional property but also the survival of intrahepatic T_{reg}.

Keywords: autoimmune liver disease, Bcl-2, CTLA-4, interleukin-2, regulatory T cells, STAT-5

could provide long-term immune-regulation without daily medications and global immunosuppression.

T_{reg} survival and function is dependent upon interleukin (IL)–2 [10], which is required for the maintenance of effective levels of functional T_{reg} in autoimmune diseases [11–13]. The importance of IL-2 for T_{reg} function has not been studied closely in autoimmune liver diseases. The cell surface receptor for IL-2 (IL-2R) is composed of three subunits, alpha (IL-2RA, CD25), beta (IL-2RB, CD122) and common gamma (IL-2RG, CD132). All leucocytes express IL-2RG constitutively. Natural killer (NK), NK T cells (NK T) and memory CD8⁺ T cells also express IL-2RB and T_{reg} express IL-2RA constitutively. IL-2RA is required for high-affinity IL-2 binding, while IL-2RB and IL-2RG transduce the IL-2 signal [14]. Two major signalling pathways

conduct IL-2-induced responses: signalling from IL-2RB leads to the activation of the serine/threonine kinase, AKT, and to up-regulation of anti-apoptotic molecules such as Bcl-2, which is required for T cell survival [15]. Signalling from IL-2RG via Janus kinase 3 (JAK3) leads to signal transducer and activator of transcription-5 (STAT-5) activation [16], and is needed for T cell proliferation and differentiation and expression of anti-apoptotic molecules [17,18]. Owing to their high levels of high-affinity CD25, T_{reg} consume IL-2 competitively, thereby maintaining their survival and function while suppressing bystander effector cells [19,20]. Where IL-2 availability is low, such as in the inflamed hepatic microenvironment, T_{reg} function may be compromised and may be inadequate to counteract the activated immune infiltrate.

A number of studies have indicated that treatment with IL-2 could improve immune-mediated diseases. In rodents, type 1 diabetes mellitus could be prevented by *in-vivo* IL-2 administration [21,22]. In humans, low-dose IL-2 therapy enhanced T_{reg} frequencies and improved outcome in graft-versus-host disease, vasculitis and type 1 diabetes [11,23–25]. We recently reported very low levels of IL-2 in the inflamed human liver [4]. Thus, we considered that clinical grade IL-2 (Proleukin) therapy might be effective in AILD.

In this study, we examined the effect of very low dose Proleukin (VLDP) on the biology of both peripheral and intrahepatic T_{reg} , focusing upon regulatory phenotype and function. Successful T_{reg} therapy in AILD would require not only enhancing T_{reg} phenotype and function, but also recruitment of peripheral T_{reg} to the inflamed autoimmune livers. We demonstrate for the first time, to our knowledge, that VLDP selectively enhances T_{reg} STAT-5 phosphorylation and subsequently up-regulates functional molecules cytotoxic T lymphocyte antigen-4 (CTLA-4), CD25, FoxP3 and T_{reg} anti-apoptosis marker Bcl-2 in AILD. VLDP also maintains liver homing chemokine receptor CXCR3 expression on T_{reg} . Importantly, VLDP enhances the suppressive function of T_{reg} via CTLA-4, and anti-CTLA-4 can block this effect. Thus, we demonstrated both phenotype and mechanistic effects of VLDP on blood and intrahepatic T_{reg} from AILD patients suggesting that VLDP therapy may enhance immune-regulatory restoration in AILD.

Materials and methods

Ethics statement

Written informed consent was obtained from all subjects in this study. Local Research Ethics Committees (LREC) and the University of Birmingham approved all experimental protocols (South Birmingham LREC reference: 98 CA/5192; Walsall LREC reference: 06/Q2708/11).

Blood and liver tissue

Venous blood, collected in ethylenediamine tetraacetic acid (EDTA) tubes, was obtained from healthy (control) individuals and individuals with AILD, including AIH, PBC and PSC. Explanted diseased liver tissue was obtained from patients undergoing liver transplantation for end-stage AILD including PBC, PSC and AIH/PBC and AIH/PSC overlap diseases.

Isolation of liver infiltrating leucocytes (LIL) and peripheral blood mononuclear cells (PBMC)

LIL were isolated from fresh liver tissue. Briefly, explanted liver tissue was diced into 5 mm³ cubes, washed with phosphate-buffered saline (PBS) and then homogenized in a Seward stomacher 400 circulator (260 rpm, 5 min). The homogenate was filtered through fine (63 µm) mesh (John Staniar and Co., Manchester, UK) and the lymphocytes isolated by density gradient separation using Lympholyte (VH Bio, Gateshead, UK) at 800 g for 20 min. The lymphocyte layer was collected and washed with PBS. Cell viability was assessed by trypan blue exclusion. Peripheral blood lymphocytes were isolated similarly from whole blood by density gradient separation using Lympholyte.

Culture of PBMC and LIL

PBMC and LIL were cultured in 24-well plates at a density of 1×10^6 cells/ml in RPMI-1640 with L-glutamine medium containing penicillin (100 IU/ml), streptomycin (100 IU/ml), additional glutamine (2 mM) (GIBCO, Carlsbad, CA, USA) and 10% human AB serum (TCS Biosciences, Buckingham, UK) and supplemented with 0 or 5 IU/ml Proleukin (Aldesleukin) (Novartis, Camberley, UK).

Surface phenotyping of freshly isolated intrahepatic and peripheral blood lymphocytes with or without Proleukin treatment

Cell phenotypes were examined by flow cytometry. Dead cells were identified by staining with the Zombie NIRTM fixable viability dye (BioLegend, San Diego, CA, USA) or e506 viability dye (eBioscience, San Diego, CA, USA) prior to staining with antibodies. To analyse expression of surface antigens, cells were incubated on ice for 30 min with antibodies against CD3, CD4, CD8, CD25, CD127 and markers of interest or isotype-matched control antibodies in 2% fetal bovine serum (FBS) (Sigma Aldrich, Poole, UK) diluted in PBS. After washing with 2% FBS (Sigma Aldrich), cells were fixed for 10 min with 3% formaldehyde solution (Sigma Aldrich). To analyse expression of intracellular proteins, cells were fixed and stained using the FoxP3/transcription factor staining set (eBioscience), according to the manufacturer's instructions. Antibodies against surface markers (CD3, CD4, CD8, CD25, CD127) were generally added together with antibodies against intracellular

markers of interest during the permeabilization and intracellular staining steps. Data were acquired using a CyAN ADP flow cytometer (Dako, Glostrup, Denmark). Single fluorophore-labelled anti-mouse immunoglobulin (Ig)G κ /negative control (FBS) compensation particles (BD Biosciences, Franklin Lakes, NJ, USA) were used for compensation. Data were analysed offline using FlowJo (TreeStar Inc., Ashland, OR, USA).

The anti-human antibodies used in flow cytometric analysis of marker expression included: anti-CD3-phycoerythrin-cyanin 7 (PE-Cy7) (SK7; BD Biosciences), anti-CD4-peridinin chlorophyll (PerCP/Cy5.5 (RPA-T4; eBioscience), anti-CD4-Viogreen (VIT4; Miltenyi Biotec), anti-CD8-PE-CF594 (RPA-T8; BD Biosciences), anti-CD25-BV421 (M-A251; BD Biosciences), anti-CD45R α -allophycocyanin (APC)-Vio770 (T6D11; Miltenyi Biotec), anti-CD127-fluorescein isothiocyanate (FITC) (eBioRDR5; eBioscience), anti-CCR7-PE-CF594 (150503; BD Biosciences), anti-granzyme B-PE (GB11; eBioscience), anti-CTLA-4-PE (BN13; BD Biosciences), anti-Bcl-2-PE (100; BioLegend), anti-FoxP3-APC (PCH101; eBioscience) anti-CD39-PE (A1; eBioscience), anti-T cell immunoglobulin and mucin domain-containing-3 (TIM3)-PE (F38-2E2; eBioscience), anti-OX40-PE (ACT35; BD Biosciences), anti-CD69-PE (FN50; Miltenyi Biotec), anti-2B4-PE (REA112; Miltenyi Biotec), anti-CD73-APC (AD2; eBioscience), anti-CD137-APC (4B4-1; Miltenyi Biotec), anti-glucocorticoid-induced tumour necrosis factor receptor (GITR)-APC (DT5D3; Miltenyi Biotec), anti-lymphocyte-activation gene 3 (LAG3)-APC (3DS223H; eBioscience) and anti-PD-1-APC (PD1.3.1.3; Miltenyi Biotec).

Analysis of STAT-5 phosphorylation in response to IL-2

To examine responsiveness of PBMC or LIL to IL-2, cells in RPMI were stained with anti-CD127-FITC (eBioRDR5; eBioscience), anti-CD20-Viogreen (LT20; Miltenyi Biotec) and anti-CD56-pacific blue (HCD56; BioLegend) for 10 min at room temperature. Cells were then stimulated for 10 min at 37°C with Proleukin (0–1000 IU/ml). Cells were fixed and permeabilized with BD Biosciences Phosflow buffers I and III according to the manufacturer's instructions then stained for 1 h at room temperature in 2% FBS (Sigma) with anti-pSTAT-5 (Y694)-AlexaFluor 647 (47/STAT-5), anti-CD3-PeCy7 (SK7), anti-CD8-PE-CF594 (RPA-T8) (all from BD Biosciences), anti-CD4-PerCPCy5.5 (RPA-T4; eBioscience) and anti-CD25-PE (3G10; Miltenyi Biotec). All data were acquired using a CyAn ADP (Dako) flow cytometer and analysed using FlowJo (Tree Star) software.

Analysis of suppression of autologous T responder cell division by T_{reg} in response to IL-2

CD4⁺CD25⁺CD127⁻ T_{reg} and CD4⁺CD25⁻CD127⁺ responder T cells were isolated by flow sorting following

prior enrichment of total CD4⁺ T cells from PBMC using magnetic negative selection (Biolegend). T_{reg} were cultured overnight with or without Proleukin (5 IU/ml). Responder T cells were labelled with cell trace violet (molecular probes; ThermoFisher, Waltham, MA, USA) and cultured overnight without stimulation. Dendritic cells (DC) were derived from monocytes that were isolated from healthy donor PBMCs by magnetic negative selection (StemCell Technologies, Vancouver, Canada) and cultured for 5–7 days in IL-4 (500 IU/ml; Miltenyi Biotec) and granulocyte-macrophage colony-stimulating factor (GM-CSF) (800 IU/ml; Berlex Laboratories, Richmond, CA, USA). DCs and T_{reg} were washed to remove cytokines and co-cultured for 5 days with responder T cells in the presence of 0.5 μ g/ml anti-CD3 (clone OKT3; Biolegend) with or without 40 μ g/ml anti-CTLA-4. Division of responder T cells under 50 μ g/ml CTLA-4-Ig (abatacept) was also monitored. Cells were cultured at a ratio of 1 DC : 20 T cells with a 1 T_{reg} : 2.5 responder T cell ratio. At 5 days, cell trace violet dilution was measured by flow cytometry and the statistics percentage division and division index (the average number of cell divisions undergone by a cell in the original population) calculated using the FlowJo proliferation analysis platform.

Statistical analysis

Statistical significance between two variables across multiple subsets was tested by two-way two-tailed analysis of variance (ANOVA) with Bonferroni's *post-hoc* analysis. Significance between two populations was tested by paired *t*-test and between multiple treatments by one way ANOVA with Bonferroni's *post-hoc* analysis. Analysis and graphical representation was performed using GraphPad Prism version 5 (GraphPad software, San Diego, CA, USA).

Results

Phosphorylation of STAT-5 occurs selectively in T_{reg} at very low doses of Proleukin in normal blood

To analyse the responsiveness of different leucocyte populations to IL-2 (Proleukin) and identify a dose that would phosphorylate STAT-5 selectively in T_{reg}, we administered Proleukin at doses of between 0 and 1000 IU/ml to PBMC from healthy control bloods and examined phosphorylation of STAT-5 in T_{reg} and other leucocyte populations: (Fig. 1a and Supporting information, Fig. 1). With the exception of B cells, all cell types responded to Proleukin at 10 IU/ml and above (Fig. 1a). However, at 1 IU/ml, 25% of T_{reg} showed measurable pSTAT-5 expression, while there was no up-regulation in pSTAT-5 in other subsets. More than 75% of T_{reg} demonstrated pSTAT-5 induction at 10 IU/ml; however, this dose also led to phosphorylation of




Fig. 1. Very low dose interleukin (IL)-2 up-regulates phosphorylated signal transducer and activator of transcription-5 (pSTAT-5) selectively in peripheral and liver-infiltrating regulatory T cells (T_{reg}) from patients with autoimmune liver disease. Peripheral blood mononuclear cells (PBMCs) from control bloods (a, $n = 3$) or autoimmune liver disease patient bloods [b, autoimmune hepatitis (AIH) remission, $n = 5$; AIH relapse, $n = 4$; primary biliary cholangitis (PBC), $n = 3$; and primary sclerosing cholangitis (PSC), $n = 3$] and liver-infiltrating lymphocytes from explanted livers of patients with autoimmune liver disease (c, $n = 4$) were stimulated for 10 min with IL-2 (Proleukin) at doses in the range 0–1000 IU/ml and the expression of p(Y694) STAT-5 determined by flow cytometry for each leucocyte population. Graphs summarize pSTAT-5 expression, quoted as percentage of positive cells or median fluorescence intensity (MFI). Data are mean \pm standard error of the mean (s.e.m.).

STAT-5 in other $CD4^+$ T cells and $CD56^{bright}$ NK cells. Nevertheless, this pSTAT-5 induction in non- T_{reg} s at this concentration was barely above baseline, as indicated by pSTAT-5 median fluorescence intensity (MFI) (Fig. 1a). Most importantly, at 2–4 IU/ml, T_{reg} pSTAT-5 MFI was 2.3–3.6-fold above baseline, but there was no increase in

pSTAT-5 MFI for other subsets (Fig. 1a). Thus, we observed that with a very low dose of Proleukin (VLDP) (< 5 IU/ml), selective phosphorylation of STAT-5 in T_{reg} could be achieved. The characteristics of the control cohort used in these and subsequent phenotyping studies in this paper are given in Table 1.

Table 1. Demographic, laboratory, histology and treatment details of patients in the peripheral blood cohort

Disease	Age (years)	Sex	BR	AST	ALT	ALP	Na+	CR	IgM	IgG	INR	UKELD	Histology	Treatment
AIH F	28	M	19	78	23	65	139	68	0.6	17	1.2	49	Moderate hepatitis, established cirrhosis	Azathioprine + tacrolimus + prednisolone
AIH F	61	F	10	129	125	120	139	71	0.5	25	1.2	47	Moderate-severe activity, F4	Mycophenolate
AIH F	57	F	15	76	53	114	140	61	1.4	18	1.1	48	Moderate hepatitis, F3	Azathioprine + prednisolone
AIH F	42	M	37	68	120	104	142	86	1.3	17	1.0	49	Moderate hepatitis, established cirrhosis	Azathioprine, prednisolone
AIH F	64	M	7	72	127	93	141	87	0.7	17	1.0	45	Moderate lobular hepatitis, F2	Mycophenolate + prednisolone
AIH F	24	M	59	167	253	120	138	49	0.7	21	1.2	53	Moderate hepatitis, F3	Mycophenolate + prednisolone
AIH R	58	M	19	42	30	82	142	86	1	16	0.9	47	Moderate lobular inflammatory activity, F2	Azathioprine
AIH R	46	F	4	17	8	76	137	67	1.2	13	1.0	44	Mild hepatitis, F2	Azathioprine
AIH R	51	F	11	26	21	58	143	72	1	10	1.0	45	Quiescent AIH, cirrhotic	None
AIH R	58	F	15	39	40	90	144	58	0.7	12	1.1	45	Quiescent AIH, F1	Azathioprine + prednisolone
AIH R	45	F	10	16	11	86	142	67	0.9	9	1.0	45	Mild hepatitis, F1	Azathioprine
AIH R	35	F	14	22	19	75	139	58	1	11	1.1	48	Moderate hepatitis, F3	Azathioprine
AIH R	54	F	7	21	28	76	143	66	1.2	9	1.0	43	Moderate hepatitis, F3	Mycophenolate + prednisolone
AIH R	45	F	13	43	36	84	138	64	1	20	1.0	47	Mild hepatitis, F2	Azathioprine
AIH R	21	F	17	30	35	70	143	63	1	8	1.0	46	Mild hepatitis, Established cirrhosis	Azathioprine + prednisolone
AIH R	57	F	3	25	28	65	142	82	0.8	12	1.0	42	Mild hepatitis, F2	Mycophenolate + prednisolone
AIH R	33	F	7	28	24	46	139	65	0.6	11	1.0	46	Mild lobular, hepatitis, F2	Mycophenolate + prednisolone
AIH R	51	F	8	33	33	93	140	60	1.3	12	1.0	46	Mild hepatitis, F1	Mycophenolate + prednisolone
AIH R	36	M	11	18	22	67	141	58	0.8	14	1.2	45	Mild hepatitis, established cirrhosis	Azathioprine
AIH R	21	F	11	17	19	67	141	61	0.8	8	1.0	46	Mild hepatitis, F1	Mercaptopurine
PBC	72	F	8	36	34	210	144	105	2.9					
PBC	63	F	5	32	20	160	141	83	3.5	9.37	0.9	43	Non-cirrhotic	Ursodeoxycholic acid
PBC	51	F	39	113	76	184	142	93	5.6	16.81	1.1	49	Non-cirrhotic	Ursodeoxycholic acid
PBC	47	F	13	29	21	150	142	52	1.71	10.64	1.1	46	Chronic biliary disease, paucity of bile ducts, cirrhosis	Ursodeoxycholic acid
PBC	58	F	11	n.a.	88	653	138	66	7.15	23.63	1.0	47	2007 - PBC with FNH; no cirrhosis	UDCA
PBC	41	F	5	20	23	74	145	68	1.31	12.75	1.1	42	n.a.	UDCA
PBC	47	F	32	112	129	446	139	57	7.78	13.32	1.0	50	n.a.	UDCA
PBC	54	F	4	14	19	297	142	78	0.6	9	0.8	42	PSC, F2	UDCA
PBC	43	M	70	151	68	575	139	60	0.8	25	1.2	54	PSC cirrhotic	UDCA
PBC	77	M	3	21	10	64	136	118	0.6	10	0.9	45	PSC, F2	UDCA
Control	45	M	8	12	18	n.a.	133	69	n.a.	n.a.	0.9	49	n.a.	None
Control	56	F	4	8	20	n.a.	140	74	n.a.	17.07	1.0	43	n.a.	None
Control	60	M	10	41	55	n.a.	145	105	n.a.	15.75	1.1	44	n.a.	None
Control	49	M	18	16	13	n.a.	141	85	n.a.	10.11	1.1	48	n.a.	None
Control	60	M	3	12	16	n.a.	139	76	n.a.	N/A	n.a.	n.a.	n.a.	None

Table 1. Continued

Disease	Age (years)	Sex	BR	AST	ALT	ALP	Na+	CR	IgM	IgG	INR	UKELD	Histology	Treatment
Control	67	F	9	33	28	n.a.	140	64	n.a.	n.a.	1.1	46	n.a.	None
Control	42	M	6	24	21	n.a.	135	72	n.a.	n.a.	n.a.	n.a.	n.a.	None
Control	76	M	6	32	27	n.a.	143	79	n.a.	12.71	n.a.	n.a.	n.a.	None
Control	41	M	63	26	12	n.a.	140	93	n.a.	9.19	n.a.	n.a.	n.a.	None

AIH R = autoimmune hepatitis in remission; AIH F = autoimmune hepatitis in flare up/relapse; PSC = primary sclerosing cholangitis; PBC = primary biliary cholangitis; AST = aspartate transaminase; ALT = alanine transaminase; ALP = alkaline phosphatase BR = bilirubin; Na+ = sodium; CR = creatinine; UKELD = United Kingdom End Stage Liver Disease Scoring; UDCA = ursodeoxycholic acid; n.a. = not available.

Very low doses of Proleukin induce selective phosphorylation of STAT-5 in intrahepatic and peripheral blood T_{reg} of patients with autoimmune liver diseases

PBMC from AIH, PBC and PSC patients were treated with Proleukin in the range of 0–100 IU/ml. Similar to controls, T_{reg} of AIH patients [both remission and flare-up (relapse)], PBC patients and PSC patients demonstrated a selective pSTAT-5 induction at doses below 10 IU/ml, especially at 5 IU/ml (Fig. 1b and Supporting information, Fig. 2). We then treated human LIL isolated from explanted autoimmune liver disease tissues with 0–100 IU/ml Proleukin, and again selective pSTAT-5 enhanced induction was seen in T_{reg} at doses below 10 IU/ml (Fig. 1c). Descriptions of all AILD blood and liver explant donors whose PBMC and LIL were used in these and subsequent studies in this paper to address the effect of IL-2 on phenotype are given in Tables 1 and 2 [blood donors were: AIH = 20 (aged 44.4 ± 13.8 years, disease in remission = 14, disease in flare up/relapse = 6); PSC = 3 (aged 58.0 ± 17.3 years); PBC = 10 (aged 54.0 ± 9.6 years); control = 9 (aged 55.1 ± 11.9 years)]. Liver donors were: PBC = 5 (aged 59.4 ± 6.4 years); PSC = 2 (aged 51.5 ± 23.3 years); AIH overlap = 2 (aged 38.0 ± 24.0 years), PBC/AIH overlap = 1; and PSC/AIH overlap = 1).

Very low doses of Proleukin induce T_{reg} functional markers CTLA-4, CD25 and FoxP3 selectively in both blood and intrahepatic T_{reg}

Having found 5 IU/ml Proleukin by dose titration to be the optimal concentration to induce a selective STAT-5 response in blood and liver T_{reg} of AILD patients, we determined the effect of 5 IU/ml Proleukin on T cell phenotype and function. PBMC and LIL from AILD patients were exposed to 0 IU/ml or 5 IU/ml Proleukin for 18 h and effects on markers of T cell activation and T_{reg} function by T cell subsets were assessed; 5 IU/ml Proleukin increased significantly the levels of CD25, CTLA-4 and FoxP3 on peripheral blood T_{reg} of AILD patients (Fig. 2a and Supporting information, Fig. 3). In liver, basal expressions of CTLA-4, CD25 and FoxP3 on T_{reg} were higher than in blood and similar to those observed in blood following Proleukin exposure (Fig. 2a). There were clear trends towards increased expression of the T_{reg} markers CD25 and FoxP3 with VLDP in liver T_{reg} from PBC, PSC and AIH, supporting potentiation of T_{reg} in the liver in settings of the AILDs (Fig. 2a and Supporting information, Fig. 3). These observations are consistent with and also reflective of upstream pSTAT-5 induction by VLDP in liver T_{reg} (Fig. 1c). Frequencies of expression of CD25, CTLA-4 and FoxP3 were largely not altered in blood or liver T_{reg} by VLDP (Supporting information, Fig. 4).

Because we noticed that 18 h exposure to Proleukin led to STAT-5 phosphorylation in CD25⁺CD127⁺CD4⁺ T cells that mirrored that in T_{reg} (CD25⁺CD127⁺CD4⁺) (both frequency and MFI) (Supporting information, Fig. 5), we evaluated the

Table 2. Demographic, histology and treatment details of patients in the liver explant cohort

Disease	Age (years)	Sex	Explant history	Treatment before transplantation
PBC	51	M	Moderate fibrosis	Ursodeoxycholic acid
PBC	63	F	Established cirrhosis	Ursodeoxycholic acid
PBC	65	F	Established cirrhosis	Ursodeoxycholic acid
PBC	64	F	Established cirrhosis	Ursodeoxycholic acid
PBC	54	F	Established cirrhosis	Ursodeoxycholic acid
PSC	68	M	Established cirrhosis	
PSC	35	M	Established cirrhosis	
PBC/AIH overlap	55	F	Established cirrhosis	Ursodeoxycholic acid
PSC/AIH overlap	21	F	Established cirrhosis	

AIH = autoimmune hepatitis; PBC = primary biliary cholangitis; PSC = primary sclerosing cholangitis.

effect of 18-h VLDP on the expression of CD25, CTLA-4 and FoxP3 by the CD25⁺CD127⁺CD4⁺ T effector cell population in the blood of AILD patients (Fig. 2b). Importantly, despite phosphorylation of STAT-5, this subset, unlike T_{reg}, did not up-regulate CD25, CTLA4 or FoxP3 (Fig. 2b).

We then investigated the effect of VLDP on the expression by peripheral and liver-infiltrating T_{reg} of tumour necrosis factor (TNF) receptor superfamily members CD137, OX40 and GITR. Culture in 5 IU/ml Proleukin helped to maintain expression frequencies of CD137 and GITR on peripheral T_{reg} over 18 h and promoted an increase in the frequency of OX40 expressing peripheral T_{reg} (Fig. 3a and Supporting information, Fig. 3). Exposure to VLDP did not influence the frequency of expression of these markers on liver-infiltrating T_{reg} (Fig. 3a). We also investigated the impact of VLDP on T_{reg}-associated surface markers, including CD39, CD73 and LAG3. Proleukin 5 IU/ml had no effect on the expression frequency of any of these molecules (Fig. 3b and Supporting information, Fig. 3): high expression of CD39 was maintained, CD73 remained low-expressed and TIM3 and LAG3, which were not detected at baseline, were not induced by Proleukin treatment on T_{reg}. Also, VLDP did not alter expression of the activation marker, CD69 or the immune checkpoint and programmed cell death receptor-1 (PD-1) on CD4⁺ or CD8⁺ T cells or T_{reg} (Fig. 3c and Supporting information, Fig. 3). Consistent with a lack of pSTAT-5 induction in CD8⁺ T cells under VLDP, 5 IU/ml Proleukin did not affect expression of granzyme B or activation marker 2B4 by CD8⁺ T cells (Fig. 3c and Supporting information, Fig. 3).

Very low dose Proleukin enhances T_{reg} suppressive potential in a CTLA-4-dependent manner across all autoimmune liver diseases

Having established that VLDP 5 IU/ml enhances T_{reg} expression of regulatory functional molecules we sought to verify that these changes in phenotype associate with improvements in T_{reg} function. We tested whether exposure to VLDP could increase the ability of T_{reg} to suppress the division of autologous CD25⁺CD4⁺ T effector cells. We used a system of anti-CD3 together with monocyte-derived DC to

supply co-stimulatory ligands. This system was chosen over conventional anti-CD3/CD28 bead activation of T cell proliferation in order that the functional impact of antigen-presenting cell-T_{reg} contact-dependent mechanisms of suppression such as involving CTLA-4 might be studied. Pre-treatment of T_{reg} with Proleukin reduced significantly the division of T responder cells from control individuals cultured 2.5 : 1 with autologous T_{reg} (Fig. 3d,e). This elevated suppressive potential of VLDP-treated T_{reg} was overcome in the presence of anti-CTLA-4 (Fig. 3d,f). Consistent with our observations that CTLA-4-Ig can reduce T responder division (Supporting information, Fig. 6a), anti-CTLA-4 also tended to increase T responder division in the presence of untreated T_{reg} (Fig. 3f). We observed the same trends of improved T_{reg} function involving contribution from CTLA-4 with VLDP treatment of T_{reg} in both cases of AILD patients tested (Supporting information, Fig. 6b).

Very low dose Proleukin treatment does not alter liver homing CXCR3 expression by peripheral blood T_{reg} in autoimmune liver diseases

Regulatory T cell homing to the site of liver inflammation is crucial for the success of T_{reg} therapies, thus we examined the crucial liver-homing chemokine receptor, CXCR3, on blood T_{reg} of autoimmune liver disease patients. *De-novo* expression levels (MFI and frequency) of CXCR3 were similar in patients with AIH compared to controls (Fig. 4a). Importantly, VLDP 5 IU/ml did not have any impact upon CXCR3 expression in AIH patients over 18 h or up to 3 days (Fig. 4b,c). Peripheral T_{reg} from PSC and PBC patients also showed no change in their CXCR3 expression with sustained culture in VLDP (Fig. 4b,d), thus VLDP therapy may be applicable for all three autoimmune liver diseases without impairing the recruitment capacity of T_{reg}.

Very low dose Proleukin treatment increases total T_{reg} frequencies and promotes the strongest T_{reg} suppressive phenotype in the CD45RA⁺CCR7⁻ population of T_{reg}

We noted that the CD25⁺CD127⁻FoxP3⁺ T_{reg} population, as a proportion of total CD4⁺ T cells, increased




Fig. 2. Effect of very low dose interleukin (IL)–2 on expression of IL-2-regulated regulatory T cells (T_{reg}) functional markers CD25, cytotoxic T lymphocyte antigen-4 (CTLA-4) and forkhead box protein 3 (FoxP3) by T_{reg} cells from blood and liver. Peripheral blood mononuclear cells (PBMCs) from patients with autoimmune hepatitis (AIH) ($n = 5$), primary sclerosing cholangitis (PSC), ($n = 3$) and primary biliary cholangitis (PBC), ($n = 6$) and liver-infiltrating leucocytes from autoimmune liver diseases (AILD) livers were exposed to 0 or 5 IU/ml IL-2 (Proleukin) for 18 h and the median fluorescence intensity of CD25, CTLA-4 and FoxP3 examined by flow cytometry for (a) CD4, CD8 and T_{reg} cells and (b) CD4 subsets defined by CD25 versus CD127 expression. Data are mean \pm standard error of the mean (s.e.m.). Significant effects of IL-2 analysed by paired t -tests (a) and two-way analysis of variance (ANOVA) with Bonferroni's *post-hoc* test (b) are shown.




Fig. 3. (a–c) Effect of very low dose interleukin (IL)–2 on regulatory T cells (T_{reg}) and T effector functional and activation phenotypes. Peripheral blood mononuclear cells (PBMCs) from patients with autoimmune hepatitis (AIH) and liver infiltrating leucocytes from autoimmune liver diseases (AILD) livers were exposed to 0 or 5 IU/ml IL-2 (Proleukin) for 18 h and the expression of functional and activation markers by T cell subsets analysed by flow cytometry. (a) Expression of tumour necrosis factor (TNF) receptor superfamily members by blood and liver T_{reg} . (b) Expression of CD39, CD73, T cell immunoglobulin and mucin domain-containing-3 (TIM3) and lymphocyte-activation gene 3 (LAG3) by blood T_{reg} . (c) Expression of CD69, programmed death 1 (PD-1), 2B4 and granzyme B by blood CD4, CD8 and T_{reg} . Data are mean \pm standard error of the mean (s.e.m.) for 2–6 donors. (d–f) Very low dose IL-2 increases T_{reg} suppressive ability in a mechanism involving cytotoxic T lymphocyte antigen-4 (CTLA-4). $CD4^+CD25^+CD127^-$ T_{reg} and autologous $CD4^+CD25^-$ T responder cells were isolated from PBMC of control individuals. T responders were labelled with cell trace violet and following overnight exposure of T_{reg} to 0 or 5 IU/ml Proleukin were co-cultured with the T_{reg} in the presence of anti-CD3 and dendritic cells, with or without CTLA-4 blockade. Cell trace violet dilution indicating T responder cell division was analysed by flow cytometry at 5 days. (d) Representative flow cytometry histograms of T responder division showing percentage division. (e) Division index summary data ($n = 3$) of T responders in the presence of T_{reg} pretreated with 0 or 5 IU/ml Proleukin. (f) Division index summary data ($n = 3$) of T responders in the presence of T_{reg} pretreated with 0 or 5 IU/ml Proleukin with or without anti-CTLA-4. Data are mean \pm standard error of the mean (s.e.m.). Significant effects of IL-2 and CTLA-4 blockade analysed by paired t -tests (e) and one-way analysis of variance (ANOVA) with Bonferroni *post-hoc* test (f) are shown.




Fig. 4. Very low dose interleukin (IL)-2 does not down-regulate liver-homing CXCR3 receptor on regulatory T cells (T_{reg}). (a) Peripheral blood mononuclear cells (PBMCs) were isolated from controls and autoimmune hepatitis (AIH) patients and T_{reg} phenotyped for CXCR3 by flow cytometry *ex vivo*. (b,c) CXCR3 expression was measured in PBMC T_{reg} from patients with AIH, primary sclerosing cholangitis (PSC) or primary biliary cholangitis (PBC) *ex vivo* and after culture for 18 h and/or 3 days in 5 IU/ml IL-2 (Proleukin). Expression of CXCR3 by $CD4^+CD25^+CD127^- T_{reg}$ of one representative donor from each disease cohort is shown in (b) and expression summarized for AIH, PSC donors and PBC donors in (c). Data are mean \pm standard error of the mean (s.e.m.).

significantly with culture in VLDP up to 3 days in AILD patients and controls (Fig. 5a). No differences in frequency were seen between any of the patient or control groups at either condition. The naive ($CD45RA^+CCR7^+$) subset of

T_{reg} has been reported to maintain the T_{reg} -specific demethylated region (TSDR) following expansion, making them an attractive option for T_{reg} cell immunotherapy [26,27]; thus, we evaluated the effect of 3 days VLDP on frequencies




Fig. 5. Low dose Proleukin treatment increases total regulatory T cell (T_{reg}) frequencies and promotes the strongest T_{reg} phenotype in the CD45Ra⁻CCR7⁻ effector memory population of T_{reg} . (a) Peripheral blood mononuclear cells (PBMCs) from controls, autoimmune hepatitis (AIH), primary sclerosing cholangitis (PSC) and primary biliary cholangitis (PBC) patients were exposed to 0 or 5 IU/ml IL-2 (Proleukin) for 3 days and the frequency of CD4⁺CD25⁺CD127⁻forkhead box protein 3 (FoxP3)⁺ T_{reg} assessed by flow cytometry. Significant effects of IL-2 were analysed by two-way analysis of variance (ANOVA) with Bonferroni's *post-hoc* analysis. Frequencies of memory and naive subsets of CD4⁺CD25⁺CD127⁻FoxP3⁺ T_{reg} as defined by CD45Ra and CCR7 expression were also determined at 3 days by flow cytometry. (b) Representative flow cytometry density plot of CCR7 *versus* CD45Ra for one PSC donor after 3 days of culture in 0 or 5 IU/ml Proleukin and summary data for all donors (c). Significant effects of IL-2 on the frequencies of the different memory/naive subsets were compared by two-way ANOVA with Bonferroni's *post-hoc* analysis, but no significant differences were identified. (D) Expression of cytotoxic T lymphocyte antigen-4 (CTLA-4), CD25 and FoxP3 by each memory or naive subset of T_{reg} . Data are mean \pm standard error of the mean (s.e.m.) (control ($n = 3$), AIH ($n = 4$), PSC ($n = 3$) and PBC ($n = 6$), central memory (CM) (CD45Ra⁻CCR7⁺); naive (CD45Ra⁺CCR7⁺); terminally differentiated effector memory (TEMRA) (CD45Ra⁺CCR7⁻); effector memory (EM) (CD45Ra⁻CCR7⁻). Effects of 5 IU/ml Proleukin on the expression of these markers by each subset were analysed by two-way ANOVA with Bonferroni's *post-hoc* tests. Stars indicate where there was a significant effect of IL-2 upon the expression of the marker by the subset (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$). Braces with stated P -values indicate significant differences in expression by the subsets under 5 IU/ml Proleukin conditions. With the exception of CTLA-4 expression by naive *versus* EM cells in PSC patients, or of FoxP3 expression by CN *versus* naive, CM *versus* TEMRA or TEMRA *versus* effector memory in PBC patients, there were no significant differences between the subsets for expression of any marker in the absence of IL-2 treatment.

of each of the maturation subsets of CD25⁺CD127⁻FoxP3⁺ T_{reg} defined by CD45RA and CCR7 expression [28]. VLDP did not alter the proportions of the maturation subsets in any AILD patient cohorts (AIH, PSC, PBC) or controls (Fig. 5b,c). None the less, as before, VLDP increased significantly the expression of IL-2 regulated markers CTLA-4, CD25 and FoxP3 on T_{reg} from all subsets. Overall, across AIH, PSC, PBC and control, the effector memory (CD45RA⁻CCR7⁻) subset of CD25⁺CD127⁻FoxP3⁺ T_{reg} cells showed the greatest induction of T_{reg} functional markers with VLDP treatment (Fig. 5d).

Liver-infiltrated T_{reg} are restricted predominantly to the CD45RA⁻CCR7⁻ population

Blood CD25⁺CD127⁻FoxP3⁺ T_{reg} subsets in AILD patients are comprised of approximately 60% effector memory (CD45RA⁻CCR7⁻), 15–20% central memory (CD45RA⁻CCR7⁺) and 15–20% naive (CD45RA⁺CCR7⁺) populations. Because VLDP increased significantly the expression of T_{reg} functional markers CTLA-4, CD25 and FoxP3, especially in the CD45RA⁻CCR7⁻ subset of T_{reg} , we investigated the frequency of these subsets in the inflamed human liver. We found that the effector memory CD45RA⁻CCR7⁻ subset of CD25⁺CD127⁻FoxP3⁺ T_{reg} population is increased significantly (95%) in the inflamed human livers compared to peripheral blood (60%) (Fig. 6a,b).

Very low dose Proleukin up-regulates expression of the anti-apoptotic molecule Bcl-2 on T_{reg} compared to effector T cells

The anti-apoptotic protein Bcl-2 is regulated differentially in effector T cells *versus* T_{reg} [29]. To assess whether VLDP alters Bcl-2 expression, lymphocytes from AILD patients and controls were treated with 5 IU/ml Proleukin for 18 h or 3 days and their Bcl-2 expression was evaluated. While expression by total CD4⁺ and CD8⁺ T cells was not altered up to 3 days, VLDP tended to increase blood T_{reg}

expression of Bcl-2 by 18 h (Fig. 7a). By 3 days, a significant twofold up-regulation of Bcl-2 expression in T_{reg} of patients with AIH was seen (Fig. 7b), and the same trend was observed in PSC and PBC (Fig. 7c). We also verified the effect of VLDP on Bcl-2 in AILD liver T_{reg} at 3 days. Consistent with the response found in blood, we observed selective up-regulation of Bcl-2 in liver T_{reg} (Fig. 7d), which were of PBC and PSC disease backgrounds. In view of the STAT-5 response by the CD4⁺CD25⁺CD127⁺ subset with sustained exposure to VLDP, we analysed the effect of VLDP on Bcl-2 expression by each CD4⁺ T cell subset, defined by CD25 *versus* CD127 expression at 3 days in blood (Fig. 7e,f) and liver (Fig. 7g), and saw that in blood the CD4⁺CD25⁺CD127⁺ subset also showed increased Bcl-2 expression with Proleukin; however, the relative fold increase in Bcl-2 expression by this effector subset was less compared to that observed for T_{reg} .

Discussion

IL-2 is a crucial cytokine for survival and function of T cells, including T_{reg} . We demonstrated in this study that very low dose clinical grade IL-2 (Proleukin) induces STAT-5 phosphorylation selectively in CD4⁺CD25⁺CD127⁻ T_{reg} from blood and liver of patients with AILD and is accompanied not only by a series of phenotypic and functional changes, but also up-regulates Bcl-2 to support T_{reg} survival. These data support existing evidence in autoimmune conditions such as diabetes and vasculitis that VLDIL-2 administration contributes to the maintenance of self-tolerance by increasing T_{reg} frequency [23,25,30]. Our study extends the potential of VLDP therapy to the treatment of autoimmune liver diseases.

Administration of the IL-2 or the IL-2 : anti-IL-2 monoclonal antibody complex reduces autoimmune disease in rodent models [31,32]. Studies in vasculitis, graft-*versus*-host disease, systemic lupus erythematosus and type 1




Fig. 6. Comparison of the frequencies of memory and naive regulatory T cell (T_{reg}) populations in blood and liver. (a) Frequencies of memory and naive subsets of $CD4^+CD25^+CD127^- T_{reg}$ as defined by CD45Ra and CCR7 expression were determined by flow cytometry for peripheral blood mononuclear cells (PBMCs) and liver infiltrating lymphocytes from patients with autoimmune liver diseases (AILD). (a) Representative flow cytometry density plot for expression of CD45Ra and CCR7 by blood and liver infiltrating T_{reg} showing the four subsets including: central memory (CM) ($CD45Ra^-CCR7^+$); naive ($CD45Ra^+CCR7^+$); terminally differentiated effector memory (TEMRA) ($CD45Ra^+CCR7^-$); effector memory (EM) ($CD45Ra^-CCR7^-$). (b) Summary frequencies for each memory and naive subset in blood [$n = 6$ donors with autoimmune hepatitis (AIH)] and liver [$n = 4$ donors with AILDs including primary sclerosing cholangitis (PSC) and primary biliary cholangitis (PBC)]. Data are mean \pm standard error of the mean (s.e.m.). Significance was tested by non-matched two-way analysis of variance (ANOVA) and showed significant interaction for subset *versus* tissue $P < 0.0001$. Bonferroni's *post-hoc* tests identified significant differences between blood and liver in the frequencies of CM and EM cells, as indicated. There was no significant effect of tissue on subset distribution but there was a significant effect of subset $P < 0.0001$.

diabetes suggest that an IL-2 dose range of $0.3\text{--}3.0 \times 10^6$ IU/m² is well tolerated, and achieves a preferential increase in the percentage of T_{reg} with no induction of effector T cell activation [23,25,33,34]. The dose range we have

identified working with clinical grade IL-2 (Proleukin) *in vitro* is similar to other investigators, who have applied VLDP in autoimmune diseases [11,23,25,34]. High-dose IL-2 is approved in cancer therapy, where it promotes tumour killing by activating NK cells and $CD4^+$ and $CD8^+$ effector T cells [12]; however, pulmonary vascular leakage occurs as a side effect to this treatment [35]. By avoiding the activation of effector cells, VLDP should be safe for clinical application in general. Indeed, a former dose escalation study of low-dose IL-2 for treatment of graft-*versus*-host disease reported very few adverse events except skin induration in a minority of study patients. Other potentially linked rare side effects include fatigue, malaise, fever, thrombocytopenia and raised serum creatinine [23]. Despite this, it is prudent that low-dose Proleukin therapy in a hospital setting be restricted to patients with normal cardiac and pulmonary function.

IL-2 is crucial for T_{reg} function. Binding of IL-2 to IL-2RA (CD25) on T_{reg} leads to phosphorylation of STAT-5, resulting in up-regulation of FoxP3 [36] and functional surface markers. FoxP3 is essential for T_{reg} development and function, as evident from the occurrence of immunodysregulation polyendocrinopathy enteropathy X-linked (IPEX) syndrome in patients with mutations in FoxP3 [37,38]. Although T_{reg} do not secrete IL-2, their high expression of CD25 allows them to consume and respond to low concentrations of exogenous IL-2 [39,40]. $CD4^+CD25^-CD127^+$ effector T cells express intermediate-affinity IL-2 receptors, IL-2RB and IL-2RG constitutively; thus, they require higher concentrations of IL-2 for activation in the absence of T cell receptor (TCR) engagement [41]. Importantly, we observed that applying VLDP of less than 5 IU/ml could enhance FoxP3 selectively in T_{reg} compared to $CD8^+$, $CD4^+CD25^-CD127^{+/-}$ and $CD4^+CD25^+CD127^+$ effector T cells from the blood and liver of patients with AILD. There was no noticeable activation of these effector subsets based on expression of CD69, granzyme B or 2B4.

CTLA-4 is an essential functional marker on T_{reg} [42]. It functions by removing its co-stimulatory ligands, CD80/CD86, from antigen-presenting cells by transendocytosis [43]. Mice deficient in CTLA-4 display an autoimmune phenotype and immune dysregulation is observed in patients with CTLA-4 polymorphisms [44]. Our data suggest that VLDP up-regulates CTLA-4 expression by T_{reg} , but has no impact on $CD4^+CD25^+CD127^+$ effector T cells. Furthermore, we identify that CTLA-4 is a significant mediator in the mechanisms by which VLDP confers enhanced suppressive potential to T_{reg} ; VLDP also maintains T_{reg} functional markers such as GITR and OX40.

We have reported recently that the inflamed liver micro-environment is deficient in IL-2 protein and that activated intrahepatic T cells are the main source of IL-2 in the human liver [4]. Local consumption of IL-2 by immune cells, including T_{reg} , creates a microenvironment that is




Fig. 7. Very low dose interleukin (IL)-2 up-regulates Bcl-2 expression in regulatory T cells (T_{reg}) from blood and liver. Peripheral blood mononuclear cells (PBMCs) from autoimmune hepatitis (AIH) patients (a,b,e), primary sclerosing cholangitis (PSC) patients (c,f), primary biliary cholangitis (PBC) patients (c,f) and liver-infiltrating lymphocytes from autoimmune liver diseases (AILD) livers (d,g) were exposed to 0 or 5 IU/ml IL-2 (Proleukin) for 18 h (a) or 3 days (b–g) and the median fluorescence intensity (MFI) of Bcl-2 on CD4, CD8 and T_{reg} examined by flow cytometry (a–d). (E–G) Day 3 Bcl-2 expression by blood (e,f) and liver (g) CD4⁺ T cell subsets defined by CD25 versus CD127 expression. Data are mean \pm standard error of the mean (s.e.m.). Significant effects of IL-2 were assessed by paired *t*-tests (a–d) and two-way analysis of variance (ANOVA) (e–g).

deficient in IL-2 to the level that it is unable to support T_{reg} function. This finding is consistent with our previous observation that only some of the FoxP3⁺ T_{reg} in the liver undergo STAT-5 phosphorylation [45]. We observed that liver T_{reg} had higher baseline expressions of CTLA-4, CD25 and FoxP3 than peripheral T_{reg} and these were equivalent

to levels seen on peripheral blood T_{reg} after IL-2 stimulation. A higher baseline expression probably reflects the more activated, effector state of liver *versus* blood lymphocytes. Liver T_{reg} also expressed higher levels of the TNF receptor superfamily members CD137, OX40 and GITR than peripheral T_{reg} , and comparing our current data in

peripheral blood with our previous phenotyping of intrahepatic T_{reg} in AILD we identify that CD69 (40%), LAG3 (20%), Tim3 (5%) and cytolytic granzyme B (5%) are also all up-regulated on T_{reg} in the liver compared to periphery (< 5% in peripheral blood) [4]. This is in contrast to CD39 and PD-1, which are expressed similarly on T_{reg} at both sites [4].

The effectiveness of T_{reg} function within the inflamed tissue setting might be compromised due to the action of inflammatory cytokines such as IL-6, IL-8, IL-12, IFN- γ and IL-1 β [4], which can enhance effector T cell activation and proliferation at the expense of T_{reg}. Thus, exogenous IL-2 such as VLDP might help to counteract these inflammatory signals [46,47]. CTLA-4-mediated T_{reg} suppressive function is via depletion of CD80/86 through transendocytosis [43]. Excitingly, we observed that VLDP enhances T_{reg} suppressive function via CTLA-4.

Recruitment of peripheral blood T_{reg} to the site of hepatic inflammation is dependent upon expression of the tissue-homing CXCR3 chemokine receptor [45,48,49]. We found expression of CXCR3 on approximately 50% of peripheral blood T_{reg} from AILD patients, comparable to controls, suggesting that cells from patients, even on immunosuppression, retain the liver-homing receptors for recruitment. Moreover, we report that CXCR3 expression was unaffected by VLDP, implying that T_{reg} of patients on VLDP therapy would retain the capability to home to inflamed hepatic lobules and portal tracts. We have reported recently that human liver-infiltrating T_{reg} are mainly of CD45RA⁻CCR7⁻ phenotype [4]. Our findings now demonstrate that exogenous VLDIL-2 up-regulates T_{reg} functional markers predominantly in this effector (suppressor) population in all types of AILD. A recent study also described high expression of CXCR3 in CD4⁺ T cells in early-stage PBC, which was associated with increased demethylation of the CXCR3 promoter. This up-regulation was most striking in the activated memory CD45RO⁺ population [50]. Taken together, these data suggest that CD4⁺ T_{reg} from patients with AILD will recruit to the human liver and be potentiated functionally by IL-2 therapy.

T_{reg} depend upon IL-2 for expansion and survival and cells undergo apoptosis upon IL-2 deprivation [51]. Bcl-2 is a critical target in IL-2 signalling [18,52], helping to protect responding cells from apoptosis. A recent study also confirmed that use of a pan-Bcl-2 inhibitor leads to profound down-regulation of FoxP3 and CTLA-4 and a reduction in the suppressive function of T_{reg} [53]. We demonstrated that VLDP enhances Bcl-2 expression selectively in T_{reg} but not CD8⁺ T cells or CD4⁺ T effector cells from both peripheral blood and liver. This is an important observation because enhanced survival of immune-regulatory T_{reg} is critical to the maintenance of tolerance.

In conclusion, our findings provide compelling evidence to support the design of T_{reg}-directed Phase I and II clinical

trials administering VLDP as a cytokine monotherapy or in combination with autologous T_{reg} cell therapy in AILD. Given its short half-life, regular doses of VLDP may be required to maintain efficacy.

Acknowledgements

We would like to thank Matthew Graeme MacKenzie at the University of Birmingham Technology Hub Services for his help with cell sorting. We would also like to thank medical, surgical and anaesthetic staff from the UHB NHS Foundation Trust for their help with tissue acquirement and patients for their donations.

Author contributions

Y. H. O. and H. C. J. designed the experiments. H. C. J. L. E. J. and P. L. performed the experiments, analysed the data and prepared the figures. H. C. J., L. E. J., P. L., D. H. A. and Y. H. O. wrote the manuscript. G. W., M. C. and G. H. consented patients and collected clinical data. All authors reviewed the manuscript.

Disclosure

Y. H. O and H. C. J. are funded by Clinician Scientist Award from the Medical Research Council, Queen Elizabeth Hospital Charity and National Institute for Health Research Liver Biomedical Research Unit, Birmingham. All authors declare that there are no financial conflicts of interest associated with this work. This paper presents independent research supported by the Birmingham NIHR Liver Biomedical Research Unit based at the University of Birmingham and University Hospitals Birmingham NHS Foundation Trust. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

References

- 1 Sakaguchi S, Sakaguchi N, Asano M, Itoh M, Toda M. Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor alpha-chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. *J Immunol* 1995; **155**:1151–64.
- 2 Liu W, Putnam AL, Xu-Yu Z *et al*. CD127 expression inversely correlates with FoxP3 and suppressive function of human CD4⁺ Treg cells. *J Exp Med* 2006; **203**:1701–11.
- 3 Sebode M, Peiseler M, Franke B *et al*. Reduced FOXP3(+) regulatory T cells in patients with primary sclerosing cholangitis are associated with IL2RA gene polymorphisms. *J Hepatol* 2014; **60**: 1010–6.
- 4 Chen YY, Jeffery HC, Hunter S *et al*. Human intrahepatic Tregs are functional, require IL-2 from effector cells for survival and are susceptible to Fas ligand mediated apoptosis. *Hepatology* 2016; **64**:138–50.

- 5 Lan RY, Cheng C, Lian ZX *et al.* Liver-targeted and peripheral blood alterations of regulatory T cells in primary biliary cirrhosis. *Hepatology* 2006; **43**:729–37.
- 6 Beuers U, Gershwin ME, Gish RG *et al.* Changing nomenclature for PBC: from ‘cirrhosis’ to ‘cholangitis’. *Hepatology* 2015; **62**: 1620–2.
- 7 Oo YH, Adams DH. Regulatory T cells and autoimmune hepatitis: defective cells or a hostile environment? *J Hepatol* 2012; **57**:6–8.
- 8 Oo YH, Adams DH. Regulatory T cells and autoimmune hepatitis: what happens in the liver stays in the liver. *J Hepatol* 2014; **61**:973–5.
- 9 Longhi MS, Hussain MJ, Mitry RR *et al.* Functional study of CD4+CD25+ regulatory T cells in health and autoimmune hepatitis. *J Immunol* 2006; **176**:4484–91.
- 10 Wing K, Sakaguchi S. Regulatory T cells exert checks and balances on self tolerance and autoimmunity. *Nat Immunol* 2010; **11**: 7–13.
- 11 Hartemann A, Bensimon G, Payan CA *et al.* Low-dose interleukin 2 in patients with type 1 diabetes: a phase 1/2 randomised, double-blind, placebo-controlled trial. *Lancet Diabetes Endocrinol* 2013; **1**:295–305.
- 12 Yang JC, Sherry RM, Steinberg SM *et al.* Randomized study of high-dose and low-dose interleukin-2 in patients with metastatic renal cancer. *J Clin Oncol* 2003; **21**:3127–32.
- 13 Klatzmann D, Abbas AK. The promise of low-dose interleukin-2 therapy for autoimmune and inflammatory diseases. *Nat Rev Immunol* 2015; **15**:283–94.
- 14 Taniguchi T, Miyazaki T, Minami Y *et al.* IL-2 signaling involves recruitment and activation of multiple protein tyrosine kinases by the IL-2 receptor. *Ann N Y Acad Sci* 1995; **766**:235–44.
- 15 Van Parijs L, Refaeli Y, Lord JD, Nelson BH, Abbas AK, Baltimore D. Uncoupling IL-2 signals that regulate T cell proliferation, survival, and Fas-mediated activation-induced cell death. *Immunity* 1999; **11**:281–8.
- 16 Kirken RA, Rui H, Malabarba MG *et al.* Activation of JAK3, but not JAK1, is critical for IL-2-induced proliferation and STAT5 recruitment by a COOH-terminal region of the IL-2 receptor beta-chain. *Cytokine* 1995; **7**:689–700.
- 17 Moriggl R, Topham DJ, Teglund S *et al.* Stat5 is required for IL-2-induced cell cycle progression of peripheral T cells. *Immunity* 1999; **10**:249–59.
- 18 Lord JD, McIntosh BC, Greenberg PD, Nelson BH. The IL-2 receptor promotes lymphocyte proliferation and induction of the c-myc, bcl-2, and bcl-x genes through the trans-activation domain of Stat5. *J Immunol* 2000; **164**:2533–41.
- 19 Hofer T, Krichevsky O, Altan-Bonnet G. Competition for IL-2 between regulatory and effector T cells to chisel immune responses. *Front Immunol* 2012; **3**:268.
- 20 Busse D, de la Rosa M, Hobiger K *et al.* Competing feedback loops shape IL-2 signaling between helper and regulatory T lymphocytes in cellular microenvironments. *Proc Natl Acad Sci USA* 2010; **107**:3058–63.
- 21 Gutierrez-Ramos JC, Andreu JL, Revilla Y, Vinuela E, Martinez C. Recovery from autoimmunity of MRL/lpr mice after infection with an interleukin-2/vaccinia recombinant virus. *Nature* 1990; **346**:271–4.
- 22 Zielasek J, Burkart V, Naylor P, Goldstein A, Kiesel U, Kolb H. Interleukin-2-dependent control of disease development in spontaneously diabetic BB rats. *Immunology* 1990; **69**:209–14.
- 23 Koreth J, Matsuoka K, Kim HT *et al.* Interleukin-2 and regulatory T cells in graft-versus-host disease. *N Engl J Med* 2011; **365**:2055–66.
- 24 Matsuoka K, Koreth J, Kim HT *et al.* Low-dose interleukin-2 therapy restores regulatory T cell homeostasis in patients with chronic graft-versus-host disease. *Sci Transl Med* 2013; **5**: 179ra43.
- 25 Saadoun D, Rosenzweig M, Joly F *et al.* Regulatory T-cell responses to low-dose interleukin-2 in HCV-induced vasculitis. *N Engl J Med* 2011; **365**:2067–77.
- 26 Sakaguchi S, Vignali DA, Rudensky AY, Neeb RE, Waldmann H. The plasticity and stability of regulatory T cells. *Nat Rev Immunol* 2013; **13**:461–7.
- 27 Canavan JB, Scotta C, Vossenkamper A *et al.* Developing *in vitro* expanded CD45RA+ regulatory T cells as an adoptive cell therapy for Crohn’s disease. *Gut* 2015; **65**:584–94.
- 28 Sallusto F, Lenig D, Forster R, Lipp M, Lanzavecchia A. Two subsets of memory T lymphocytes with distinct homing potentials and effector functions [see comments]. *Nature* 1999; **401**: 708–12.
- 29 Pandiyan P, Zheng L, Ishihara S, Reed J, Lenardo MJ. CD4+CD25+Foxp3+ regulatory T cells induce cytokine deprivation-mediated apoptosis of effector CD4+ T cells. *Nat Immunol* 2007; **8**:1353–62.
- 30 Setoguchi R, Hori S, Takahashi T, Sakaguchi S. Homeostatic maintenance of natural Foxp3(+) CD25(+) CD4(+) regulatory T cells by interleukin (IL)–2 and induction of autoimmune disease by IL-2 neutralization. *J Exp Med* 2005; **201**:723–35.
- 31 Wilson MS, Pesce JT, Ramalingam TR, Thompson RW, Cheever A, Wynn TA. Suppression of murine allergic airway disease by IL-2:anti-IL-2 monoclonal antibody-induced regulatory T cells. *J Immunol* 2008; **181**:6942–54.
- 32 Webster KE, Walters S, Kohler RE *et al.* In vivo expansion of T reg cells with IL-2-mAb complexes: induction of resistance to EAE and long-term acceptance of islet allografts without immunosuppression. *J Exp Med* 2009; **206**:751–60.
- 33 von Spee-Mayer C, Siegert E, Abdirema D *et al.* Low-dose interleukin-2 selectively corrects regulatory T cell defects in patients with systemic lupus erythematosus. *Ann Rheum Dis* 2015; **75**:1407–15.
- 34 Rosenzweig M, Churlaud G, Mallone R *et al.* Low-dose interleukin-2 fosters a dose-dependent regulatory T cell tuned milieu in T1D patients. *J Autoimmun* 2015; **58**:48–58.
- 35 Rosenstein M, Ettinghausen SE, Rosenberg SA. Extravasation of intravascular fluid mediated by the systemic administration of recombinant interleukin 2. *J Immunol* 1986; **137**:1735–42.
- 36 Fontenot JD, Gavin MA, Rudensky AY. Foxp3 programs the development and function of CD4+CD25+ regulatory T cells. *Nat Immunol* 2003; **4**:330–6.
- 37 Bennett CL, Christie J, Ramsdell F *et al.* The immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) is caused by mutations of FOXP3. *Nat Genet* 2001; **27**: 20–1.
- 38 d’Hennezel E, Ben-Shoshan M, Ochs HD *et al.* FOXP3 forkhead domain mutation and regulatory T cells in the IPEX syndrome. *N Engl J Med* 2009; **361**:1710–3.
- 39 Stauber DJ, Debler EW, Horton PA, Smith KA, Wilson IA. Crystal structure of the IL-2 signaling complex: paradigm for a heterotrimeric cytokine receptor. *Proc Natl Acad Sci USA* 2006; **103**:2788–93.

- 40 Wang X, Rickert M, Garcia KC. Structure of the quaternary complex of interleukin-2 with its alpha, beta, and gamma receptors. *Science* 2005; **310**:1159–63.
- 41 Grant AJ, Roessler E, Ju G, Tsudo M, Sugamura K, Waldmann TA. The interleukin 2 receptor (IL-2R): the IL-2R alpha subunit alters the function of the IL-2R beta subunit to enhance IL-2 binding and signaling by mechanisms that do not require binding of IL-2 to IL-2R alpha subunit. *Proc Natl Acad Sci USA* 1992; **89**:2165–9.
- 42 Walker LS, Sansom DM. The emerging role of CTLA4 as a cell-extrinsic regulator of T cell responses. *Nat Rev Immunol* 2011; **11**:852–63.
- 43 Qureshi OS, Zheng Y, Nakamura K *et al.* Trans-endocytosis of CD80 and CD86: a molecular basis for the cell-extrinsic function of CTLA-4. *Science* 2011; **332**:600–3.
- 44 Schubert D, Bode C, Kenefick R *et al.* Autosomal dominant immune dysregulation syndrome in humans with CTLA4 mutations. *Nat Med* 2014; **20**:1410–6.
- 45 Oo YH, Weston CJ, Lalor PF *et al.* Distinct roles for CCR4 and CXCR3 in the recruitment and positioning of regulatory T cells in the inflamed human liver. *J Immunol* 2010; **184**:2886–98.
- 46 Jeffery LE, Wood AM, Qureshi OS *et al.* Availability of 25-hydroxyvitamin D(3) to APCs controls the balance between regulatory and inflammatory T cell responses. *J Immunol* 2012; **189**:5155–64.
- 47 Liaskou E, Jeffery LE, Trivedi PJ *et al.* Loss of CD28 expression by liver-infiltrating T cells contributes to pathogenesis of primary sclerosing cholangitis. *Gastroenterology* 2014; **147**:221–32 e7.
- 48 Curbishley SM, Eksteen B, Gladue RP, Lalor P, Adams DH. CXCR3 activation promotes lymphocyte transendothelial migration across human hepatic endothelium under fluid flow. *Am J Pathol* 2005; **167**:887–99.
- 49 Erhardt A, Wegscheid C, Claass B *et al.* CXCR3 deficiency exacerbates liver disease and abrogates tolerance in a mouse model of immune-mediated hepatitis. *J Immunol* 2011; **186**:5284–93.
- 50 Lleo A, Zhang W, Zhao M *et al.* DNA methylation profiling of the X chromosome reveals an aberrant demethylation on CXCR3 promoter in primary biliary cirrhosis. *Clin Epigenetics* 2015; **7**:61.
- 51 Downward J. Signal transduction. A target for PI(3) kinase. *Nature* 1995; **376**:553–4.
- 52 Miyazaki T, Liu ZJ, Kawahara A *et al.* Three distinct IL-2 signaling pathways mediated by bcl-2, c-myc, and lck cooperate in hematopoietic cell proliferation. *Cell* 1995; **81**:223–31.
- 53 Kim PS, Jochems C, Grenga I *et al.* Pan-Bcl-2 inhibitor, GX15-070 (obatoclax), decreases human T regulatory lymphocytes while preserving effector T lymphocytes: a rationale for its use in combination immunotherapy. *J Immunol* 2014; **192**:2622–33.

Supporting information

Additional Supporting Information may be found in the online version of this article at the publisher's website.

Fig. S1. Flow cytometry gating strategy for the analysis of signal transducer and activator of transcription-5 (STAT-5) expression by regulatory T cells (T_{reg}) and eight other immune cell subsets.

Fig. S2. Very low dose interleukin (IL)–2 selectively up-regulates signal transducer and activator of transcription-5 (STAT-5) in peripheral regulatory T cells (T_{reg}) from autoimmune hepatitis (AIH) and primary biliary cholangitis (PBC) patient bloods. Peripheral blood mononuclear cells (PBMCs) from patients with AIH, PBC or primary sclerosing cholangitis (PSC) were stimulated for 10 min with IL-2 (Proleukin) 0–100 IU/ml and the percentage expression of phosphorylated (Y694)STAT5 by each leucocyte population assessed by flow cytometry at each IL-2 dose. Data are mean \pm standard error of the mean (s.e.m.) for five donors (AIH remission), four donors (AIH relapse) and three donors (PBC and PSC).

Fig. S3. Flow cytometry gating strategy (a) to define CD4, CD8 and regulatory T cell (T_{reg}) populations and representative overlays (b,d) for markers of T cell activation and function on CD4, CD8 and T_{reg} in autoimmune liver disease (AILD) blood [autoimmune hepatitis (AIH)] and liver [primary sclerosing cholangitis (PSC)] following 18 h exposure to Proleukin 5 IU/ml.

Fig. S4. Effect of very low dose interleukin (IL)–2 on expression of IL-2-regulated regulatory T cell (T_{reg}) functional markers CD25, cytotoxic T lymphocyte antigen-4 (CTLA-4) and forkhead box protein 3 ($FoxP3^+$) by CD4, CD8 and T_{reg} cells from blood and liver. Peripheral blood mononuclear cells (PBMCs) from patients with autoimmune hepatitis (AIH), primary sclerosing cholangitis (PSC) and primary biliary cholangitis (PBC) and liver infiltrating lymphocytes from autoimmune liver disease (AILD) livers were exposed to 0 or 5 IU/ml IL-2 (Proleukin) for 18 h and the percentage expression of CD25, CTLA-4 and $FoxP3$ by CD4⁺, CD8⁺ and T_{reg} cells examined by flow cytometry. Data are mean \pm standard error of the mean (s.e.m.) for five donors (AIH), three donors (PSC) and five donors (liver).

Fig. S5. Sustained exposure to low dose interleukin (IL)–2 up-regulates signal transducer and activator of transcription-5 (STAT-5) selectively in CD4⁺CD25⁺CD127⁺ T cells as well as in regulatory T cells (T_{reg}). Peripheral blood mononuclear cells (PBMCs) from patients with autoimmune liver disease (AILD) were stimulated for 18 h with IL-2 (Proleukin) doses of 0, 1, 5 and 10 IU/ml and expression of phosphorylated (Y694)STAT-5 by each leucocyte population assessed by flow cytometry. (a) Percentage of each leucocyte population expressing pSTAT-5 at each IL-2 dose. (b) median fluorescence intensity (MFI) for pSTAT-5.

Fig. S6. Effect of very low dose interleukin (IL)–2 on the suppressive ability of regulatory T cells (T_{reg}) in autoimmune liver diseases. (a) CD4⁺CD25[−] T responder cells were isolated from peripheral blood mononuclear cells (PBMCs) of controls ($n = 3$) and the effect of blocking CD28-mediated co-stimulation with Abatacept [cytotoxic T lymphocyte antigen-4-immunoglobulin (CTLA-4-Ig)] on division index at 5 days examined. (b)

CD4⁺CD25⁺CD127⁻ T_{reg} and autologous CD4⁺CD25⁻ T responder cells were isolated from PBMC of two patients with autoimmune hepatitis (AIH) (one in remission and one in relapse). T responders were labelled with cell trace violet, and following overnight exposure of T_{reg} to 0 or 5

IU/ml Proleukin were co-cultured with the T_{reg} in the presence of anti-CD3 and dendritic cells, with or without CTLA-4 blockade. Cell trace violet dilution indicating T responder cell division was analysed by flow cytometry at 5 days.