

UNIVERSITY OF BIRMINGHAM

Research at Birmingham

An integrative literature review exploring the clinical management of delirium in patients with advanced cancer

Lawley, Hayley ; Hewison, Alistair

DOI:

[10.1111/jocn.13960](https://doi.org/10.1111/jocn.13960)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Lawley, H & Hewison, A 2017, 'An integrative literature review exploring the clinical management of delirium in patients with advanced cancer', *Journal of Clinical Nursing*, vol. 26, no. 23-24, pp. 4172-4183. <https://doi.org/10.1111/jocn.13960>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the peer reviewed version of the following article: Lawley H, Hewison A. An integrative literature review exploring the clinical management of delirium in patients with advanced cancer. *J Clin Nurs*. 2017;00:1–12. <https://doi.org/10.1111/jocn.13960>, which has been published in final form at [10.1111/jocn.13960](https://doi.org/10.1111/jocn.13960). This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy


While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



An integrative literature review exploring the clinical management of delirium in advanced cancer patients.

Journal:	<i>Journal of Clinical Nursing</i>
Manuscript ID	JCN-2017-0051.R2
Manuscript Type:	Review
Keywords:	Adult Nursing, Cancer, Death and Dying, End of Life Care, Management



[Type text]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

An integrative literature review exploring the clinical management of delirium in patients with advanced cancer.

For Peer Review

[Type text]

Abstract

Aim: The aim of this paper is to present the findings of an integrative literature review of the evidence for the clinical management of delirium in patients with advanced cancer.

Background: Patients with advanced cancer frequently experience delirium which can be distressing for both patients and their families. Current guidelines recommend that underlying causes of the delirium be addressed and a course of antipsychotics considered. However the research into the effectiveness of treatments for delirium in people with advanced cancer is limited.

Design: Integrative literature review

Data sources: Systematic searches of the MEDLINE, CINAHL, ProQuest Nursing and Allied Health and PsychInfo databases were conducted in April 2016 to include papers published in 2000 and later. The returns were screened using inclusion and exclusion criteria and the seven studies found to be suitable were subject to review.

Review Methods: Findings of the seven papers were extracted, appraised critically and reviewed using a narrative approach.

Results: A number of interventions, including the use of atypical antipsychotics, opioid rotation, methylphenidate hydrochloride and celiac plexus block were reported however there was limited evidence of their effectiveness. One study reported the use of exercise therapy as a non-pharmacological intervention.

Conclusion: A variety of interventions to treat delirium in patients with advanced cancer have been tested through non-blinded, non-randomised trials which has not produced a clear evidence-base for practice. There is a need for further research (particularly randomised control trials) to determine the most effective treatments for patients with advanced cancer experiencing delirium.

Key Words: literature review, delirium, palliative care, end of life, advanced cancer, neoplasm, management, nursing.

[Type text]

What does this paper contribute to the wider global clinical community?

Why is this review needed?

- There is a lack of evidence to inform the management of delirium in patients with advanced cancer. .
- The lack of evidence for the management of delirium in patients with advanced cancer means that patients, their families and carers, may not be receiving the most appropriate care.

Key Findings:

- Available research on the clinical management of delirium in advanced cancer is limited and of variable quality.
- A number of pharmacological and non-pharmacological interventions have been used to treat delirium in patients with advanced cancer however the evidence base for practice is limited.

How should these findings be used?

- These findings highlight the need for further investigation of the treatment of delirium in advanced cancer.
- Provide insights for nurses and other professionals on the effectiveness of the treatments currently in use.

[Type text]

The Review

Introduction

Patients with advanced cancer often experience delirium, yet strategies for its early detection, prevention, and management are limited (Caraceni and Simonetti 2009). Delirium is one of the most difficult syndromes to diagnose and treat and has an adverse impact on the quality of life for the patient and their family in the final days of life (Centeno et al 2004). There are various definitions of advanced cancer, however for the purpose of this paper, advanced cancer is that which cannot be cured and is likely to cause death (American Cancer Society, 2014). This means that End of Life concerns, such as preferred place of death, are real and valid for those diagnosed with advanced cancer. For example, although most people would prefer to die at home rather than in hospital (National Audit Office, 2008), delirium can result in longer hospital stays (Ljubisavljevic and Kelly, 2003), meaning patients may not be able to die in their preferred place. Moreover family members and carers generally provide the most care for their loved ones experiencing delirium which causes distress (Namba et al, 2007), which is often exacerbated as delirious patients are more likely to be non-compliant and risk putting themselves or others in danger (for example by pulling out IV lines/catheters) and being aggressive.

In view of this a need to examine the evidence for the clinical management of delirium in patients with advanced cancer was identified, in order to determine how it can be best managed. The aim of this review was to examine evidence for pharmacological and non-pharmacological treatments for delirium in patients with advanced cancer to determine the evidence base for practice and to highlight gaps in knowledge

[Type text]

Background

One of the most challenging aspects of end of life care, and a reason why many patients stay in and die hospital, is poor symptom control. Effective amelioration of symptoms including delirium, pain and nausea can help a patient to have a 'good death' (de Jong and Clarke, 2009). Although guidelines exist for the general and more specific management of delirium in cancer patients (NICE 2015, CCO 2010, Western Australia Cancer & Palliative Care Network 2010), the guidance concerning treatment in patients with advanced cancer is limited, in part because of the limited evidence available. The treatment and medication patients may receive as part of their cancer treatment can mean the standard approach to treating delirium is less effective.

Delirium is defined as a neurocognitive disorder characterised by a disturbance in attention and cognition, which can result from another medical condition (or from multiple aetiologies) for example pyrexia and malnutrition. The condition develops over a short period of time and cannot be explained by any another neurocognitive disorder (Diagnostic and Statistical Manual of Mental Disorders - V, 2013). Delirium can manifest in a number of ways (see table 1) and in practice, it is often nurses who recognise the symptoms when they first present. A variety of tools exist which can be used to help identify and diagnose the condition. For example the National Institute for Health and Care Excellence (NICE 2010) recommends the use of the Diagnostic and Statistical Manual of Mental Disorders – IV (DSM-IV) criteria or the Confusion Assessment Method (CAM), although other tools including the MDAS (Memorial Delirium Assessment Scale) and the MMSE (Mini-Mental State Examination) are available and often used in screening. Despite the availability of these tools however, the rate of detection of delirium in patients in practice is low. Ryan et al, 2013 found that nurses identified 63.6% of patients with delirium as confused or delirious,

[Type text]

1
2
3 however only 43.6% of those patients had the observation they were confused recorded in
4
5 their notes. In a study of patents with advanced cancer it was found that the overall detection
6
7 rate in patients with terminal cancer by any member of the palliative team was 44.9% and the
8
9 detection rate of the hypoactive subtype was only 20.5%, significantly lower than that of the
10
11 hyperactive/mixed subtypes (see below) (Fang et al 2008).
12
13

14
15 (Table 1 here)
16
17

18
19 Although the term delirium is widely used, there are distinct subtypes of delirium which are
20
21 based on their clinical presentation. They are: hyperactive, hypoactive and mixed, and
22
23 delirium with catatonic symptoms. For example, a comparison of symptoms of delirium in a
24
25 range of patients found that 50.15% had the hyperactive subtype, 24.61% the mixed subtype
26
27 and 19.93% the hypoactive subtype (Grover et al 2014). In other studies, the mixed type of
28
29 delirium was most common (Peterson et al 2006; Kim et al 2015). Patients with hyperactive
30
31 delirium are often restless, agitated and hyper-vigilant and experience perceptual disturbances
32
33 and delusions, whereas patients with hypoactive delirium demonstrate lethargy and minimal
34
35 spontaneous movement (Fang et al, 2009; Grover et al, 2014). The identification of different
36
37 subtypes raises the issue of whether they should be treated as separate conditions, especially
38
39 noting that patients experiencing hypoactive and mixed types of delirium have a poorer
40
41 prognosis than those with hyperactive delirium (Kim et al, 2015), suggesting different
42
43 treatment approaches may be needed.
44
45
46
47

48
49 Estimates for total cancer deaths in 2012 were 8.2 million (about 22,000 cancer deaths a day),
50
51 and by 2030 the global burden is expected to grow to 21.7 million new cancer cases and 13
52
53 million cancer deaths (American Cancer Society 2015). The prevalence of delirium in
54
55 patients with terminal cancer ranges from 28% to 85% (Minagawa et al 1996; de la Cruz et
56
57 al, 2015; Massie et al 1983). This large variation may be a result of the difficulties involved
58
59
60

[Type text]

1
2
3 in accurately identifying the symptoms of the different subtypes of delirium (particularly
4 hypoactive delirium), misdiagnosis, or a lack of the training staff need to make an
5
6
7 assessment. Patients with cancer are at particular risk of developing delirium for a number of
8
9
10 reasons, including exposure to certain drugs (opioids, corticosteroids, and benzodiazepines)
11
12 (Gaudreau, 2005), and the presence of bone metastases and haematological malignancies
13
14 (Ljubisavljevic and Kelly, 2003). In a recent literature review it was concluded that there
15
16 needs to be a greater understanding of delirium in palliative care, including its
17
18 pathophysiology and causation, as well as its treatment (Grassi et al 2015).
19
20
21
22
23
24

25 Current Clinical Management

26
27
28 Current approaches to the care of people with delirium involve non-pharmacological and
29
30 pharmacological management. Non-pharmacological interventions are recommended such as
31
32 nurses ensuring there is a clock visible in clinical areas and explaining to patients where they
33
34 are and introducing themselves and other staff members, helping to keep patients orientated
35
36 to time and place (NICE, 2010). Preventing precipitating factors such as constipation and
37
38 dehydration can also help (NICE 2010), and this entails nursing staff planning care to ensure
39
40 patients are mobilised, encouraged to drink and that bowel movements are monitored.
41
42
43 Pharmacologically, the recommended initial treatment is a short term, low dose course of
44
45 haloperidol or olanzapine. Hui et al (2010) for example found olanzapine was given to 17%
46
47 of delirious patients and haloperidol to 72% of the patients, perhaps reflecting the evidence
48
49 that Haloperidol has been shown to reduce symptoms of delirium in patients with cancer with
50
51 no significant side effects (Akechi et al 1996). Delirium is also one of the most common
52
53 reasons patients with advanced cancer are prescribed sedation (Alonso-Babarro et al, 2010;
54
55
56
57
58
59
60

[Type text]

1
2
3 Beller et al, 2014), which has been criticised on the basis that its use should be decided on an
4
5 individual basis (National Ethics Committee, Veterans Health Administration, 2007).
6
7

8
9 The management of delirium requires an interdisciplinary approach (Milisen et al, 2001;
10
11 Meagher, 2001) with nurses holding a key role in its prevention, detection and management
12
13 (Siddiqi et al 2006), as they are at the front line of care and provide the majority of screening
14
15 on admission to hospital. Tools available for nurses to use to identify the presence of
16
17 delirium include the Mini-Mental State Exam (MMSE) which is comprised of five sections
18
19 assessing orientation, registration, attention, recall and language. Its use has been effective in
20
21 detecting delirium and as a diagnostic aid (Anthony 1982; Faught 2014). On the basis of the
22
23 outcome nurses can refer patients to the appropriate healthcare professionals for further
24
25 assessment and treatment.
26
27

28
29
30
31 Despite the increasing amount of research into delirium experienced at the end of life in the
32
33 past decade (Close and Long, 2012; Harris, 2007), there are very few studies that focus on the
34
35 treatment of delirium in patients with advanced cancer., A review of the evidence exploring
36
37 the management of delirium in patients post-surgery and those being cared for in intensive
38
39 care settings, found delirium can be improved in around half of patients by removing
40
41 precipitating factors (Kang et al 2013). Although this highlights some useful evidence for
42
43 practice, the study did not specifically address the needs of patients with advanced cancer,
44
45 where it is often not possible to eliminate some of the precipitating factors identified.
46
47

48
49 Patients who are at the end of their life as a result of advanced cancer are often receiving
50
51 chemotherapy and opioid pain relief, which may mean the delirium experienced and the
52
53 treatment for that delirium is different to that experienced by patients who do not have
54
55 advanced cancer.
56
57
58
59
60

[Type text]

Design

The design of this study is an integrative literature review. The purpose of nursing research is to solve problems or answer questions that are relevant to nursing practice (Polit and Beck, 2014). The clinical management of delirium in patients with advanced cancer is one such problem. Although it is a common phenomenon (Minagawa et al 1996; de la Cruz et al, 2015; Massie et al 1983), there is little guidance in practice concerning how best to manage it. The review question was developed using the Population, Intervention and Outcome elements of the PICO model (Polit and Beck, 2014). This was used to guide the development of the question and because the focus was on studies reporting any intervention to alleviate delirium in patients with advanced cancer the C (comparator) element was not required. This process is summarised in Table 2.

(Table 2 here)

An initial search revealed there was little research reporting the treatment of delirium in patients with advanced cancer. Consequently although a systematic review is generally the recommended approach (Cullinan, 2005), it was not appropriate for this question because of the variety of interventions explored in the relevant literature. In view of this an integrative review was undertaken. In contrast to a systematic reviews, which aim to systematically search for, appraise and synthesise research evidence, aiming for exhaustive and comprehensive searching (Grant and Booth, 2009) and adhere to a strict design based on explicit, pre-specified, and reproducible methods (Gopalakrishnan and Ganeshkumar 2013) and generally focus on randomised controlled trials (Akenborg 2005), in an integrative of review, studies using diverse methodologies can be included and the approach has the potential to play a greater role in evidence-based practice for nursing (Whittemore and Knaf 2005). It is a review method which summarises previous empirical or theoretical literature, to

[Type text]

1
2
3 provide a more comprehensive understanding of a healthcare problem (Broome 1993 [cited in
4
5 Whittemore and Knafl, 2005]). The primary sources included in an integrative review need
6
7 to be logically grouped to facilitate analysis (Whittemore and Knafl, 2005). Ryan (2013)
8
9 recommends that the similarities and differences of the findings of different studies and
10
11 patterns in the data should be explored and in the review reported here there were
12
13 interventions that could be considered together.
14
15
16
17

18 Search Methods

19
20 Four databases; MEDLINE, CINAHL, ProQuest Nursing and Allied Health and PsychInfo
21
22 were searched because they were likely to include sources addressing the review question as
23
24 they focus on medicine, nursing and psychology. The searches were conducted in April 2016
25
26 to include papers published in 2000 and later. Table 3 identifies the search terms used. The
27
28 search terms were derived from the research question and the databases were searched using
29
30 those key words or phrases identified. Boolean operators (AND, OR, NOT) were used to
31
32 further refine the search. A 'wildcard' was also used, which allows multiple words to be
33
34 searched for which have the same truncation, for example 'deliri*' would return results for
35
36 'delirium' and 'delirious' (Aveyard, 2010). One article was found through hand-searching
37
38 through the reference lists of relevant articles (Horsely et al, 2001). Papers were excluded if
39
40 they were unavailable in English, but not if they were conducted in other countries and
41
42 reported in English. The inclusion and exclusion criteria used to determine the selection of
43
44 papers for review are shown in Table 4.
45
46
47
48
49

50
51 (Table 3 here)

52
53
54
55 (Table 4 here).
56
57
58
59
60

[Type text]

Search Outcomes

The results of the screening process are shown in the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) in figure 1 below. The selection of the papers for inclusion in the review was made following independent review of titles and abstracts by both authors. Titles and abstracts were identified by the systematic search of the databases noted earlier, guided by the application inclusion and exclusion criteria (Table 4). Papers found not to be relevant to the aims of study by their titles and abstracts were excluded. The full text versions of the twenty seven papers which met the inclusion criteria were obtained and read. On the basis of this second stage of independent review seven were found to be suitable for full critical review (see figure 1).

(Figure 1 here)

Data Abstraction

All the articles retrieved were read several times by one author and reviewed by the second author to gain a deeper understanding of the studies. A data abstraction form was used to record the key content of each paper in preparation for analysis (Polit and Beck, 2014). The design, aim sample, tool, results, limitations identified and recommendations made in the papers are summarised in table 5.

(Table 5 here)

Summary of the Studies included

Arai at al, 2013: This prospective study investigated terminal delirium in patients with pancreatic cancer and found that those who underwent a celiac plexus block had terminal delirium for a shorter time than those who did not.

[Type text]

1
2
3 Boettger and Breitbart, 2011: An open label naturally assigned study of patients with cancer
4 (46.7% had advanced cancer) that investigated whether aripiprazole could be used to treat
5 delirium. It found that aripiprazole may be safe and effective in treating delirium.
6
7

8
9 Moryl et al, 2005: Investigated 'switching' to methadone as part of an opioid rotation strategy
10 in patients with advanced cancer and found that methadone may be considered in treating
11 terminal delirium before the introduction of sedation.
12
13

14
15 Morita et al, 2005: Examined opioid rotation (morphine to fentanyl) in patients with cancer
16 receiving palliative care who were experiencing morphine induced delirium, and found that
17 the switch to fentanyl may alleviate the delirium.
18
19

20
21 Tatematsu et al, 2011: Investigated exercise therapy as an approach to reducing delirium in
22 patients with cancer and found that the antipsychotic dose was lower in those in who received
23 exercise therapy concluding that exercise therapy had potential as a non-pharmacological
24 intervention for the management of delirium.
25
26

27
28 Gagnon et al, 2005: Investigated the use of methylphenidate in treating hypoactive delirium
29 in patients with advanced cancer, finding that hypoactive delirium with no known aetiology
30 may be improved by prescribing methylphenidate.
31
32

33
34 Breitbart et al, 2002: Demonstrated the efficacy and safety of olanzapine for use in the
35 management of delirium patients with cancer in hospital.
36
37
38
39
40
41
42
43
44

45 Critical Appraisal

46
47 The articles were then subjected to critical appraisal which involved assessing the strengths
48 and weaknesses of the selected papers in order to assess their relevance to the review
49 question (Aveyard, 2010). The papers were reviewed to assess trustworthiness and quality
50 using a set of recognised criteria (Mhaskar et al, 2009; Polit and Beck, 2014). The
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

[Type text]

1
2
3 studies employing a quantitative design (Hek and Moule 2006), supplemented with key
4
5 questions developed by Polit and Beck (2014). These were combined in an appraisal form
6
7 which can be seen in figure 2.
8

9
10 (Figure 2 here).
11

12
13
14 The appraisal of the specific statistical tests used in the studies can be seen in table 6 (seen in the
15
16 statistical analysis section below), and the overall quality ratings of the papers resulting from the
17
18 application of the appraisal tool in figure 2 can be found in table 7.
19

20 (Table 7 here)
21
22
23

24 Synthesis

25
26
27 There is a lack of research which investigates delirium experienced by patients with advanced
28
29 cancer and that which exists is of variable quality. No randomised control trials of
30
31 interventions developed to treat delirium could be located. The review of the 7 articles
32
33 suitable for inclusion demonstrated that a range of pharmacological and non-pharmacological
34
35 interventions have been explored. It was not possible to combine and analyse the results
36
37 using statistics because the papers investigated the use of different interventions using
38
39 different methods. In view of this, the results are grouped on the basis of the main findings as
40
41 a basis for further discussion of their implications. The common main findings were
42
43 identified by reading through the included studies, identifying the types of interventions that
44
45 were used and considering the implications of these approaches. The interventions identified
46
47 were use of: atypical antipsychotics, opioid rotation, other pharmacological interventions and
48
49 non-pharmacological interventions. However before considering these further, a number of
50
51 general points about the studies are presented below.
52
53
54
55
56
57
58
59
60

[Type text]

Results

Design

All of the studies lacked the rigour of RCTs, as they were open-label, non-blinded and non-randomised (Sedgwick 2014), although most did report statistically significant results (apart from Moryl et al, 2005 which does not include any statistical analysis). All of the studies excepting Tatematsu et al (2011) were prospective in nature.

Sampling

The studies all used purposive sampling, a technique often used in qualitative research, where researchers select participants who can best address the purpose of the study (Aveyard, 2010). This type of sampling can introduce bias because of the absence of randomisation (Clifford, 1997). However in the studies reviewed here, the sample characteristics were so specific that it may have been difficult to recruit the patients in any other way. Excepting Breitbart et al (2002) sample sizes were less than 30 (the 2010 study by Tatematsu used 31 patients for the control group and 17 for the intervention group) which affects the statistical significance of the findings (Krithikadatta, 2014). Power calculations to determine the sample size required to test their hypothesis (Polit and Beck, 2014), were not carried out, (or were not reported on), in the seven studies.

Tools

A number of tools were used to collect data about the presence and severity of delirium. The MDAS was the commonest (Boettger and Breitbart 2011, Moryl et al 2005, Breitbart et al 2002). The Japanese version, which omits some the original items, was used by Morita et al (2005). The CAM, DOS and MMSE were used in other studies (Tatematsu et al 2011; Arai et al 2013; Gagnon et al 2005). The use of different tools means identification of delirium varied between the studies. Those using MDAS applied different thresholds for the definition

[Type text]

of delirium, for example Breitbart et al (2002), Boettger and Breitbart (2011) and Morita et al (2005) considered an MDAS < 10 to indicate delirium resolution, whereas a score of less than 13 is generally accepted as accurate (Alici and Breitbart, n.d).

Statistical Analysis

All of the papers include statistical analysis of the results (producing inferential statistics) apart from Moryl et al (2005), which includes descriptive statistics. The statistical test used depends on the type of data, whether it is parametric or non-parametric, and then the type of data generated; nominal, ordinal, interval or ratio. Parametric data follows a normal distribution (usually a sample of 30 or more is required for this) and non-parametric data does not follow normal distribution (Krithikadatta, 2014). A variety of statistical tests were used to analyse the data. Table 6 summarises the tests used and their appropriateness for the type of data they were analysing (Institute for Digital Research and Education, n.d.).

(Table 6 here)

Main Treatment Approaches

Atypical Antipsychotics

The use of atypical antipsychotics was explored in two of the papers, olanzapine (Breitbart et al 2002) and aripiprazole (Boettger and Breitbart 2011). Both report statistically significant results, with Boettger and Breitbart (2011) recording the mean MDAS at T1 was 18.0 which reduced to a mean MDAS at a stable dose of aripiprazole at T3 of 8.3 ($p < 0.001$). Breitbart et al (2002) found 76% of the patients had resolution of their delirium at T3 based on an MDAS of < 10, meaning the MDAS score improved from 19.85 (± 3.79) to 10.78 (± 7.31), $P = 0.001$. The statistical significance of these findings needs to be treated with caution because of the open-label non-randomised study designs. For example Breitbart et al (2002)

[Type text]

1
2
3 analysed confounding factors and found that treatment outcome was affected by designation
4 of the delirium subtype. This suggests that the management of subtypes could be different.
5
6

7 8 Opioid Rotation 9

10 Two studies explored opioid rotation in the management of delirium. The change to
11 Methadone for patients with morphine induced delirium (Moryl et al, 2005), and the
12 replacement of morphine with Fentanyl (Morita et al, 2005). Moryl et al (2005) found
13 'average' MDAS improved from 23.6 to 10.6, but it is not specified if this is referring to the
14 mean, mode, or median. When Fentanyl was introduced the mean MDAS decreased from 14
15 at T1 to 3.6 at T3 ($p < 0.001$) (Morita et al 2005). However, the tool used to collect this data
16 was the Japanese version of the MDAS, with some of the original items excluded, so again
17 the results need to be regarded with caution.
18
19
20
21
22
23
24
25
26
27
28
29
30
31

32 Other Pharmacological Interventions 33

34 Two studies reporting other pharmacological interventions were located. The first
35 investigated the use of a Celiac Plexus Block (CPB), and it was found that the duration of
36 Terminal Delirium (TD) was lower in the CPB group than the control group (1.8 ± 2.9 vs
37 10.4 ± 7.5 days respectively) (Arai et al, 2013). In the second Methylphenidate
38 hydrochloride (Gagnon et al 2005) was prescribed for patients with hypoactive delirium
39 resulting in an improvement from 20.9 ± 4.9 pre-treatment to 27.8 ± 2.4 on the MMSE when
40 the patients were on a stable dose of methylphenidate (Gagnon et al 2005). Its use for the
41 treatment of hypoactive delirium in the study limits its generalisability to other forms of
42 delirium.
43
44
45
46
47
48
49
50
51
52
53
54

55 Non-Pharmacological Intervention 56 57 58 59 60

[Type text]

The only report of a non-pharmacological intervention located, investigated the use of exercise therapy (Tatematsu et al, 2011). It found that antipsychotic dose was lower in the experimental group (2.198mg) than the control group (5.533mg) ($P < 0.036$), however the sample size was too small for the results to be generalizable. This study indicates there may be potential in non-pharmacological interventions as a wider range of staff would be able to deliver them.

Discussion

The findings of the research demonstrate that although there has been some investigation of the treatment of delirium in patients with advanced cancer, relatively few rigorous studies have been conducted. In terms of pharmacological interventions, the use of atypical antipsychotics reduced mean MDAS scores (Boettger and Breitbart, 2011; Breitbart et al, 2002), as did opioid rotation (Moryl et al, 2005 and Morita et al, 2005), a celiac plexus block reduced the duration of terminal delirium (Arai et al, 2013), and methylphenidate was found to be effective in patients with hypoactive delirium (Gagon et al, 2005). The one report of a non-pharmacological intervention found that the dose of antipsychotics could be reduced for patients participating in exercise therapy (Tatematsu et al, 2011).

The findings of this literature review are inconclusive in terms of identifying a single safe and effective treatment for delirium experienced by patients with advanced cancer. Haloperidol is often the first line treatment for delirium (NICE, 2010), however, a conference paper identified in the literature search reported that haloperidol alone was insufficient to control delirium in patients with advanced cancer because while 128 of the 167 patients required therapy with only haloperidol, 39 required a second neuroleptic to control symptoms (Susman, 2014).

Another first line treatment for delirium is olanzapine, which Breitbart et al (2002) investigated and even though the rigour of the study is not comparable with an RCT, the

[Type text]

1
2
3 results were still deemed to be significant. The use of aripiprazole has not been explored to a
4
5 great extent although small studies such as the one by Boettger and Breitbart (2011) indicate
6
7 it has helped resolve delirium. Further work is needed to determine if it can be added to the
8
9 drugs licenced to treat delirium.
10

11
12 The substitution of one opioid for another can improve therapeutic response and/or reduce
13
14 side effects (Moryl et al 2005; Morita et al 2005; Knotkova et al, 2009). Cancer patients
15
16 often experience pain and are prescribed opioids for pain relief even though this increases the
17
18 risk of delirium (Gaudreau et al, 2007; Morrison et al 2003). Further investigation of the
19
20 effects of opioid rotation is important for the treatment of delirium in patients with advanced
21
22 cancer. The final pharmacological intervention studied was the use of methylphenidate
23
24 hydrochloride in patients with hypoactive delirium (Gagnon et al, 2005). Methylphenidate is
25
26 licenced to treat attention-deficit hyperactivity disorder (ADHD) (BNF, 2014). It works by
27
28 increasing the amount of dopamine available in areas of the brain responsible for reward and
29
30 motivation (Volkow et al, 2012) and causes a 'calming' effect on patients with ADHD.
31
32 Because the pathophysiology of delirium is still not fully understood (Bush and Bruera 2009),
33
34 it is not clear how methylphenidate works to improve symptoms. This suggests further trials
35
36 of this agent would be of value.
37
38
39

40
41
42 The one non-pharmacological intervention studied was exercise therapy (Tatematsu et al,
43
44 2011), which was found to be a useful adjunct to pharmacological treatment in mental health
45
46 settings and is of relatively low cost (Daley, 2002). Studies which explore non-
47
48 pharmacological interventions are of particular interest to nurses, because if they are proven
49
50 to be effective, they can be readily incorporated into the nursing management of patients with
51
52 delirium. They are also useful because they can be investigated without the concomitant risk
53
54 of adverse effects such as side effects or serious complications, anaphylactic shock for
55
56 example, that can occur with pharmacological interventions (Cancer Research UK, 2015).
57
58
59
60

[Type text]

1
2
3 There remains limited information from randomised controlled trials of pharmacological
4 agents to guide practice in evidence-based neuroleptic administration to cancer and palliative
5 care patients (Bush and Bruera 2009). Further research is needed to determine efficacious
6 and safe drugs and dosages for the different delirium subtypes and aetiologies, as well as the
7 role of nonpharmacological and environment management strategies in improving improve
8 the comprehensive multifaceted management of this distressing syndrome (Bush and Bruera
9 2009).

10
11 With regard to the implications of the findings for clinical practice it is important to consider
12 options available to alleviate the distressing symptoms of delirium. Although treatments
13 unlicensed for use in delirium such as aripiprazole and olanzapine cannot be introduced,
14 interventions including exercise therapy could be considered. Also the findings suggest that
15 initiating discussions with doctors and pharmacists about the potential of opioid rotation is
16 worthy of further exploration. In addition practicing in accordance with current guidelines
17 for the care and treatment of delirium, including the provision of aides to orientation (for
18 example having clocks and calendars visible in clinical areas), should remain a priority of
19 care (NICE 2010).

Limitations

20
21 As with any literature review this one has limitations. For example, unpublished or 'grey'
22 literature was not included which has the potential to introduce publication bias, in that
23 journals tend to favour articles which report positive results, leading to over-representation of
24 significant findings (Cullinan, 2005; Polit and Beck, 2014). Also in view of the type of
25 interventions used in the treatment of delirium the nature of the studies reported varied, an
26 integrated approach was taken because a meta-analysis was not possible.

[Type text]

Conclusion

Delirium is a poorly understood condition which is frequently experienced by patients with advanced cancer, however there is limited knowledge of its pathophysiology, particularly of its subtypes. The findings from the seven studies examined in this review do not provide a conclusive evidence base for the treatment of delirium for patients with advanced cancer, although the interventions seemed to have some impact in terms of reducing the severity or the duration of the delirium.

However despite the lack of evidence on which to make definitive recommendations about treatment and care the current gap in knowledge has been identified, and there appear to be some promising areas for further research. More work is needed to build understanding of the pathophysiology of delirium in patients with advanced cancer and once this is more widely understood there can be more informed research into its treatment. Ideally this should take the form of Randomised Controlled Trials that can take account of confounding factors including age, co-treatments and cancer diagnosis (stage and tumour site) so that the effect of these factors can be analysed, so the true reason for delirium resolution can be identified.

[Type text]

References

Agency for Health Care Research and Quality (2011) Palliative care for the patient with incurable cancer or advanced disease. Part 2: pain and symptom management. Available at <https://www.guideline.gov/summaries/summary/38885/palliative-care-for-the-patient-with-incurable-cancer-or-advanced-disease-part-2-pain-and-symptom-management> Accessed 4th August 2016.

Akechi T., Uchitomi Y., Okamura H., Fukue M., Kagaya A., Nishida A., Oomori N., Yamawaki S. 1996. Usage of haloperidol for delirium in cancer patients. *Supportive Care in Cancer*. 4:390-392.

Akobeng A.K. (2005) Principles of evidence based medicine. *Archive of Diseases in Childhood* 90, 837-840.

Alici Y., Breitbart W., n.d. MDAS. Encyclopedia of psycho-pharmacology [online]. Available at: http://download.springer.com/static/pdf/325/prt%253A978-3-540-68706-1%252F13.pdf?originUrl=http%3A%2F%2Flink.springer.com%2Freferenceworkentry%2F10.1007%2F978-3-540-68706-1_1159&token2=exp=1470056650~acl=%2Fstatic%2Fpdf%2F325%2Fprt%25253A978-3-540-68706-1%25252F13.pdf%3ForiginUrl%3Dhttp%253A%252F%252Flink.springer.com%252Freferenceworkentry%252F10.1007%252F978-3-540-68706-1_1159*~hmac=85c0c3880267abb98ef947d7a78b3604765f4fcb86c8e8e06539f35c4c6190de [Accessed 01/08/16].

Alao A.O., and Moskowitz L., 2006. Aripiprazole and delirium. *Annals of Clinical Psychiatry* 18(4):267-9.

Alonso-Babarro, M. Varela-Cerdeira, I. Torres-Vigil, R. Rodriguez-Barrientos, E. Bruera, 2010. At-home palliative sedation for end-of-life cancer patients. *Palliative Medicine*, 24:5 pp. 486–492

American Cancer Society, 2014. What is advanced cancer? Available at: <http://www.cancer.org/treatment/understandingyourdiagnosis/advancedcancer/advanced-cancer-what-is> [Accessed 18/10/16].

American Cancer Society. (2015). *Global Cancer Facts & Figures* (Third Edition) Edition. American Cancer Society, Atlanta.

American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders: DSM-V* [online]. Available at: [http://reader.eblib.com/\(S\(xstn1pgsjtq1mnsvb4lplfce\)\)/Reader.aspx#](http://reader.eblib.com/(S(xstn1pgsjtq1mnsvb4lplfce))/Reader.aspx#) [Accessed 03/03/15].

Anthony J.C., LeResche L., Niaz U., Von Korff M.R., and Folstein M.F., (1982). Limits of the 'Mini-Mental State' as a screening test for dementia and delirium among hospital patients. *Psychological Medicine*, 12:2, pp 397-408.

[Type text]

1
2
3 Arai Y-C. P., Nishihara M., Kobayashi K., Kanazawa T., Hayashi N., Tohyama Y., Nishida
4 K., Arakawa M., Suzuki C., Kinoshita A., Kondo M., Masturbara S., Yokoe N., Hayashi R.,
5 Ohta A., Sato J., Ushida T., 2013. Neurolytic celiac plexus block reduces occurrence and
6 duration of terminal delirium in patients with pancreatic cancer. *Japanese Society of*
7 *Anaesthesiologists*. 27:1 p 88 – 92.

8
9 Averyard H., 2010. *Doing a Literature Review in Health and social care; a practical guide*.
10 3rd edition. Berkshire; Open University Press.

11
12 Bekelman J.E., Halpern S.D., Blankart C.R., Bynum J.P., Cohen J., Fowler R., Kaasa S.,
13 Kwietniewski L., Melberg H.O., Onwuteaka-Philipsen B., Oosterveld-Vlug M., Pring A.,
14 Schreyögg J., Ulrich C.M., Verne J., Wunsch H., Emanuel E.J. Comparison of Site of Death,
15 Health Care Utilization, and Hospital Expenditures for Patients Dying With Cancer in 7
16 Developed Countries. *JAMA*. 19;315(3):272-83.

17
18 Beller E.M., van Driel M.L., McGregor L., Truong S., Mitchell G., 2014. Palliative
19 pharmacological sedation for terminally ill adults. The Cochrane Library [online]. Available
20 at:
21 [http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD010206.pub2/abstract;jsessionid=18](http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD010206.pub2/abstract;jsessionid=18B5F99B1C059A767E0D25E110F473CB.f02t02)
22 [B5F99B1C059A767E0D25E110F473CB.f02t02](http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD010206.pub2/abstract;jsessionid=18B5F99B1C059A767E0D25E110F473CB.f02t02). Accessed 01/08/16.

23
24 Bettany-Saltikov J., 2012. How to do a systematic Literature Review in Nursing: A step-by-
25 step guide. New York; Open University Press.

26
27 BNF, 2014. *Methylphenidate hydrochloride* [online]. Available at:
28 [https://www.evidence.nhs.uk/formulary/bnf/current/4-central-nervous-system/44-cns-](https://www.evidence.nhs.uk/formulary/bnf/current/4-central-nervous-system/44-cns-stimulants-and-drugs-used-for-attention-deficit-hyperactivity-disorder/methylphenidate-hydrochloride)
29 [stimulants-and-drugs-used-for-attention-deficit-hyperactivity-disorder/methylphenidate-](https://www.evidence.nhs.uk/formulary/bnf/current/4-central-nervous-system/44-cns-stimulants-and-drugs-used-for-attention-deficit-hyperactivity-disorder/methylphenidate-hydrochloride)
30 [hydrochloride](https://www.evidence.nhs.uk/formulary/bnf/current/4-central-nervous-system/44-cns-stimulants-and-drugs-used-for-attention-deficit-hyperactivity-disorder/methylphenidate-hydrochloride) [Accessed 08/03/16].

31
32 Boettger S., Breitbart W., 2011. An open trial of aripiprazole for the treatment of delirium in
33 hospitalized cancer patients. *Palliative and Supportive Care*. 9:4 pp 351-357

34
35 Breitbart W., Tremblay A., Gibson C., 2002. An Open Trial of Olanzapine for the Treatment
36 of Delirium in Hospitalized Cancer Patients. *Psychosomatics*. 43:3 p 175–182.

37
38 Bush S.H., and Bruera E. (2009). The assessment and management of delirium in cancer
39 patients. *The Oncologist* 14 (10), 1039-1049.

40
41 Cancer Care Ontario (CCO)(2010) Cancer Care Ontario's Symptom Management Guide-to-
42 Practice: Delirium. Available at:
43 https://www.cancercare.on.ca/CCO_DrugFormulary/Pages/FileContent.aspx (accessed 4th
44 August 2016)

45
46 Cancer Research UK (2015) Advantages & drawbacks of taking part in a clinical trial
47 Available at: [http://www.cancerresearchuk.org/about-cancer/find-a-clinical-trial/what-you-should-be-](http://www.cancerresearchuk.org/about-cancer/find-a-clinical-trial/what-you-should-be-told-about-a-clinical-trial/advantages-and-drawbacks#C6cDTBRg3CYmJI7o.99)
48 [told-about-a-clinical-trial/advantages-and-drawbacks#C6cDTBRg3CYmJI7o.99](http://www.cancerresearchuk.org/about-cancer/find-a-clinical-trial/what-you-should-be-told-about-a-clinical-trial/advantages-and-drawbacks#C6cDTBRg3CYmJI7o.99) (accessed 4th August
49 2016)

50
51 Caraceni, A. and Simonetti, F. (2009) Palliating delirium in patients with cancer. *The Lancet*
52 *Oncology* 10, 164-72.

53
54 Centeno C., Bruera E. and Sanz, Á. (2004) Delirium in advanced cancer patients. *Palliative*
55 *Medicine* 18, 184-194.

[Type text]

1
2
3 Clifford C., 1997. Nursing and Health Care Research: A skills-based Introduction.
4 Hertfordshire; Prentice Hall Europe.

5
6 Close J.F, and Long C.O., Delirium: Opportunity for Comfort in Palliative Care. *Journal of*
7 *Hospice and Palliative Nursing.* 14:6 pp386-394

8
9 Cullinan P., 2005. Evidence-based healthcare: systematic reviews. In: Bowling A., and
10 Ebrahim S., eds 2005. *Handbook of Health Research Methods; Investigation, Measurement*
11 *and Analysis.* England; Open University Press.

12
13 Daley A.J., 2002. Exercise therapy and mental health in clinical populations: is exercise
14 therapy a worthwhile intervention? *Advances in Psychiatric Treatment.* 8:4 p 262-270

15
16 de Jong, J.D; Clarke L.E, 2009. What Is a Good Death? Stories from Palliative Care. *Journal*
17 *of Palliative Care.* 25:1 PP. 61-7.

18
19 de la Cruz M; Noguera A; San Miguel-Arregui MT; Williams J; Chisholm G; Bruera E,
20 2015. Delirium, agitation, and symptom distress within the final seven days of life among
21 cancer patients receiving hospice care. *Palliative & Supportive Care.* 13:2 p211-216.

22
23 Fang C-K., Chen H-W., Liu S.I., Lin C-J., Tsai L-Y., and Lai y-l. 2008. Prevalence,
24 Detection and Treatment of Delirium in Terminal Cancer Inpatients: A Prospective Survey.
25 *Japanese Journal of Clinical Oncology* 38(1)56–63.

26
27 Faight, D.D., 2014. Delirium: The Nurse's Role in Prevention, Diagnosis, and
28 Treatment. *Medsurg Nursing* 23:5 301-305.

29
30 Fong, T. G., Tulebaev, S. R., Inouye, S. K., 2009. Delirium in elderly adults: diagnosis,
31 prevention and treatment. *Nature Reviews. Neurology,* 5(4), 210–220.

32
33 Gagnon B., Graeme L., Schreier G., 2005. Methylphenidate hydrochloride improves
34 cognitive function in patients with advanced cancer and hypoactive delirium: a prospective
35 clinical study. *Journal of Psychiatry and Neurosciences.* 30:2 p 100 – 107.

36
37 Gaudreau J.D., Gagnon P., Roy M.A., Harel F., Tremblay A. 2007. Opioid medications and
38 longitudinal risk of delirium in hospitalized cancer patients. *Cancer.* 109(11):2365-73

39
40 Gaudreau J-D., Gagnon P., Harel F., Roy M-A., Tremblay A., 2005. Psychoactive
41 Medications and Risk of Delirium in Hospitalized Cancer Patients. *American Society of*
42 *Clinical Oncology.* 23:27 pp6712-6718.

43
44 Gopalakrishnan S. and Ganeshkumar P. (2013) Systematic Reviews and Meta-analysis:
45 Understanding the Best Evidence in Primary Healthcare. *Journal of Family Medicine and Primary*
46 *Care* 2 (1), 9-14.

47
48 Grant M.J., Booth A., 2009. A typology of reviews: an analysis of 14 review types and associated
49 methodologies. *Health Information and Libraries Journal.* 26: pp.91–108

50
51 Grassi L., Caraceni A., Mitchell A.J, Nanni M.G., Berardi M.A., Caruso R., Riba M., 2015
52 *Management of delirium in palliative care: a review.* *Current Psychiatry reports* 17 (13) pp 550.

[Type text]

1
2
3 Grover S., Sharma A., Aggarwal M., Mattoo S.K, Chakrabarti S., Malhotra S., Avasthi
4 A., Kulhara P., and Basu D., 2014. Comparison of symptoms of delirium across various
5 motoric subtypes. *Psychiatry and Clinical Neurosciences*. 68: 283–291

6
7 Harris D., 2007. *Delirium in advanced disease*. *Postgraduate Medical Journal*. 83(982): 525–
8 528.

9
10 Hek G., and Moule P., 2006. *Making sense of research; an introduction for health and social*
11 *care practitioners*. 3rd edition. London; SAGE publication ltd.

12
13
14 Horsley T., Dingwall O., Sampson M. 2011. Checking reference lists to find additional
15 studies for systematic reviews. *Cochrane Database of Systematic Reviews*. 10:8.

16
17 Hui D., Bush S.H., Gallo L.E., Palmer J.L., Yennurajalingam S., and Bruera E., 2010.
18 Neuroleptic dose in the management of delirium in advanced cancer patients. *Journal of pain*
19 *and symptom management*. 39:2 186 – 196.

20
21 Husebo B.S., Ballard C., Sandvik, R, Nilsen O.B, Aarsland D, 2011. Efficacy of treating pain
22 to reduce behavioural disturbances in residents of nursing homes with dementia: cluster
23 randomised clinical trial. *British Medical Journal*. 343.

24
25
26
27 Institute for Digital Research and Education, n.d. *What statistical Analysis Should I Use?*
28 [online]. Available at <http://www.ats.ucla.edu/stat/sas/whatstat/> [Accessed 02/02/16].

29
30 Kang J.H., Shin S.H., Bruera E. 2013. Comprehensive approaches to managing delirium in
31 patients with advanced cancer. *Cancer Treatment Reviews*. 39(1):105-12.

32
33 Kennedy JS, Jeste D, Kaiser CJ, Golshan S, Maguire GA, Tollefson G, Sanger T, Bymaster
34 FP, Kinon BJ, Dossenbach M, Gilmore JA, Breier A., 2003. Olanzapine vs haloperidol in
35 geriatric schizophrenia: analysis of data from a double-blind controlled trial. *International*
36 *Journal of Geriatric Psychiatry*. 18(11):1013-20.

37
38 Kim S-Y., Kim S-W., Kim J-M., Shin I-S., Bae K-Y., Shim, H-J., Bae, W-K., Cho S-H.,
39 Chung I-J., Yoon J-S., 2015. Differential Associations between Delirium and Mortality
40 According to Delirium Subtype and Age: A Prospective Cohort Study. *Psychosomatic*
41 *Medicine*. 77:8. p 903–910.

42
43
44 Knotkova H., Fine .PG., Portenoy R.K., 2009. Opioid rotation: the science and the limitations
45 of the equianalgesic dose table. *Journal of Pain and Symptom Management* 38(3): p426-39.

46
47
48 Krithikadatta J., 2014. Normal Distribution. *Journal of Conservative Dentistry*. 17(1): 96–97.

49
50 Ljubisavljevic V., Kelly B., 2003. Risk factors for development of delirium among oncology
51 patients. *General Hospital Psychiatry* 25:5 pp 345- 352.

52
53 Massie MJ, Holland J, Glass E., 1983. Delirium in terminally ill cancer patients. *American*
54 *Journal of Psychiatry* 140:8 1048–1050

55
56 Meagher D.J., 2001. Delirium: Optimising Management. *British Medical Journal*. 322:114

[Type text]

1
2
3 Meier D.E., 2012. Pain as a Cause of Agitated Delirium. *Journal of the American Medical*
4 *Association*. 172:15 p 1130.

5
6 Mhaskar R., Emmanuel P, Mishra S, Patel S., Naik E., Kumar A., 2009. Critical appraisal
7 skills are essential to informed decision-making. *Indian Journal of Sexually Transmitted*
8 *Diseases*. 30(2): 112–119.

9
10 Milisen, K., Foreman, M. D., Abraham, I. L., De Geest, S., Godderis, J., Vandermeulen, E.,
11 Fischler, B., Delooz, H. H., Spiessens, B. and Broos, P. L. O. (2001), A Nurse-Led
12 Interdisciplinary Intervention Program for Delirium in Elderly Hip-Fracture Patients. *Journal*
13 *of the American Geriatrics Society*, 49:5 p 523–532

14
15
16 Minagawa H., Uchitomi Y., Yamawaki S., Ishitani K., 1996. Psychiatric Morbidity in
17 Terminally 111 Cancer Patients: A Prospective Study. *Cancer*. 78: 5 p 1131 – 1137.

18
19 Moher, D., Liberati, A., Tetzlaff, J. and Altman, D.G. (2009) *Preferred Reporting for*
20 *systematic reviews and meta-analysis: The PRISMA Statement* [online].
21 <http://www.plosmedicine.org/article/info%3Adoi%2F10.1371%2Fjournal.pmed.1000097>
22 [Accessed 30/01/2016]

23
24
25 Morita T., Takigawa C., Onishi H., Tajima T., Tani K., Matsubara T., Miyoshi M, Ikenaga
26 M., Akechi T, and Uchitomi Y., 2005. Opioid Rotation from Morphine to Fentanyl in
27 Delirious Cancer Patients: An Open-Label Trial. *Journal of Pain and Symptom Management*.
28 30:1 p 96 - 103

29
30
31 Morrison R.S., Magaziner J., Gilbert M., Koval K.J., McLaughlin M.A., Orosz G., Strauss E.,
32 and Siu A.L., 2003. Relationship Between Pain and Opioid Analgesics on the Development
33 of Delirium Following Hip Fracture. *Journal of Gerontology*. 58:1, P 76–78

34
35 Moryl N., Kogan M., Comfort C., Obbens E., 2005. Methadone in the treatment pf pain and
36 terminal delirium in advanced cancer patients. *Palliative and Supportive Care*. 3:4 p 311-317.

37
38 Namba M, Morita T, Imura C, Kiyohara E, Ishikawa S, Hirai K, 2007. Terminal delirium:
39 families' experience. *Journal of Palliative Medicine*. 21(7):587-94

40
41 National Audit Office, 2008. *End of Life care* [online]. Available at
42 <https://www.nao.org.uk/wp-content/uploads/2008/11/07081043.pdf> [Accessed 04/03/16].

43
44
45 National Ethics Committee, Veterans Health Administration, 2007. The Ethics of Palliative
46 Sedation as a Therapy of Last Resort. *American Journal of Hospice & Palliative Medicine*.
47 23:6 pp 483-491.

48
49 NICE, 2010. *Delirium: prevention, diagnosis and management [CG103]* [online]. Available
50 at: <https://www.nice.org.uk/guidance/cg103> [Accessed 07/03/16].

51
52
53 NICE, 2015. Care of dying adults in the last days of life [NG31] [online]. Available at:
54 <https://www.nice.org.uk/guidance/ng31/chapter/Recommendations> [Accessed 25/07/16].

[Type text]

Peterson J.F, Pun B.T, Dittus R.S, Thomason J.W.W., Jackson J.C., Shintani A.K., Ely E.W., 2006. Delirium and Its Motoric Subtypes: A Study of 614 Critically Ill Patients. *The Journal of the American Geriatrics Society*. 54: 3 pp 479 – 484.

Polit D.F., and Beck C.T., 2014. *Essentials of Nursing Research: Appraising Evidence for Nursing Practice*. 8th edition. London: Wolters Kluwer Health/Lippincott Williams & Wilkins.

Ryan D.J., O'Regan N.A., Ó Caoimh R., Clare J., O'Connor M, Leonard M., McFarland J., Tighe S, O'Sullivan K, Trzepacz P.T., Meagher D., Timmons S., 2012. Delirium in an adult acute hospital population: predictors, prevalence and detection. *BMJ Open*. 3:1.

Ryan R; Cochrane Consumers and Communication Review Group. 'Cochrane Consumers and Communication Review Group: data synthesis and analysis'. <http://cccrg.cochrane.org>, June 2013 (accessed 02/03/2017).

Sedgwick P. (2014) What is an open label trial? *British Medical Journal* 348:g3434 doi: 10.1136/bmj.g3434.

Siddiqi N., House A.O., and Holmes J.D., 2006. Occurrence and outcome of delirium in medical in-patients: a systematic literature review. *Age and Ageing*. 35 (4):350-364.

Susman E., 2014. Haloperidol alone found sufficient to control delirium in advanced cancer patients. *Oncology times* 36:15 pp 20-20.

Tatematsu N., Hayashi A., Narita K., Tamaki A., Tsuboyama T., 2001. The effect of exercise therapy on delirium in cancer patients: a retrospective study. *Supportive Care in Cancer*. 19:6 p 765 – 770.

The Economist Intelligence Unit, 2010. *The quality of death: Ranking end-of-life care across the world* [online]. Available at <http://graphics.eiu.com/upload/eb/qualityofdeath.pdf> [accessed 04/03/15].

Volkow N. D., Wang, G.-J., Tomasi, D., Kollins, S.H., Wigal, T.L., Newcorn, J.H., Telang, F.W., Fowler, J.S., Logan, J., Wong, C.T., Swanson, J.M., 2012. Methylphenidate-Elicited Dopamine Increases in Ventral Striatum Are Associated with Long-Term Symptom Improvement in Adults with Attention Deficit Hyperactivity Disorder. *Journal of Neuroscience*. 32(3) 841-849.

Western Australia Cancer & Palliative Care Network (2010) Evidence based clinical guidelines for adults in the terminal phase, Palliative Care for People Living at Home Initiative. Government of Western Australia, Department of Health. Available at: http://www.healthnetworks.health.wa.gov.au/cancer/docs/Evidenced_based_guidelines_flipbook.pdf (accessed 4th August 2016)

Whittemore R., and Knafl K, 2005. Methodological issues in nursing research: The integrative review: Updated Methodology. *Journal of Advanced Nursing*. 52:5. 546

Table 1: Symptoms of delirium by subtype.

Delirium Subtype	Symptoms
Hyperactive	Restlessness Agitation Hallucinations Delusions
Hypoactive	Lethargy Minimal spontaneous movement Slow in responding to questions Thought process abnormality
Mixed	A combination of hyperactive and hypoactive symptoms.

For Peer Review

Table 2: Research Question Formulation, PIO

Population	Patients with advanced cancer
Interventions	Pharmacological and non-pharmacological treatments of delirium
Outcome	The management of delirium; whether the interventions improve, worsen or result in no change to delirium symptoms.

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3: Database Search Terms

Population	Population	Intervention	Outcome
Oncology	Supportive	Treatment	Deliri*
Cancer	“end of life”	Management	
Neoplasms	Palliative	Therapeutics	“acute confusion”


For Peer Review

Table 4: Inclusion and exclusion criteria and rationale

Inclusion Criteria	Rationale
Patients with advanced cancer must make up the full or partial sample group	The research question focuses on advanced cancer patients, studies on those with without cancer or cancer that is not advanced
Primary research	Less chance of bias than with secondary research.
Treatment of delirium must be the main focus of the article	The article must focus on treatment of delirium (treatment can be pharmacological or non-pharmacological) rather than prevention to address the research question
Exclusion Criteria	Rationale
Studies published before the year 2000	The studies are less likely to be relevant to todays practice
Secondary research	More chance of bias
Case Reports	The lack of generalisability was viewed as too great as the cases were often very complex.
Unpublished/grey literature	The researcher is novice and has limited time to search for data
Studies not published in English	The researcher is novice and only speaks fluent English, so does not have the time or resources to translate the articles.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1: PRISMA (adapted from Moher et al, 2009)



Paper	Design	Sample & Setting	Intervention	Delirium Screening tool(s)	Are tools reliable?	Analysis	Results	Conclusion
Boettger and Breitbart, 2011	Open Label Naturally Assigned prospective	N = 21 46.7% had advanced cancer Memorial Sloan-Kettering Cancer Centre – US	Aripiprazole	MDAS	Yes	Freidman	P < 0.001 mean MDAS T1 = 18.0, mean MDAS T3 = 8.3	Aripiprazole may be safe and effective in treating delirium.
Moryl et al, 2005	Open Label, prospective	N = 20 Advanced cancer patients. Tertiary cancer palliative care hospital -US	Methadone 'switch'	MDAS	Yes	None	'Average' MDAS. Improved from 23.6 to 10.6	A switch to methadone may be considered before sedation
Morita et al, 2005	Open Label Prospective	N = 21 Palliative Care patients, morphine induced delirium 7 palliative care units - Japan	Opioid Rotation; morphine to fentanyl.	MDAS (Japanese version). Omitted items	Original – yes Modified - no	Freidman	P < 0.001. Mean MDAS decreased from 14 at T1 to 3.6 at T3	Opioid Rotation; morphine to fentanyl may be effective in alleviating morphine induced delirium
Tatematsu et al, 2011	Open-label Retrospective	N = 48, EG = 17 NG = 31. Prognosis < 6 months' survival in EG = 35.5%; NG=41.9%. Kyoto University Hospital – Japan	Exercise therapy	CAM	Yes	Students t test	P < 0.036. Antipsychotic dose lower in EG (2.198mg) than NG (5.533mg).	Exercise therapy could be as an environmental intervention for delirium
Arai et al, 2013	Open-label Prospective CPB group,	N = 36 Control Group N = 17 CPB Group N = 19. Pancreatic cancer. TD. Palliative-care team - Japan	Neuroleptic Celiac Plexus Block	DOS	Yes	Unclear	P < 0.003. Duration of TD lower in CPB than control (1.8 ± 2.9 vs 10.4 ± 7.5 days respectively).	Duration of TD significantly less in patients who underwent CPB
Gagnon et al, 2005.	Open-label Prospective	N= 14 hypoactive delirium, advanced cancer. Montreal General Hospital – Canada.	Methylphenidate hydrochloride	MMSE	Yes	Wilcoxon signed rank test	P<0.001. Pre-treatment MMSE mean=20.9 ± 4.9, stable treatment MMSE mean=27.8±2.4	Hypoactive delirium with no known aetiology may be improved by methylphenidate
Breitbart et al, 2002	Open-Label, prospective	N = 79 Stage: terminal = 5%, Memorial Sloan-Kettering Cancer - US	Olanzapine	MDAS	Yes	ANOVA	MDAS score improved from a mean of 19.85 to 10.78, P = 0.001. Subtype can affect outcome. 76% delirium resolution.	Study begins to demonstrate the efficacy and safety of olanzapine for use in the management of delirium among hospitalized medically ill patients

Table 5: Summary of Papers

Abbreviations;

MDAS = Memorial Delirium Assessment Scale
 MMSE = Mini-Mental State Examination.
 DOS = Delirium Observation Screening Scale
 ANOVA = Analysis of variance

CAM = Confusion Assessment Method
 T1 = Time 1; baseline measurement
 T3 = Time 3; 4-7 days after intervention

± = standard deviation
 TD = Terminal Delirium
 N = number of participants

EG = Exercise Group
 NG = Non-exercise group
 CPB = celiac plexus block

Figure 2 - Critical Appraisal Tool

Abstract	Are key elements included? Are the Aims clearly stated?
Literature Review	Is the literature review up to date? Does the literature review establish a focus for the study due to a gap in knowledge?
Method: Research Design	Was the most rigorous possible design used? Was bias minimised? Were the researchers blinded? Were the participants randomised?
Population and sample	Was the population identified and described? Was the best possible sampling design used to enhance the samples representativeness? Was the sample size adequate? Was sample bias present? Was there a control group? Were the inclusion/exclusion criteria stated?
Data collection and Measurement	Are the tools used adequately described and was it properly implemented? Does the report provide evidence that the data collection methods yielded data high on reliability and validity? Was data collected in a manner that minimised bias?
Procedures	If there was an intervention was it clearly described and properly implemented? Did most participants allocated to the intervention group receive it?
Results: Data Analysis	Were appropriate statistical methods used? Was $p < 0.05$? Did the analysis control for confounding variables?
Findings/Conclusions	Was information about statistical significance presented? Are limitations identified and discussed? Are the results generalizable? Does the researcher discuss the results in context of the previous literature? Does the researcher suggest implications for practice and further research?

Adapted from

Polit D.F., and Beck C.T., 2014. Essentials of Nursing Research: Appraising Evidence for Nursing Practice 8th edition. London: Wolters Kluwer Health/Lippincott Williams & Wilkins

And

Hek G., and Moule P., 2006. Making sense of research; an introduction for health and social care practitioners. 3rd edition. London; SAGE publication ltd.

Table 6: Statistical tests used

Article	Data	Normal Distribution	Type of Data	Statistical Test Used	Was the test appropriate?
Moryl, 2005	Change in MDAS score	No	Interval	None Used	N/A
Morita, 2005	Change in MDAS score	No	Interval	Friedman	Yes
Tatematsu, 2010	Difference in antipsychotic dose	Unclear	Ratio	Students t test	Unclear
Arai et al, 2013	Duration of terminal delirium	No	Interval	Unclear	Unclear
Boettger and Breitbart, 2011	Change in MDAS score	No	Interval	Friedman	Yes
Gagnon, 2005	Change in MMSE score	No	Interval	Matched-paired Wilcoxon signed rank test	Yes
Breitbart, 2002.	Change in MDAS score	Yes	Interval	ANOVA	Yes

Table 7 - Results of the Quality Appraisal

Article	Design (n/5)	Sample/Sampling (n/5)	Tools (n/5)	Analysis (n/5)	Overall Quality (n/20)
Moryl et al, 2005	1	2	5	0	9
Morita et al, 2005	1	2	4	4	11
Tatematsu et al, 2011	2	2	2	2	8
Arai et al, 2013	2	2	4	2	10
Boettger and Breitbart, 2011	1	2	5	3	11
Gangon et al, 2005	1	2	4	4	11
Breitbart et al, 2002.	3	3	5	4	15

Key: 0 (very poor) 5 (excellent)

Or Peer Review