

# UNIVERSITY OF BIRMINGHAM

## Research at Birmingham

### Weight status, cardiorespiratory fitness and high blood pressure relationship among 5-12 years old Chinese primary school children

Pallan, Miranda; Griffin, Tania; Cheng, Kar; Li, Bai; Adab, Peymane

DOI:

[10.1038/jhh.2017.67](https://doi.org/10.1038/jhh.2017.67)

#### Document Version

Peer reviewed version

#### Citation for published version (Harvard):

Pallan, M, Griffin, T, Cheng, K, Li, B & Adab, P 2017, 'Weight status, cardiorespiratory fitness and high blood pressure relationship among 5-12 years old Chinese primary school children', *Journal of Human Hypertension*. <https://doi.org/10.1038/jhh.2017.67>

[Link to publication on Research at Birmingham portal](#)

#### Publisher Rights Statement:

Checked for eligibility: 04/08/2017

doi: 10.1038/jhh.2017.67

#### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

1 **Title:** Weight status, Cardiorespiratory fitness and high blood pressure relationship  
2 among 5-12 years old Chinese primary school children

3  
4

5 **Authors:**

6 WJ Liu<sup>1</sup>, LH Xiong<sup>1</sup>, CS Guo<sup>1</sup>, B Li<sup>2</sup>, M Pallan<sup>2\*</sup>, T Griffin<sup>2</sup>, KK Cheng<sup>2</sup>, and P Adab<sup>2</sup>

7 <sup>1</sup>Faculty of School Health, Guangzhou Centre for Disease Control and Prevention,  
8 China;<sup>2</sup>Public Health, Epidemiology and Biostatistics, Institute of Applied Health  
9 Research, College of Medical and Dental Sciences, University of Birmingham UK

10  
11

12 **Running title:** Weight status, fitness and blood pressure in children

13  
14  
15  
16

17 **\*Corresponding author:**

18 Dr Miranda Pallan, Senior Clinical Lecturer in Public Health, Institute of Applied  
19 Health Research, The University of Birmingham, Edgbaston, Birmingham, B15 2TT,  
20 UK.

21 Tel: +441214147990  
22 e-mail: M.J.Pallan@bham.ac.uk

23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71

## Summary Table

### What is known about this topic

- Cardiorespiratory fitness (CRF) and adiposity contribute to high blood pressure (HBP), but their relative importance is unknown.
- Data analyzed these relationships among Chinese children is lacking.

### What this study adds

- Hypertension is common in Chinese children (prevalence 15.3%), with higher prevalence in obese (40.5% and 45.9% in boys and girls respectively) and overweight (27.6% and 30.2%).
- Weight status is strongly associated with likelihood of high blood pressure in Chinese primary school aged children.
- There was no evidence that increased CRF modified the risk of HBP in overweight and obese children.

72 **Abstract**

73 Cardiorespiratory fitness (CRF) and adiposity contribute to high blood pressure (HBP)  
74 in adults and children. However, their relative importance as risk factors is unknown.  
75 We examined the relationships between weight status, CRF and HBP among Chinese  
76 primary school children. A cross-sectional study was conducted with 4926 school  
77 children aged 5–12 years. CRF was estimated from a modified Cooper test, body  
78 mass index z-scores and weight categories were calculated from objective height and  
79 weight measurements, and BP was measured using an electronic  
80 sphygmomanometer. HBP was defined as >95th percentile based on reference  
81 cut-offs for Chinese boys and girls. Generalized Linear Mixed Models, adjusting for  
82 age, pubertal status and height, were developed for boys and girls to explore the  
83 independent and combined associations between fitness, weight status and HBP. 752  
84 (15.3%) children had HBP, with a higher prevalence in obese (40.5% and 45.9% in  
85 boys and girls respectively) and overweight (27.6% and 30.2%) compared with  
86 non-overweight (9.0% and 13.8%) children. HBP prevalence was lower in boys with  
87 higher CRF (OR for highest vs. lowest CRF quartile in boys 0.64; 95%CI 0.46-0.89).  
88 This association was not seen in girls. With weight status and CRF in the same model,  
89 weight status, but not CRF, remained significantly associated with HBP (obesity in  
90 boys: OR 4.19; 95%CI 2.63-6.67; in girls: OR 2.49; 95%CI 1.19-5.19). The interaction  
91 effect for CRF and weight status was non-significant. Overweight/obesity was  
92 significantly associated with HBP among children. There was no evidence of  
93 modification of this relationship by CRF.

94

95 **Keywords:** Blood pressure; Cardiorespiratory fitness; Obesity, School children, China  
96

## 97 **Introduction**

98 High blood pressure (HBP) is increasing in prevalence among children and youth, and  
99 there is evidence that HBP is present in Chinese children as young as 5 years old<sup>1</sup>.  
100 Serial cross-sectional studies in China indicate that the prevalence in adults has  
101 increased by 83.4% since 1991, affecting over a quarter of the population in  
102 2007-2008<sup>2-5</sup>. There is increasing evidence of tracking of blood pressure (BP) levels  
103 from childhood to adult life and hypertension in childhood is associated with higher  
104 risk of cardiovascular disease and mortality in adulthood<sup>6-10</sup>. Two of the major  
105 modifiable risk factors for hypertension include obesity and low cardiorespiratory  
106 fitness (CRF)<sup>11-14</sup>. However, the exact pathophysiological mechanisms through which  
107 obesity and low CRF contribute to hypertension are not fully understood. Obesity is  
108 likely to act through inter-related complex mechanisms related to diet, and  
109 activation of the sympathetic nervous system causing vascular and renal injury<sup>15</sup>.  
110 Similarly there are multiple hypotheses to explain the mechanisms by which  
111 increased CRF alters blood pressure. Animal studies suggest that exercise training,  
112 which increases CRF, alters vasomotor tone through endothelial and smooth muscle  
113 adaptations<sup>16</sup>. In addition, a study in young adults has suggested that the  
114 relationship between CRF and arterial stiffness is mediated by resting heart rate<sup>17</sup>.  
115 There is some debate about the relative importance of CRF versus weight status as  
116 risk factors for HBP and whether they act independently. There is a wealth of  
117 evidence to suggest that low CRF is a potentially more important risk factor for

118 premature all-cause mortality and cardiovascular mortality than obesity, and that  
119 amongst individuals with obesity, physical activity and CRF can significantly positively  
120 impact on obesity and the subsequent risk of cardiovascular disease<sup>18-23</sup>. Trials that  
121 compare the effects of diet only versus diet and physical activity interventions in  
122 obese adolescents, suggest that whilst both interventions result in reduction of BP,  
123 the addition of exercise results in normalization of forearm vascular conductance  
124 (similar to that seen in non-overweight children), whereas this is not seen in the diet  
125 only group<sup>24</sup>. Thus CRF may have effects on vascular health independently of weight  
126 status, although the exact mechanisms are not fully understood. Other studies  
127 suggest that higher levels of physical activity or CRF do not compensate for obesity in  
128 relation to cardiovascular risk<sup>25-26</sup>. Therefore, there is a need to further understand  
129 the relative importance of these two risk factors and their relationship with one  
130 another. A better understanding of the relationship between weight status, CRF and  
131 HBP in this age group would help to shape recommendations for interventions to  
132 improve the health of children and adolescents.

133 The prevalence of overweight and obesity in school-aged children in China (aged  
134 5-12 years) has increased more rapidly than that seen in most Western countries<sup>27-28</sup>,  
135 with the prevalence in some urban regions approaching those in developed  
136 countries<sup>29-30</sup>. In addition a report on the Physical Fitness and Health Surveillance of  
137 Chinese School Students in 2010 revealed that CRF in children has been decreasing  
138 during the past twenty five years<sup>31</sup>. Given the significant increases in these two  
139 major cardiovascular risk factors in Chinese children, we aimed to explore the

140 prevalence of HBP and its relationship with these two risk factors, and also to  
141 examine the potential modification effect of CRF on the association between weight  
142 status and HBP. We hypothesized that both obesity and CRF would be associated  
143 with HBP in this population of children, and that CRF would attenuate the  
144 association between obesity and HBP.

145

## 146 **Methods**

147 We included a sub-group of participants who had the relevant measurements from a  
148 larger cross-sectional study that aimed to determine the prevalence and risk factors  
149 for childhood obesity in primary school-aged children in Guangzhou<sup>32</sup>. A multi-stage  
150 stratified cluster random sampling procedure was used to obtain a representative  
151 sample of primary school children in grades 1-5 (aged 5-12 years). Five of the ten  
152 urban districts were selected using a random number generator. Within each of  
153 these, all primary schools were stratified by public (residents) or private (migrant)  
154 status, and six were randomly chosen with a 2:1 ratio from each stratum. Permission  
155 for the study was not obtained for one of the private schools, leaving 29  
156 participating schools. Within each school two classes per grade were randomly  
157 selected. The exclusion criteria included: children with invalid anthropometric  
158 measurements or questionnaire information and no parental consent.

159 Written informed consent was sought from the parents of 11445 eligible children  
160 through schools, resulting in 9917 (86.6%) participants (1403 children with no  
161 parental consent or invalid questionnaire; 125 children with invalid anthropometric

162 measurements). Data collection took place from April to June 2014, with  
163 anthropometric, CRF (reduced Cooper test) and blood pressure measurements  
164 undertaken in school, by trained research staff using standardized procedures and  
165 instruments (see details below). All consented children had measurements of their  
166 height and weight, but more detailed measurements, including blood pressure and  
167 the reduced Cooper test, were undertaken on a subsample (children from one of the  
168 two classes per grade were randomly selected; n=4926 including 2725 boys and  
169 2201 girls).

170 The study was approved by the Ethical Committee of the Guangzhou Center for  
171 Disease Control and Prevention and the University of Birmingham Research Ethics  
172 Committee. Permissions to conduct the study were granted by the Departments of  
173 Education and Health.

#### 174 **Anthropometric and BP measurements**

175 Height and weight were measured with subjects wearing light clothing and without  
176 shoes. Weight was measured to the nearest 0.1 kg using an electronic scale  
177 (JH-1993T, weighing Apparatus Co. Ltd. Dalian). Height was recorded to the nearest  
178 0.1 cm with a TGZ height tester (Dalian) according to the following protocol: no shoes,  
179 heels together, and student's heels, buttocks, shoulders, and head touching the  
180 vertical surface with line of sight aligned horizontally.

181 Blood pressure was measured by the same trained nurse using an electronic  
182 sphygmomanometer (Omron HEM-7211, Dalian) at the right arm with students in  
183 the seated position after at least 5 minutes of rest in a quiet classroom. The cuff size



184 was based on the length and circumference of the upper arm and was chosen to be  
185 as large as possible without having the elbow skin crease obstruct the stethoscope.  
186 Two consecutive readings were taken on the same arm with a two-minute interval  
187 between each reading; the mean of the 2 measures was used for analysis. Systolic  
188 blood pressure (SBP) was defined by the first Korotkoff sound (appearance of sounds),  
189 and diastolic blood pressure (DBP) was defined by the fourth Korotkoff sound (sound  
190 muffling).

#### 191 **CRF measurements**

192 CRF was assessed using a reduced Cooper test<sup>33</sup>. The full 12-minute Cooper test has  
193 been shown to be a very good predictor of maximum oxygen intake<sup>34</sup>, but is  
194 inappropriate to administer to young children within a school setting<sup>35</sup>. We therefore  
195 opted for the reduced Cooper test (6-minute) which has been successfully used in  
196 primary school aged children and can be easily administered within school settings<sup>33</sup>.  
197 Children were asked to run counter clockwise along a track of fixed size (marked  
198 rectangle measuring 9×18m) as many times as they could within 6 minutes. The  
199 exercise was undertaken outdoors on level ground. Outdoor climate varies little  
200 from day to day in Guangzhou, where it is generally warm and humid. We  
201 incentivized children to make a maximal effort by explaining that the test results  
202 would be included in their school report. However, if they lacked the physical  
203 strength to run at any point, they were allowed to walk. The distance covered  
204 (measured in meters) was timed by a trained physical education teacher within  
205 school, using a stopwatch (CASIO, HS-70W stopwatch) to the nearest 0.1s. The

206 physical education teacher recorded the number of complete laps done by each child,  
207 and estimated the distance for any incomplete laps at the end of the reduced Cooper  
208 test.

### 209 **Other measurements**

210 All children over the age of 9 years were assessed for whether or not they were  
211 pubertal by self-report, by asking girls if they had reached menarche and boys if they  
212 had experienced a first nocturnal emission. These questions were asked by a trained  
213 physician when the physical measurements were being undertaken.

### 214 **Statistical Analysis**

215 Body mass index (BMI; [weight (kg)]/[height (m)]<sup>2</sup>), was calculated and standard  
216 deviation scores (BMI z-score) derived using the age and sex specific WHO growth  
217 reference for school-aged children<sup>36</sup>. BMI z-scores were used to classify participants  
218 as non-overweight ( $\leq 1SD$ ), overweight ( $> 1SD, \leq 2SD$ ) or obese ( $> 2SD$ ). HBP was  
219 defined as systolic or diastolic BP above the 95th percentile for age and gender  
220 specific reference cut-offs for Chinese children and adolescents<sup>37</sup>. As we were using a  
221 reduced Cooper test, we were unable to calculate the  $VO_2$  max from the distance run  
222 using reference tables, therefore the distance covered was categorized into quartiles  
223 based on the child's age and sex, using the study data as the reference, with further  
224 categorization into higher (3rd and 4th quartiles) or lower (1st and 2nd quartiles)  
225 CRF.

226 Summary statistics (mean  $\pm$ SD and percentages) were used to describe participant  
227 characteristics, prevalence of hypertension and the proportion of participants in the

228 different CRF categories. Differences in characteristics between the sexes or weight  
229 status groups were determined using t-tests, analysis of covariance and Chi-square  
230 tests, where appropriate. Generalized linear mixed models, with school as a random  
231 effect to account for clustering, were used to examine the relationships between HBP  
232 as a dependent variable and CRF or weight status as independent variables by sex,  
233 both adjusted for age, pubertal status, and height (adjusted model I)<sup>38</sup>. A further  
234 model included both weight status, CRF and weight status  $\times$  CRF as covariates  
235 (adjusted model II) by sex. We also examined differences in prevalence of HBP, mean  
236 SBP and mean DBP in those with high CRF compared to those with low CRF in each of  
237 the weight status subgroups to further explore the potential modification effect of  
238 CRF on the relationship between weight status and HBP. Finally, we carried out  
239 sensitivity analyses using the criteria introduced by international obesity taskforce  
240 (IOTF). Data were analyzed using SPSS 21.0 statistical software package (SPSS Inc.,  
241 Chicago, IL). A 2-tailed P value less than 0.05 was considered statistically significant.

242

## 243 **Results**

### 244 **Characteristics of the Study Sample**

245 Among 4926 children in the sub-group who had BP measurements, 4,726 (2,725  
246 boys and 2,201 girls) had complete data for the reduced Cooper test, BP, and weight  
247 and height, and so were included in the analysis. Descriptive characteristics of the  
248 sample are shown in Table 1. Overall 10.9% of children were overweight and an  
249 additional 6.9% were obese, with rates being higher in boys compared with girls.

250 Almost all children (100% of boys and 96.5% of girls) were pre-pubertal. Boys  
251 covered a greater distance in the reduced Cooper test (899.4 vs 864.8 m) and had a  
252 higher SBP compared with girls (107.6 vs 106.1 mmHg). SBP and DBP increased with  
253 increasing age in boys and girls. Around 15% of children had HBP, with prevalence  
254 rates being similar in boys and girls and no significant difference in prevalence by  
255 age.

#### 256 **Relationship between weight status and CRF among school children**

257 There was a significant association between CRF and weight status (Table 2), with a  
258 higher proportion of non-overweight children being in the highest quartile for CRF  
259 (28.5% boys and 25.8% girls) compared with those who were obese (6.1% boys and  
260 6.6% girls). Significant differences were clearly seen in the proportions of children in  
261 the high and low CRF categories across weight status groups, with 54.7% of  
262 non-overweight boys, 42.8% of overweight boys and 20.1% of obese boys in the high  
263 CRF group. The corresponding percentages of girls in the high CRF group were 51.5%,  
264 39.2% and 19.7% for the non-overweight, overweight and obese categories,  
265 respectively.

#### 266 **Association between weight status, CRF and HBP among school children**

267 There was a clear relationship between weight status and HBP, with prevalence of  
268 9.0% (boys) and 13.8% (girls) in non-overweight children, 27.6% (boys) and 30.2%  
269 (girls) in overweight children, and 40.5% (boys) and 45.9% (girls) in obese children.  
270 Similarly there was a relationship between lower CRF and higher risk of hypertension  
271 in boys (prevalence of HBP 18.2% in the lowest versus 11.8% in the highest fitness  
272 quartile), although no clear relationship between fitness levels and BP was seen in

273 girls (Table 3).

274 In the adjusted models (model I; adjusted for age, pubertal status, height), the  
275 likelihood of having HBP was significantly higher for boys and girls who were  
276 overweight (OR 3.51; 95%CI 2.62-4.70 and 1.93; 1.34-2.78 respectively) or obese (OR  
277 5.55; 4.07-7.57 and 4.11; 2.37-7.13 respectively) compared with those who were  
278 non-overweight. There was also a statistically significant trend for reduced risk of  
279 HBP with increasing quartile of CRF in boys (OR for 4th vs 1th quartile CRF 0.64;  
280 95%CI 0.46-0.89,  $p < 0.05$ ), but not in girls (OR for 4th vs 1th quartile CRF 0.70; 95%CI  
281 0.43-1.13,  $p > 0.05$ ). In the combined weight status and CRF model, simultaneously  
282 adjusted for age, pubertal status, height and weight status  $\times$  CRF (adjusted model II),  
283 weight status but not CRF remained significantly associated with HBP in both boys  
284 and girls. The likelihood of having HBP was significantly higher for boys and girls who  
285 were overweight (OR 2.96; 95%CI 1.71-5.11 and 1.75; 1.00-3.06 respectively;  $p < 0.05$ )  
286 or obese (OR 4.19; 2.63-6.67 and 2.49; 1.19-5.19 respectively;  $p < 0.05$ ) compared  
287 with those who were non-overweight (Table 3). The interaction term for weight  
288 status and CRF was non-significant in the models.

289 Table 4 shows the mean SBP, DBP and prevalence of hypertension among those with  
290 high or low CRF within each weight status category (non-overweight, overweight,  
291 obese). All measures of BP were similar within the weight status categories  
292 irrespective of CRF level, and BP and prevalence of HBP were greater with increasing  
293 weight status in both boys and girls.

#### 294 **Sensitivity Analyses**

295 Repeating the analyses for factors associated with HBP, using the IOTF categorization  
296 of weight status, did not alter the findings. The magnitude and direction of effect of  
297 all variables reported above remained similar to the main analysis, although the  
298 absolute values differed (likelihood of HBP for those in obese compared with  
299 non-overweight category in model II was 5.06 (95%CI 2.94 to 8.73,  $p<0.05$ ) in boys  
300 and 2.34 (95%CI 1.38 to 4.09,  $p<0.05$ ) in girls, whilst likelihood of HBP for those in  
301 highest vs lowest quartile for reduced Cooper test was 0.89 (95%CI 0.59 to 1.33,  
302  $p>0.05$ ) and 0.75 (95%CI 0.46 to 1.22,  $p>0.05$ ) respectively).

303

#### 304 **Discussion**

305 We found that weight status is strongly associated with the likelihood of HBP in  
306 primary school aged children, and that in boys, but not in girls, the level of CRF was  
307 also inversely associated with HBP. However, when both CRF and weight status were  
308 included in the same model, only the association between weight status and HBP  
309 remained significant. In contrast to our hypothesis, there was no evidence in our  
310 analyses that higher CRF attenuated the association between obesity and HBP. Our  
311 finding of a strong association between weight status and blood pressure in children  
312 is in line with reports from other studies and systematic reviews, and suggests that  
313 tackling obesity in childhood may help reduce the burden of cardiovascular disease  
314 in adulthood<sup>39</sup>.

315 Previous studies have suggested that higher CRF could attenuate the effects of  
316 obesity on cardiovascular health and there is evidence from longitudinal studies to

317 suggest that higher CRF is associated with lower cardiovascular mortality among  
318 those who are obese<sup>21,40</sup>. A recent large longitudinal study of adolescents in the US  
319 reported a significant interaction between CRF and weight status in predicting risk of  
320 hypertension, and an association between lower CRF and HBP, even among those  
321 who were not overweight<sup>41</sup>. Among the Chinese children in our study, we did not  
322 find any evidence that increasing fitness levels were associated with lower BP once  
323 weight status was taken into account. Our findings are similar to those from another  
324 study in 9-12 year old children in the USA<sup>42</sup>. These differences may be due to the  
325 cross-sectional nature of these two studies, with the effects of CRF on blood  
326 pressure becoming apparent in the longer term. There is some limited evidence to  
327 suggest that CRF in childhood predicts later increases in blood pressure<sup>43-44</sup>. It may  
328 also be related to the younger age of the children in these studies, suggesting that  
329 weight status is the predominant predictor of HBP in childhood<sup>45</sup>, with CRF becoming  
330 more important in later life. The younger age of participants in this study sample  
331 may also potentially explain the observed difference between boys and girls in  
332 relation to the association between CRF and HBP. The majority of children in this  
333 study were pre-pubertal, and there is some evidence that pre-pubertal girls have  
334 stiffer large arteries compared with boys<sup>46</sup>. This could make them less responsive to  
335 blood pressure changes irrespective of CRF levels. However, even without the  
336 moderating effect of CRF on obesity and the risk of cardiovascular disease, CRF is an  
337 important factor in increasing longevity<sup>18-21</sup>, and may positively impact on the  
338 prognosis of hypertension. Therefore improving CRF should be prioritized as an

339 intervention target alongside obesity. Another important consideration is the role of  
340 physical activity, as this contributes to both CRF<sup>47</sup> and reduction in obesity,  
341 (particularly vigorous intensity physical activity<sup>48-49</sup>. Physical activity has also been  
342 shown to be important in protecting against all cause mortality in epidemiological  
343 studies<sup>50</sup>.

344 Strengths of our study include the large representative sample of children, the  
345 inclusion of objective standardized anthropometric, blood pressure and CRF  
346 measurements, and adjustment for school level clustering, which accounts for  
347 potential confounding from important socioeconomic factors.

348 Limitations include the use of only one measure of physical fitness and the lack of a  
349 validated method of estimating VO2 max from the reduced Cooper test. We also did  
350 not have information on family history of HBP or on salt and dietary intake, which  
351 could be important confounding factors and ideally should be adjusted for within the  
352 analyses. Finally, the cross-sectional nature of the study limits interpretation of  
353 causal associations.

354 In conclusion, our results demonstrated that overweight/obesity is strongly  
355 associated with HBP in both boys and girls. This supports recent evidence in the US  
356 where the increasing prevalence of HBP is attributed to the rise in overweight and  
357 obesity among young children and youth<sup>51-54</sup>. Given the rising prevalence of  
358 childhood obesity in China, it is imperative that comprehensive strategies are put  
359 into place to tackle obesity in order to reduce the future burden of cardiometabolic  
360 disease.



361

362 **Conflict of Interest**

363 The authors declare that they have no competing interests.

364 **Acknowledgements**

365 The data used for this analysis came from a study funded by the Guangzhou Medical  
366 Foundation(20131A031001).The authors gratefully acknowledge the contribution of  
367 Yuexiu, Haizhu, Liwan, Luogang and Nansha Education Bureau and 29 elementary  
368 schools. None of the authors had any personal or financial conflicts of interest.

369

370

371 **References**

372 1) Zhou Y, Qian Z, Vaughn MG, Boutwell BB, Yang M, Zeng XW, et al. Epidemiology of elevated blood  
373 pressure and associated risk factors in Chinese children: the SNEC study. J Hum Hypertens  
374 2016;30(4):231-236.

375 2) Ostchega Y, Carroll M, Prineas RJ, McDowell MA, Louis T, Tilert T. Trends of elevated blood pressure  
376 among children and adolescents: data from the National Health and Nutrition Examination  
377 Survey1988-2006. Am J Hypertens 2009;22:59-67.

378 3) Juonala M, Magnussen CG, Berenson GS, Venn A, Burns TL, Sabin MA, et al. Childhood adiposity,  
379 adult adiposity, and cardiovascular risk factors. N Engl J Med 2011;365(20):1876-1885.

380 4) Zhao D, Wu ZS, Wang W, Liu J, Wu GX, Zeng ZC, et al. Association between blood pressure level and  
381 risk of cardiovascular diseases in Chinese: a cohort study in 11 provinces of China. Chin J Cardiol  
382 2001;29(10):612-617.

383 5) Gao Y, Chen G, Tian HM, Wang J,Sun Y, Ding R,et al. Prevalence of hypertension in China:

384 across-sectional study. Plos one 2013;8(6):e65938.

385 6) Goon D, Amusa L, Mhlongo D, Khoza L, Any-Anwu F. Elevated blood pressure among Rural South  
386 African Children in Thohoyandou, South Africa. Iran J Pub Health 2013;42(5):489-96.

387 7) Jafar TH, Islam M, Poulter N, Hatcher J, Schmid CH, Levey AS, et al. Children in South Asia have  
388 higher body mass-adjusted blood pressure levels than white children in the United States: a  
389 comparative study. Circulation 2005;111(10):1291-7.

390 8) Du SF, Lv B, Wang ZH, Zhai FY, Barry MP. Transition of dietary pattern in China. J Hygiene Res 2001;  
391 30:221-225.

392 9) Franks PW, Hanson RL, Knowler WC, Sievers ML, Bennett PH, Looker HC. Childhood obesity, other  
393 cardiovascular risk factors, and premature death. N Engl J Med 2010;362:485-493.

394 10) Mahoney LT, Burns TL, Stanford W, Thompson BH, Witt JD, Rost CA, et al. Coronary risk factors  
395 measured in childhood and young adult life are associated with coronary artery calcification in young  
396 adults: the Muscatine Study. J Am Coll Cardiol 1996;27:277-284.

397 11) Carnethon MR, Gulati M, Greenland P. Prevalence and cardiovascular disease correlates of low  
398 cardiorespiratory fitness in adolescents and adults. JAMA 2005;294:2981-8.

399 12) Diaz KM, Shimbo D. Physical activity and the prevention of hypertension. Curr Hypertens Rep  
400 2013;15(6):659-668.

401 13) Kelly RK, Magnussen CG, Sabin MA, Cheung M, Juonala M. Development of hypertension in  
402 overweight adolescents: a review. Adolesc Health Med Ther 2015;6:171-87.

403 14) U.S. Department of Health and Human Services. Physical activity and health: a report of the  
404 Surgeon General. U.S. Department of Health and Human Services, 1997. Atlanta: Centers for Disease  
405 Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion.

406 15) Hall JE, da Silva AA, do Carmo JM, Dubinion J, Hamza S, Munusamy S, et al. Obesity-induced  
407 Hypertension: Role of Sympathetic Nervous System, Leptin, and Melanocortins. *J Biol Chem*  
408 2010;285(23):17271-6.

409 16) Delp MD, McAllister RM, Laughlin MH. Exercise training alters endothelium-dependent  
410 vasoreactivity of rat abdominal aorta. *J Appl Physiol*(1985) 1993;75(3):1354-63.

411 17) Quan HL, Blizzard CL, Sharman JE, Magnussen CG, Dwyer T, Raitakari O, et al. Resting heart rate  
412 and the association of physical fitness with carotid artery stiffness. *Am J Hypertens* 2014; 27(1):65-71.

413 18) Myers J, McAuley P, Lavie CJ, Despres JP, Arena R, Kokkinos P. Physical Activity and  
414 Cardiorespiratory Fitness as Major Markers of Cardiovascular Risk: Their Independent and Interwoven  
415 Importance to Health Status. *Progress in Cardiovascular Diseases. Prog Cardiovasc Dis*  
416 2015;57(4):306-14.

417 19)DeFina LF, Haskell WL, Willis BL, Barlow CE, Finley CE, Levine BD, et al. Physical activity versus  
418 cardiorespiratory fitness: two (partly) distinct components of cardiovascular health? *Prog Cardiovasc*  
419 *Dis* 2015;57(4):324-9.

420 20)Lavie CJ, De Schutter A, Parto P, Jahangir E, Kokkinos P, Ortega FB, et al. Obesity and Prevalence of  
421 Cardiovascular Diseases and Prognosis-The Obesity Paradox Updated. *Prog Cardiovasc*  
422 *Dis* 2016;58(5):537-47.

423 21) Barry VB, Baruth M, Beets MW, Durstine JL, Liu JH, Blair SN. Fitness vs. fatness on all-cause  
424 mortality: a meta-analysis. *Progress in Cardiovascular Disease* 2014; 56(4):382-390.

425 22) Fogelholm M. Physical activity, fitness and fatness: relations to mortality, morbidity and disease  
426 risk factors. A systematic review. *Obesity Reviews* 2010;11(3):202–221.

427 23) Lee DC, SuiX, Blair SN. Does physical activity a meliorate health hazards of obesity? *Br J Sports*

428 Med 2009;43:49-51.

429 24) Ribeiro MM, Silva AG, Santos NS, Guazzelle I, Matos LN, Trombetta IC, et al. Diet and exercise  
430 training restore blood pressure and vasodilatory responses during physiological maneuvers in obese  
431 children. *Circulation* 2005; 111(15):1915-23.

432 25) Hu G, Barengo NC, Tuomilehto J, Lakka TA, Nissinen A, Jousilahti P. Relationship of physical activity  
433 and body mass index to the risk of hypertension: a prospective study in Finland. *Hypertension*  
434 2004;43(1):25-30

435 26) Chen J, Das S, Barlow CE, Grundy S, Lakoski SG. Fitness, fatness, and systolic blood pressure: Data  
436 from the Cooper Center Longitudinal Study. *Am Heart J* 2010, 160 (1), pp. 166-170.

437 27) Ji CY, Chen TJ. Empirical changes in the prevalence of overweight and obesity among Chinese  
438 students from 1985 to 2010 and corresponding preventive strategies. *Biomed Environ Sci* 2013; 26(1):  
439 1-12.

440 28) Wang Y, Lobstein T. Worldwide trends in childhood overweight and obesity. *Int J Pediatr Obes*  
441 2006; 1(1): 11-25.

442 29) Ji CY, Cheng TO. Prevalence and geographic distribution of childhood obesity in China in 2005. *Int J*  
443 *Cardiol* 2008; 131(1): 1-8.

444 30) Ji CY. Report on childhood obesity in China (4) prevalence and trends of overweight and obesity in  
445 Chinese urban school-age children and adolescents, 1985-2000. *Biomed Environ Sci* 2007; 20(1): 1-10.

446 31) The research group on Chinese school students physical fitness and health. Report on the physical  
447 fitness and health surveillance of Chinese school students in 2010, Higher Education Press, 2012.

448 32) Liu WJ, Liu W, Lin R, Li B, Pallan M, Cheng KK, et al. Socioeconomic determinants of childhood  
449 obesity among primary school children in Guangzhou, China. *BMC Public Health* 2016;16(1):482.

450 33) Fjørtoft I, Pedersen AV, Sigmundsson H, Vereijken B. Measuring physical fitness in children who are  
451 5 to 12 years old with a test battery that is functional and easy to administer. *Phys Ther*  
452 2011;91(7):1087-95.

453 34) Grant S, Corbett K, Amjad AM, Wilson J, Aitchison T. A comparison of methods of predicting  
454 maximum oxygen uptake. *Br J Sports Med* 1995;29:147-152

455 35) Cooper KH, Kenneth H. A means of assessing maximal oxygen intake: correlation between field and  
456 treadmill testing. *JAMA* 1968;203(3):201-204.

457 36) de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth  
458 reference for school-aged children and adolescents. *Bulletin of the World Health Organization* 2007;  
459 85(9): 660-7.

460 37) Mi J, Wang TY, Meng LH, Zhu GJ, Han SM, Zhong Y, et al. Development of blood pressure  
461 reference standards for Chinese children and adolescents. *Chin J Evid Based Pediatr* 2010;5(1):4-14.

462 38) Jones MA, Hitchen PJ, Stratton G. The importance of considering biological maturity when assessing  
463 physical fitness measures in girls and boys aged 10 to 16 years. *Ann Hum Biol* 2000;27(1):57-65.

464 39) Torrance B, McGuire KA, Lewanczuk R, McGavock J. Overweight, physical activity and high blood  
465 pressure in children: a review of the literature. *Vasc Health Risk Manag* 2007;3(1):139-149.

466 40) Lavie CJ, Parto P, Archer E. Obesity, Fitness, Hypertension, and Prognosis Is Physical Activity the  
467 Common Denominator? *JAMA Intern Med* 2016;176(2):217-218.

468 41) Crump C, Sundquist J, Winkleby MA, Sundquist K. Interactive effects of physical fitness and body  
469 mass index on the risk of hypertension. *JAMA Intern Med* 2016;176(2):210-6.

470 42) Hayes HM, Eisenmann JC, Pfeiffer K, Carlson JJ. Weight Status, physical activity, and vascular health  
471 in 9- to 12-Year-old children. *J Phys ActHealth* 2013;10(2):205-10.

472 43) Lambrechtsen J, Rasmussen F, Hansen HS, Jacobsen IA. Tracking and factors predicting rising  
473 in 'tracking quartile' in blood pressure from childhood to adulthood: Odense Schoolchild Study. *J Hum*  
474 *Hypertens* 1999; 13(6): 385-391.

475 44) Kelly RK, Thomson R, Smith KJ, Dwyer T, Venn A, Magnussen CG. Factors affecting tracking of  
476 blood pressure from childhood to adulthood: the Childhood Determinants of Adult Health Study. *J*  
477 *Pediatr* 2015; 167(6): 1422-1428.

478 45) Jago R, Drews KL, McMurray RG, Thompson D, Volpe SL, Moe EL, et al. Fatness, fitness, and  
479 cardiometabolic risk factors among sixth-grade youth. *Med Sci Sports Exerc* 2010;42(8): 1502-10.

480 46) Ahimastos AA, Formosa M, Dart AM, Kingwell BA. Gender differences in large artery stiffness  
481 pre-and post puberty. *J Clin Endocrinol Metab* 2003; 88(11):5375-80.

482 47) Bai Y, Chen S, Laurson KR, Kim Y, Saint-Maurice PF, Welk GJ. The associations of youth physical  
483 activity and screen time with fatness and fitness: The 2012 NHANES National Youth Fitness Survey."  
484 *PLoS One* 2016;11(1): e0148038.

485 48) Ruiz JR, Rizzo NS, Hurtig-Wennlöf A, Ortega FB, Wärnberg J, Sjöström M. Relations of total  
486 physical activity and intensity to fitness and fatness in children: the European Youth Heart Study. *Am J*  
487 *Clin Nutr* 2006; 84(2): 299-303.

488 49) Carson V, Rinaldi RL, Torrance B, Maximova K, Ball GD, Majumdar SR, et al. Vigorous physical  
489 activity and longitudinal associations with cardiometabolic risk factors in youth. *Int J Obes(Lond)* 2014;  
490 38(1): 16-21.

491 50) Samitz G, Egger M, Zwahlen M. Domains of physical activity and all-cause mortality: systematic  
492 review and dose-response meta-analysis of cohort studies. *Int J Epidemiol* 2011;40(5):1382-400.

493 51) Muntner P, He J, Cutler JA, Wildman RP, Whelton PK. Trends in blood pressure among children and

494 adolescents. JAMA 2004;291(17):2107-13.

495 52) Tu W, Eckert GJ, DiMeglio LA, Yu Z, Jung J, Pratt JH. Intensified effect of adiposity on blood pressure  
496 in overweight and obese children. Hypertension 2011; 58(5):818-824.

497 53) Ataei N, Hosseini M, Iranmanesh M. The relationship of body mass index and blood pressure in  
498 Iranian children <7 years old. J Trop Pediatr 2009;55:313-317.

499 54) Makgae PJ, Monyeki KD, Brits SJ, Kemper HC, Mashita J. Somatotype and blood pressure of rural  
500 South African children aged 6-13 years: Ellisras longitudinal growth and health study. Ann Hum Biol  
501 2007;34:240-251.

502

503  
504

**Table 1: Anthropometric and physiological parameters in Chinese boys and girls aged 5 to 12 years (n=4926).**

<b>Variables</b>	<b>Boys (2725)</b>	<b>Girls (2201)</b>	<b>Total (4926)</b>
Mean age, years (SD)	9.3±1.5	9.3±1.5	9.3±1.5
Pre-Pubertal, n (%)	2725 (100)	2125 (96.5)	4850 (98.5)
Overweight, n (%)	348 (12.8)	189 (8.6)*	537 (10.9)
Obese, n (%)	279 (10.2)	61 (2.8)*	340 (6.9)
BMI-z score	-0.01±1.37	-0.45±1.15*	-0.21±1.30
Reduced Cooper test (distance run in m)	899.4±133.4	864.8±163.0*	884.0±148.4
<b>SBP, mmHg (Mean ± SD)</b>			
	107.6±9.6 <sup>§</sup>	106.1±9.7 <sup>§</sup>	106.9±9.7
5-6 years (n=320)	101.8±8.8	101.9±8.1	101.9±8.5
7-8 years (n=1868)	105.6±9.3	103.3±9.0	104.6±9.2
9-10 years (n=1977)	109.0±9.1	107.6±9.7	108.3±9.4
11-12 years (n=761)	111.4±9.7	110.8±9.3	111.15±9.6
<b>DBP, mmHg (Mean ± SD)</b>	64.7±7.4 <sup>#</sup>	65.0±7.4 <sup>#</sup>	64.9±7.4
5-6 years (n=320)	61.1±7.1	61.9±6.4	61.5±6.8
7-8 years (n=1868)	63.3±7.2	63.3±7.2	63.3±7.2
9-10 years (n=1977)	65.8±7.1	66.2±7.4	66.0±7.2
11-12 years (n=761)	67.0±7.3	67.5±6.8	67.2±7.1
<b>High SBP, n (%)</b>			
Yes	367 (13.5)	327 (14.9)	694 (14.1)
<b>High DBP, n (%)</b>			
Yes	100 (3.7)	102 (4.6)	202 (4.1)
<b>HBP, n (%)</b>			
Yes	398 (14.6)	354 (16.1)	752 (15.3)

505 **Note:** Continuous variables were described by means ± standard deviation. Categorical variables  
506 are described by frequency (%). \* Statistical significant between boys and girls, P<0.05. § SBP  
507 increased with increasing age in boys and girls, P<0.05. # DBP increased with increasing age in  
508 boys and girls, P<0.05.



**Table 2: Relationship between weight status and cardiorespiratory fitness among school children in Guangzhou, China.**

	N	Cardiorespiratory fitness (Reduced Cooper test )							
		1 <sup>st</sup> Quartile	2 <sup>nd</sup> Quartile	3 <sup>rd</sup> Quartile	4 <sup>th</sup> Quartile	Lower Fitness	Higher Fitness		
		%	%	%	%	P-value	%	%	P-value
<b>Boys</b>									
Non-overweight	2098	21.5	23.8	26.2	28.5		45.3	54.7	
Overweight	348	27.9	29.3	24.7	18.1	<0.001	57.2	42.8	<0.001
Obese	279	49.8	30.1	14	6.1		79.9	20.1	
<b>Girls</b>									
Non-overweight	1951	24.7	23.9	25.6	25.8		48.5	51.5	
Overweight	189	31.7	29.1	20.6	18.5	0.001	60.8	39.2	<0.001
Obese	61	44.3	36.1	13.1	6.6		80.3	19.7	
		Mean distance (m) + SD	Mean distance (m) + SD	Mean distance (m) + SD	Mean distance (m) + SD		Mean distance (m) + SD	Mean distance (m) + SD	
<b>Boys</b>									
Non-overweight	2098	761.0±85.0	868.8±39.3	951.0±49.3 <sup>§</sup>	1060.8±76.4		816.2±84.9 <sup>§</sup>	997.4±82.5	
Overweight	348	750.3±86.4	865.5±38.5	941.5±44.8	1047.9±95.7		810.8±97.3	993.0±99.9	
Obese	279	740.3±85.7	863.3±41.7	933.4±43.0	1034.0±54.2		786.6±93.7	969.6±63.9	
<b>Girls</b>									
Non-overweight	1951	737.0±80.2	838.3±24.6	905.4±41.7	1151.6±98.2 <sup>§</sup>		785.4±78.7	1021.9±65.8 <sup>§</sup>	
Overweight	189	734.3±85.0	837.7±32.4	896.3±33.7	986.9±68.5		784.2±81.6	941.8±70.5	
Obese	61	735.8±73.2	835.7±29.4	896.0±34.7	976.0±79.1		781.6±77.4	922.7±63.4	

**Note:** weight status (defined using WHO 2007 reference standard); 1<sup>st</sup> Quartile: 1%-25% percentiles, 2<sup>nd</sup> Quartile: 26%-50% percentiles; 3<sup>rd</sup> Quartile: 51%-75% percentiles, 4<sup>th</sup> Quartile: 76%-100% percentiles. § Reduced Cooper test decreased with weight status in boys and girls, P<0.05. Lower Fitness (1st and 2nd Quartile); Higher Fitness (3rd and 4th Quartile).

**Table 3: Generalized linear mixed model analysis of the association between weight status, cardiorespiratory fitness and HBP among school children in Guangzhou, China.**

Characteristics	HBP	Likelihood of HBP		
	Number (%)	Unadjusted model OR(95% CI)	Adjusted model I OR(95% CI)	Adjusted model II OR(95% CI)
<b>Boys(n=2725)</b>				
<b>Children's weight status</b>				
Non-overweight	2098(9.0) **	Reference	Reference	Reference
Overweight	348(27.6)	3.94(2.97-5.22) **	3.51(2.62-4.70) **	2.96(1.71-5.11) **
Obese	279(40.5)	7.03(5.28-9.37) **	5.55(4.07-7.57) **	4.19(2.63-6.67) **
<b>Reduced Cooper test</b>				
1 <sup>st</sup> quartile	688(18.2) **	Reference	Reference	Reference
2 <sup>nd</sup> quartile	685(15.5)	0.81(0.61-1.09)	0.71(0.56-1.02)	0.69(0.44-1.07)
3 <sup>rd</sup> quartile	675(12.9)	0.64(0.47-0.88) *	0.64(0.47-0.89) *	0.88(0.58-1.33)
4 <sup>th</sup> quartile	677(11.8)	0.59(0.43-0.81) *	0.64(0.46-0.89) *	0.87(0.56-1.35)
<b>Girls (n=2201)</b>				
<b>Children's weight status</b>				
Non-overweight	1951(13.8) **	Reference	Reference	Reference
Overweight	189(30.2)	2.66(1.89-3.73) **	1.93(1.34-2.78) **	1.75(1.00-3.06) **
Obese	61(45.9)	5.16(3.04-8.75) **	4.11(2.37-7.13) **	2.49(1.19-5.19) **
<b>Reduced Cooper test</b>				
1 <sup>st</sup> quartile	568(15.3)	Reference	Reference	Reference
2 <sup>nd</sup> quartile	543(17.3)	1.14(0.83-1.59)	1.14(0.85-1.54)	1.14(0.83-1.57)
3 <sup>rd</sup> quartile	547(18.6)	1.25(0.90-1.74)	0.92(0.66-1.29)	1.01(0.71-1.43)
4 <sup>th</sup> quartile	543(13.1)	0.81(0.57-1.17)	0.70(0.43-1.13)	0.71(0.43-1.18)

**Note:** Adjusted model I: Adjusted for age, height and pubertal status. Adjusted model II: Model includes age, height, pubertal status, weight status (WHO 2007 categories), CRF quartiles and weight status×CRF quartiles.\*\* P<0.001, \* P<0.05.

**Table 4: The mean SBP, DBP and prevalence of hypertension across weight status and cardiorespiratory fitness groups among Chinese schoolboys and girls in Guangzhou.**

Characteristics	Boys(n=2725)				Girls(n=2201)			
	Hypertension		Mean ± SD		Hypertension		Mean ± SD	
	Number	%	SBP	DBP	Number	%	SBP	DBP
<b>Reduced Cooper test</b>								
Non-overweight								
Higher fitness	1147	8.7	105.49±8.77	63.28±6.99	1004	14.1	105.28±9.36	64.32±7.25
Lower fitness	951	9.4	105.86±8.75	63.81±6.81	947	13.4	105.12±9.36	64.52±7.18
P-value		0.61	0.33	0.08		0.64	0.71	0.54
Overweight								
Higher fitness	149	28.2	113.62±9.42	67.70±7.19	74	31.1	111.85±9.69	67.96±7.00
Lower fitness	199	27.1	111.92±9.55	67.65±7.27	115	29.6	111.79±9.19	69.28±6.50
P-value		0.83	0.10	0.95		0.83	0.97	0.19
Obese								
Higher fitness	56	44.6	116.46±9.96	70.62±6.80	12	66.7	117.63±5.96	72.33±6.22
Lower fitness	223	39.5	115.68±9.10	70.17±7.54	61	49.5	115.78±10.65	71.43±7.97
P-value		0.48	0.57	0.69		0.11	0.43	0.72

**Note:** Weight status (defined using WHO 2007 reference standard); Lower fitness refer to 1<sup>st</sup> Quartile and 2<sup>nd</sup> Quartile fitness and Higher fitness refer to 3<sup>rd</sup> Quartile and 4<sup>th</sup> Quartile fitness.