

UNIVERSITY OF BIRMINGHAM

Research at Birmingham

Resolvin D1 stimulates epithelial wound repair and inhibits TGF- induced EMT whilst reducing fibroproliferation and collagen production

Zheng, Shenxing; Wang, Qian; D'Souza, Vijay; Bartis, Domokos; Dancer, Rachel; Parekh, Dhruv; Gao Smith, Fang; Lian, Qingquan; Jin, Shengwei; Thickett, David

DOI:

[10.1038/labinvest.2017.114](https://doi.org/10.1038/labinvest.2017.114)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Zheng, S, Wang, Q, D'Souza, V, Bartis, D, Dancer, R, Parekh, D, Gao Smith, F, Lian, Q, Jin, S & Thickett, D 2017, 'Resolvin D1 stimulates epithelial wound repair and inhibits TGF- induced EMT whilst reducing fibroproliferation and collagen production', Laboratory investigation. <https://doi.org/10.1038/labinvest.2017.114>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

Resolvin D1 stimulates epithelial wound repair and inhibits TGF- β induced EMT whilst reducing fibroproliferation and collagen production

Shenxing Zheng^{1,2}, Qian Wang^{1,2}, Vijay D'Souza², Dom Bartis², Rachel Dancer², Dhruv Parekh², Fang Gao², Qingquan Lian¹, Shengwei Jin¹, David R Thickett²

1. Department of Anesthesia and Critical Care, The Second Affiliated Hospital and Yuying Children's Hospital of Wenzhou Medical University, Zhejiang 325027, China.

2. Institute of Inflammation and Aging, University of Birmingham, Edgbaston Birmingham B15 2TT UK.

Shengxing Zheng (zsx0106014@163.com)

Qian Wang (wangqian84@aliyun.com)

Vijay K D'Souza (V.K.DSouza@bham.ac.uk)

Domokos Bartis (d.bartis@outlook.com)

Rachel CA Dancer (r.dancer@bham.ac.uk)

Dhruv Parekh (d.parekh@bham.ac.uk)

Babu Naidu (babu.naidu@heartofengland.nhs.uk)

Fang Gao-Smith (f.g.smith@bham.ac.uk)

ShengweiJin (jinshengwei69@163.com)

QingquanLian (lianqingquanmz@163.com)

David R Thickett (d.thickett@bham.ac.uk)

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

Correspondence: Dr David Thickett d.thickett@bham.ac.uk

telephone – sec+44 121 371 4841 fax +44 121 627 8245.

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

ABSTRACT

Acute and chronic inflammatory lung diseases are often associated with epithelial cell injury/loss and fibroproliferative responses. ResolvinD₁ (RvD1) is biosynthesized during the resolution phase of inflammatory response and exerts potent anti-inflammatory and promotes resolution of inflammatory lung diseases. The aim of this study was to investigate whether RvD1 exerts protective effects on alveolar epithelial cell function / differentiation and protects against fibroproliferative stimuli. Primary human alveolar type II cells were used to model the effects of RvD1 in vitro upon wound repair, proliferation, apoptosis, transdifferentiation and epithelial mesenchymal transition (EMT). Effects of RvD1 upon primary human lung fibroblast proliferation, collagen production, and myofibroblast differentiation were also examined. RvD1 promoted alveolar type II (ATII) cell wound repair and proliferation. RvD1 protected ATII cells against sFas-ligand/TNF- α -induced apoptosis and inhibition on cell proliferation and viability. RvD1 promoted ATII cells transdifferentiation. Moreover, we demonstrate that RvD1 inhibited EMT in response to TGF- β . Furthermore RvD1 inhibited HLF proliferation, collagen production and myofibroblast differentiation induced by both TGF- β and bronchoalveolar lavage fluid (BALF) from ARDS patients. The effects of RvD1 were PI3-kinase dependent and mediated via the resolvin receptor. RvD1 seems to

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

promote alveolar epithelial repair by stimulating AII cells wound repair, proliferation, reducing apoptosis and inhibiting TGF- β induced EMT. While RvD1 reduced fibroproliferation, collagen production and myofibroblast differentiation. Together, these results suggest a potential new therapeutic strategy for preventing and treating chronic diseases (such as IPF) as well as the fibroproliferative phase of ARDS by targeting RvD1 actions that emphasizes natural resolution signalling pathways.

Key words: idiopathic pulmonary fibrosis, acute respiratory distress syndrome, resolvins, alveolar type II cells, lung fibroblast.

INTRODUCTION

Chronic lung diseases such as idiopathic pulmonary fibrosis (IPF) are associated with loss of alveolar epithelial cells due to apoptosis, excessive fibroproliferation, aberrant deposition of extracellular matrix (EMC), inflammation and dysregulated repair of lung tissue.^{1, 2} Many Studies suggest alveolar epithelial cells play a central role in the pathogenesis of IPF.³⁻⁶ In IPF, alveolar epithelial cells may undergo increased apoptosis through a Fas ligand-mediated pathway.⁷ Injury to the alveolar epithelial barrier is an early event during the development of pulmonary fibrosis.⁶ Timely repair of lung injury is essential for proper restoration of function. The repair of a damaged alveolar epithelial barrier is a complex and poorly understood process that includes transdifferentiation of type II epithelial cells into type I epithelial cells, as well as regeneration of epithelial cells from stem cells.⁸ Dysregulation of repair mechanisms such as epithelial to mesenchymal transition, may contribute to the generation of numerous ECM-producing fibroblasts/myofibroblasts,^{9, 10} that result in clinically significant pulmonary fibrosis. Similarly, in acute lung diseases such as acute respiratory distress syndrome (ARDS), the degree of the epithelial injury and the loss of alveolar epithelial cells due to Fas ligand-mediated apoptosis is an important predictor of outcome.¹¹⁻¹³ In some cases of ARDS, a marked fibroproliferative response is

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

associated with bad outcome.¹⁴ Therefore, a therapy that promotes epithelial repair and inhibits EMT could be useful in both acute and chronic respiratory disease, but only if it wasn't also a stimulus for fibro-proliferation.

ResolvinD₁ (RvD1) is a lipid mediator derived from both eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) which acts to dampen excessive PMN infiltration and transmigration.¹⁵ Previous studies have suggested that RvD1 attenuates lung inflammation and can maintain the integrity of lung epithelium.¹⁶ Furthermore, RvD1 has recently been shown to reduce interstitial fibrosis,¹⁷ inhibit cytokines release at sites of inflammation,^{18, 19} and is protective after ischemia-reperfusion second organ injury.^{17, 20} RvD1 directly activates the lipoxinA₄ receptor/formyl peptide receptor 2 (ALX/FPR2) with high affinity.²¹

Currently what is not known is whether RvD1 has a direct role in modulating human lung epithelial cell or primary human lung fibroblast proliferation and function.

Our results indicate that RvD1 promotes epithelial wound repair and inhibited TGF- β induced EMT in human adult type II alveolar epithelial cells, whilst inhibiting fibroproliferation and reducing the effects of TGF- β on primary human lung fibroblast (HLF) collagen production and myofibroblast differentiation.

MATERIAL AND METHODS

Reviewers these are the short methods – more detailed methods are in the online supplement.

Reagents

ResolvinD1 was purchased from Cayman chemicals (Cayman Chemical Company, USA). Recombinant human TGF- β was purchased from R&D (R&D Systems, Abingdon, UK). Antibody against caspase-8, AKT and phospho-AKT were obtained from Cell Signal Technology (Cell signal Technology, Boston, USA). Antibody against E-cadherin, N-cadherin and α -SMA were obtained from Abcam (Abcam, Cambridge, UK). Antibody against β -actin was purchased from Santa Cruz Biotechnology Inc (Santa Cruz, CA, USA).

Primary lung cell culture

ATII cells were isolated from peripheral normal lung tissue distal from the tumour in patients undergoing lung cancer resection. The cells were isolated in accordance with approval from the local research ethics committees at the University of Birmingham (Birmingham, UK). Primary human alveolar type II (AT II) cells were extracted according to methods described previously (see

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

online supplement).²² Average yields of primary human alveolar type II cells were 30.2 million cells per resection with an average purity of 92% ATII-like cells. Cells were tested for primary human alveolar type II (AT II) cell phenotype by alkaline phosphatase staining, lysotracker lamellar body staining and by PCR expression of surfactant protein C—a type II cell marker with negative expression of aquaporin V (a type I cell marker) (data not shown).

Primary human lung fibroblasts (HLF) from Lonza™ were similarly cultured in dulbecco modified Eagle medium culture media (ECACC, Sigma, Poole, UK) supplemented with 10% FCS (sigma, Poole, UK) at 37 °C and 5% CO₂. Cells were subcultured at 60-80% confluence using trypsin/EDTA. Cells were obtained from three separate donors, and all experiments were repeated in triplicate.

Stimuli and Inhibitors. ATII cells and fibroblasts were treated with resolvinD₁ (10nM, 25nM or 100nM) (Cayman Chemical Company, USA). Appropriate vehicle controls were used for all experiments with inhibitors. Inhibitors were used at the following concentrations according to manufacturers' instructions: LY294002, a PI3-kinase inhibitor (Calbiochem, Nottingham, UK) at 10 μM; and the ALXR antagonist, Boc-2 (N-t-Boc-Phe-Leu-Phe-Leu-Phe; GenScript USA Inc), at 10μM. Inhibitors were added to cells 1 hour prior to every treatment.

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

Bronchoalveolar Lavage Fluid (BALF) Collection

BALF from ARDS patients is known to stimulate epithelial repair in the scratch wound assay in an IL-1 β dependent fashion.²³ To test whether resolvin D1 could augment or synergise with this effect, the BALF from patients with ARDS were mixed 50:50 with appropriate culture media for each cell type as a positive control stimulus. We used BALF from patients enrolled into the BALTI-1 trial, demographics for whom have been published previously.²⁴

In Vitro Alveolar Epithelial Wound Repair Assay. Epithelial repair was determined using an in vitro epithelial wound repair assay as described before.²⁵ Briefly, primary human alveolar type II (AT II) cells were grown to confluent monolayers before wounding with a 1-mL pipette tip. Digital images of the same point on the wound were taken at time 0 and at time 36 hours. To control for the inconsistencies in wound size, only monolayers in which the original wound areas varied by 10% of the mean were analyzed. Repair is expressed as the percentage of the original wound area covered by cells relative to control media. To allow for variability between cell types and batches, data are expressed as the mean (SE) percentage of control).

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

BRDU cell proliferation assay and cell viability assays: BrdU incorporation was assessed according to manufacturers' instructions (BRDU Cell Proliferation Assay, Promega, UK). Cell Viability after 24 hours was assessed adding 20µL of Cell Titer 96 aqueous one solution cell proliferation solution (Promega, UK) to cells for 1.5 hours at 37°C and 5% CO₂ as described.²⁶

Flow Cytometry: Apoptosis of epithelial cells was assessed as described previously using flow cytometry.²² Cells were left in serum-free media for 24 hours before exposure to 100 ng/ml Fas-ligand (R&D Systems, Abingdon, UK). Apoptosis was determined by flow cytometry using the Annexin V and SyTOX antibody according to the manufacturer's recommendations (Molecular Probes, Eugene, OR) after 24 hours exposure.

Quantitative PCR: Quantitative PCR was performed using commercially obtained primers as outlined in supplementary methods.

Western Blot Analysis: Western blot analyses from cells homogenates were performed as described previously. After equal amounts of protein were electrophoresed on 10/12% sodium dodecyl sulfate-polyacrylamide gels and then transferred to polyvinylidene difluoride membranes (Millipore, Billerica MA01821).

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

Western blot analysis was performed using the Image Quant LAS 4000 mini (GE).

HLF proliferation assay in response to ARDS BALF: HLF were plated out at 2500 cells per well. A standard curve of cell counts was pipetted from 1250 to 15000 per well. BALF mixed 50:50 with media was added from 10 ARDS patients. After 24 hours BRDU incorporation was assessed according to manufacturers' instructions (BRDU Cell Proliferation Assay, Promega, UK). Results were extrapolated from the standard curve. Each patient sample was run with 6 replicates upon a single batch of HLF at passage 3.

Statistical Analysis: Data were normally distributed and analyzed by analysis of variance with Tukey's test for *post hoc* comparisons using Minitab 14.0 (Minitab, State College, PA). A p value equal or less than 0.05 was considered significant. Data are expressed as mean (SEM).

RESULTS

ResolvinD₁ stimulates ATII cell wound repair and proliferation in vitro.

RvD1 increased AT II cell wound closure after 36 hours compared with control media. ARDS BALF increased ATII cell wound closure

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

compared with media control²³. RvD1 + ARDS BALF further enhanced the wound repair response (Fig. 1A).

Scratch wound repair can occur due to either spreading of cells and /or proliferation. Cell proliferation studies confirmed that RvD1 stimulated proliferation of AT II cell in a dose dependent manner (Fig. 1B).

ResolvinD₁ promotes ATII cell proliferation through activation of ALX receptor and the PI3K/AKT signaling pathway.

The PI3K/AKT signaling pathway plays an important role in cell proliferation. To determine if PI3-kinase signalling was involved in the RvD1 proliferative response, LY294002 (PI3K inhibitor) was incubated with ATII cells at 10 μ M for 1 hour before RvD1 treatment of ATII cells. LY294002 treatment reversed the effects of RvD1 (100nM) on the proliferation of ATII cells compared with control media-treated cells (Fig 2A). To investigate whether RvD1 can activate AKT phosphorylation in ATII cells, ATII cells were stimulated with different concentrations of RvD1 (10, 25, and 100 nmol/ml) for 24h. We found that RvD1 activated AKT phosphorylation (Fig 2B). In addition, Pre-treatment of cells with Boc-2 (the resolvin ALX receptor antagonist) inhibited the effects of RvD1 on the proliferation of ATII cells (Fig. 2A).

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

ResolvinD₁ protects against Fas-ligand (sFasL) and TNF- α actions on ATII cells.

sFasL and TNF- α inhibited cellular proliferation compared with control media-treated cells. This effect was attenuated by 100nM RvD1 pretreatment (see online supplement Fig.s1A). The addition of 100 ng/ml soluble Fas-ligand (sFasL) or 100 ng/ml TNF- α , as expected, significantly reduced cellular viability compared with control group. Co-treatment with 100nM RvD1 significantly increased cellular viability compared with sFasL ($p < 0.01$) (see online supplement Fig.s1B)- an effect that was also reduced by RvD1 rescue therapy added 30 mins after treatment with sFasL (data not shown). sFasL treatment of ATII cells (which are known to be resistant to apoptosis) increased the number of apoptotic cells from $3.29 \pm 0.11\%$ in control cells to $10.34 \pm 0.33\%$ ($p = 0.01$). Rescue treatment with RvD1 reduced the number of apoptotic cells to $4.12 \pm 0.52\%$ ($p = 0.01$) (Fig. 3A).

ResolvinD₁ inhibited Fas-ligand- induced caspase-8 activation in ATII cells.

To research the impact of RvD1 on Fas-ligand induced ATII cells apoptosis, we preliminarily detected the caspase-8 levels in vitro. ATII cells were treated with Fas-ligand or/and RvD1. The protein levels of caspase-8 were quantified and analyzed in the indicated

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

groups (Fig. 3B). Fas-ligand promoted caspase-8 activation. RvD1 inhibited Fas-ligand induced caspase-8 activation in ATII cells.

ResolvinD₁ promotes Aquaporin 5 gene expression (a type I epithelial cell marker) whilst inhibiting surfactant protein C (SP-C, a type II epithelial cell marker) gene expression on ATII cells.

Aquaporin 5 (AQP 5) is a membrane protein that mainly facilitates osmotic water transport.²⁸ AQP5 may promote alveolar fluid clearance or maintain integrity of epithelial barrier.²⁹ Roles of AQP5 other than fluid transport have been explored in animal lung injury models, results from these studies show that lung injury is associated with down-regulation of AQP5 expression.^{30, 31} AQP5 expression has been used as a type I epithelial cell marker. To observe the effect of RvD1 on AQP5 and SP-C gene expression, ATII cells were treated by RvD1 for 24 hours. RvD1 increased AQP5 gene expression (2.42 ± 0.34 fold) relative to control group, $p=0.001$ (Fig. 4A). RvD1 induced AQP5 expression was blocked by BOC-2 (the ALXR antagonist) (Fig. 4A). In contrast, RvD1 down-regulated mRNA expression of SP-C suggesting that RvD1 may promote transdifferentiation of ATII cells into ATI like cells (Fig. 4B).

ResolvinD₁ inhibit TGF- β induced EMT in Primary human alveolar type II cells

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

The epithelial-mesenchymal transition (EMT) of alveolar epithelial cells is a phenotype conversion, which is one of the main mechanisms of pulmonary fibrosis.³² The EMT process of epithelial cells is often stimulated with TGF- β .³² In our study, EMT was induced in ATII cells with TGF- β treatment. TGF- β treated ATII cells showed a mesenchymal morphology (fibroblast-like), and RvD1 restored the epithelial morphology of the cells to a certain extent (Fig. 5A). RvD1 blocked the expression of mRNA of mesenchymal markers including N-cadherin, vimentin, type I collagen, S100A4, and α -SMA, while RvD1 restored the expression of mRNA of E-cadherin (Fig.5B). The effects of RvD1 on the TGF- β -treated E-cadherin, α -SMA, N-cadherin of ATII cells were confirmed by Western blot. (Fig.5C).

To elucidate the mechanism involved in the effects of RvD1 on EMT, pre-treatment of cells with Boc-2 (the ALX receptor antagonist), inhibited the effects of RvD1 on EMT of ATII cells. TGF- β -induced N-cadherin expression was suppressed by RvD1, and pre-treatment of cells with Boc-2 incapacitated the effects of RvD1 on N-cadherin expression (Fig. 6). Reduced expression of E-cadherin in TGF- β -treated ATII cells was restored by RvD1, but pre-treatment of cells with Boc-2 abolished the effects of RvD1 (Fig. 6).

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

ResolvinD₁ inhibits proliferation of Primary HLF Induced by TGF- β and this effect was PI3-Kinase dependent and blocked by BOC-2.

Cell proliferation studies confirmed that 100 nM RvD1 inhibited proliferation of primary HLF Induced by TGF- β . Cells were treated with TGF- β for 24 h with or without preincubation with the PI3-kinase inhibitor LY294002 (10 μ M) or BOC-2 (10 μ M). RvD1 inhibited the effects of TGF- β on HLF proliferation, and these effects were blocked by both LY294002 and BOC-2 (Fig. 7).

ResolvinD₁ inhibits proliferation of primary HLF induced bronchoalveolar lavage fluid (BALF) from patients with ARDS.

ARDS BALF has previously been shown to promote fibroblast proliferation in vitro.³³ To model the in vivo stimulus for fibroproliferation in ARDS, HLF were treated with a 50:50 mix of BALF from patients with ARDS. RvD1 inhibited ARDS BALF induced proliferation in HLF (Fig 8).

ResolvinD₁ reduces primary HLF collagen production and α -smooth muscle actin (α -SMA) induced by TGF- β and BALF from patients with ARDS.

We investigated the mRNA expression of type I collagen, type IV collagen and α -SMA in HLF induced by TGF- β 10ng/ml with quantitative PCR. Gene expression of type I collagen, type IV

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

collagen and α -SMA were increased in HLF induced by 24 hours of treatment with TGF- β relative to control group. Treatment with RvD1 significantly inhibited gene expression of type I collagen, type IV collagen and α -SMA in HLF induced by TGF- β compared to TGF- β group respectively (Table 1). We also investigated the effect of ARDS BALF upon type I collagen, type IV collagen and α -SMA mRNA expression. Gene expression of type I collagen, type IV collagen and α -SMA were increased in HLF induced by ARDS BALF relative to control group (see table 1). RvD1 significantly inhibited gene expression of type I collagen, type IV collagen and α -SMA in HLF induced by by ARDS BALF (see Table 1).

DISCUSSION

Epithelial injury is one of the hallmarks of ARDS and fibrotic lung diseases.^{1, 34} Progressive pulmonary fibrosis occurs due to recurrent injury to AECs followed by aberrant repair/regeneration of epithelial barrier, persistence of activated fibroblasts, and alterations in extracellular matrix (ECM).^{1, 35} It has been suggested that resolvinD1 (RvD1) exerts potent anti-inflammatory and proresolution effects, without causing immunosuppression. Furthermore, we previously reported that RvD1 improved alveolar fluid clearance, decreased pulmonary edema and maintained the

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

integrity of lung epithelia in LPS-induced murine lung injury.³⁶ Aspirin-triggered RvD1 also improved epithelial and endothelial barrier integrity in a murine model of hydrochloric acid-induced ALI.³⁷ RvD1 promoted macrophage phagocytosis of zymosan and apoptotic PMNs signaling through the human ALX and GPR32 receptors.²¹ Based on this background, our purpose was to evaluate whether RvD1 stimulated physical wound repair, promoted cellular proliferation, and inhibited apoptosis in AII cells processes that are dysregulated repair in ARDS and fibrotic lung diseases.

The restoration of the alveolar epithelial barrier is a critical aspect of alveolar repair,²⁵ and our study clearly demonstrated that RvD1 stimulated alveolar repair promoting physical wound closure by inducing proliferation of primary alveolar epithelial cells. The PI3K-AKT signaling pathway regulates proliferation. Recombinant mouse osteopontin induces the proliferation of human bronchial smooth muscle cells via the PI3K/AKT signaling pathway.³⁸ Our study also showed that the mitogenic response of AII cells to RvD1 is mediated through activation of ALX receptor and the PI3K/AKT signaling pathway.

However, the precise role of RvD1 as modulators of apoptosis remains elusive and it is also unclear whether the effects relate to increasing lung epithelial survival. Previous studies reported that RvD1 can stimulate apoptosis of T cells and PMNs, while our study

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

demonstrated that RvD1 inhibited apoptosis in sFasL treated cells. One study has also indicated that RvD1 decreased apoptosis induced by ER stress in HepG2 cells.³⁹ Our study demonstrates that RvD1 reduced cell death in response to sFasL and /or TNF- α even when given after the onset of injury and, therefore, may have potential as a rescue therapy post-injury. Furthermore, these effects seemed to relate to caspase-8 activation as caspase-8 levels were elevated in the sFasL treated cells, and RvD1 suppressed sFasL induced caspase-8 activation in ATII cells.

The normal alveolar epithelium is composed of two types of cells (AT I and ATII). The repair of the epithelial barrier is believed to involve the transdifferentiation of type II cells into type I epithelial cells.⁸ The inability of type II AECs to transdifferentiate into type I AECs have also been observed in human lung fibrosis.⁴⁰ Our study showed that RvD1 promoted gene expression of AQP5 (type I marker) in ATII cells via activation of ALX supporting a potential role for RvD1 in promoting fluid transport. while RvD1 reduced gene expression of surfactant protein C (type II markers) in ATII cells suggesting that it may promote transdifferentiation of ATII cells towards the ATI phenotype. Otherwise, recurrent alveolar epithelial cell (AEC) injury that leads to aberrant activation of AEC (such as epithelial to mesenchymal transition, EMT), producing fibroblasts and myofibroblasts.^{9, 10}

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

Epithelial mesenchymal transition (EMT) has been increasingly proposed as one of the causative mechanisms of lung fibrosis.^{41, 42}

In this study, we demonstrated that RvD1 restored the epithelial morphology of the cells to a certain extent. Our results also revealed that RvD1 inhibited TGF- β induced expression of the mesenchymal markers N-cadherin, vimentin, (FSP)-1 (also called S100A4), type I collagen and α -SMA (the mesenchymal cell markers) mRNA expression, while maintaining the epithelial marker E-cadherin (the epithelial cell marker) mRNA expression in ATII cells. Effects that were confirmed by Western blot. These results suggest that RvD1 may be a suppressor of TGF- β -induced EMT in ATII cells. These effects were inhibited by pre-treatment of ATII cells with Boc-2 (the resolvin ALX receptor antagonist). The role of RvD1 on EMT has also been documented in A549 cells lung cancer cells.⁴³

Recent studies have indicated that 17(R)-RvD1 attenuated bleomycin-induced pulmonary fibrosis by promoting the resolution of neutrophilic inflammation in mice.⁴⁴ Treatment with 17(R)-RvD1 attenuated neutrophil alveolar infiltration, lung collagen content, and type I collagen mRNA expression, which was inhibited by an antagonist of ALX/FPR2 receptor.⁴⁴ Resolvin D1 (RvD1) can suppress renal fibrosis in the obstructed kidney via inhibiting fibroblast proliferation and production of fibronectin and collagen I.^{45, 46} We therefore studied the effects of RvD1 upon primary HLF.

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

RvD1 inhibited TGF- β induced proliferation in primary HLF via activation of ALX/FPLR-1 and this effect was mediated through the PI3/Akt signalling pathway.

It is established that TGF- β , as a potent inducer of fibroblast differentiation into myofibroblasts, can stimulate fibroblast proliferation and collagen production.^{47, 48} We therefore addressed the possibility that RvD1 may inhibit their differentiation into myofibroblasts. RvD1 significantly inhibited gene expression of type I collagen, type IV collagen and α -SMA (a reliable myofibroblast marker) in HLF induced by TGF- β . We also showed ARDS BALF stimulated the expression of markers of myofibroblast differentiation when incubated with normal HLF; an effect that was blocked by RvD1. Our data therefore suggests that RvD1 has differential effects upon ATII and HLF cells in vitro; both promoting epithelial repair and inhibiting TGF- β induced EMT whilst reducing fibroproliferation. This differential effect is potentially vitally important for RvD1 if used as a novel therapy in both acute and chronic inflammatory lung diseases such as ARDS and IPF.

In summary, these data provide evidence for a new mechanism by which RvD1 may contribute to alveolar repair promoting physical wound closure by inducing proliferation of primary human ATII cells in vitro. RvD1 protected ATII cells from pro-apoptotic stimuli even when given after the initial injury. RvD1 increased the expression of

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

the type I marker, AQP5, with reduction in SP-C by ATII like cells, potentially promoting transdifferentiation. Moreover, RvD1 inhibited EMT in response to TGF- β . Intriguingly RvD1 also inhibited HLF proliferation, collagen production and α -SMA expression induced by both TGF- β and ARDS BALF. The potential key role for RvD1 during primary human ATII cells and HLF in vitro is summarized in Fig. 9. Our next step is to determine whether RvD1 has a role in regulating repair and EMT in animals models.

In conclusion, these results suggest a potential new therapeutic strategy for preventing and treating chronic diseases (such as IPF) as well as the fibroproliferative phase of ARDS by targeting RvD1 actions that emphasizes natural resolution signaling pathways. Further experiments are necessary to understand the basic mechanism underlying the anti-fibrotic and anti- apoptotic effects of RvD1, which are currently under investigation.

ACKNOWLEDGMENTS

We would like to thank the patients and relatives who agreed to both tissue donation during elective surgery and participation in translational studies when in intensive care. We would like to thank Teresa Melody and Dawn Hill as nurse/coordinator (both Heart of England NHS Foundation Trust, Birmingham, UK).

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

Support statement: D.R. Thickett was funded by the Medical Research Council (MRC G1100196/1) and the Wellcome Trust. D. Bartis was funded by a European Respiratory Society long-term training fellowship. D. Parekh (MR/J011266/1) and R.C.A.Dancer (G1100196/1) were funded by the MRC. This work was supported by the National Natural Science Foundation of China (grants 81401579) and the Wenzhou Science & Technology Bureau (grants Y20160370).

DISCLOSURE/CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

1. Thannickal VJ, Toews GB, White ES et al. Mechanisms of pulmonary fibrosis. *Annu Rev Med* 2004; 55: 395-417.
2. Wipff PJ, Rifkin DB, Meister JJ et al. Myofibroblast contraction activates latent TGF-beta1 from the extracellular matrix. *J Cell Biol* 2007; 179(6): 1311-1323.
3. Thomas AQ, Lane K, Phillips J et al. Heterozygosity for a surfactant protein C gene mutation associated with usual interstitial pneumonitis and cellular nonspecific interstitial pneumonitis in one kindred. *Am J Respir Crit Care Med* 2002; 165(9): 1322-1328.
4. Garcia, C.K. Idiopathic pulmonary fibrosis: update on genetic discoveries. *Proc Am Thorac Soc* 2011; 8(2): 158-162.

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

5. Alder JK, Barkauskas CE, Limjunyawong N et al. Telomere dysfunction causes alveolar stem cell failure. *Proc Natl Acad Sci U S A* 2015; 112(16):5099-5104.
6. Chen H, Matsumoto K, Brockway BL et al. Airway epithelial progenitors are region specific and show differential responses to bleomycin-induced lung injury. *Stem Cells* 2012; 30(9): 1948-60.
7. Maeyama T, Kuwano K, Kawasaki M et al. Upregulation of Fas-signalling molecules in lung epithelial cells from patients with idiopathic pulmonary fibrosis. *Eur Respir J* 2001; 17(2): 180-9.
8. Thickett, D.R., A.R. Poole, and A.B. Millar et al. The balance between collagen synthesis and degradation in diffuse lung disease. *Sarcoidosis Vasc Diffuse Lung Dis* 2001; 18(1): 27-33.
9. Andersson-Sjöland A, de Alba CG, Nihlberg K et al. Fibrocytes are a potential source of lung fibroblasts in idiopathic pulmonary fibrosis. *Int J Biochem Cell Biol* 2008; 40(10): 2129-40.
10. Omenetti A, Porrello A, Jung Y et al. Hedgehog signaling regulates epithelial-mesenchymal transition during biliary fibrosis in rodents and humans. *J Clin Invest* 2008; 118(10): 3331-42.
11. Martin TR, Hagimoto N, Nakamura M et al. Apoptosis and epithelial injury in the lungs. *Proc Am Thorac Soc* 2005; 2(3): 214-20.
12. Matthay, M.A., L. Robriquet, and X. Fang et al. Alveolar epithelium: role in lung fluid balance and acute lung injury. *Proc Am Thorac Soc* 2005; 2(3): 206-13.

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

13. Zemans, R.L. and M.A. Matthay. Bench-to-bedside review: the role of the alveolar epithelium in the resolution of pulmonary edema in acute lung injury. *Crit Care* 2004; 8(6): 469-77.
14. Martin C, Papazian L, Payan MJ et al. Pulmonary fibrosis correlates with outcome in adult respiratory distress syndrome. A study in mechanically ventilated patients. *Chest* 1995; 107(1): 196-200.
15. Serhan CN, Hong S, Gronert K et al. Resolvins: a family of bioactive products of omega-3 fatty acid transformation circuits initiated by aspirin treatment that counter proinflammation signals. *J Exp Med* 2002; 196(8): 1025-37.
16. Wang B, Gong X, Wan JY et al. Resolvin D1 protects mice from LPS-induced acute lung injury. *Pulm Pharmacol Ther* 2011; 24(4): 434-41.
17. Duffield JS, Hong S, Vaidya VS et al. Resolvin D series and protectin D1 mitigate acute kidney injury. *J Immunol* 2006; 177(9): 5902-11.
18. Recchiuti A, Krishnamoorthy S, Fredman G et al. MicroRNAs in resolution of acute inflammation: identification of novel resolvin D1-miRNA circuits. *FASEB J* 2011; 25(2): 544-60.
19. Tian H, Lu Y, Sherwood AM et al. Resolvins E1 and D1 in choroid-retinal endothelial cells and leukocytes: biosynthesis and mechanisms of anti-inflammatory actions. *Invest Ophthalmol Vis Sci* 2009; 50(8): 3613-20.
20. Kasuga K, Yang R, Porter TF et al. Rapid appearance of resolvin precursors in inflammatory exudates: novel mechanisms in resolution. *J Immunol* 2008; 181(12): 8677-87.

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

21. Krishnamoorthy S, Recchiuti A, Chiang N et al. Resolvin D1 binds human phagocytes with evidence for proresolving receptors. *Proc Natl Acad Sci U S A* 2010; 107(4): 1660-5.
22. Richter AG, McKeown S, Rathinam S et al. Soluble endostatin is a novel inhibitor of epithelial repair in idiopathic pulmonary fibrosis. *Thorax* 2009; 64(2): 156-61.
23. Perkins, G.D., F. Gao, and D.R. Thickett et al. In vivo and in vitro effects of salbutamol on alveolar epithelial repair in acute lung injury. *Thorax* 2008; 63(3): 215-20.
24. Perkins GD, McAuley DF, Thickett DR et al. The beta-agonist lung injury trial (BALTI): a randomized placebo-controlled clinical trial. *Am J Respir Crit Care Med* 2006; 173(3): 281-7.
25. Geiser T, Atabai K, Jarreau PH et al. Pulmonary edema fluid from patients with acute lung injury augments in vitro alveolar epithelial repair by an IL-1beta-dependent mechanism. *Am J Respir Crit Care Med* 2001; 163(6): 1384-8.
26. Zolnai A, Tóth EB, Wilson RA et al. Comparison of 3H-thymidine incorporation and CellTiter 96 aqueous colorimetric assays in cell proliferation of bovine mononuclear cells. *Acta Vet Hung* 1998; 46(2): 191-7.
27. Livak, K.J. and T.D. Schmittgen. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{(-Delta Delta C(T))} Method. *Methods* 2001; 25(4): 402-8.
28. King, L.S., S. Nielsen, and P. Agre. Respiratory aquaporins in lung inflammation: the night is young. *Am J Respir Cell Mol Biol* 2000; 22(1): 8-10.

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

29. Song Y, Fukuda N, Bai C et al. Role of aquaporins in alveolar fluid clearance in neonatal and adult lung, and in oedema formation following acute lung injury: studies in transgenic aquaporin null mice. *J Physiol* 2000; 525 (3): 771-9.
30. Gropper, M.A. and J. Wiener-Kronish. The epithelium in acute lung injury/acute respiratory distress syndrome. *Curr Opin Crit Care* 2008; 14(1): 11-5.
31. Towne JE, Harrod KS, Krane CM et al. Decreased expression of aquaporin (AQP)1 and AQP5 in mouse lung after acute viral infection. *Am J Respir Cell Mol Biol* 2000; 22(1): 34-44.
32. Zhou W, Mo X, Cui W et al. Nrf2 inhibits epithelial-mesenchymal transition by suppressing snail expression during pulmonary fibrosis. *Sci Rep* 2016; 6: 38646.
33. Marshall RP, Bellingan G, Webb S et al. Fibroproliferation occurs early in the acute respiratory distress syndrome and impacts on outcome. *Am J Respir Crit Care Med* 2000; 162(5): 1783-8.
34. Matthay, M.A., H.G. Folkesson, and C. Clerici. Lung epithelial fluid transport and the resolution of pulmonary edema. *Physiol Rev* 2002; 82(3): 569-600.
35. Zoz, D.F., W.E. Lawson, and T.S. Blackwell. Idiopathic pulmonary fibrosis: a disorder of epithelial cell dysfunction. *Am J Med Sci* 2011; 341(6): 435-8.
36. Wang Q, Zheng X, Cheng Y et al. Resolvin D1 stimulates alveolar fluid clearance through alveolar epithelial sodium channel, Na,K-ATPase via ALX/cAMP/PI3K pathway in lipopolysaccharide-induced acute lung injury. *J Immunol* 2014; 192(8): 3765-77.

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

37. Eickmeier O, Seki H, Haworth O et al. Aspirin-triggered resolvin D1 reduces mucosal inflammation and promotes resolution in a murine model of acute lung injury. *Mucosal Immunol* 2013; 6(2): 256-66.
38. Simoes DC, Xanthou G, Petrochilou K et al. Osteopontin deficiency protects against airway remodeling and hyperresponsiveness in chronic asthma. *Am J Respir Crit Care Med* 2009;179:894–902.
39. Jung TW, Hwang HJ, Hong HC et al. Resolvin D1 reduces ER stress-induced apoptosis and triglyceride accumulation through JNK pathway in HepG2 cells. *Mol Cell Endocrinol* 2014; 391(1-2): 30-40.
40. Kasper, M. and G. Haroske. Alterations in the alveolar epithelium after injury leading to pulmonary fibrosis. *Histol Histopathol* 1996; 11(2): 463-83.
41. Noguchi, S., Y. Yamauchi, and H. Takizawa. Novel therapeutic strategies for fibrotic lung disease: a review with a focus on epithelial-mesenchymal transition. *Recent Pat Inflamm Allergy Drug Discov.* 8(1): p. 9-18.
42. Bartis D, Mise N, Mahida RY et al. Epithelial-mesenchymal transition in lung development and disease: does it exist and is it important? *Thorax* 2014; 69(8): 760-5.
43. Lee HJ, Park MK, Lee EJ et al. Resolvin D1 inhibits TGF-beta1-induced epithelial mesenchymal transition of A549 lung cancer cells via lipoxin A4 receptor/formyl peptide receptor 2 and GPR32. *Int J Biochem Cell Biol* 2013; 45(12): 2801-7.
44. Yatomi M, Hisada T, Ishizuka T et al. 17(R)-resolvin D1 ameliorates bleomycin-induced pulmonary fibrosis in mice. *Physiol Rep* 2015; 3(12): 12628.

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

45. Sun YB, Qu X, Li X et al. Endothelial dysfunction exacerbates renal interstitial fibrosis through enhancing fibroblast Smad3 linker phosphorylation in the mouse obstructed kidney. PLoS One 2013; 8(12): e84063.
46. Qu X, Zhang X, Yao J et al. Resolvins E1 and D1 inhibit interstitial fibrosis in the obstructed kidney via inhibition of local fibroblast proliferation. J Pathol 2012; 228(4): 506-19.
47. Laurent GJ, McAnulty RJ, Hill Met al. Escape from the matrix: multiple mechanisms for fibroblast activation in pulmonary fibrosis. Proc Am Thorac Soc 2008; 5(3): 311-5.
48. Caraci F, Gili E, Calafiore M et al. TGF-beta1 targets the GSK-3beta/beta-catenin pathway via ERK activation in the transition of human lung fibroblasts into myofibroblasts. Pharmacol Res 2008; 57(4): 274-82.

Figure legends

Figure 1 ResolvinD1 stimulates ATII cells wound repair and proliferation in vitro. **A:** RvD1 at different concentrations was added to monolayers of ATII cells physically wounded with a 1-mL pipette tip. To allow for variability between cell batches, data are expressed as the mean (SE) percentage of the baseline wound size for each separate set of experiments for each culture condition. **B:** RvD1 stimulated the proliferation of primary human Alveolar Type II

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

cell (ATII cell). Values of >1 fold of control reflect increased proliferation. N=6 for each culture condition, repeated using cells from 3 donors.

Figure 2 RvD1 promotes ATII cells proliferation through activation of ALX receptor and the PI3K/AKT signaling pathway.

A: 100 nM RvD1 promoted proliferation of ATII cells. Pretreatment with 10 μ M LY294002, a phosphatidylinositol 3'-kinase/Akt inhibitor inhibited the effects of resolvinD1 on ATII cells proliferation of suggesting that the pro-proliferation effects of resolvinD1 are PI3-kinase dependent. BOC-2, the ALX receptor antagonist, was re-incubated with primary human Alveolar Type II cells at 10 μ M for 1 hour before resolvinD1 treatment of AT II cells. BOC-2 treatment inhibited the effects of resolvinD1 on the proliferation of primary human Alveolar Type II cells (ATII cells) suggesting that the promoting proliferation effects of resolvinD1 are ALX receptor dependent. **B:** To investigate whether RvD1 can activate AKT phosphorylation in ATII cells, ATII cells were stimulated with different concentrations of RvD1 (10, 25, and 100 nmol/ml) for 24h. We found that RvD1 activated AKT phosphorylation in a dose-dependent manner.

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

Figure 3 Effect of RvD1 upon effects of soluble Fas ligand and

TNF-alpha on apoptosis and caspase-8 activation .

A: Flow cytometry analysis of annexin-positive cells 24 hours after treatment with 100 ng/mL sFasL. Co-treatment with resolvinD1 at 100 nM reduced annexin binding. sFasL treatment of ATII cells increased the number of apoptotic cells from $3.29 \pm 0.11\%$ in control cells to $10.34 \pm 0.33\%$ ($p=0.01$). Rescue treatment with RvD1 reduced the number of apoptotic cells to $4.12 \pm 0.52\%$ ($p=0.01$) **B:** RvD1 inhibited sFasL induced caspase-8 activation on ATII cells. Western blots showing caspase-8 protein in ATII cells treated with 100 ng/mL sFasL or/and RvD1 at 100 nM for 24 h. Caspase-8 protein was quantified and analyzed in the indicated groups. ** $p < 0.01$ relative to control group, ## $p < 0.01$ compared to FasL group.

Figure 4 RvD1 up-regulates Aquaporin V and down-regulates

SPC.

To observe the effect of RvD1 on AQP5 and surfactant protein C gene expression, ATII cells were treated by resolvinD1 for 24 hours.

A: Aquaporin5 gene expression: resolvinD1 only (2.42 ± 0.34 fold) relative to control group, $p=0.001$. The effect that can be blocked by BOC-2 (the FPR antagonist) (resolvinD1+BOC-2, mean,

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

0.54±0.11 fold, p=0.001). **B:** Surfactant protein C gene expression: resolvinD1 only (0.54±0.05 fold) relative to control group, p=0.001. The effect that can be blocked by BOC-2 (the FPR antagonist) (resolvinD1+BOC-2, mean, 0.82±0.03 fold, p=0.01).

Figure 5 ResolvinD₁ inhibits TGF-β induced EMT in Primary human alveolar type II cells

A: TGF-β treated ATII cells showed a mesenchymal morphology (fibroblast-like), and RvD1 restored the epithelial morphology of the cells to a certain extent. **B:** ATII cells were pre-treated with RvD1 for 2 h. The cells were then cultured with TGF-β (10 ng/ml) for 48h. β-Actin was used here as an internal control. EMT was induced with TGF-β treatment. TGF-β treatment induced the expression of mRNA of mesenchymal markers including N-cadherin, vimentin, type I collagen, S100A4, and α-SMA, and reduced the expression of epithelial markers such as E-cadherin. RvD1 blocked the expression of mRNA of mesenchymal markers including N-cadherin, vimentin, type I collagen, S100A4, and α-SMA, while RvD1 restored the expression of mRNA of E-cadherin, **p<0.01 relative to control group respectively, ##p<0.01 compared to TGF-β group respectively. **C:** The effects of RvD1 on the TGF-β-treated E-cadherin, α-SMA, N-cadherin of ATII cells were confirmed by

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

Western blot. ** $p < 0.01$ relative to control group respectively, ## $p < 0.01$ compared to TGF- β + RvD1 group respectively.

Figure 6 RvD1 inhibits TGF- β induced EMT in Primary human alveolar type II cells via ALX/FPR2 receptor.

To elucidate the mechanism involved in the effects of RvD1 on EMT, pre-treatment of cells with Boc-2 (the ALX receptor antagonist), inhibited the effects of RvD1 on EMT of ATII cells. TGF- β -induced N-cadherin expression was suppressed by RvD1, and pre-treatment of cells with Boc-2 incapacitated the effects of RvD1 on N-cadherin expression. Reduced expression of E-cadherin in TGF- β -treated ATII cells was restored by RvD1, but pre-treatment of cells with Boc-2 abolished the effects of RvD1. ** $p < 0.01$ relative to control group respectively, ## $p < 0.01$ compared to TGF- β +RvD1 group respectively, && $p < 0.01$ compared to TGF- β +RvD1 group respectively.

Figure 7 Effect of RvD1 on primary HLF proliferation in response to TGF- β

Cell proliferation studies confirmed that resolvin D1 inhibited proliferation of primary HLF induced by TGF- β . Cultured and serum-deprived cells were treated with 10 ng/ml TGF- β for 24 h with or without pre-incubation with LY294002 (10 μ M) for 1h, BOC-2 (10 μ M)

ResolvinD₁ stimulates epithelial wound repair in HAECs

S Zheng et al

for 1 h. Data are mean \pm SEM of three independent experiments. $P < 0.05$, compared with no treatment group; $**P < 0.01$, compared with TGF- β only group.

Figure 8 Effect of RvD1 on primary human lung fibroblast (HLF) proliferation in response to acute respiratory distress syndrome (ARDS) bronchoalveolar lavage fluid (BALF).

Bronchoalveolar lavage fluid (BALF) from patients with ARDS stimulated proliferation of primary HLF. Resolvin D1 inhibited the proliferation of primary HLF induced bronchoalveolar lavage fluid (BALF) from patients with ARDS. Data are mean \pm SEM of three independent experiments.

Figure 9 The key role for RvD1 during primary human ATII cells and HLF in vitro

Table 1 Summary of the different types gene expression in response to treatment of different stimulations