

UNIVERSITY OF BIRMINGHAM

Research at Birmingham

Avoidance of Harm From Treatment for ANCA-Associated Vasculitis

King, Catherine; Harper, Lorraine

DOI:

[10.1007/s40674-017-0082-y](https://doi.org/10.1007/s40674-017-0082-y)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

King, C & Harper, L 2017, 'Avoidance of Harm From Treatment for ANCA-Associated Vasculitis', *Current Treatment Options in Rheumatology*, vol. 3, no. 4, pp. 230-243. <https://doi.org/10.1007/s40674-017-0082-y>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 09/01/2018

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Avoidance of Harm From Treatment for ANCA-Associated Vasculitis

Catherine King, MRCP, MBChB
Lorraine Harper, PhD, MRCP, MBChB*

Address

*Centre for Translational Inflammation Research University of Birmingham Research Laboratories, Queen Elizabeth Hospital, Mindelsohn Way, Edgbaston, Birmingham, B15 2WB, UK
Email: L.harper@bham.ac.uk

Published online: 13 November 2017

© The Author(s) 2017. This article is an open access publication

This article is part of the Topical Collection on *Vasculitis*

Keywords Anti-neutrophil cytoplasm antibody-associated vasculitides (AAV) · Adverse events · Cyclophosphamide · Rituximab

Opinion statement

Purpose of review With established immunosuppressant treatment regimens for anti-neutrophil cytoplasm antibody-associated vasculitides (AAV), prognosis has significantly improved. The mainstay of treatment still comprises high-dose corticosteroids and cyclophosphamide for severe forms, although rituximab is being increasingly utilised instead of cyclophosphamide as induction therapy. AAV patients experience an excess of infections, malignancies and cardiovascular events as compared to the general population, which is a combination of the systemic inflammatory process associated with vasculitis and the adverse events from treatment.

Recent findings Successful therapy should focus on suppressing disease activity and minimising treatment-related toxicity. Infection is the largest contributor to morbidity and mortality in the first year of treatment, and annual pneumococcal and influenza vaccinations, *Pneumocystis jiroveci* prophylaxis and tuberculosis (TB) and Hepatitis B virus screening are advised. Patients on high-dose corticosteroid treatment should have regular blood sugar monitoring, a FRAX assessment with vitamin D and calcium supplementation, consideration of prophylaxis for gastric ulcers and a cardiovascular risk assessment. Patients who are treated with cyclophosphamide could also receive MESNA to reduce the risk of chemical cystitis. Cyclophosphamide, methotrexate and azathioprine all require blood monitoring schedules due to the risk of bone marrow suppression, liver and renal toxicity. Hypogammaglobulinaemia is a recognised risk of rituximab treatment. Patients of reproductive age need to be counselled on the infertility risks with cyclophosphamide and the teratogenicity associated with it, methotrexate and mycophenolate mofetil.

Summary A greater focus on identifying clinical and biological markers that will help identify those patients at greatest risk of relapse, e.g. GPA and PR3-ANCA specificity, from

those patients at greatest risk of toxicity, e.g. increasing age and declining GFR, is required to allow treatment to be tailored accordingly.

Introduction

The introduction of effective immunosuppressive therapy to treat granulomatosis with polyangiitis (GPA) and microscopic polyangiitis, commonly grouped as anti-neutrophil cytoplasm antibody-associated vasculitis (AAV), has significantly changed prognosis. Untreated, these conditions are almost universally fatal [1]. Following the introduction of cyclophosphamide and corticosteroid-based immunosuppressive regimens, survival rates approach 80% at 5 years [2], with infection, cardiovascular disease and malignancy being the commonest causes of death. Although improved, this remains 2.6 times worse than an age and sex-matched healthy population, with death predicted by age, glomerular filtration rate < 15 ml/min and higher disease severity [2]. The greatest risk to patients with AAV is in the first year of diagnosis and most commonly from the adverse events associated with treatment rather than active vasculitis [3]. A study of patients with GPA using primary care data noted a 9-fold increase in mortality compared with a control population in the first year following diagnosis [4].

The improvement in survival has changed our perception of vasculitides from acutely fatal diseases to chronic conditions with a relapsing remitting nature and high morbidity due to disease damage and the side effects associated with treatment. There is a complex relationship between the severity of vasculitis and the risk of developing adverse events, with the estimated glomerular filtration rate (eGFR) being consistently found to be the greatest overall predictor. This is probably due to reduced excretion of active drug metabolites, as well as the general immunosuppressive effect of kidney failure. Damage due to disease and treatment is common and increases with time. In a long-term follow-up study of patients recruited to four European AAV trials, over 30% of patients reported greater than 5 damage items at long-term follow-up using the Vasculitis Damage Index (VDI) [5]. Patients with damage are also at increased risk of death [6].

The insight we now have into morbidity and mortality in AAV highlights the need for a change in our treatment approach to maximise efficacy and minimise toxicity.

Treatment

Treatment overview

The treatment of AAV is usually divided into two phases comprised of induction of remission and maintenance of remission. The latest 2015 recommendations from European league against rheumatism, European Renal Association and European Vasculitis Study (EULAR/ERA-EDTA) state that with organ or life-threatening disease, treatment typically involves pulsed cyclophosphamide or rituximab with high-dose corticosteroids (e.g. 1 mg/kg/day oral prednisolone, or intravenous methylprednisolone 0.5–1.0 g/day for 3 days) [7]. For those patients with pulmonary haemorrhage or rapidly progressive renal failure, plasma exchange should be considered. Patients with non-organ threatening disease may be successfully treated with a less toxic induction regimen of either methotrexate or mycophenolate mofetil alongside corticosteroids [7].

Maintenance therapy then consists of azathioprine, methotrexate or rituximab with corticosteroid tapering. There is great variation in the approach to corticosteroid tapering, but a target of between 7.5 and 10 mg of prednisolone (or equivalent) after 3 months of treatment is considered appropriate [7]. The duration of maintenance therapy is less clear.

Treatment must take into account quality of life (QOL) for patients with AAV. QOL among AAV patients is frequently poor and determined by multiple clinical and bio-psychosocial factors, of which fatigue appears to be of principal importance [8]. Whilst optimising disease control can improve QOL, non-pharmacological interventions targeting bio-psychosocial determinants may be more beneficial. AAV therapy must focus on person-centred outcomes such as QOL and ascertaining what a patient's priorities are for the management of their chronic disease.

Infection

Infection is a considerable cause of morbidity and mortality associated with AAV, and the immunosuppressive agents used to treat vasculitis are an undoubtable contributor to this. Overall, approximately 30% of patients will suffer from a serious infection requiring hospital admission, with respiratory infections being most common [9, 10]. The greatest risk to patients is in the first year, reflecting the intensity of immunosuppression [3, 11]. Higher doses of corticosteroids are also associated with infectious risk [9, 12, 13]. During infection, consideration should be given to pausing cytotoxic therapies especially if severe.

Advancing age and declining GFR are associated with increased risk of infection and mortality [3, 14, 15]. Immune dysfunction induced by AAV therapy predisposes to infection through neutropenia, reduced B and T cell numbers, aberrant cell function and reduced immunoglobulin levels [16]. Close monitoring of the peripheral blood count is recommended during AAV therapy as all cytotoxic therapies are associated with bone marrow suppression [3] (Table 1). Granulocyte colony-stimulating factor (G-CSF), commonly used in oncology for patients with neutropenia following chemotherapy, has been shown to be effective at reducing the duration of neutropenia in AAV and may be used safely, if necessary, without increasing risk of relapse [17].

Bacterial infections are common in AAV patients, accounting for up to 62% of major infections, in particular those causing pneumonia. Opportunistic *Pneumocystis jiroveci* pneumonia (PJP) has an incidence of 0.85–12% with increased mortality in AAV [18–20]. Although PJP is generally associated with a defect in T cell-mediated immunity, B cells have also been shown to play an important role in the defence against pneumocystis [21]. Prophylaxis with cotrimoxazole (800/160 mg on alternate days or 400/80 mg daily) is recommended for all patients undergoing cyclophosphamide or rituximab treatment [7].

GPA patients have higher *Staphylococcus aureus* colonisation of the upper respiratory tract which has been associated with a higher rate of relapse [22–24]. In patients with nasal disease, treatment with topical antibiotics such as mupirocin may be considered in the presence of chronic carriage of nasal *Staphylococcus aureus* [7, 25].

Vaccination should be used to reduce the risk of respiratory infections. The EULAR guidelines conclude that pneumococcal and seasonal influenza vaccinations are safe and effective and recommended for all patients with autoimmune inflammatory disease [11, 26]. Not all patients with AAV will have an adequate response to vaccination due to immune dysfunction [3]. Poor immunologic function, including low IgG, B cell counts and CD4 T cell counts, predict vaccine response and mortality [16]. Current EULAR recommendations are to avoid vaccination during B cell depletion therapy, e.g. with rituximab, as

Table 1. Treatment-specific blood monitoring schedules for ANCA vasculitis therapy (compiled from [29] and [55])

Drug	Laboratory monitoring	Frequency
Methotrexate	FBC LFTs, albumin Creatinine and eGFR	<ul style="list-style-type: none"> • Pre-treatment, baseline levels • Every 2 weeks until on stable dose for 6 weeks • Then once on a stable dose, monthly for 3 months • After this at least every 12 weeks • Dose increases should be monitored with FBC every 2 weeks until on stable dose for 6 weeks, then return to normal schedule • More frequent monitoring is required for patients at high risk for toxicity, i.e. chronic kidney disease and other risk factors for liver disease
Azathioprine	FBC LFTs, albumin Creatinine and eGFR	*as above
Pulsed cyclophosphamide	FBC LFTs, albumin Creatinine and eGFR	<ul style="list-style-type: none"> • Pre-treatment baseline levels • 7–10 days after first pulse • Then prior to giving next pulse • Unless dose changed, only require FBC monitoring prior to subsequent pulses • If WBC prior to pulse is $< 4 \times 10^9/L$ and neutrophils $< 2 \times 10^9/L$, postpone treatment until above these counts and reduce next dose by 25%

this has been consistently shown to impair humoral responses to influenza, pneumococcal and tetanus toxoid immunisations in patients with rheumatoid arthritis [27, 28]. Live attenuated vaccinations should be avoided in immunocompromised patients as live vaccines can, in some situations, cause severe or fatal infections due to extensive replication of the vaccine strain.

Mycobacterium tuberculosis infection has only rarely been reported in AAV patients, but can be fatal; the majority of cases show reactivation of latent TB. Current British Society of Rheumatology (BSR) and British Health Professionals in Rheumatology (BHPR) guidance recommends that all AAV patients require a thorough history, clinical examination and chest X-ray to screen for TB before commencing immunosuppressive therapy [25, 29]. TB prophylaxis is then indicated if latent TB is detected [25].

Viral infections are common, responsible for 35.8% of all major infectious episodes seen in one long-term analysis [9]. The occurrence of herpes zoster virus (HZV), shingles, occurs more often in AAV patients than a healthy population, with a frequency of 4.5 episodes of HZV/100 patient years compared to 0.40/100 patient years in the general population [12, 30]. Prophylaxis against varicella zoster virus (VZV) is not routinely recommended due to the non-life-threatening nature of VZV alongside the high risk of drug interactions and adverse events especially with renal dysfunction [25]. The varicella vaccine is a live attenuated vaccine and guidance for its use in immunosuppressed patients is mixed; if possible the vaccine should be administered prior to starting immunosuppression. Recently, a vaccine using recombinant VZV glycoprotein E has been shown to have benefit in reducing the risk of herpes zoster and post-herpetic

neuralgia in those over 70 and would be suitable for use in immunosuppressed populations when available [31].

Latent infection with cytomegalovirus (CMV) is less frequent in AAV patients than in solid-organ transplantation, accounting for only 7.5% of major infectious episodes seen in long-term follow-up studies [9]. Whilst CMV polymerase chain reaction (PCR) monitoring and prophylaxis with valganciclovir are established in solid-organ transplantation follow-up, these are not routinely recommended in AAV [25].

Hepatitis B virus (HBV) reactivation occurs early in patients receiving rituximab or cyclophosphamide therapy for immune-mediated inflammatory disorders and is an important cause of hepatic failure [32]. All patients receiving significant immunosuppression should be screened for HBV infection and to perform HBV vaccination in patients with a negative result. Guidelines recommend that hepatitis B surface antigen (HBsAg)-positive patients receive preemptive antiviral therapy before starting therapy and that hepatitis B core antibody (HBcAb)-positive patients should be considered for preemptive antiviral therapy if rituximab or cyclophosphamide are given [32, 33].

Invasive fungal infections are rare in AAV patients, so systemic anti-fungal prophylaxis is not indicated. A Cochrane review noted that oral nystatin was no better than placebo at preventing candida infections in immunosuppressed patients. Consideration should be given to using fluconazole if prophylaxis or treatment is required [34].

Progressive multifocal leucoencephalopathy (PML) is a rare demyelinating infection of the central nervous system caused by John Cunningham virus which can result in irreversible neurological damage and death. Cases of PML have been reported in patients receiving cyclophosphamide and/or rituximab [35]. Clinicians must consider PML in their differentials for neurological presentations in patients who are heavily immunosuppressed as earlier diagnosis and restorations of immune function with withdrawal of immunosuppression may improve outcomes [36].

Corticosteroids

Other adverse events are treatment specific. The extensive side effect profile associated with corticosteroids is well established. Data collected from four European trials showed that of AAV patients treated with corticosteroids within the first year, 8.2% developed new onset diabetes, 50% of which occurred within 1.7 months [3]. 29% of patients gained over 10 kg weight, 2.6% developed peptic ulceration, 2.5% had fractures, 2% developed cataracts and 0.4% developed avascular necrosis [3]. The adverse effects from corticosteroids are even more pronounced with increasing length of exposure. At long-term follow-up (median 5 years) of 270 patients, 41% had hypertension, 38% osteoporosis, 28% diabetes mellitus and 25% had developed cataracts [5].

Corticosteroid administration modulates carbohydrate metabolism particularly in adipose tissue as well as inducing insulin resistance, with impaired glucose uptake stimulating gluconeogenesis [37]. Risk factors for diabetes include previous corticosteroid-induced hyperglycaemia, pre-diabetes, obesity or a family history of diabetes [37]. Capillary blood glucose should be closely monitored whilst patients are on corticosteroids; if persistently over 12 mmol/L, treatment should be commenced.

Osteoporosis is a concern in patients with AAV. Corticosteroids promote osteoporosis through direct cellular effects stimulating a positive effect on osteoclasts, leading to increased bone resorption, and a negative effect on osteoblasts and osteocytes, causing reduced bone formation and reduced repair of micro-damaged bone. AAV alone is also associated with reduced BMD, although the mechanisms for this are not yet fully understood. In a cross-sectional study involving 99 patients with AAV treated with corticosteroids and cyclophosphamide, 78% of patients were found to have abnormal dual-energy X-ray absorptiometry (DEXA) measurements in one of three sites (the lumbar spine, proximal femur or radius); 21% of these patients had osteoporosis and 57% osteopenia [38]. Cumulative dosage of corticosteroid therapy was inversely related to Z-score; significant correlations were also seen between bone mineral density (BMD) and renal function [38]. Current UK guidelines developed by the National Osteoporosis Guideline Group (NOGG) advise assessment with FRAX® tool for all patients identified to be at risk of osteoporosis, to help guide further investigations and treatment.

Both the American College of Rheumatology and NOGG guidance suggests lifestyle advice should be given to all patients on commencement of long-term corticosteroid therapy including smoking cessation, alcohol limitation and regular exercise [39, 40]. Oral calcium and vitamin D supplementation is recommended for all patients undergoing long-term corticosteroid therapy [29, 41]. A Cochrane meta-analysis of 274 patients on corticosteroids taking calcium and vitamin D compared to placebo showed a clinically and statistically significant increase in bone mineral density of 2.6% at 2 years [41]. Bone-protective treatment with oral bisphosphonates should be started at the onset of corticosteroid therapy in individuals at moderate or high risk of fracture [40]. Bisphosphonate therapy is contraindicated in patients with significant renal impairment (GFR < 30 ml/min). Special consideration should be given to women of child-bearing potential due to the risk of foetal harm [40].

There is evidence that there is an increased risk of cardiovascular disease (CVD) in AAV patients [42]. Outcomes from the EUVAS trials showed that 14% of AAV patients have a cardiovascular event within 5 years of diagnosis; risk factors included MPO-ANCA positivity, increasing age and diastolic hypertension [43]. Whilst the mechanisms behind this association are not fully understood, it is believed that inflammation accelerates atherosclerosis through endothelial dysfunction [44]. Corticosteroid use may also increase cardiovascular risk [45]. Corticosteroids are known to cause multiple metabolic side effects including hypertension, hyperlipidaemia, weight gain and diabetes which are all significant cardiovascular risk factors. Due to this established increased risk of CVD in AAV patients, EULAR guidelines advise annual review of traditional Framingham risk factors [7].

It is also advisable to consider prophylactic gastric protection for AAV patients receiving a prolonged course of high-dose corticosteroids due to the risk of mucosal irritation.

Whilst it is appreciated that steroids are associated with significant toxicity, there is limited data available on the most appropriate duration of treatment or dose. Early withdrawal has been associated with increased disease relapse [46]. Data from EUVAS trials demonstrated that approximately half (47.8%) of the patients were on glucocorticoids beyond 2 years, with a mean duration of

corticosteroid use of 40.4 months [5]. New studies are addressing the need and dose of corticosteroids required to treat AAV. PEXIVAS (plasma exchange and glucocorticoid dosing in the treatment of ANCA-associated vasculitis) addresses initial corticosteroid dosage (<http://clinicaltrials.gov/ct2/show/NCT00987389>). The CLEAR (C5aR inhibitor on leukocytes exploratory ANCA-associated renal vasculitis) study has just been published investigating a new C5aR inhibitor avacopan as a replacement for high-dose glucocorticoids in treating newly diagnosed or relapsing AAV [47].

Cyclophosphamide

Cyclophosphamide is an alkylating agent, which is metabolised by cytochrome P450 to acrolein and phosphoramidate mustard. Phosphoramidate mustard forms covalent bonds with DNA inducing lymphopenia, particularly of B cells and suppressing humoral responses. Cyclophosphamide metabolites are primarily excreted in the urine unchanged, resulting in increased systemic drug exposure in those with renal dysfunction. Dosing should be reduced for both age and renal function to reduce the risk of bone marrow suppression (Table 2). These metabolites are toxic to the bladder and can cause haemorrhagic cystitis and bladder cancer [48, 49]. Consideration should be given to the use of oral or intravenous 2-mercaptoethanesulfonate sodium (MESNA) for patients receiving intravenous cyclophosphamide alongside adequate hydration [7, 29]. MESNA binds to acrolein, a toxic metabolite of cyclophosphamide, rendering it non-toxic [49]. Current guidelines recommend routine surveillance with urinalysis; this should be continued indefinitely following initiation of any cyclophosphamide therapy. If non-glomerular haematuria is confirmed on urine microscopy, a urology referral should be sought [7].

The cumulative dose of cyclophosphamide, with a threshold of 36 g, increases the risk of malignancy [50]. The overall incidence of cancer in AAV patients was previously reported as being up to two times higher than that of general population, with bladder cancer, non-melanoma skin cancer (NMSC), leukaemia and lymphoma being the most common malignancies [12, 14]. Recent studies suggest that with reduction in cyclophosphamide dosing, only NMSC is increased [51]. AAV patients should be educated on the risk of skin malignancy and should be strongly encouraged to attend the national cancer screening programs.

Table 2. Dose modification of pulsed cyclophosphamide as used in the CYCLOPS trial and stated in the EULAR ANCA vasculitis guidelines [79]

Pulsed dose cyclophosphamide reductions for age and renal function		
Age (years)	Creatinine ($\mu\text{mol/L}$)	
	< 300	300–500
< 60	15 mg/kg/pulse	12.5 mg/kg/pulse
60–70	12 mg/kg/pulse	10 mg/kg/pulse
> 70	10 mg/kg/pulse	7.5 mg/kg/pulse

Methotrexate

Methotrexate is an anti-folate drug with anti-proliferative and anti-inflammatory effects, due to the inhibition of several key enzymes involved in folate, methionine, adenosine and de novo nucleotide synthesis pathways [52]. Side effects are common with 10–37% of patients discontinuing treatment due to toxicity. GI toxicity is frequent although usually mild, associated with diarrhoea, nausea and vomiting; 30% of patients will develop abnormal LFTs although this may be transient. Regular monitoring of liver function tests is recommended especially in those with other risk factors for liver disease [53] (Table 1). Patients are advised to limit alcohol intake whilst taking methotrexate to reduce the risk of liver toxicity.

Interstitial pneumonitis, although rare, remains a concern with methotrexate use; a low threshold for investigations is required for patients on methotrexate presenting with new or increasing dyspnoea or a dry cough. If detected early and methotrexate is discontinued, most patients recover without progression to lung fibrosis [53]. Renal failure from low-dose methotrexate is rare; however, as it is primarily excreted by the kidneys, it is contraindicated in severe renal impairment and should be used with caution if creatinine > 150 µmol/L.

All patients should be co-prescribed folic acid supplementation at a minimal dose of 5 mg once weekly, as adverse effects from methotrexate are thought to be mediated by folate antagonism [54, 55].

Azathioprine

Azathioprine is a prodrug of mercaptopurine, which is subsequently metabolised by several pathways, one of which involves the enzyme thiopurine methyltransferase (TPMT). Azathioprine may cause a toxic reaction, which results in hepatotoxicity and myelosuppression and is related to variation in TPMT levels. Up to 14% of the population is known to have reduced levels of this enzyme as a result of genetic mutations [56]. The positive predictive value of measuring TPMT levels is widely debated, but British Society of Rheumatology guidelines state that patients should have baseline TPMT status assessed [55]. Allopurinol, a xanthine oxidase inhibitor, an enzyme involved in metabolising 6-mercaptopurine, should not be used with azathioprine due to the high risk of bone marrow suppression.

Rituximab

The safety of rituximab has become a topic of great interest with its increasing use in both remission induction and maintenance therapy in AAV. Rituximab was first used successfully to treat rheumatological disease without significant rates of infection at long-term follow-up, so its promise in reducing the toxicity from cumulative doses of cyclophosphamide and ongoing maintenance therapy was particularly appealing [57, 58]. Whilst its efficacy has been demonstrated, its side effect profile remains in question as the trials demonstrating efficacy failed to show any difference in adverse events, such as infection, in patients with AAV [59, 60].

The advantages of rituximab in reducing infections rates as demonstrated in patients with rheumatoid arthritis, may not have materialised within these trials as the follow-up periods may not have been long enough to have incorporated the benefit from reduced cumulative cyclophosphamide exposure and prolonged maintenance therapy [60].

Hypoimmunoglobulinaemia is associated with the use of cyclophosphamide and rituximab [61]. A recent retrospective study of 243 patients with multi-system disease treated with rituximab demonstrated that 56% had IgG hypoimmunoglobulinaemia during follow-up; although in 50%, this was transient. Only 4.2% of these patients required serum IgG replacement for infection [62]. EULAR guidelines state that surveying patients with AAV is warranted post cyclophosphamide and rituximab treatment for serum immunoglobulin concentrations and persisting hypoimmunoglobulinaemia [7].

Late-onset neutropenia (LON), defined as an absolute neutrophil count (ANC) $< 1.5 \times 10^9/L$ at least 4 weeks after the last infusion, has been observed in patients with rheumatological disease, including AAV treated with rituximab [63]. A recent study involving 59 patients with AAV treated with rituximab demonstrates that 7 patients (11.9%) developed LON after a median time of 86 days (range 56–168 days) since their latest rituximab treatment [64]. LON is idiosyncratic in its development; it may occur after both single and repeated rituximab courses and does not recur after subsequent doses of rituximab. No predisposing factors for LON have been identified.

Fertility and pregnancy

Disease activity and its therapy are threats to the fertility of patients with vasculitis. The cumulative dose of cyclophosphamide and increasing age at initiation of treatment both increase the risk of primary ovarian failure [65, 66]. Data from a lupus trial suggests that a total cyclophosphamide exposure of 14–20 g causes infertility in $> 50\%$ of women aged over 32 years, with a lower risk in younger women [67]. The long-term effects of rituximab on fertility and teratogenicity during pregnancy have not been studied, but no concerns have been reported suggesting rituximab should be used for young patients with AAV wishing to preserve fertility [7].

Cyclophosphamide, methotrexate and mycophenolate mofetil are teratogenic. Patients taking these agents should be counselled on this and advised to use effective contraception for the duration of therapy and at least 3 months thereafter. Azathioprine is considered low risk in pregnancy without a significant increase in major congenital malformations [68, 69]. There is less guidance available on the risk of paternal immunosuppressant exposure. The British Association of Dermatologists' guidelines for the use of azathioprine state that studies do confirm male patients receiving azathioprine father healthy children and standard dose azathioprine does not appear to affect fertility [70].

There is still little research on pregnancy outcomes in patients with vasculitis. Two recent reports suggest that relapse during pregnancy is uncommon and that outcomes when disease is controlled prior to conception are good [71, 72].

Thromboembolic disease

Although microthrombosis is a component of vasculitic pathology, the frequency of deep venous thrombosis and pulmonary embolism is also increased, being highest during periods of disease activity and inflammation. A rate of 15% was reported in a prospective trial; and in retrospective surveys, rates of 1.8%/year for all AAV patients rising to 6.7–9%/year for those with active disease were reported [73]. Autoantibodies to plasminogen and tissue

plasminogen activator occur in the sera of 25 and 14% of AAV patients [74]. Their presence has been associated with more severe renal outcomes and increased risk for thromboembolic events. Anti-coagulants have been used historically in the treatment of renal vasculitis, but thromboprophylactic strategies have not been developed for AAV in the current era. Until further data emerges of the clinical utility of anti-plasminogen autoantibody testing, it is advisable to address conventional risk factors and maintain a high threshold for suspicion of thromboembolism.

Risk of relapse versus prolonged maintenance therapy

Relapse in AAV occurs in 30–50% of patients by 5 years, with an increased risk associated with GPA, PR3-ANCA specificity, ear nose and throat involvement, persisting ANCA positivity after induction therapy, a lower serum creatinine at diagnosis and early withdrawal of corticosteroids or immunosuppressives [75]. However, prolonged duration of maintenance therapy is associated with increased exposure to the toxicities of immunosuppressives and corticosteroids. Current recommendations suggest at least 24 months of induction therapy once remission has been obtained, although, the optimal duration of remission-maintenance therapy remains unknown [7].

The recent REMAIN trial (prolonged REmission-MAINtenance therapy in systemic vasculitis) by EUVAS looked at whether continued azathioprine/prednisolone was more effective in preventing relapse in AAV than their withdrawal at 24 months. It demonstrated that continuation of corticosteroids and azathioprine beyond 2 years reduced the relapse risk and improved renal outcome. Continuation of immunosuppressive therapy was associated with a trend towards more frequent adverse events such as malignancy, infections and cardiovascular complications ($p = 0.07$) [76]. However, another study investigating the use of prolonged azathioprine to prevent disease relapse failed to show benefit [77].

Further research is required to identify clinical or biological markers that will help to identify high-risk patients who require prolonged therapy, until then, the treating clinician must weigh up the individual risk of relapse compared with the risk of treatment toxicity.

Conclusion

Over the last 20 years, careful use and monitoring of cyclophosphamide and corticosteroid-based treatment regimens have improved outcomes [78]. This may be improved further with use of rituximab as an alternative to cyclophosphamide and new therapies reducing the need for corticosteroids.

It is the responsibility of the treating clinician to assess every patient as an individual, considering the risks associated with AAV therapy before commencement of induction therapy, using the risk factors identified for adverse events and mortality including increasing age and declining kidney function [3]. Individual patient risk of relapse must be balanced against risk from treatment toxicity when deciding duration of maintenance therapy.

The impact of this chronic relapsing disease and the burden of therapy on patients' physical and emotional wellbeing are significant, with fatigue being a considerable concern. Many patients experience reduced quality of life despite

successful therapy. We must identify patient priorities at the start of treatment with regular reassessment, alongside patient education on both the nature of their disease and the considerable risks from disease and therapy. This will aid clinicians and patients with shared decision-making to achieve the best outcomes for patients with AAV.

Until the development of treatments without the significant side effects that still remain from current regimens, we need to focus on prophylaxis to prevent the known toxicity from AAV therapies.

Compliance with Ethical Standards

Conflict of Interest

Catherine King declares that she has no conflict of interest.

Lorraine Harper declares that she has no conflict of interest.

Human and Animal Rights and Informed Consent

This article does not contain any studies with human or animal subjects performed by any of the authors.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References and Recommended Reading

- Walton EW. Giant-cell granuloma of the respiratory tract (Wegener's granulomatosis). *Br Med J*. 1958;2(5091):265–70.
- Flossmann O, Berden A, de Groot K, Hagen C, Harper L, Heijl C, et al. Long-term patient survival in ANCA-associated vasculitis. *Ann Rheum Dis*. 2011;70(3):488–94.
- Little MA, Nightingale P, Verburgh CA, Hauser T, De Groot K, Savage C, et al. Early mortality in systemic vasculitis: relative contribution of adverse events and active vasculitis. *Ann Rheum Dis*. 2010;69(6):1036–43.
- Luqmani R, Suppiah R, Edwards CJ, Phillip R, Maskell J, Culliford D, et al. Mortality in Wegener's granulomatosis: a bimodal pattern. *Rheumatology (Oxford)*. 2011;50(4):697–702.
- Robson J, Doll H, Suppiah R, Flossmann O, Harper L, Hoggund P, et al. Damage in the anca-associated vasculitides: long-term data from the European vasculitis study group (EUVAS) therapeutic trials. *Ann Rheum Dis*. 2015;74(1):177–84.
- Exley AR, Bacon PA, Luqmani RA, Kitas GD, Gordon C, Savage CO, et al. Development and initial validation of the Vasculitis Damage Index for the standardized clinical assessment of damage in the systemic vasculitides. *Arthritis Rheum*. 1997;40(2):371–80.
- Yates M, Watts RA, Bajema IM, Cid MC, Crestani B, Hauser T, et al. EULAR/ERA-EDTA recommendations for the management of ANCA-associated vasculitis. *Ann Rheum Dis*. 2016;75(9):1583–94.
- Basu N, McClean A, Harper L, Amft EN, Dhaun N, Luqmani RA, et al. The characterisation and determinants of quality of life in ANCA associated vasculitis. *Ann Rheum Dis*. 2014;73(1):207–11.
- Charlier C, Henegar C, Launay O, Pagnoux C, Berezne A, Bienvenu B, et al. Risk factors for major infections in Wegener granulomatosis: analysis of 113 patients. *Ann Rheum Dis*. 2009;68(5):658–63.
- Reinhold-Keller E, Beuge N, Latza U, de Groot K, Rudert H, Nolle B, et al. An interdisciplinary approach to the care of patients with Wegener's granulomatosis: long-term outcome in 155 patients. *Arthritis Rheum*. 2000;43(5):1021–32.
- van Assen S, Agmon-Levin N, Elkayam O, Cervera R, Doran MF, Dougados M, et al. EULAR recommendations for vaccination in adult patients with autoimmune inflammatory rheumatic diseases. *Ann Rheum Dis*. 2011;70(3):414–22.

12. Hoffman GS, Leavitt RY, Kerr GS, Fauci AS. The treatment of Wegener's granulomatosis with glucocorticoids and methotrexate. *Arthritis Rheum.* 1992;35(11):1322-9.
13. McGregor JG, Falk RJ. Vasculitis: the elusive optimal induction strategy for vasculitis. *Nat Rev. Nephrol.* 2012;8(4):195-6.
14. Wong L, Harper L, Little MA. Getting the balance right: adverse events of therapy in anti-neutrophil cytoplasm antibody vasculitis. *Nephrol Dial Transplant.* 2015;30(Suppl 1):i164-70.
15. Harper L, Savage CO. ANCA-associated renal vasculitis at the end of the twentieth century—a disease of older patients. *Rheumatology (Oxford).* 2005;44(4):495-501.
16. David Morgan M, Richter A, Al-Ali S, Flint J, Yiannakis C, Drayson M, et al. Association of low B cell count and IgG levels with infection, and poor vaccine response with all-cause mortality in an immunosuppressed vasculitis population. *Arthritis Care Res (Hoboken).* 2016;68(6):853-60.
17. Hellmich B, Schnabel A, Gross WL. Granulocyte colony-stimulating factor treatment for cyclophosphamide-induced severe neutropenia in Wegener's granulomatosis. *Arthritis Rheum.* 1999;42(8):1752-6.
18. Ognibene FP, Shelhamer JH, Hoffman GS, Kerr GS, Reda D, Fauci AS, et al. Pneumocystis carinii pneumonia: a major complication of immunosuppressive therapy in patients with Wegener's granulomatosis. *Am J Respir Crit Care Med.* 1995;151(3 Pt 1):795-9.
19. Falagas ME, Manta KG, Betsi GI, Pappas G. Infection-related morbidity and mortality in patients with connective tissue diseases: a systematic review. *Clin Rheumatol.* 2007;26(5):663-70.
20. Ward MM, Donald F. Pneumocystis carinii pneumonia in patients with connective tissue diseases: the role of hospital experience in diagnosis and mortality. *Arthritis Rheum.* 1999;42(4):780-9.
21. Lund FE, Hollifield M, Schuer K, Lines JL, Randall TD, Garvy BAB. cells are required for generation of protective effector and memory CD4 cells in response to Pneumocystis lung infection. *J Immunol.* 2006;176(10):6147-54.
22. Stegeman CA, Tervaert JW, Sluiter WJ, Manson WL, de Jong PE, Kallenberg CG. Association of chronic nasal carriage of Staphylococcus aureus and higher relapse rates in Wegener granulomatosis. *Ann Intern Med.* 1994;120(1):12-7.
23. Laudien M, Gadola SD, Podschun R, Hedderich J, Paulsen J, Reinhold-Keller E, et al. Nasal carriage of Staphylococcus aureus and endonasal activity in Wegener's granulomatosis as compared to rheumatoid arthritis and chronic rhinosinusitis with nasal polyps. *Clin Exp Rheumatol.* 2010;28(1 Suppl 57):51-5.
24. Salmela A, Rasmussen N, Tervaert JWC, Jayne DRW, Ekstrand A. European Vasculitis Study G. Chronic nasal Staphylococcus aureus carriage identifies a subset of newly diagnosed granulomatosis with polyangiitis patients with high relapse rate. *Rheumatology (Oxford).* 2017;56(6):965-72.
25. Moosig F, Holle JU, Gross WL. Value of anti-infective chemoprophylaxis in primary systemic vasculitis: what is the evidence? *Arthritis Res Ther.* 2009;11(5):253.
26. Holvast A, Stegeman CA, Benne CA, Huckriede A, Wilschut JC, Palache AM, et al. Wegener's granulomatosis patients show an adequate antibody response to influenza vaccination. *Ann Rheum Dis.* 2009;68(6):873-8.
27. van Assen S, Holvast A, Benne CA, Posthumus MD, van Leeuwen MA, Voskuyl AE, et al. Humoral responses after influenza vaccination are severely reduced in patients with rheumatoid arthritis treated with rituximab. *Arthritis Rheum.* 2010;62(1):75-81.
28. Bingham CO 3rd, Looney RJ, Deodhar A, Halsey N, Greenwald M, Coddling C, et al. Immunization responses in rheumatoid arthritis patients treated with rituximab: results from a controlled clinical trial. *Arthritis Rheum.* 2010;62(1):64-74.
29. Lapraik C, Watts R, Bacon P, Carruthers D, Chakravarty K, D'Cruz D, et al. BSR and BHPR guidelines for the management of adults with ANCA associated vasculitis. *Rheumatology (Oxford).* 2007;46(10):1615-6.
30. Wung PK, Holbrook JT, Hoffman GS, Tibbs AK, Specks U, Min YI, et al. Herpes zoster in immunocompromised patients: incidence, timing, and risk factors. *Am J Med.* 2005;118(12):1416.
31. Cunningham AL, Lal H, Kovac M, Chlibek R, Hwang SJ, Diez-Domingo J, et al. Efficacy of the herpes zoster subunit vaccine in adults 70 years of age or older. *N Engl J Med.* 2016;375(11):1019-32.
32. Droz N, Gilardin L, Cacoub P, Berenbaum F, Wendling D, Godeau B, et al. Kinetic profiles and management of hepatitis B virus reactivation in patients with immune-mediated inflammatory diseases. *Arthritis Care Res (Hoboken).* 2013;65(9):1504-14.
33. Sebastiani M, Atzeni F, Milazzo L, Quartuccio L, Scire C, Gaeta GB, et al. Italian consensus guidelines for the management of hepatitis B virus infections in patients with rheumatoid arthritis. *Joint Bone Spine.* 2017;84(5):525-30.
34. Gotzsche PC, Johansen HK. Nystatin prophylaxis and treatment in severely immunodepressed patients. *Cochrane Database Syst Rev.* 2014;9:CD002033.
35. Bharat A, Xie F, Baddley JW, Beukelman T, Chen L, Calabrese L, et al. Incidence and risk factors for progressive multifocal leukoencephalopathy among patients with selected rheumatic diseases. *Arthritis Care Res (Hoboken).* 2012;64(4):612-5.
36. Molloy ES, Calabrese CM, Calabrese LH. The risk of progressive multifocal leukoencephalopathy in the biologic era: prevention and management. *Rheum Dis Clin North Am.* 2017;43(1):95-109.
37. Tamez-Perez HE, Quintanilla-Flores DL, Rodriguez-Gutierrez R, Gonzalez-Gonzalez JG, Tamez-Pena AL. Steroid hyperglycemia: prevalence, early detection and therapeutic recommendations: a narrative review. *World J Diabetes.* 2015;6(8):1073-81.

38. Boomsma MM, Stegeman CA, Kramer AB, Karsijns M, Piers DA, Tervaert JW. Prevalence of reduced bone mineral density in patients with anti-neutrophil cytoplasmic antibody associated vasculitis and the role of immunosuppressive therapy: a cross-sectional study. *Osteoporos Int*. 2002;13(1):74–82.
39. Compston J, Cooper A, Cooper C, Gittoes N, Gregson C, Harvey N, et al. UK clinical guideline for the prevention and treatment of osteoporosis. *Arch Osteoporos*. 2017;12(1):43.
40. Buckley L, Guyatt G, Fink HA, Cannon M, Grossman J, Hansen KE, et al. 2017 American College of Rheumatology guideline for the prevention and treatment of glucocorticoid-induced osteoporosis. *Arthritis Care Res (Hoboken)*. 2017;69(8):1095–110.
41. Homik J, Suarez-Almazor ME, Shea B, Cranney A, Wells G, Tugwell P. Calcium and vitamin D for corticosteroid-induced osteoporosis. *Cochrane Database Syst Rev*. 2000;2:CD000952.
42. Morgan MD, Turnbull J, Selamet U, Kaur-Hayer M, Nightingale P, Ferro CJ, et al. Increased incidence of cardiovascular events in patients with antineutrophil cytoplasmic antibody-associated vasculitides: a matched-pair cohort study. *Arthritis Rheum*. 2009;60(11):3493–500.
43. Suppiah R, Judge A, Batra R, Flossmann O, Harper L, Høglund P, et al. A model to predict cardiovascular events in patients with newly diagnosed Wegener's granulomatosis and microscopic polyangiitis. *Arthritis Care Res (Hoboken)*. 2011;63(4):588–96.
44. Filer AD, Gardner-Medwin JM, Thambyrajah J, Raza K, Carruthers DM, Stevens RJ, et al. Diffuse endothelial dysfunction is common to ANCA associated systemic vasculitis and polyarteritis nodosa. *Ann Rheum Dis*. 2003;62(2):162–7.
45. Souverein PC, Berard A, Van Staa TP, Cooper C, Egberts AC, Leufkens HG, et al. Use of oral glucocorticoids and risk of cardiovascular and cerebrovascular disease in a population based case-control study. *Heart*. 2004;90(8):859–65.
46. Walsh M, Merkel PA, Mahr A, Jayne D. Effects of duration of glucocorticoid therapy on relapse rate in antineutrophil cytoplasmic antibody-associated vasculitis: a meta-analysis. *Arthritis Care Res (Hoboken)*. 2010;62(8):1166–73.
47. Jayne DR, Bruchfeld AN, Harper L, Schaefer M, Venning MC, Hamilton P, et al. Randomized trial of C5a receptor inhibitor avacopan in ANCA-associated vasculitis. *J Am Soc Nephrol*. 2017;28(9):2756–67.
48. Grosse Y, Baan R, Straif K, Secretan B, El Ghissassi F, Bouvard V, et al. A review of human carcinogens—part A: pharmaceuticals. *Lancet Oncol*. 2009;10(1):13–4.
49. Knight A, Askling J, Granath F, Sparen P, Ekblom A. Urinary bladder cancer in Wegener's granulomatosis: risks and relation to cyclophosphamide. *Ann Rheum Dis*. 2004;63(10):1307–11.
50. Faurischou M, Sorensen IJ, Mellemkjaer L, Loft AG, Thomsen BS, Tvede N, et al. Malignancies in Wegener's granulomatosis: incidence and relation to cyclophosphamide therapy in a cohort of 293 patients. *J Rheumatol*. 2008;35(1):100–5.
51. Heijl C, Harper L, Flossmann O, Stucker I, Scott DG, Watts RA, et al. Incidence of malignancy in patients treated for antineutrophil cytoplasmic antibody-associated vasculitis: follow-up data from European Vasculitis Study Group clinical trials. *Ann Rheum Dis*. 2011;70(8):1415–21.
52. Inoue K, Yuasa H. Molecular basis for pharmacokinetics and pharmacodynamics of methotrexate in rheumatoid arthritis therapy. *Drug Metab Pharmacokin*. 2014;29(1):12–9.
53. Romao VC, Lima A, Bernardes M, Canhao H, Fonseca JE. Three decades of low-dose methotrexate in rheumatoid arthritis: can we predict toxicity? *Immunol Res*. 2014;60(2-3):289–310.
54. Whittle SL, Hughes RA. Folate supplementation and methotrexate treatment in rheumatoid arthritis: a review. *Rheumatology (Oxford)*. 2004;43(3):267–71.
55. Ledingham J, Gullick N, Irving K, Gorodkin R, Aris M, Burke J, et al. BSR and BHPR guideline for the prescription and monitoring of non-biologic disease-modifying anti-rheumatic drugs. *Rheumatology (Oxford)*. 2017;56(6):865–8.
56. Asadov C, Aliyeva G, Mustafayeva K. Thiopurine S-methyltransferase as a pharmacogenetic biomarker: significance of testing and review of major methods. *Cardiovasc Hematol Agents Med Chem*. 2017;15(1). <https://doi.org/10.2174/1871525715666170529091921>.
57. Emery P, Fleischmann R, Filipowicz-Sosnowska A, Schechtman J, Szczepanski L, Kavanaugh A, et al. The efficacy and safety of rituximab in patients with active rheumatoid arthritis despite methotrexate treatment: results of a phase IIB randomized, double-blind, placebo-controlled, dose-ranging trial. *Arthritis Rheum*. 2006;54(5):1390–400.
58. van Vollenhoven RF, Emery P, Bingham CO 3rd, Keystone EC, Fleischmann RM, Furst DE, et al. Long-term safety of rituximab in rheumatoid arthritis: 9.5-year follow-up of the global clinical trial programme with a focus on adverse events of interest in RA patients. *Ann Rheum Dis*. 2013;72(9):1496–502.
59. Jones RB, Tervaert JW, Hauser T, Luqmani R, Morgan MD, Peh CA, et al. Rituximab versus cyclophosphamide in ANCA-associated renal vasculitis. *N Engl J Med*. 2010;363(3):211–20.
60. Stone JH, Merkel PA, Spiera R, Seo P, Langford CA, Hoffman GS, et al. Rituximab versus cyclophosphamide for ANCA-associated vasculitis. *N Engl J Med*. 2010;363(3):221–32.
61. Venhoff N, Effelsberg NM, Salzer U, Warnatz K, Peter HH, Lebrecht D, et al. Impact of rituximab on immunoglobulin concentrations and B cell numbers after cyclophosphamide treatment in patients with ANCA-associated vasculitides. *PLoS One*. 2012;7(5):e37626.
62. Roberts DM, Jones RB, Smith RM, Alberici F, Kumaratne DS, Burns S, et al. Rituximab-associated hypogammaglobulinemia: incidence, predictors and

- outcomes in patients with multi-system autoimmune disease. *J Autoimmun.* 2015;57:60–5.
63. Wolach O, Bairey O, Lahav M. Late-onset neutropenia after rituximab treatment: case series and comprehensive review of the literature. *Medicine (Baltimore).* 2010;89(5):308–18.
 64. Knight A, Sundstrom Y, Borjesson O, Bruchfeld A, Malmstrom V, Gunnarsson I. Late-onset neutropenia after rituximab in ANCA-associated vasculitis. *Scand J Rheumatol.* 2016;45(5):404–7.
 65. McDermott EM, Powell RJ. Incidence of ovarian failure in systemic lupus erythematosus after treatment with pulse cyclophosphamide. *Ann Rheum Dis.* 1996;55(4):224–9.
 66. Huong DL, Amoura Z, Duhaut P, Sbai A, Costedoat N, Wechsler B, et al. Risk of ovarian failure and fertility after intravenous cyclophosphamide. A study in 84 patients. *J Rheumatol.* 2002;29(12):2571–6.
 67. Hickman RA, Gordon C. Causes and management of infertility in systemic lupus erythematosus. *Rheumatology (Oxford).* 2011;50(9):1551–8.
 68. Machen L, Clowse ME. Vasculitis and pregnancy. *Rheum Dis Clin North Am.* 2017;43(2):239–47.
 69. Goldstein LH, Dolinsky G, Greenberg R, Schaefer C, Cohen-Kerem R, Diav-Citrin O, et al. Pregnancy outcome of women exposed to azathioprine during pregnancy. *Birth Defects Res A Clin Mol Teratol.* 2007;79(10):696–701.
 70. Garritsen FM, van den Broek MPH, van Zuilen AD, Fidler HH, de Bruin-Weller MS, Spuls PI. Pregnancy and fetal outcomes after paternal exposure to azathioprine, methotrexate or mycophenolic acid: a critically appraised topic. *Br J Dermatol.* 2017;176(4):866–77.
 71. Croft AP, Smith SW, Carr S, Youssouf S, Salama AD, Burns A, et al. Successful outcome of pregnancy in patients with anti-neutrophil cytoplasm antibody-associated small vessel vasculitis. *Kidney Int.* 2015;87(4):807–11.
 72. Clowse ME, Richeson RL, Pieper C, Merkel PA. Vasculitis Clinical Research C. Pregnancy outcomes among patients with vasculitis. *Arthritis Care Res (Hoboken).* 2013;65(8):1370–4.
 73. Merkel PA, Lo GH, Holbrook JT, Tibbs AK, Allen NB, Davis JC Jr, et al. Brief communication: high incidence of venous thrombotic events among patients with Wegener granulomatosis: the Wegener's Clinical Occurrence of Thrombosis (WeCLOT) Study. *Ann Intern Med.* 2005;142(8):620–6.
 74. Berden AE, Nolan SL, Morris HL, Bertina RM, Erasmus DD, Hagen EC, et al. Anti-plasminogen antibodies compromise fibrinolysis and associate with renal histology in ANCA-associated vasculitis. *J Am Soc Nephrol.* 2010;21(12):2169–79.
 75. Walsh M, Flossmann O, Berden A, Westman K, Hoglund P, Stegeman C, et al. Risk factors for relapse of antineutrophil cytoplasmic antibody-associated vasculitis. *Arthritis Rheum.* 2012;64(2):542–8.
 76. Karras A, Pagnoux C, Haubitz M, Groot K, Puechal X, Tervaert JWC, et al. Randomised controlled trial of prolonged treatment in the remission phase of ANCA-associated vasculitis. *Ann Rheum Dis.* 2017;76(10):1662–68.
 77. Sanders JS, de Joode AA, DeSevaux RG, Broekroelofs J, Voskuyl AE, van Paassen P, et al. Extended versus standard azathioprine maintenance therapy in newly diagnosed proteinase-3 anti-neutrophil cytoplasmic antibody-associated vasculitis patients who remain cytoplasmic anti-neutrophil cytoplasmic antibody-positive after induction of remission: a randomized clinical trial. *Nephrol Dial Transplant.* 2016;31(9):1453–9.
 78. Rhee RL, Hogan SL, Poulton CJ, McGregor JA, Landis JR, Falk RJ, et al. Trends in long-term outcomes among patients with antineutrophil cytoplasmic antibody-associated vasculitis with renal disease. *Arthritis Rheumatol.* 2016;68(7):1711–20.
 79. Mukhtyar C, et al. EULAR recommendations for the management of primary small and medium vessel vasculitis. *Annals of the Rheumatic Diseases.* 2009;68:310–7.