

Affinity proteomics reveals extensive phosphorylation of the Brassica chromosome axis protein ASY1 and a network of associated proteins at prophase I of meiosis

Osman, Kim; Yang, Jianhua; Roitinger, Elisabeth; Lambing, Christophe; Heckmann, Stefan; Howell, Elaine; Cuacos, Maria; Imre, Richard; Dürnberger, Gerhard; Mechtler, Karl; Armstrong, Susan; Franklin, Chris

DOI:

[10.1111/tpj.13752](https://doi.org/10.1111/tpj.13752)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Osman, K, Yang, J, Roitinger, E, Lambing, C, Heckmann, S, Howell, E, Cuacos, M, Imre, R, Dürnberger, G, Mechtler, K, Armstrong, S & Franklin, F. C. (2018), Affinity proteomics reveals extensive phosphorylation of the Brassica chromosome axis protein ASY1 and a network of associated proteins at prophase I of meiosis', *The Plant journal : for cell and molecular biology*, vol. 93, no. 1, pp. 17-33. <https://doi.org/10.1111/tpj.13752>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Osman, K. , Yang, J. , Roitinger, E. , Lambing, C. , Heckmann, S. , Howell, E. , Cuacos, M. , Imre, R. , Dürnberger, G. , Mechtler, K. , Armstrong, S. and Franklin, F. C. (2018), Affinity proteomics reveals extensive phosphorylation of the Brassica chromosome axis protein ASY1 and a network of associated proteins at prophase I of meiosis. *Plant J*, 93: 17-33. doi:10.1111/tpj.13752

Published in *The Plant Journal* on 27/10/2017

DOI: 10.1111/tpj.13752

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Affinity proteomics reveals extensive phosphorylation of the Brassica chromosome axis protein ASY1 and a network of associated proteins at prophase I of meiosis

Kim Osman^{1,‡}, Jianhua Yang^{1,†,‡}, Elisabeth Roitinger², Christophe Lambing^{1,‡}, Stefan Heckmann^{1,§}, Elaine Howell¹, Maria Cuacos^{1,§}, Richard Imre², Gerhard Dürnberger^{2,3}, Karl Mechtler², Susan Armstrong¹ and F. Christopher H. Franklin^{1,*}

¹School of Biosciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK,

²IMP-IMBA, 1030, Vienna, Austria, and

³Gregor Mendel Institute of Molecular Plant Biology, Dr. Bohr-Gasse 3, 1030, Vienna, Austria

Received 5 June 2017; revised 10 October 2017; accepted 17 October 2017; published online 27 October 2017.

*For correspondence (e-mail f.c.h.franklin@bham.ac.uk)

‡These authors contributed equally to this work.

†Present address: Faculty of Engineering and Computing, Coventry University, Coventry, CV1 5FB, UK.

§Present address: Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EA, UK.

§Present address: Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), OT Gatersleben, Correnstrasse 3, D-06466, Stadt Seeland, Germany.

SUMMARY

During meiosis, the formation of crossovers (COs) generates genetic variation and provides physical links that are essential for accurate chromosome segregation. COs occur in the context of a proteinaceous chromosome axis. The transcriptomes and proteomes of anthers and meiocytes comprise several thousand genes and proteins, but because of the level of complexity relatively few have been functionally characterized. Our understanding of the physical and functional interactions between meiotic proteins is also limited. Here we use affinity proteomics to analyse the proteins that are associated with the meiotic chromosome axis protein, ASY1, in *Brassica oleracea* anthers and meiocytes. We show that during prophase I ASY1 and its interacting partner, ASY3, are extensively phosphorylated, and we precisely assign phosphorylation sites. We identify 589 proteins that co-immunoprecipitate with ASY1. These correspond to 492 Arabidopsis orthologues, over 90% of which form a coherent protein–protein interaction (PPI) network containing known and candidate meiotic proteins, including proteins more usually associated with other cellular processes such as DNA replication and proteolysis. Mutant analysis confirms that affinity proteomics is a viable strategy for revealing previously unknown meiotic proteins, and we show how the PPI network can be used to prioritise candidates for analysis. Finally, we identify another axis-associated protein with a role in meiotic recombination. Data are available via ProteomeXchange with identifier PXD006042.

Keywords: meiosis, chromosome axis, phosphorylation, LC-MS/MS, protein–protein interaction, *Brassica oleracea*, *Arabidopsis thaliana*.

INTRODUCTION

During meiosis, homologous recombination (HR) generates crossovers (COs) that provide genetic variation and promote accurate chromosome segregation at the first meiotic division. The HR pathway has been studied extensively in *Saccharomyces cerevisiae*, and is thought to be broadly similar in plants (Osman *et al.*, 2011). HR occurs within the context of profound changes in chromosome organization (Kleckner, 2006). Following replication, sister chromatids are linked by the cohesin complex (Haering and Jessberger, 2012). At the leptotene stage of prophase I, the sister chromatids become organized into linear looped arrays that are

conjoined at the loop bases by a proteinaceous axis running along their length. At zygotene, the pairs of homologous chromosomes begin to align and become tightly linked by the synaptonemal complex (SC). This is a highly conserved tripartite structure comprising the chromosome axes with transverse filament (TF) proteins bridging the region between the axes (Page and Hawley, 2004). In many organisms, including plants, mutations leading to defects in axis/SC proteins often have a profound effect on CO formation, whereas in turn recombination pathway mutants can disrupt chromosome morphogenesis (Couteau *et al.*, 1999;

Grelon *et al.*, 2001; Armstrong *et al.*, 2002; Li *et al.*, 2004; Higgins *et al.*, 2005; Ferdous *et al.*, 2012).

CO formation is highly coordinated, such that chromosome pairs receive at least one, termed the 'obligate' CO (Jones and Franklin, 2006). CO designation is thought to occur early in prophase I, and reduces the probability that another CO will occur in an adjacent region, a phenomenon known as CO interference (reviewed in Berchowitz and Copenhaver, 2010). Precisely how these outcomes are achieved remains to be fully elucidated. Nevertheless, proteins associated with the chromosome axis and SC clearly play an important role (Zickler and Kleckner, 2016). The most extensively studied plant meiotic chromosome axis protein is ASY1 (PAIR2 in rice). Arabidopsis *asy1* mutants fail to synapse and have severely reduced CO formation (Ross *et al.*, 1997). In the absence of ASY1, the DMC1 recombinase fails to become stably established on the chromosomes, with the result that interhomologue recombination is severely compromised (Sanchez-Moran *et al.*, 2007). Although not required for axis formation *per se*, ASY1 association with the chromatin is concurrent with axis morphogenesis. Immunolocalization of male meocytes indicates that it first appears as punctate foci in G2 before progressing to a more linear signal along the entire length of the chromosome axes by leptotene (Armstrong *et al.*, 2002). ASY1 remains detectable throughout prophase I, but remodelling of the axis by the AAA+ ATPase, PCH2, during zygotene appears to progressively deplete it from the axis, such that its signal is more obviously associated with the chromatin loops, a process necessary for the normal extension of the SC and the patterned formation of COs (Lambing *et al.*, 2015). Taken together, these studies indicate that ASY1 plays important roles in the coordination of axis/SC morphogenesis and recombination to produce the meiosis-specific bias that favours interhomologue recombination and the maturation of CO-designated recombination intermediates.

To date, insight into meiosis in plants has largely derived from mutant analysis of individual genes (Mercier *et al.*, 2015), identified either through sequence conservation with other species or from mutant or suppressor genetic screens (De Muyt *et al.*, 2009; Crismani *et al.*, 2012; Girard *et al.*, 2014). Global approaches to identify plant meiotic genes and proteins have also been adopted. The transcriptomes of developing anthers undergoing meiosis and of isolated meocytes have been analysed using microarrays and RNAseq (Chen *et al.*, 2010; Tang *et al.*, 2010; Aya *et al.*, 2011; Deveshwar *et al.*, 2011; Libeau *et al.*, 2011; Yang *et al.*, 2011 and Dukowic-Schulze *et al.*, 2014). These studies reveal a highly complex picture, identifying in the order of 1000–2000 meiotically implicated genes. Moreover, the fact that the relationship between mRNA transcription and the cellular level of the corresponding proteins is nonlinear, the possibility of alternatively spliced

meiotic protein variants (Kalsotra and Cooper, 2011; Schmid *et al.*, 2013; Sprink and Hartung, 2014 and Wang *et al.*, 2014), plus evidence from budding yeast and other species that post-translational modifications of meiotic proteins play a key role in their function (for example, Rockmill and Roeder, 1991; Lin *et al.*, 2010; Attner *et al.*, 2013), increases the complexity still further. Proteomic studies present a similarly complex picture (Zhang *et al.*, 2017). Analysis of the proteome and phosphoproteome of *Oryza sativa* (rice) anthers identified 4984 proteins and 3203 phosphoproteins associated with early anther development and meiosis (Ye *et al.*, 2015), whereas a further study focusing on rice meocytes identified 1316 proteins (Collado-Romero *et al.*, 2014).

Here we aimed to reduce complexity by using affinity-based proteomics to immuno-target the key meiotic axis protein, ASY1, to enrich for associated protein complexes using *Brassica oleracea*, which we have previously shown can be used to provide an enriched source of meiotic tissue for proteomic analysis (Sánchez-Morán *et al.*, 2005; Osman *et al.*, 2009). We anticipated that this strategy might also begin to reveal the physical interactions that occur between proteins during prophase I of meiosis. We identify the BoASY1 co-immunoprecipitating proteins and show that their Arabidopsis counterparts form a coherent protein–protein interaction (PPI) network that, importantly, can be used to prioritize candidates for verification of a meiotic role. We present ICU2, the DNA polymerase α subunit, as proof of principle of this approach. We also describe the discovery of another axis-associated protein with an apparent role in meiotic recombination and identify multiple phosphorylation sites in BoASY1 and its interacting partner, BoASY3, providing further insights into meiosis in higher plants.


RESULTS

Identification of BoASY1 co-immunoprecipitating proteins

We carried out co-immunoprecipitation (co-IP) of ASY1 from *B. oleracea* to enrich for associated protein complexes (Figure 1a). Anthers ($n = 200$) were used either intact or their contents were extruded to further enrich for meiotic cells (hereafter, these samples are referred to as 'anthers' or 'meocytes', respectively). Proteins were extracted under non-denaturing conditions to preserve meiotic complexes. With the high level of sequence identity between BoASY1 and AtASY1 (83.6%) we could target BoASY1 using an anti-AtASY1 antibody (Figure S1; Armstrong *et al.*, 2002). Parallel control co-IPs were carried out using non-specific IgG. Proteins were analysed by in-solution mass spectrometry (MS) and identified using a combined database comprising *Brassica rapa* sequences (Wang *et al.*, 2011) and Brassica sequences obtained from the National Center for Biotechnology Information (NCBI,

Figure 1. Identification of BoASY1 co-immunoprecipitating proteins.

(a) Summary of workflow for co-immunoprecipitation (co-IP) experiments. (b) Numbers of ASY1 sample-specific proteins identified in Brassica meiotic tissues, with corresponding numbers of putative Arabidopsis orthologues in parentheses. (c) Molecular function of putative Arabidopsis orthologues indicated by gene ontology classification. (d) ASY1 co-IP network, with nodes representing proteins and edges representing interactions. ASY1 (green) and cluster of cohesin, histone and replication-related proteins (orange) are highlighted. (e) Detailed view of cluster. Proteins are named according to TAIR.



<https://www.ncbi.nlm.nih.gov>) in 2010. Putative orthologues in *Arabidopsis thaliana* were identified using the best BLASTP score. Five anther and four meiocyte data sets were collected. ASY1 was identified in all data sets with 55 peptides in total and up to 75% sequence coverage (Figure S2a), confirming that targeting was successful. In addition, each data set contained several hundred BoASY1 sample-specific proteins. These ranged from proteins present in all data sets and identified by a relatively large number of unique peptides to those that appeared only once with two unique peptides. To obtain an indication of protein reliability, the raw data from all experiments were searched together and peptide/protein label-free quantification was carried out using the in-house tool PEAKJUGGLER (unpubl. data; Andersen *et al.*, 2017), to determine peak area. Six of the nine data sets were obtained using three technical replicates of each sample, allowing the statistical significance of proteins to be determined using LIMMA analysis (Smyth Gordon, 2004). Proteins showing a fold-change of ≥ 5 in sample relative to control and $P < 0.01$ in

at least one data set were considered significant in label-free quantification. In addition, to avoid inadvertently excluding any meiotically relevant proteins, we decided to retain all ASY1 sample-specific proteins satisfying the minimum identification threshold of two peptides while excluding all proteins identified in any of the control samples. This also allowed us to consider data from the three remaining data sets that lacked replicates. Any proteins that were accepted purely on this qualitative basis were considered less reliable than the quantitatively significant group. Nevertheless, our decision to retain them appeared justified when several were subsequently confirmed as having a meiotic role (see below). Details of both sets of accepted proteins are presented in Table S1.

The group taken forward for further analysis therefore comprised 589 Brassica proteins corresponding to 492 Arabidopsis gene loci. Note that all but one Brassica protein could be assigned a putative Arabidopsis orthologue; the discrepancy between the number of Brassica and Arabidopsis proteins is explained partly by the database

containing sequences from several different Brassica species and partly by an ancient *Brassicaceae* lineage-specific whole-genome triplication event (Liu *et al.*, 2014; Parkin *et al.*, 2014), such that in some cases several Brassica proteins indicate the same Arabidopsis orthologue. As expected, there was some overlap between anther and meiocyte data sets, with 421 of the total 589 Brassica proteins (71.5%) identified in both tissue types (336 of 492 for Arabidopsis; Figure 1b); however, 125 Brassica proteins (21.2%) were identified only in meiocytes despite ASY1 being detected equally well in both tissues (117 proteins, 23.8%, for Arabidopsis).

Chromosome axis and SC-associated proteins co-immunoprecipitate with ASY1

Gene ontology (GO) classification of the 492 Arabidopsis orthologues using The Arabidopsis Information Resource website (TAIR, <https://www.arabidopsis.org>) indicated that the group of ASY1 co-IP proteins covered a range of molecular functions (Figure 1c). Analysis of GO enrichment relative to the Arabidopsis genome was conducted using PANTHER accessed through the GO consortium website. Further analysis was carried out using the 453 orthologues identified from meiocytes. In both cases, a large number of GO terms were found to be enriched, but notable amongst the Biological Process terms showing the highest fold enrichment were several relating to DNA processing and nucleus organisation (Table S2a), and 'DNA-dependent ATP-ase activity' was one of the most highly enriched terms for Molecular Function (Table S2b). Several large protein complexes and functional pathways or families were well represented in the ASY1 co-IP data, so where appropriate we used a combination of the KEGG pathway database (<http://www.kegg.jp/kegg/pathway.html>) and examination of the relevant literature to group Arabidopsis orthologues accordingly (Table S3).

We identified 12 proteins with a prior confirmed meiotic role in Arabidopsis, including several axis and SC-associated proteins (Table S3). From the cohesin complex we identified sub-units SMC1, SMC3 and SCC3 and one of the five Arabidopsis SPO76 cohesin cofactor proteins, PDS5C (Chelysheva *et al.*, 2005; Lam *et al.*, 2005; Pradillo *et al.*, 2015). The SC transverse filament protein, ZYP1a, and the condensin I subunit, CAP-D2, were also detected (Higgins *et al.*, 2005; Smith *et al.*, 2014), as were the axis protein, ASY3, and PCH2, an AAA+ ATPase with a role in prophase I axis remodelling, both of which we characterized and published during the course of this study (Ferdous *et al.*, 2012; Lambing *et al.*, 2015). Given the functional relationship of the HR pathway and the developing axis and SC, it was encouraging that several meiotic recombination proteins immunoprecipitated with ASY1, notably PRD3, required for DNA double-strand break (DSB) formation (De Muyt *et al.*, 2009), and the

recombinase DMC1 (Klimyuk and Jones, 1997; Doutriaux *et al.*, 1998). Finally, we identified two peptides of the CDK1 homologue, CDKA;1, previously implicated as having a role in meiotic progression (Cromer *et al.*, 2012). Most of the previously confirmed meiotic proteins were identified either from both tissue types or solely from meiocyte samples; however, CAP-D2 was identified only from intact anthers.

Other proteins that have (or are predicted to have) a close association with chromatin were present in the ASY1 co-IP data (Table S3), including proteins involved in DNA replication and repair, chromatin remodelling proteins, putative transcription factors and regulators, and histone proteins. There were several proteins implicated in the RNA-dependent DNA methylation (RdDM) pathway, including AGO4 (Table S3). Argonaute proteins have been shown to have important pre-meiotic and meiotic roles in a range of organisms, including several plant species (Nonomura *et al.*, 2007; Olmedo-Monfil *et al.*, 2010; Singh *et al.*, 2011; Oliver *et al.*, 2014, 2016; Liu and Nonomura, 2016).

Twenty 26S proteasome and 11 ubiquitination-related proteins were identified (Table S3), suggesting a close association between these proteins and the meiotic chromosome axis. In animals and yeast, the importance of the ubiquitin–proteasome system (UPS) in regulating key aspects of meiosis, such as recombination and meiotic progression, is well established (Bose *et al.*, 2014), and is now also beginning to be elucidated in plants (Wang and Yang, 2006; Zhao *et al.*, 2006; He *et al.*, 2016).

ASY1 co-IP proteins form a coherent protein–protein interaction network

As the ASY1 co-IP proteins covered a wide range of protein types and GO terms, we carried out network analysis to determine whether they were predicted to interact based on existing data in the public domain. We used an open-source database of known and predicted protein interactions: STRING (Szklarczyk *et al.*, 2015). The STRING network was created using the 492 putative Arabidopsis orthologues of BoASY1 co-IP proteins (Table S1). The network was visualized using CYTOSCAPE (Appendix S1; Figure 1d): 92.7% of proteins (456) formed a single network, with relatively few 'orphans' (36), suggesting that they had immunoprecipitated as a coherent group, providing further evidence that the co-IP approach was successful. It is worth noting that six proteins in the 'orphans' group lacked annotation or were otherwise 'unknown', and a further five uncharacterized proteins were linked to the main network only by virtue of co-expression or by being co-mentioned in public text collections (Appendix S1; Table S3). Given that axis and SC proteins tend to be poorly conserved, these proteins were interesting candidates for further study (see below).

The ASY1 co-IP network can be used to prioritize candidates for functional analysis

Candidate proteins were investigated for a meiotic role by cytological examination of chromosome spreads of male meiocytes from homozygous Arabidopsis T-DNA insertion lines. Initially, we chose candidates based largely on confidence of identification (Table S1) and absence of a previously published role, but we also considered their potential function as inferred from conserved domains, etc. We found several with a strong meiotic mutant phenotype, including ASY3 and PCH2 (Ferdous *et al.*, 2012; Lambing *et al.*, 2015). In addition, from a sample of 10 candidates exhibiting only a modest or no reduction in fertility, four displayed a relatively minor mutant phenotype, where meiotic defects were clearly observed at the cytological level but were present in only a subset of meiocytes (<10%). Defects included chromosome fragmentation, unresolved interlocks, interbivalent connections, univalency and chromosome bridges at the division stages. Results from the four candidates are summarized in Figures S3 and S4, and Table S4. Interestingly, a phospho-modified peptide was detected in the N terminus of the Brassica orthologue of one of the candidates (gi257685916; At5g59210), an structural maintenance of chromosomes (SMC) domain protein, and other phosphopeptides were identified in the N- and C-terminal regions of Bra004279 (At1g68060, MAP70-1) a microtubule-associated protein (Figure S2b, c). Data from IntAct (Arabidopsis Interactome Mapping Consortium, 2011; Orchard *et al.*, 2014), revealed a two-hybrid array interaction between At5g59210 and At1g68060, thus supporting a direct physical interaction between the *B. oleracea* orthologues of these two proteins in our study.

As mentioned above, many of the ASY1 co-IP proteins were identified with few unique peptides, appearing in only one or two data sets, and as such might be considered relatively low-confidence candidates (Table S1). We therefore investigated whether we could use the ASY1 co-IP STRING network to prioritize candidates for analysis, particularly as the process of identifying and screening homozygous mutants is labour and time intensive. In the network, ASY1 and PCH2 are located in a cluster that contains several histone-related proteins, cohesin complex components and proteins associated with DNA replication, including RFC complex and MCM family proteins (Figure 1e; Table S3). A member of the MCM family, MCM8, is involved in DMC1-independent DSB repair in Arabidopsis (Crismani *et al.*, 2013), whereas the large subunit of the heteropentameric RFC complex, RFC1, is required for meiotic DSB repair (Liu *et al.*, 2013) and interference-sensitive COs (Wang *et al.*, 2012b). Topoisomerase II is necessary for resolving heterochromatic DNA entanglements during female meiosis I in *Drosophila melanogaster* (Hughes and

Hawley, 2014), for meiotic chromosome condensation and separation in mice (Li *et al.*, 2013), and has a role in mediating CO interference in budding yeast (Zhang *et al.*, 2014). Given the clear link between proteins in this cluster and aspects of meiotic DNA metabolism, we decided to investigate other cluster members and, indeed, mutant analysis of several of these, for example ICU2, did suggest a meiotic role.

ICU2 is required for normal progression through meiosis

ICU2 (the catalytic subunit of DNA polymerase α) is involved in mediating epigenetic states in Arabidopsis (Barrero *et al.*, 2007; Liu *et al.*, 2010 and Hyun *et al.*, 2013), and has a potential role in HR (Liu *et al.*, 2010). To determine whether ICU2 is involved in meiosis we analysed *icu2-1*, homozygous for a non-lethal missense allele of the gene (Barrero *et al.*, 2007). *icu2-1* has a pleiotropic phenotype, including early flowering, leaf incurvature, homeotic transformations of some floral parts, reduced plant height and reduced fertility (Barrero *et al.*, 2007). In our hands, the fertility of *icu2-1* was 45.4% (seed count per silique, $n = 50$). Analysis of male meiosis in *icu2-1* indicated that during most of prophase I the mutant was indistinguishable from wild-type (WT) Arabidopsis (background En2, $2n = 10$; Figure 2). Chromosomes appeared as thin threads in leptotene (Figure 2a, d), with homologues becoming fully paired and synapsed by pachytene (Figure 2b, e). ZYP1 immunolocalisation at this stage suggested synapsis was complete (Figure 2c, f). Chromosomes then began to desynapse and condense, and at metaphase I five aligned bivalents were observed in the WT (Figure 2g). Separation of the homologues at anaphase I (Figure 2h) followed by separation of sister chromatids at the second division then resulted in a tetrad of the four haploid products of meiosis (Figure 2i). In *icu2-1*, however, 44.4% of first divisions appeared aberrant ($n = 30$); nuclei did not exhibit five normal bivalents at metaphase I (Figure 2j), and as homologues began to separate at anaphase I, fragmentation and abnormal chromosomal connections were observed (Figure 2k). This resulted in unbalanced nuclei with fragmented chromosomes at the tetrad stage (Figure 2l). The programmed formation of meiotic DSBs and subsequent recombination and synapsis is dependent on the activity of the SPO11 complex (Bergerat *et al.*, 1997). To determine whether the fragmentation/connections observed in *icu2-1* resulted from unrepaired breaks incurred during pre-meiotic replication or resulted from defective repair of SPO11-induced DSBs during recombination, we generated a *spo11-1-4 icu2-1* double mutant. Chromosome spreads of *spo11-1-4 icu2-1* indicated that most meiocytes had a similar phenotype to the *spo11-1-4* single mutant, with 10 achiasmatic univalents rather than five bivalents at metaphase I (Figure 2m, n). Only 10.0% of nuclei exhibited fragmentation and/or




Figure 2. Cytological analysis of *icu2-1* showing male meiotic chromosome spreads stained with 4',6-diamidino-2-phenylindole (DAPI).

(a–c, g–i) Wild type (WT). (d–f, j–l) *icu2-1*. (c, f) Immunolocalization of ZYP1 (green), DAPI (blue). (m–p) First division in single and double mutants of *SPO11-1* and *ICU2*. (m) *spo11-1-4* and (n) *spo11-1-4 icu2-1* nucleus with 10 univalents. (o) *spo11-1-4 icu2-1* and (p) *icu2-1* nucleus with unresolved chromosomal connections (yellow arrows) and fragmentation (orange arrows). Scale bars: 10 μ m.

unresolved connections in the double mutant (Figure 2o) compared with 48.3% of nuclei in the *icu2-1* single mutant (Figure 2p; $\chi^2_{(1)} = 19.52$, $P < 0.0001$, $n = 60$). Therefore the *spo11-1-4* mutation can largely rescue the phenotype of the *icu2-1* single mutant, indicating that *ICU2* has a role in meiotic recombination.

Analysis of *ICU2* therefore provides 'proof of principle' of using the co-IP networks to prioritize particular proteins or key interactions for verification and analysis, and appears to justify our choice of acceptance level in the co-IP analysis as the protein was identified in the lower confidence qualitative group with only three peptides (see above and Table S1).

We also analysed a homozygous mutant of *MCM2*, albeit in less detail than *icu2-1*, and observed mild meiotic defects and a 10% reduction in fertility (Figure S5; Table S4).

Identification of an axis-associated protein

As axis and SC proteins tend to be poorly conserved at the primary sequence level, we were interested to note that the ASY1 co-IP data included several uncharacterized proteins lacking known functional domains (Table S3). As mentioned above, one such protein was subsequently characterized as ASY3 (Ferdous *et al.*, 2012). Another protein, encoded by At2g33793, was found to share 23.9% identity and 40.1% similarity with the C-terminal predicted coiled-coil region of AtASY3 (Figure S6a). At2g33793 formed links with several other STRING network proteins on the basis of co-expression, including TOPII, MCM5, CAP-D2 and PRD3 (Appendix S1). During this project, At2g33793 was independently identified by Mathilde Grelon's group (INRA, France), and was subsequently referred to as ASY4.


Preliminary characterization of a weak mutant allele of ASY4 (Figure S6b, c) indicated normal vegetative growth and silique length, but a slight reduction in fertility based on seed count per silique (mean of 57.50 compared with 60.54 in the WT, $P < 0.001$, $n = 50$). Analysis of meiocytes from *asy4* confirmed a meiotic role (Figure 3). During prophase I, *asy4* appeared similar to the WT, and by pachytene homologues appeared paired and synapsed based on ZYP1 immunolocalization (Figure 3a–f), although we cannot rule out the possibility of short stretches of chromosomes remaining unsynapsed in some nuclei. Most *asy4* nuclei completed meiosis apparently normally; however, at metaphase I, unlike the situation in the WT where five aligned bivalents were invariably observed, a small proportion of *asy4* nuclei contained univalents (2.3%, $n = 130$; Figure 3g, h). Abnormal inter-bivalent connections were also apparent (Figure 3i). As homologues separated at anaphase I, chromosome bridges were observed in

15.6% ($n = 32$) of nuclei (Figure 3k, l; $\chi^2_{(1)}$ for aberrant nuclei at the first division = 6.28, $P = 0.012$, $n = 162$).

We then explored the interaction between ASY4 and the other axis proteins, ASY1 and ASY3. Previously we showed that ASY1 and ASY3 directly interact via the C-terminal predicted coiled-coil region of ASY3 (Ferdous *et al.*, 2012). Given the high sequence similarity between this region of ASY3 and ASY4, we investigated whether ASY4 could also directly interact with ASY1 or, indeed, with ASY3. Yeast two-hybrid analysis found no direct interaction between ASY1 and ASY4 (Figure 4a). In contrast, in an analysis of full-length cDNAs from ASY3 and ASY4, yeast growth was enabled even under high-stringency selection, demonstrating a direct physical interaction between their encoded proteins. Furthermore, as in the interaction between ASY3 and ASY1 (Ferdous *et al.*, 2012), the predicted coiled-coil region of ASY3 (residues 623–793) was sufficient for interaction with ASY4 (Figure 4b). These

Figure 3. Cytological analysis of *asy4* showing male meiotic chromosome spreads.

(a–f) During early meiotic stages *asy4* appears similar to the wild type (WT), with chromosomes becoming paired and synapsed by pachytene. (g) WT metaphase I. (h–i) *asy4* metaphase I with (h) univalents and (i) inter-bivalent connections. (j) WT anaphase I. (k–l) *asy4* anaphase I, with (k) chromosome bridges and (l) stray chromosome or large fragment and chromosome bridge. DNA is stained with 4',6-diamidino-2-phenylindole (DAPI). In (c) and (f) the immunolocalization of ZYP1 (green) marks the synaptonemal complex. Arrows indicate relevant features. Scale bars: 10 μ m.






Figure 4. Yeast 2-hybrid analysis of ASY4. Plasmid constructs were co-transformed into yeast cells and plated on SD-Leu/Trp (-LT), SD-Leu/Trp/His (-LTH) and SD-Leu/Trp/His/Ade (-LTHA).

(a) ASY4 and ASY1: absence of growth on -LTH and -LTHA, but growth on the control medium, -LT, suggested that there was no direct interaction between ASY4 and ASY1. (b) ASY4 and ASY3: growth on -LTH and -LTHA confirmed that the predicted coiled coil-containing region of ASY3 (amino-acid residues 623–793) is sufficient for interaction with ASY4.

results confirm that ASY4 is axis-associated with a potential role in meiotic recombination.

BoASY1 and BoASY3 are phosphorylated at multiple sites

In other organisms the phosphorylation of chromosome axis proteins is important in regulating their activity during meiosis (Rogers *et al.*, 2002; Brar *et al.*, 2006; Carballo *et al.*, 2008; Katis *et al.*, 2010; Fukuda *et al.*, 2012; Penedos *et al.*, 2015; Sakuno and Watanabe, 2015). We were therefore interested in whether MS analysis would enable us to identify phospho-modified residues in BoASY1 and its interacting partner BoASY3. This proved to be the case. For BoASY1, phospho-modified forms of 13 different peptides were identified with a total of 18 distinct phospho-modified Serine (S) or Threonine (T) sites (Table 1). For two of the peptides the precise position of the phosphate group was unclear, but in the majority of cases the position of the phosphorylated residue could be unambiguously determined. Several peptides had doubly phosphorylated forms. Of the 18 phospho-modified residues, four corresponded to S/TQ motifs, the preferred sites of phosphorylation for the ATM/ATR family of DNA

damage response serine/threonine kinases. Notably, all of the phospho-modified S/TQ sites were located within two S/TQ cluster domains (SCDs), defined as a region where three or more S/TQ motifs occur within a span of up to 100 residues (Traven and Heierhorst, 2005; Figure 5). SCDs are known targets of ATM/ATR. SCD1 is located near the centre of the protein, between HORMA and SWIRM domains, and comprises four S/TQ sites (S267, T272, T294 and S300), with phospho-modification detected at T294 and S300. AtASY1 also has a SCD in this region, although it differs slightly from the BoASY1 SCD, containing only three S/TQ motifs (S267, T269 and T295) and lacking a site corresponding to S300 in BoASY1 (Figure 5). BoASY1 T294 is conserved, however, corresponding to T295 in AtASY1. Both BoASY1 and AtASY1 also contain a second SCD consisting of three S/TQ motifs close to the C terminus. Two of the S/TQ motifs in this SCD (S569 and S572) lie in close proximity on the same BoASY1 peptide (Figure 5; Table 1), and we identified phospho-modification at S568, S569 and S572.

The remaining phospho-modified sites in BoASY1 also tend to occur in clusters. Of particular note are S260, S262

Table 1 Phosphorylation sites identified in BoASY1 and BoASY3

| Protein | Site | Phosphopeptide(s) | Tissue | ptmRS: Best site probabilities |
|---------------|-----------------------|--|---|--|
| BoASY1 | S17 | EAEITEQD(S)LLLTR | A | S9(Phospho), 100.00 |
| | S253 and S260 | STGPN(S)VHDEQP(S)DSDSEISQTK | M | S6 (Phosho), 97.33; S13 (Phosho), 99.82 |
| | S260 | STGPNSVHDEQP(S)DSDSEISQTK | A and M | S13(Phospho), 99.99 |
| | S260 and S262 | STGPNSVHDEQP(S)D(S)DSEISQTK | | S13(Phospho), 100.00; S15(Phospho), 100.00 |
| | S262 | STGPNSVHDEQPSD(S)DSEISQTK | | S15(Phospho), 99.85 |
| | S262 and S264 | STGPNSVHDEQPSD(S)D(S)EISQTK (with S264) | | S15(Phospho), 82.82; S17(Phospho), 90.01 |
| | T294 | ETQFLVAAVEKQEDDDGEVDEDN(T)QDPVESQQQLER | A and M | T24(Phospho), 100.00 |
| | | QEDDDGEVDEDN(T)QDPVESQQQLER | | T13(Phospho), 100.00 |
| | S300 | QEDDDGEVDEDNTQDPVE(S)QQQLER | A and M | S19(Phospho), 100.00 |
| | | QEDDDGEVDEDN(T)QDPVE(S)QQQLER (with T294) | | T13(Phospho), 100.00; S19(Phospho), 100.00 |
| | S442 or S443 | MVQEGYVED(S)SNRR or MVQEGYVEDS(S)NRR | A | S10(Phospho), 50.00; S11(Phospho), 50.00 |
| | T493 | TNGQDAKL(T)PDVSTR | A and M | T9(Phospho), 100.00 |
| | | L(T)PDVSTR | | T2(Phospho), 100.00 |
| | S504 | GGIH(S)IGSDLTR | | S5(Phospho), 98.97 |
| | S504 and S507 | GGIH(S)IG(S)DLTR | A and M | S5(Phospho), 100.00; S8(Phospho), 99.88 |
| | S507 | GGIHSIG(S)DLTR | | S8(Phospho), 100.00 |
| | S526 | SAMHQNGSVL(S)EQTISK | M | S11(Phospho), 99.98 |
| | T536 | ANN(T)PMSSNAQPVASR | A and M | T4(Phospho), 100.00 |
| | S539 | ANNTPM(S)SNAQPVASR | A and M | S7 (Phospho), 99.39 |
| | S547 or S550 | ANNTPMSSNAQPVA(S)RESFAVK or ANNTPMSSNAQPVASRE(S)FAVK | M | S15(Phospho), 50.00; S18(Phospho), 50.00 |
| S568 and S569 | ICTDAGTD(S)(S)QASQDRR | A and M | S9 (Phospho), 99.89; S10 (Phospho), 91.24 | |
| S569 | ICTDAGTDS(S)QASQDR | A and M | S10(Phospho), 96.10 | |
| S572 | ICTDAGTDSSQA(S)QDRR | A and M | S13(Phospho), 98.72 | |
| BoASY3 | S15 | SFGSNFHPS(S)QPR | M | S10(Phospho), 94.53 |
| | S156 | GNEMDK(S)PER | A and M | S7(Phospho), 100.00 |
| | S205 | A(S)PEYNEDVNSETPEVVK | M | S2(Phospho), 99.67 |
| | T231 or S232 | LNQDK(T)SNDDPLTK or LNQDKT(S)NDDPLTK | M | T6(Phospho), 50.00; S7(Phospho), 50.00 |
| | S251 or S253 | HHSDTIETD(S)E(S)PEVATR | M | S10(Phospho), 49.72; S12(Phospho), 49.72 |
| | S432 and S441 | EK(S)VEPENDFQ(S)PTFGYK | A | S3(Phospho), 100.00; S12(Phospho), 92.49 |

Phospho-modified residues are indicated by parentheses. Some peptides were confirmed as doubly phosphorylated. For a few peptides a phospho-modification could be confirmed, but the precise location within the peptide could not be determined. All sites were identified by ptmRS and manually verified. A (anther) and M (meiocyte) indicate the tissue(s) from which the phosphopeptides were identified.

and S264, located immediately upstream of SCD1 (Figure 5; Table 1). Phosphorylation at these sites was complex, with both singly and various doubly phosphorylated peptides observed. Interestingly, the multiple acidic residues surrounding the phosphoserines matches the hallmark motif of casein kinase 2 (CK2; Pinna, 2002). Two more loose clusters, each consisting of three phospho-modified sites within a stretch of up to 15 residues, occur between the SWIRM domain and SCD2. One site in each cluster (T493 and T536) is at a consensus minor CDK1 motif (S/TP) (Figure 5; Table S1).

In BoASY3 we identified phospho-modified forms of six different peptides (Table 1). Most carried a single modification, but one was doubly phosphorylated at S432 and S441. The positions of most sites could be unequivocally determined, but in two cases was ambiguous (T231 or S232; S251 or S253). Four of the sites are at consensus CDK1 motifs: S205, S253 and S441 are at minor motifs and S156 is at a full motif (S/TPXK/R; Figure 6). Only one site, S15, is at an S/TQ motif.

DISCUSSION

In plants, as in other sexually reproducing organisms, the frequency and distribution of COs during meiosis is governed by the functional inter-relationship between the recombination machinery and the proteinaceous structures that organize the chromosomes during prophase I of meiosis. Thus far our understanding of plant meiosis largely derives from the analysis of around 90 plant meiotic genes, primarily identified through mutant analysis. Although effective, this approach is hampered by several factors. For instance, many of the proteins that are crucial for meiosis are likely to be involved in essential processes in somatic cells. Some genes are duplicated, functionally redundant or, when mutated, produce only subtle phenotypes with little impact on fertility (at least under standard glasshouse growth conditions). Here we have demonstrated that affinity proteomics can be used as an additional approach to identify proteins that play a role in meiosis, and that by targeting a specific component of the meiotic machinery it is




Figure 5. Phosphorylation sites identified in BoASY1. The full-length sequence of BoASY1 is shown aligned with AtASY1 and its budding yeast orthologue ScHop1 (CLUSTAL OMEGA). BoASY1 phospho-modified residues are highlighted in blue (note that two phospho-sites could not be precisely determined, as indicated above the sequence). ScHop1 phospho-sites (Carballo *et al.*, 2008) are highlighted in green. S/TQ cluster domains in BoASY1 and AtASY1 are indicated by red lines above the sequence. Predicted HORMA (residues 13-222) and SWIRM (residues 389-438) domains are highlighted in yellow in AtASY1. [Colour figure can be viewed at wileyonlinelibrary.com].

possible to begin to define the protein–protein interaction network in which the protein participates.

Proteins that co-immunoprecipitate with ASY1 can be organized into a coherent interaction network

Previously, global analyses of the proteomes of developing rice anthers during meiosis and isolated rice meiocytes have identified several thousand proteins, highlighting the complexity of the meiotic proteome (Collado-Romero *et al.*, 2014; Ye *et al.*, 2015). Nevertheless, it is likely that the picture is incomplete as Ye *et al.* (2015) identified peptides corresponding to just 10 of at least 28 characterized rice meiotic proteins (Luo *et al.*, 2014), and homologues of only 14 and seven Arabidopsis and budding yeast meiotic proteins, respectively.

Adopting a strategy based on affinity purification of meiotic complexes has provided a viable alternative approach,

in that focusing on proteins associated with a key meiotic protein, in this case ASY1, substantially reduces complexity and facilitates functional analysis. This approach has enabled us to define a PPI network of 492 nodes that incorporates ASY1. That a substantial proportion of the network proteins are likely to have a meiotic role has been validated using a combination of prior functional knowledge (for example, the presence of CAP-D2, ZYP1 and PRD3; Higgins *et al.*, 2005; De Muyt *et al.*, 2009; Smith *et al.*, 2014) and functional analysis (this study; Ferdous *et al.*, 2012; Lambing *et al.*, 2015). It is apparent from this analysis that although some network proteins such as ASY3, PCH2, ICU2 and PRD3 have major meiotic roles, this is not the case for a significant proportion. Mutant analysis of a small sample suggests that many of the network proteins may have only a minor effect on meiosis; however, this is based on a preliminary analysis, and hence we cannot rule

(Wang *et al.*, 2012b), and DNA leading-strand synthesis was found to be important for the formation of interference-sensitive COs in Arabidopsis (Huang *et al.*, 2015). The precise role of ICU2 will require further study.

Cytological analysis suggested a potentially interesting mutant phenotype for At5g59210, the protein product of which is predicted to contain extensive coiled-coil regions. Further work will be required to fully characterize the role of this protein, but the identification of a phospho-site near the N-terminus of its Brassica orthologue is interesting, particularly as IntAct indicates an interaction with MAP70-1, a plant-specific microtubule-associated protein, the Brassica orthologue of which was also found to be phosphorylated. We have not yet investigated MAP70-1 for a meiotic role. Any analysis would need to address the fact that it is part of a multigene family, sharing a high degree of identity with three other proteins (Korolev *et al.*, 2005), and that the Brassica orthologues of all four proteins were present in the ASY1 co-IP data (Tables S1 and S3). Consistent with the identification of microtubule-associated proteins, several β -tubulin and α -tubulin proteins were also present in the co-IP data.

Of the 492 Arabidopsis loci submitted for STRING analysis, 11 were unknown/uncharacterized and six could not be incorporated into the PPI network. These proteins were thought to be good candidates for investigation for a meiotic role, which led to the identification and preliminary characterization of ASY4. Although we were able to analyse only a weak mutant allele of ASY4, its cytological phenotype of impaired CO formation, together with its high degree of similarity to the C terminal of ASY3, and the confirmation of a direct Y2H interaction between the two proteins, strongly supports an axis-associated role for the protein. No direct Y2H interaction with ASY1 was detected, suggesting that ASY4 may have been co-immunoprecipitated by an indirect interaction with ASY1 via ASY3, thus illustrating a further advantage of using an affinity proteomics approach in that by targeting meiotic complexes, secondary and even higher order protein interactors might be identified. An indirect interaction with ASY1 might explain why we identified relatively few unique peptides of ASY4 compared with ASY1 and ASY3 (3, 55 and 39, respectively), although this could also have been influenced by its smaller size (ASY4 has a predicted molecular weight of 24.69 kDa compared with 67.21 kDa for ASY1 and 88.00 kDa for ASY3).

Protein phosphorylation

Mass spectrometry (MS) revealed that *in vivo* BoASY1 is extensively phosphorylated at prophase I of meiosis. Amongst the 18 identified sites there were four S/TQ motifs distributed between two SCDs. SCD1 is located near the centre of the protein, between the HORMA and SWIRM domains, and SCD2 is near the C terminus. Sequence

alignment suggests a comparable arrangement in AtASY1, although there appears to be an additional S/TQ motif in SCD1 of BoASY1 (Figure 5). Although further studies will be required to determine whether phosphorylation of the S/TQ residues in SCD1 and SCD2 is of functional significance, comparison with Hop1 in budding yeast (Figure 5) suggests that this is possible, at least for some of the S/TQ sites in SCD1 (Carballo *et al.*, 2008). Hop1 contains eight S/TQ motifs; three of them form an SCD located just downstream of the HORMA domain, and all three are phosphorylated *in vivo* during meiosis. Phosphorylation at T318 has the greatest effect in promoting Hop1-dependent interhomologue recombination (Carballo *et al.*, 2008), and phosphorylation at S298 promotes stable interaction of HOP1 and Mek1 effector kinase on the chromosomes following initial phospho-T318 mediated Mek1 recruitment (Penedos *et al.*, 2015). Similar to Hop1, SCD1 is found near the centre of the protein between the HORMA and SWIRM domains in BoASY1 and AtASY1. A full-length alignment of BoASY1 with Hop1 (Figure 5) suggests that residue T294 in SCD1 corresponds in position to Hop1 T318. As is the case for T318 in Hop1, a flanking S/TQ motif (S300) is also phosphorylated. Although it is reasonable to speculate that the S/TQ sites within SCD1 and the Hop1 SCD may be functionally comparable, this is clearly not the case for SCD2, which is absent from the budding yeast protein. SCD2 is conserved in the rice HORMA domain protein, OsPAIR2, however, and has also been reported to undergo phosphorylation (Ye *et al.*, 2015). This occurred at S579, which corresponds to S572 in BoASY1. A second OsPAIR2 phospho-site was detected nearby at S569, a non-S/TQ site. Nevertheless, the significance of phosphorylation in the ASY1 C terminal remains obscure, as it appears that this SCD is present in the orthologues of some plant species, but not in others.

The significance of the tendency for the non-S/TQ phosphosites in BoASY1 to also occur in clusters remains to be determined. The cluster S260, S262 and S264 just upstream of SCD1 (Figure 5; Table 1) is particularly interesting because the multiple acidic residues surrounding the phospho-serines matches the hallmark motif of casein kinase II (Pinna, 2002). CK2 motifs were recently identified amongst irradiation and ATM/ATR-dependent upregulated phosphorylation sites in Arabidopsis, although it remains to be seen whether these sites are actually targeted by CK2 in an ATM/ATR-dependent manner (Roitinger *et al.*, 2015). Phosphorylation at minor CDK1 sites within the two clusters situated between the SWIRM domain and SCD2 may also be of significance, particularly as CDKA;1 was amongst the proteins we identified, albeit with only two peptides.

The most striking feature of the BoASY3 phospho-sites is that all seven are located in the N-terminal region of the protein (Figure 6). This is consistent with an earlier

functional analysis of ASY3 that showed that the C-terminal coiled-coil region of the protein (residues 623–793) is involved in its interaction with ASY1 in Arabidopsis (Ferdous *et al.*, 2012), and may therefore be inaccessible for signalling. Four of the sites were at consensus CDK1 motifs: one at a full motif and three at minor motifs. A single phospho-site, at position S81, was identified in the rice ASY3 orthologue, PAIR3, and was also at a minor CDK1 motif (Ye *et al.*, 2015). Red1, the budding yeast orthologue of BoASY3, contains seven putative target sites of Cdc28 (CDK1) and at least four Cdc28-independent phosphorylation sites (Lai *et al.*, 2011); however, in a full-length alignment of BoASY3 and Red1, only S432 and S441 lie in close proximity to a Red1 phosphosite (S469; Figure 6). Of these, S441 is a minimal S/TP motif, like Red1 S469. Functional analysis of phosphorylation in Red1 suggested that it was non-essential for its functions in meiosis (Lai *et al.*, 2011), so it will be interesting to investigate any potential role for ASY3 and ASY1 phosphorylation in future studies.

Besides CDKA;1, several other kinases and phosphatases were identified in the ASY1 co-IP data, including the protein phosphatase 2A subunits PP2AA2 and PP2A-3 (Tables S1 and S3). PP2A has been implicated in a number of meiotic roles in animals and yeast (e.g. Lu *et al.*, 2002; Kitajima *et al.*, 2006; Riedel *et al.*, 2006; Nolt *et al.*, 2011; Tang *et al.*, 2016), and is regulated by the UPS during mouse oocyte maturation (Yu *et al.*, 2015). It remains to be established whether it, or any of the kinases identified in our study, has a role in plant meiosis, however.

Proteins associated with other cellular processes

The ASY1 co-IP data contained multiple components of several large complexes and functional pathways, such as the 26S proteasome, the ubiquitination system and the spliceosome (Table S3). Because of their participation in a wide range of cellular functions, one could argue that these proteins were recovered simply as a result of non-specific protein interactions. It is therefore important to emphasize that they were identified either as significant in label-free quantification or as ASY1 sample-specific (absent from all control data sets), suggesting that at least some of the complex/pathway components were isolated on the basis of a specific interaction with the ASY1 meiotic complex. Indeed, an examination of the literature provides several indications of a close association between these particular protein complexes/pathways and meiotic chromatin. For example, HEI10, a mammalian RING domain protein with E3 ubiquitin-ligase activity (Ward *et al.*, 2007), and related proteins in budding yeast (Zip3), *Sordaria macrospora*, Arabidopsis and rice (HEI10) are required for CO formation, and have been shown to localize to discrete foci along meiotic chromosomes (Agarwal and Roeder, 2000; Chelysheva *et al.*, 2012; Wang *et al.*, 2012a; De Muyt

et al., 2014; Qiao *et al.*, 2014). Furthermore, studies indicate that Zip3 in yeast and RNF212 (its mammalian orthologue) and HEI10 in mouse mediate the recruitment of proteasomes to chromosome axes to regulate axis morphogenesis, homologue pairing, synapsis and meiotic recombination (Ahuja *et al.*, 2017; Rao *et al.*, 2017). Examination of the *Caenorhabditis elegans* germline provides further evidence that proteasome recruitment to the chromosome axes is an evolutionarily conserved feature of meiosis (Ahuja *et al.*, 2017).

Around 400 splicing-related proteins have been predicted or confirmed in Arabidopsis (Wang and Brendel, 2004; Koncz *et al.*, 2012). We identified 25 spliceosome-related proteins (Table S3), including PRL1 from the spliceosome-activating NineTeen Complex (NTC) core and three NTC-associated proteins, as defined by Monaghan *et al.*, (2009). There is increasing evidence of a role for the NTC in the coordination of DNA damage responses (reviewed in Koncz *et al.*, 2012). Interestingly, Ye *et al.* (2015) found that the RNA splicing pathway was extensively phosphorylated in rice anthers at around the time of meiosis (as indeed were the DNA synthesis and RdDM pathways, which are also well represented in our data). Although any association of spliceosome factors with meiotic chromosomes or a DNA repair role in meiosis remains to be established, our analysis may be a pointer in this direction.

Technical considerations

Although we anticipated that it might be feasible to use anthers to identify proteins that co-IP with ASY1, we were concerned that meiocytes represent only a small proportion of the tissue, potentially compromising our ability to identify less abundant meiotic proteins. Our data reveal that although several of the proteins with a prior confirmed role in meiosis could be identified from anthers, most were identified exclusively from the meiocyte-enriched samples. This was also the case for ICU2 and ASY4. Other experiments carried out in our lab suggest that certain meiotic proteins, notably ZYP1, are more easily recovered from intact anthers than extruded meiocytes, however, and hence there may be technical reasons favouring their detection from this tissue. In this context, it is interesting that during native meiotic protein extraction in budding yeast, the ZYP1 orthologue Zip1 appears to be far less stable than ASY1 and ASY3 homologues Hop1 and Red1 (Lin *et al.*, 2010). Thus, although it is clear that the additional effort in preparing meiocyte-enriched material was justified, it seems that analysis of both tissue types provides the most comprehensive picture.

The decision to retain the lower confidence group of all ASY1 sample-specific proteins satisfying the minimum identification threshold of two peptides appeared justified

for two reasons. First, several of them were found to have a meiotic role, and second, over 90% of the identified proteins formed a single PPI network; however, we cannot rule out that in some cases low-confidence proteins may have appeared to be sample specific by chance. There are several biological reasons why genuine interactors might appear with low confidence. Low-abundance proteins and proteins that form only transient interactions with the ASY1 complex may be under-represented in samples (extreme examples being protein kinases and phosphatases). Indirect interactors or proteins that form only weak interactions with the complex may also be more difficult to detect. Further sample enrichment to target a substage of prophase I may begin to address some of these limitations, as would targeting other proteins in the ASY1 interaction network in order to confirm and extend the network, enabling a comprehensive picture of meiotic interactions to emerge. This, together with ongoing technical improvements in sample preparation methods and MS analysis, should help to increase our ability to identify genuine protein interactors that currently lie at the borderline of detection.

EXPERIMENTAL PROCEDURES

Plant material, nucleic acid extraction and mutation site mapping

Arabidopsis thaliana ecotypes Columbia (0) and Enkheim-2 (En-2) and *B. oleracea* var. *albobolabra* A12Dhd were used for WT analysis. *Arabidopsis* seed stocks were obtained from the Nottingham *Arabidopsis* Stock Centre (<http://arabidopsis.info>). Plants were grown, *Arabidopsis* material was harvested and nucleic acid extractions were carried out as previously described (Higgins *et al.*, 2004). T-DNA insertion sites of mutant lines were confirmed by PCR and, in the case of *asy4*, by sequencing. The missense mutation of *icu2-1* was confirmed by sequencing and tetra-primer ARMS-PCR (Ye *et al.*, 2001). Primer details are listed in Appendix S2.

Co-immunoprecipitation analysis

Brassica meiotic tissue was collected as previously described (Sánchez-Morán *et al.*, 2005). Co-IP analysis was based on a previously described method, with minor modifications (Osman *et al.*, 2013). Full details of the procedure are available in Appendix S2.

Bioinformatic analysis

Brassica proteins were used to identify putative *A. thaliana* orthologues using best BLASTP 2.6.0 score (with an acceptance threshold of an E-value of $1e^{-5}$) against TAIR 10 protein sequences (<https://www.arabidopsis.org>). GO categorization of *A. thaliana* orthologues was carried out using the TAIR website (<https://www.arabidopsis.org/tools/bulk/go/index.jsp>). GO enrichment analysis was carried out using PANTHER accessed through the GO consortium website (<http://geneontology.org>). The KEGG pathway database was used to predict functional pathways for *Arabidopsis* orthologues (<http://www.kegg.jp/kegg/pathway.html>; Kanehisa *et al.*, 2016). PPI networks of *Arabidopsis* orthologues were generated using STRING 10.5 (<http://string-db.org>; Szklarczyk *et al.*,

2015), using default settings. The resulting network and protein description files were used to produce the networks in CYTOSCAPE 3.5.1 (<http://www.cytoscape.org>). Sequence alignments were carried out using CLUSTAL OMEGA (Sievers *et al.*, 2011) or EMBOSS NEEDLE (Rice *et al.*, 2000), accessed through the EMBL-EBI website (<https://www.ebi.ac.uk>).

Antibody production

The AtZYP1B C-terminal antibody was produced using a previously described procedure (Ferdous *et al.*, 2012) with primers ZYP1B-C-F and ZYP1B-C-R (Appendix S2). Polyclonal antiserum against the recombinant protein was raised in rabbit (Orygen Antibodies Ltd.; <http://www.orygen.co.uk>).

Cytological procedures

Cytological procedures were carried out as previously described (Higgins *et al.*, 2004). Antibodies were used as follows: anti-AtASY1 (rat, 1/1000 dilution) and anti-AtZYP1B-C (rabbit, 1/500 dilution). DNA was stained with $1 \mu\text{g ml}^{-1}$ 4',6-diamidino-2-phenylindole (DAPI) in Vectashield.

Yeast 2-hybrid analysis

Yeast 2-hybrid analysis was carried out as previously described (Ferdous *et al.*, 2012). Details of primers used for plasmid construction are presented in Appendix S2.

Statistical procedures

Fertility in WT and mutant plants was compared using single-factor ANOVA. Chi-square (χ^2) tests were carried out using GRAPHPAD PRISM 7 (<https://graphpad.com>) using Yate's correction.

ACCESSION NUMBERS

The MS proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository (Vizcaíno *et al.*, 2016), with the identifier PXD006042. The following lines were used for mutant analysis: At5g46070, SALK_016366; At3g52140, SALK_046271; At5g42220, SALK_151742; At5g59210, GABI_094G05; *mcm2* (At1g44900), SALK_023429; *icu2-1* (At5g67100), N329; *spo11-1-4* (At3g13170), WiscDsLox461-464J19 (Roberts 2010); *asy4* (At2g33793), SAIL_886_D04.

ACKNOWLEDGEMENTS

Horticultural/technical support was provided by Karen Staples and Steve Price, University of Birmingham. We wish to thank Professors Graham King (Southern Cross University, Australia), Isobel Parkin (University of Saskatchewan, Canada) and Xiaowu Wang (Chinese Academy of Agricultural Sciences, China) for their help with Brassica sequences. We are grateful to Dr Eugenio Sanchez-Moran and Professor George Bassel (University of Birmingham) for helpful discussions. Research leading to these results has received funding from the European Community's Seventh Framework Programme FP7/2007-2013 under grant agreement number KBBE-2009-222883, and from the Biotechnology and Biological Sciences Research Council (BBSRC) under grant number ERA-Caps-13 BB/M004902/1.

CONFLICTS OF INTEREST

The authors declare no known conflicts of interest.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article.

Figure S1. Targeting BoASY1 using an anti-AtASY1 antibody.

Figure S2. Protein sequence coverage.

Figure S3. Mutant analysis of three meiotic candidates.

Figure S4. Mutant analysis of At5g59210.

Figure S5. Mutant analysis of meiotic candidate MCM2.

Figure S6. Alignment of ASY4 (At2g33793) with ASY3 and mapping of T-DNA insertion SAIL_886_D04 in *asy4*.

Table S1. ASY1 sample-specific Brassica proteins with their putative Arabidopsis orthologues.

Table S2. Gene ontology enrichment analysis of ASY1 sample-specific proteins.

Table S3. Functional grouping of ASY1 sample-specific proteins.

Table S4. Summary of analysis of meiotic candidates.

Appendix S1. PPI network of ASY1 sample-specific proteins in Cytoscape format.

Appendix S2. Supporting experimental procedures.

REFERENCES

- Agarwal, S. and Roeder, G.S. (2000) Zip3 provides a link between recombination enzymes and synaptonemal complex proteins. *Cell*, **102**, 245–255.
- Ahuja, J.S., Sandhu, R., Mainpal, R., Lawson, C., Henley, H., Hunt, P.A., Yanowitz, J.L. and Börner, G.V. (2017) Control of meiotic pairing and recombination by chromosomally tethered 26S proteasome. *Science*, **355**, 408–411.
- Andersen, P.R., Tirian, L., Vunjak, M. and Brennecke, J. (2017) A heterochromatin-dependent transcription machinery drives piRNA expression. *Nature*, **549**, 54–59.
- Arabidopsis Interactome Mapping Consortium. (2011) Evidence for network evolution in an Arabidopsis interactome map. *Science*, **333**, 601.
- Armstrong, S.J., Caryl, A.P., Jones, G.H. and Franklin, F.C.H. (2002) Asy1, a protein required for meiotic chromosome synapsis, localizes to axis-associated chromatin in Arabidopsis and Brassica. *J. Cell Sci.* **115**, 3645.
- Attner, M.A., Miller, M.P., Ee, L.S., Elkin, S.K. and Amon, A. (2013) Polo kinase Cdc5 is a central regulator of meiosis I. *Proc. Natl Acad. Sci.* **110**, 14278–14283.
- Aya, K., Suzuki, G., Suwabe, K. et al. (2011) Comprehensive network analysis of anther-expressed genes in rice by the combination of 33 laser microdissection and 143 spatiotemporal microarrays. *PLoS ONE*, **6**, e26162.
- Barrero, J.M., González-Bayón, R., del Pozo, J.C., Ponce, M.R. and Micol, J.L. (2007) INCURVATA2 encodes the catalytic subunit of DNA polymerase α and interacts with genes involved in chromatin-mediated cellular memory in *Arabidopsis thaliana*. *Plant Cell*, **19**, 2822–2838.
- Berchowitz, L.E. and Copenhaver, G.P. (2010) Genetic interference: don't stand so close to me. *Curr. Genom.* **11**, 91–102.
- Bergerat, A., de Massy, B., Gadelle, D., Varoutas, P.-C., Nicolas, A. and Forterre, P. (1997) An atypical topoisomerase II from archaea with implications for meiotic recombination. *Nature*, **386**, 414–417.
- Bose, R., Manku, G., Culty, M. and Wing, S.S. (2014) Ubiquitin–proteasome system in spermatogenesis. In *Posttranslational Protein Modifications in the Reproductive System* (Sutovsky, P., ed.). New York, NY: Springer New York, pp. 181–213.
- Brar, G.A., Kiburz, B.M., Zhang, Y., Kim, J.-E., White, F. and Amon, A. (2006) Rec8 phosphorylation and recombination promote the step-wise loss of cohesins in meiosis. *Nature*, **441**, 532–536.
- Carballo, J.A., Johnson, A.L., Sedgwick, S.G. and Cha, R.S. (2008) Phosphorylation of the axial element protein Hop1 by Mec1/Tel1 ensures meiotic interhomolog recombination. *Cell*, **132**, 758–770.
- Chelysheva, L., Diallo, S., Vezon, D. et al. (2005) AtREC8 and AtSCC3 are essential to the monopolar orientation of the kinetochores during meiosis. *J. Cell Sci.* **118**, 4621.
- Chelysheva, L., Vezon, D., Chambon, A., Gendrot, G., Pereira, L., Lemhendi, A., Vrielynck, N., Le Guin, S., Novatchkova, M. and Grelon, M. (2012) The arabidopsis HEI10 is a new ZMM protein related to Zip3. *PLoS Genet.* **8**, e1002799.
- Chen, C., Farmer, A.D., Langley, R.J., Mudge, J., Crow, J.A., May, G.D., Huntley, J., Smith, A.G. and Retzel, E.F. (2010) Meiosis-specific gene discovery in plants: RNA-Seq applied to isolated Arabidopsis male meiocytes. *BMC Plant Biol.* **10**, 280.
- Collado-Romero, M., Alós, E. and Prieto, P. (2014) Unravelling the proteomic profile of rice meiocytes during early meiosis. *Front. Plant Sci.* **5**, 356.
- Couteau, F., Belzile, F., Horlow, C., Grandjean, O., Vezon, D. and Doutriaux, M.-P. (1999) Random chromosome segregation without meiotic arrest in both male and female meiocytes of a *dmc1* mutant of arabidopsis. *Plant Cell*, **11**, 1623–1634.
- Crismani, W., Girard, C., Froger, N., Pradillo, M., Santos, J.L., Chelysheva, L., Copenhaver, G.P., Horlow, C. and Mercier, R. (2012) FANCM limits meiotic crossovers. *Science*, **336**, 1588.
- Crismani, W., Portemer, V., Froger, N., Chelysheva, L., Horlow, C., Vrielynck, N. and Mercier, R. (2013) MCM8 is required for a pathway of meiotic double-strand break repair independent of DMC1 in *Arabidopsis thaliana*. *PLoS Genet.* **9**, e1003165.
- Cromer, L., Heyman, J., Touati, S. et al. (2012) OSD1 promotes meiotic progression via APC/C inhibition and forms a regulatory network with TDM and CYCA1;2/TAM. *PLoS Genet.* **8**, e1002865.
- De Muyt, A., Pereira, L., Vezon, D. et al. (2009) A high throughput genetic screen identifies new early meiotic recombination functions in *Arabidopsis thaliana*. *PLoS Genet.* **5**, e1000654.
- De Muyt, A., Zhang, L., Piolot, T., Kleckner, N., Espagne, E. and Zickler, D. (2014) E3 ligase Hei10: a multifaceted structure-based signaling molecule with roles within and beyond meiosis. *Genes Dev.* **28**, 1111–1123.
- Devshwar, P., Bovill, W.D., Sharma, R., Able, J.A. and Kapoor, S. (2011) Analysis of anther transcriptomes to identify genes contributing to meiosis and male gametophyte development in rice. *BMC Plant Biol.* **11**, 78.
- Doutriaux, M.P., Couteau, F., Bergounioux, C. and White, C. (1998) Isolation and characterisation of the RAD51 and DMC1 homologs from *Arabidopsis thaliana*. *Mol. Gen. Genet.* **257**, 283–291.
- Dukowicz-Szulce, S., Sundararajan, A., Mudge, J. et al. (2014) The transcriptome landscape of early maize meiosis. *BMC Plant Biol.* **14**, 118.
- Ferdous, M., Higgins, J.D., Osman, K. et al. (2012) Inter-homolog crossing-over and synapsis in arabidopsis meiosis are dependent on the chromosome axis protein AtASY3. *PLoS Genet.* **8**, e1002507.
- Fukuda, T., Pratto, F., Schimenti, J.C., Turner, J.M.A., Camerini-Otero, R.D. and Höög, C. (2012) Phosphorylation of chromosome core components may serve as axis marks for the status of chromosomal events during mammalian meiosis. *PLoS Genet.* **8**, e1002485.
- Girard, C., Crismani, W., Froger, N., Mazel, J., Lemhendi, A., Horlow, C. and Mercier, R. (2014) FANCM-associated proteins MHF1 and MHF2, but not the other Fanconi anemia factors, limit meiotic crossovers. *Nucl. Acids Res.* **42**, 9087–9095.
- Grelon, M., Vezon, D., Gendrot, G. and Pelletier, G. (2001) AtSPO11-1 is necessary for efficient meiotic recombination in plants. *EMBO J.* **20**, 589.
- Haering, C.H. and Jessberger, R. (2012) Cohesin in determining chromosome architecture. *Exp. Cell Res.* **318**, 1386–1393.
- He, Y., Wang, C., Higgins, J.D., Yu, J., Zong, J., Lu, P., Zhang, D. and Liang, W. (2016) MEIOTIC F-BOX is essential for male meiotic DNA double-strand break repair in rice. *Plant Cell*, **28**, 1879–1893.
- Higgins, J.D., Armstrong, S.J., Franklin, F.C.H. and Jones, G.H. (2004) The Arabidopsis MutS homolog AtMSH4 functions at an early step in recombination: evidence for two classes of recombination in Arabidopsis. *Genes Dev.* **18**, 2557–2570.
- Higgins, J.D., Sanchez-Moran, E., Armstrong, S.J., Jones, G.H. and Franklin, F.C.H. (2005) The Arabidopsis synaptonemal complex protein ZYP1 is required for chromosome synapsis and normal fidelity of crossing over. *Genes Dev.* **19**, 2488–2500.
- Huang, J., Cheng, Z., Wang, C., Hong, Y., Su, H., Wang, J., Copenhaver, G.P., Ma, H. and Wang, Y. (2015) Formation of interference-sensitive meiotic cross-overs requires sufficient DNA leading-strand elongation. *Proc. Natl Acad. Sci.* **112**, 12534–12539.
- Hughes, S.E. and Hawley, R.S. (2014) Topoisomerase II is required for the proper separation of heterochromatic regions during *Drosophila melanogaster* female meiosis. *PLoS Genet.* **10**, e1004650.

- Hyun, Y., Yun, H., Park, K., Ohr, H., Lee, O., Kim, D.-H., Sung, S. and Choi, Y. (2013) The catalytic subunit of arabidopsis DNA polymerase α ensures stable maintenance of histone modification. *Development*, **140**, 156.
- Jones, G.H. and Franklin, F.C.H. (2006) Meiotic crossing-over: obligation and interference. *Cell*, **126**, 246–248.
- Kalsotra, A. and Cooper, T.A. (2011) Functional consequences of developmentally regulated alternative splicing. *Nat. Rev. Genet.* **12**, 715–729.
- Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M. and Tanabe, M. (2016) KEGG as a reference resource for gene and protein annotation. *Nucl. Acids Res.* **44**, D457–D462.
- Katis, V.L., Lipp, J.J., Imre, R., Bogdanova, A., Okaz, E., Habermann, B., Mechtler, K., Nasmyth, K. and Zachariae, W. (2010) Rec8 phosphorylation by casein kinase 1 and Cdc7-Dbf4 kinase regulates cohesin cleavage by separase during meiosis. *Dev. Cell* **18**, 397–409.
- Kitajima, T.S., Sakuno, T., Ishiguro, K.-I., Iemura, S.-I., Natsume, T., Kawashima, S.A. and Watanabe, Y. (2006) Shugoshin collaborates with protein phosphatase 2A to protect cohesin. *Nature*, **441**, 46–52.
- Kleckner, N. (2006) Chiasma formation: chromatin/axis interplay and the role(s) of the synaptonemal complex. *Chromosoma*, **115**, 175.
- Klimyuk, V.I. and Jones, J.D.G. (1997) AtDMC1, the Arabidopsis homologue of the yeast DMC1 gene: characterization, transposon-induced allelic variation and meiosis-associated expression. *Plant J.* **11**, 1–14.
- Koncz, C., deJong, F., Villacorta, N., Szakonyi, D. and Koncz, Z. (2012) The spliceosome-activating complex: molecular mechanisms underlying the function of a pleiotropic regulator. *Front. Plant Sci.* **3**, 9.
- Korolev, A.V., Chan, J., Naldrett, M.J., Doonan, J.H. and Lloyd, C.W. (2005) Identification of a novel family of 70 kDa microtubule-associated proteins in Arabidopsis cells. *Plant J.* **42**, 547–555.
- Lai, Y.J., Lin, F.M., Chuang, M.J., Shen, H.J. and Wang, T.F. (2011) Genetic requirements and meiotic function of phosphorylation of the yeast axial element protein Red1. *Mol. Cell. Biol.* **31**, 912–923.
- Lam, W.S., Yang, X. and Makaroff, C.A. (2005) Characterization of *Arabidopsis thaliana* SMC1 and SMC3: evidence that AtSMC3 may function beyond chromosome cohesion. *J. Cell Sci.* **118**, 3037.
- Lambing, C., Osman, K., Nuntasootorn, K. et al. (2015) Arabidopsis PCH2 mediates meiotic chromosome remodeling and maturation of cross-overs. *PLoS Genet.* **11**, e1005372.
- Li, W., Chen, C., Markmann-Mulisch, U., Timofejeva, L., Schmelzer, E., Ma, H. and Reiss, B. (2004) The Arabidopsis AtRAD51 gene is dispensable for vegetative development but required for meiosis. *Proc. Natl Acad. Sci. USA*, **101**, 10596–10601.
- Li, X.M., Yu, C., Wang, Z.W., Zhang, Y.L., Liu, X.M., Zhou, D., Sun, Q.Y. and Fan, H.Y. (2013) DNA topoisomerase II is dispensable for oocyte meiotic resumption but is essential for meiotic chromosome condensation and separation in mice. *Biol. Reproduct.* **89**(118), 111.
- Libeau, P., Durandet, M., Granier, F., Marquis, C., Berthomé, R., Renou, J.P., Tacconat-Soubirou, L. and Horlow, C. (2011) Gene expression profiling of Arabidopsis meiocytes. *Plant Biol.* **13**, 784–793.
- Lin, F.M., Lai, Y.J., Shen, H.J., Cheng, Y.H. and Wang, T.F. (2010) Yeast axial-element protein, Red1, binds SUMO chains to promote meiotic interhomologue recombination and chromosome synapsis. *EMBO J.* **29**, 586.
- Liu, H. and Nonomura, K.I. (2016) A wide reprogramming of histone H3 modifications during male meiosis I in rice is dependent on the Argonaute protein MEL1. *J. Cell Sci.* **129**, 3553.
- Liu, J., Ren, X., Yin, H., Wang, Y., Xia, R., Wang, Y. and Gong, Z. (2010) Mutation in the catalytic subunit of DNA polymerase α influences transcriptional gene silencing and homologous recombination in Arabidopsis. *Plant J.* **61**, 36–45.
- Liu, Y., Deng, Y., Li, G. and Zhao, J. (2013) Replication factor C1 (RFC1) is required for double-strand break repair during meiotic homologous recombination in Arabidopsis. *Plant J.* **73**, 154–165.
- Liu, S., Liu, Y., Yang, X. et al. (2014) The Brassica oleracea genome reveals the asymmetrical evolution of polyploid genomes. *Nat. Comm.* **5**, 3930.
- Lu, Q., Dunn, R.L., Angeles, R. and Smith, G.D. (2002) Regulation of spindle formation by active mitogen-activated protein kinase and protein phosphatase 2A during mouse oocyte meiosis. *Biol. Reproduct.* **66**, 29–37.
- Luo, Q., Li, Y., Shen, Y. and Cheng, Z. (2014) Ten years of gene discovery for meiotic event control in rice. *J. Genet. Genom.* **41**, 125–137.
- Mercier, R., Mézard, C., Jenczewski, E., Macaisne, N. and Grelon, M. (2015) The molecular biology of meiosis in plants. *Ann. Rev. Plant Biol.* **66**, 297–327.
- Monaghan, J., Xu, F., Gao, M., Zhao, Q., Palma, K., Long, C., Chen, S., Zhang, Y. and Li, X. (2009) Two Prp19-like U-box proteins in the MOS4-associated complex play redundant roles in plant innate immunity. *PLoS Pathog.* **5**, e1000526.
- Nolt, J.K., Rice, L.M., Gallo-Ebert, C., Bisher, M.E. and Nickels, J.T. (2011) PP2ACdc55 is required for multiple events during meiosis I. *Cell Cycle*, **10**, 1420–1434.
- Nonomura, K.I., Morohoshi, A., Nakano, M., Eiguchi, M., Miyao, A., Hirochika, H. and Kurata, N. (2007) A germ cell-specific gene of the ARGO-NAUTE family is essential for the progression of premeiotic mitosis and meiosis during sporogenesis in rice. *Plant Cell*, **19**, 2583–2594.
- Oliver, C., Santos, J. and Pradillo, M. (2014) On the role of some ARGO-NAUTE proteins in meiosis and DNA repair in *Arabidopsis thaliana*. *Front. Plant Sci.* **5**, 177.
- Oliver, C., Santos, J.L. and Pradillo, M. (2016) Accurate chromosome segregation at first meiotic division requires AGO4, a protein involved in RNA-dependent DNA methylation in *Arabidopsis thaliana*. *Genetics*, **204**, 543.
- Olmedo-Monfil, V., Duran-Figueroa, N., Arteaga-Vazquez, M., Demesa-Arevalo, E., Aufran, D., Grimanelli, D., Slotkin, R.K., Martienssen, R.A. and Vielle-Calzada, J.-P. (2010) Control of female gamete formation by a small RNA pathway in Arabidopsis. *Nature*, **464**, 628–632.
- Orchard, S., Ammari, M., Aranda, B. et al. (2014) The MIntAct project—IntAct as a common curation platform for 11 molecular interaction databases. *Nucl. Acids Res.* **42**, D358–D363.
- Osman, K., Sanchez-Moran, E., Mann, S.C., Jones, G.H. and Franklin, F.C.H. (2009) Replication protein A (AtRPA1a) is required for class I crossover formation but is dispensable for meiotic DNA break repair. *EMBO J.* **28**, 394.
- Osman, K., Higgins, J.D., Sanchez-Moran, E., Armstrong, S.J. and Franklin, F.C.H. (2011) Pathways to meiotic recombination in *Arabidopsis thaliana*. *New Phytol.* **190**, 523–544.
- Osman, K., Roitinger, E., Yang, J., Armstrong, S., Mechtler, K. and Franklin, F.C.H. (2013) Analysis of meiotic protein complexes from arabidopsis and brassica using affinity-based proteomics. In *Plant Meiosis: Methods and Protocols* (Pawlowski, W.P., Grelon, M. and Armstrong, S., eds). Totowa, NJ: Humana Press, pp. 215–226.
- Page, S.L. and Hawley, R.S. (2004) The genetics and molecular biology of the synaptonemal complex. *Ann. Rev. Cell Devel. Biol.* **20**, 525–558.
- Parkin, I.A.P., Koh, C., Tang, H. et al. (2014) Transcriptome and methylome profiling reveals relics of genome dominance in the mesopolyploid *Brassica oleracea*. *Genome Biol.* **15**, R77.
- Penedos, A., Johnson, A.L., Strong, E., Goldman, A.S., Carballo, J.A. and Cha, R.S. (2015) Essential and checkpoint functions of budding yeast ATM and ATR during meiotic prophase are facilitated by differential phosphorylation of a meiotic adaptor protein, Hop1. *PLoS ONE*, **10**, e0134297.
- Pinna, L.A. (2002) Protein kinase CK2: a challenge to canons. *J. Cell Sci.* **115**, 3873.
- Pradillo, M., Knoll, A., Oliver, C., Varas, J., Corredor, E., Puchta, H. and Santos, J.L. (2015) Involvement of the cohesin cofactor PDS5 (SPO76) during meiosis and DNA repair in *Arabidopsis thaliana*. *Front. Plant Sci.* **6**, 1034.
- Qiao, H., Prasada Rao, H.B.D., Yang, Y. et al. (2014) Antagonistic roles of ubiquitin ligase HEI10 and SUMO ligase RNF212 regulate meiotic recombination. *Nat. Genet.* **46**, 194–199.
- Rao, H.B.D.P., Qiao, H., Bhatt, S.K. et al. (2017) A SUMO-ubiquitin relay recruits proteasomes to chromosome axes to regulate meiotic recombination. *Science*, **355**, 403–407.
- Rice, P., Longden, I. and Bleasby, A. (2000) EMBOS: the European molecular biology open software suite. *Trends Genet.* **16**, 276–277.
- Riedel, C.G., Katis, V.L., Katou, Y. et al. (2006) Protein phosphatase 2A protects centromeric sister chromatid cohesion during meiosis I. *Nature*, **441**, 53–61.
- Roberts, N.Y. (2010) Investigating the control of pairing and crossover formation in meiosis of *Arabidopsis thaliana*. Ph.D. Thesis. University of Birmingham, UK. <http://etheses.bham.ac.uk/1009>
- Rockmill, B. and Roeder, G.S. (1991) A meiosis-specific protein kinase homolog required for chromosome synapsis and recombination. *Genes Dev.* **5**, 2392–2404.

- Rogers, E., Bishop, J.D., Waddle, J.A., Schumacher, J.M. and Lin, R. (2002) The aurora kinase AIR-2 functions in the release of chromosome cohesion in *Caenorhabditis elegans* meiosis. *J. Cell Biol.* **157**, 219–229.
- Roitinger, E., Hofer, M., Köcher, T., Pichler, P., Novatchkova, M., Yang, J., Schlögelhofer, P. and Mechtler, K. (2015) Quantitative phosphoproteomics of the ataxia telangiectasia-mutated (ATM) and ataxia telangiectasia-mutated and rad3-related (ATR) dependent DNA damage response in *Arabidopsis thaliana*. *Mol. Cell Proteomics*, **14**, 556–571.
- Ross, K.J., Frasz, P., Armstrong, S.J., Vizir, I., Mulligan, B., Franklin, F.C. and Jones, G.H. (1997) Cytological characterization of four meiotic mutants of *Arabidopsis* isolated from T-DNA-transformed lines. *Chromosome Res.* **5**, 551–559.
- Sakuno, T. and Watanabe, Y. (2015) Phosphorylation of cohesin Rec11/SA3 by casein kinase 1 promotes homologous recombination by assembling the meiotic chromosome axis. *Dev. Cell*, **32**, 220–230.
- Sanchez-Moran, E., Santos, J.-L., Jones, G.H. and Franklin, F.C.H. (2007) ASY1 mediates AtDMC1-dependent interhomolog recombination during meiosis in *Arabidopsis*. *Genes Dev.* **21**, 2220–2233.
- Sánchez-Morán, E., Mercier, R., Higgins, J.D., Armstrong, S.J., Jones, G.H. and Franklin, F.C.H. (2005) A strategy to investigate the plant meiotic proteome. *Cytogenet. Genome Res.* **109**, 181–189.
- Schmid, R., Grellscheid, S.N., Ehrmann, I. et al. (2013) The splicing landscape is globally reprogrammed during male meiosis. *Nucl. Acids Res.* **41**, 10170–10184.
- Sievers, F., Wilm, A., Dineen, D. et al. (2011) Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mole. Syst. Biol.* **7**, 539.
- Singh, M., Goel, S., Meeley, R.B., Dantec, C., Parrinello, H., Michaud, C., Leblanc, O. and Grimanelli, D. (2011) Production of viable gametes without meiosis in maize deficient for an ARGONAUTE protein. *Plant Cell*, **23**, 443–458.
- Smith, S.J., Osman, K. and Franklin, F.C.H. (2014) The condensin complexes play distinct roles to ensure normal chromosome morphogenesis during meiotic division in *Arabidopsis*. *Plant J.* **80**, 255–268.
- Smyth Gordon, K. (2004) Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat. Appl. Genet. Mol. Biol.* **3**, 1–25.
- Sprink, T. and Hartung, F. (2014) The splicing fate of plant SPO11 genes. *Front. Plant Sci.* **5**, 214.
- Szklarczyk, D., Franceschini, A., Wyder, S. et al. (2015) STRING v10: protein–protein interaction networks, integrated over the tree of life. *Nucl. Acids Res.* **43**, D447–D452.
- Tang, X., Zhang, Z.-Y., Zhang, W.-J., Zhao, X.-M., Li, X., Zhang, D., Liu, Q.-Q. and Tang, W.-H. (2010) Global gene profiling of laser-captured pollen mother cells indicates molecular pathways and gene subfamilies involved in rice meiosis. *Plant Physiol.* **154**, 1855–1870.
- Tang, A., Shi, P., Song, A., Zou, D., Zhou, Y., Gu, P., Huang, Z., Wang, Q., Lin, Z. and Gao, X. (2016) PP2A regulates kinetochore-microtubule attachment during meiosis I in oocyte. *Cell Cycle*, **15**, 1450–1461.
- Traven, A. and Heierhorst, J. (2005) SQ/TQ cluster domains: concentrated ATM/ATR kinase phosphorylation site regions in DNA-damage-response proteins. *BioEssays*, **27**, 397–407.
- Vizcaino, J.A., Csordas, A., del-Toro, N., Dienes, J.A., Griss, J., Lavidas, I., Mayer, G., Perez-Riverol, Y., Reisinger, F. and Ternent, T. (2016) 2016 update of the PRIDE database and its related tools. *Nucl. Acids Res.* **44**, D447–D456.
- Wang, B.-B. and Brendel, V. (2004) The ASRG database: identification and survey of *Arabidopsis thaliana* genes involved in pre-mRNA splicing. *Genome Biol.* **5**, R102.
- Wang, Y. and Yang, M. (2006) The *Arabidopsis* SKP1-LIKE1 (ASK1) protein acts predominantly from leptotene to pachytene and represses homologous recombination in male meiosis. *Planta*, **223**, 613–617.
- Wang, X., Wang, H., Wang, J. et al. (2011) The genome of the mesopolyploid crop species *Brassica rapa*. *Nat. Genet.* **43**, 1035–1039.
- Wang, K., Wang, M., Tang, D., Shen, Y., Miao, C., Hu, Q., Lu, T. and Cheng, Z. (2012a) The role of rice HEI10 in the Formation of meiotic crossovers. *PLoS Genet.* **8**, e1002809.
- Wang, Y., Cheng, Z., Huang, J., Shi, Q., Hong, Y., Copenhagen, G.P., Gong, Z. and Ma, H. (2012b) The DNA replication factor RFC1 is required for interference-sensitive meiotic crossovers in *Arabidopsis thaliana*. *PLoS Genet.* **8**, e1003039.
- Wang, H., You, C., Chang, F., Wang, Y., Wang, L., Qi, J. and Ma, H. (2014) Alternative splicing during *Arabidopsis* flower development results in constitutive and stage-regulated isoforms. *Front. Genet.* **5**, 25.
- Ward, J.O., Reinholdt, L.G., Motley, W.W. et al. (2007) Mutation in mouse Hei10, an E3 ubiquitin ligase, disrupts meiotic crossing over. *PLoS Genet.* **3**, e139.
- Yang, H., Lu, P., Wang, Y. and Ma, H. (2011) The transcriptome landscape of *Arabidopsis* male meiocytes from high-throughput sequencing: the complexity and evolution of the meiotic process. *Plant J.* **65**, 503–516.
- Ye, S., Dhillon, S., Ke, X., Collins, A.R. and Day, I.N.M. (2001) An efficient procedure for genotyping single nucleotide polymorphisms. *Nucleic Acids Res.* **29**, e88.
- Ye, J., Zhang, Z., Long, H., Zhang, Z., Hong, Y., Zhang, X., You, C., Liang, W., Ma, H. and Lu, P. (2015) Proteomic and phosphoproteomic analyses reveal extensive phosphorylation of regulatory proteins in developing rice anthers. *Plant J.* **84**, 527–544.
- Yu, C., Ji, S.-Y., Sha, Q.-Q., Sun, Q.-Y. and Fan, H.-Y. (2015) CRL4-DCAF1 ubiquitin E3 ligase directs protein phosphatase 2A degradation to control oocyte meiotic maturation. *Nat. Commun.* **6**, 8017.
- Zhang, L., Wang, S., Yin, S., Hong, S., Kim, K.P. and Kleckner, N. (2014) Topoisomerase II mediates meiotic crossover interference. *Nature*, **511**, 551–556.
- Zhang, Z., Hu, M., Feng, X., Gong, A., Cheng, L. and Yuan, H. (2017) Proteomes and phosphoproteomes of anther and pollen: availability and progress. *Proteomics*, **17**, 1770151. <https://doi.org/10.1002/pmic.201600458>
- Zhao, D., Yang, X., Quan, L., Timofejeva, L., Rigel, N.W., Ma, H. and Makaroff, C.A. (2006) ASK1, a SKP1 homolog, is required for nuclear reorganization, presynaptic homolog juxtaposition and the proper distribution of cohesin during meiosis in *Arabidopsis*. *Plant Mol. Biol.* **62**, 99–110.
- Zickler, D. and Kleckner, N. (2016) A few of our favorite things: pairing, the bouquet, crossover interference and evolution of meiosis. *Sem. Cell Dev. Biol.* **54**, 135–148.