

UNIVERSITY OF BIRMINGHAM

Research at Birmingham

Thickened endometrium in asymptomatic postmenopausal women - determining an optimum threshold for prediction of atypical hyperplasia and cancer

Ghoubara, Ahmed; Emovon, Emmanuel; Sundar, Sudha; Ewies, Ayman

DOI:

[10.1080/01443615.2018.1458081](https://doi.org/10.1080/01443615.2018.1458081)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Ghoubara, A, Emovon, E, Sundar, S & Ewies, A 2018, 'Thickened endometrium in asymptomatic postmenopausal women - determining an optimum threshold for prediction of atypical hyperplasia and cancer' *Journal of Obstetrics and Gynaecology*. <https://doi.org/10.1080/01443615.2018.1458081>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is an Accepted Manuscript of an article published by Taylor & Francis in *Journal of Obstetrics and Gynaecology* on 03/06/2018, available online: <http://www.tandfonline.com/10.1080/01443615.2018.1458081>.

Checked 03/07/2018.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.


Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



**Thickened endometrium without postmenopausal bleeding
– determining an optimum threshold for prediction of
hyperplasia and cancer**

Journal:	<i>Journal of Obstetrics and Gynaecology</i>
Manuscript ID	CJOG-2017-0907
Manuscript Type:	Original Article
Date Submitted by the Author:	19-Dec-2017
Complete List of Authors:	Ghoubara, Ahmed; University of Birmingham, Institute of Cancer and Genomic Sciences; Sandwell and West Birmingham Hospitals NHS Trust, Pan Birmingham Gynaecological Cancer Centre; Aswan University, Obstetrics and Gynaecology Emovon, Emmanuel; Sandwell and West Birmingham Hospitals NHS Trust Sundar, Sudha; University of Birmingham, Institute of Cancer and Genomic Sciences; Sandwell and West Birmingham Hospitals NHS Trust, Pan Birmingham Gynaecological Cancer Centre Ewies, Ayman; Sandwell and West Birmingham Hospitals NHS Trust, Pan Birmingham Gynaecological Cancer Centre; University of Birmingham, Institute of Cancer and Genomic Sciences
Keywords:	Endometrial cancer, Endometrial hyperplasia, Endometrial polyp, Endometrial thickness, Postmenopausal bleeding

SCHOLARONE™
Manuscripts

1
2
3 **Thickened endometrium without postmenopausal bleeding – determining an optimum**
4 **threshold for prediction of hyperplasia and cancer**
5
6
7
8
9

10 Ahmed Ghoubara^{a,b,c}, Emmanuel Emovon^c, Sudha Sundar^{a,c}, Ayman Ewies^{a,c}

11
12
13 ^a*Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham, UK*

14
15
16 ^b*Obstetrics and Gynecology department, Aswan University, Aswan, Egypt*

17
18
19 ^c*Gynaecology Department, Sandwell and West Birmingham Hospitals NHS Trust, Birmingham,*
20
21 *UK*

22
23
24
25
26 **Authors biography**

- 27
28
29 1- Ahmed Ghoubara, MSc, Research Fellow in The Institute of Cancer and Genomic Sciences,
30 University of Birmingham, UK and Sandwell and West Birmingham Hospitals NHS
31 Teaching Trust, Birmingham, UK and Assistant Lecturer in Gynecology, Aswan University,
32 Egypt. E-mail: a.ghoubara@icloud.com. ORCID: <http://orcid.org/0000-0002-9330-2344>.
33
34 Postal Address: Pan Birmingham Gynecological Cancer Center, Birmingham City Hospital,
35 Dudley Road, Birmingham B18 7QH, UK. Tel: 00441215074042.
36
37
38 2- Emmanuel Emovon, MBBS, MRCOG, Clinical Fellow in Obstetrics and Gynecology,
39 Sandwell and West Birmingham Hospitals NHS Teaching Trust, UK. E-mail:
40 Emmanuel.emovon@nhs.net. Postal Address: Pan Birmingham Gynecological Cancer
41 Center, Birmingham City Hospital, Dudley Road, Birmingham B18 7QH, UK. Tel:
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
00441215074042.

- 1
2
3- Sudha Sundar, MPhil, MRCOG, Senior Lecturer in Gynecological Oncology and Honorary
4
5 Consultant, Institute of Cancer and Genomic Sciences, University of Birmingham, UK,
6
7 Sandwell and West Birmingham Hospitals NHS Teaching Trust, Birmingham, UK. E-mail:
8
9 Sudha.sundar@nhs.net. Postal Address: Pan Birmingham Gynecological Cancer Center,
10
11 Birmingham City Hospital, Dudley Road, Birmingham B18 7QH, UK. Tel:
12
13 00441215074042.
14
15
16
17 4- Ayman A A Ewies, MD, FRCOG, Consultant in Gynecology, Sandwell and West
18
19 Birmingham Hospitals NHS Teaching Trust, UK and Honorary Senior Lecturer, The College
20
21 of Medical & Dental Sciences, University of Birmingham, UK. aymanewies@hotmail.com.
22
23
24
25

26 **Corresponding Author**

27
28 Mr. Ayman Ewies

29
30
31 Postal Address: Pan Birmingham Gynecological Cancer Center, Birmingham City Hospital,
32
33 Dudley Road, Birmingham B18 7QH, UK
34
35

36 E-mail: aymanewies@hotmail.com; ayman.ewies@nhs.net

37
38
39 Tel: 00441215074042

40
41
42 Fax: 00441215075680
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

There is lack of consensus about the significance and management of an incidental finding of endometrial thickness (ET) >4mm on transvaginal ultrasound scan in postmenopausal women without postmenopausal bleeding (PMB). The data of 1995 consecutive women attending PMB clinic were collected prospectively; of them 81 (4.1%) were referred because of ET>4mm without PMB. The prevalence of endometrial atypical hyperplasia and cancer was 4/81 (4.9%) and polyp was 20/81 (24.7%). On using Receiver Operator Characteristic curve, the diagnosis of endometrial atypical hyperplasia and cancer using ET threshold of ≥ 10 mm had sensitivity of 100% (95% CI=40-100%), specificity of 60% (95% CI=48-71%) with AUC=0.8 (95% CI=0.66-0.93), $p < 0.04$. For the 35 women with ET ≥ 10 mm, the prevalence of endometrial atypical hyperplasia and cancer was 4/35 (11.4%) and benign endometrial polyps was 16/35 (45.7%). The use of ≥ 10 mm ET threshold to prompt investigations did not miss any case of endometrial atypical hyperplasia or cancer.

Impact statement:

What is already known about this topic:

Unlike women with PMB in whom endometrial thickness (ET) ≤ 4 mm is considered as low risk for endometrial hyperplasia and cancer, in postmenopausal women without PMB, the threshold that separates normal from pathologically thickened endometrium has not been standardized. A decision-analysis study in a hypothetical cohort found that ET threshold of 11mm yields a similar separation as ≤ 4 mm in those with PMB.

What does this paper add?

The study uses prospectively collected data from consecutive patients using standardized format, thus minimizing bias from incomplete data. This study is the third prospective series in the literature to address the dilemma of management of asymptomatic postmenopausal women with incidental finding of thickened endometrium. It showed that the prevalence of endometrial atypical hyperplasia and cancer is high enough to justify investigation and for the clinical problem not to be trivialized. All cases of endometrial atypical hyperplasia and cancer had endometrial thickness of ≥ 10 mm.

Clinical implications and future research:

Our data strengthen the current body of literature to help development of clinical practice guidelines about the management work-up. However, a well-designed multi-centre large prospective study is required to confirm the findings since most studies in the literature are either retrospective or small.

1
2
3 **Keywords**
4

5 Endometrial cancer; endometrial hyperplasia; endometrial polyp; endometrial thickness;
6
7
8 postmenopausal bleeding
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Introduction

Trans-vaginal ultrasound scan (TVS) is frequently performed in postmenopausal women referred for a variety of symptoms, such as pelvic pain or bloating. Although concerns regarding endometrial pathology may not have been the indication for the test, endometrial thickness (ET) is often measured. The incidental finding of $ET > 4\text{mm}$ in the absence of postmenopausal bleeding (PMB) is a common cause of referral to secondary care, and can lead to a cascade of interventions to exclude endometrial hyperplasia or cancer (Jade et al., 2016). There is lack of consensus amongst gynecologists about its significance, and there are no evidence-based guidelines to inform clinical practice as how these women are best managed, leading to wide variations in practice (Gambacciani et al., 2004, Giannella et al., 2014).

Ninety percent of women with endometrial cancer present with PMB, and women with PMB have a 5-10% chance of having endometrial cancer (Bachmann et al., 2003, ACOG, 2009). However; it is estimated that up to 15% of endometrial cancers occur in women without PMB (NIH National Cancer Institute, 2000). It is well established that $ET \leq 4\text{mm}$ in women with PMB is associated $<1\%$ risk of endometrial cancer; therefore, these women are usually reassured without the need for further investigations (ACOG, 2009, RCOG, 2016). However; in postmenopausal women without PMB, the threshold that separates normal from pathologically thickened endometrium has not been standardized. Smith-Bindman *et al* performed a decision analysis study, in a theoretical cohort of 10 000 postmenopausal women aged ≥ 50 years using published and unpublished data, to determine the ET threshold that should be considered abnormal in asymptomatic postmenopausal women. They found that in a postmenopausal woman with PMB, the risk of endometrial cancer is approximately 0.07% if the endometrium is

1
2
3 thin (≤ 5 mm) and 7.3% if her endometrium is thick (> 5 mm). In a postmenopausal woman
4
5 without PMB, an 11mm threshold yields a similar separation between women who are at high
6
7 risk and low risk for endometrial cancer. The risk of cancer is approximately 0.002% if the
8
9 endometrium is thin (≤ 11 mm) and 6.7% if the endometrium is thick (> 11 mm). If an ET
10
11 threshold of 4 mm was used to define an abnormal test result, as in women with PMB, the
12
13 number of false-positive test results would far outnumber the true-positive test results (Smith-
14
15 Bindman et al., 2004).
16
17
18
19
20
21
22

23 In addition, two previous studies found that endometrial polyps are the commonest pathology
24
25 encountered in postmenopausal women with asymptomatic increase in ET. The reported
26
27 prevalence varied between 34%-73%, and hysteroscopy was recommended as the first line
28
29 investigation tool (Farfaras et al., 2011, Schmidt et al., 2009).
30
31
32
33
34
35

36 Given the current controversy surrounding the significance and management of thickened
37
38 endometrium of > 4 mm in postmenopausal women without PMB, this study was conducted to (1)
39
40 quantify the prevalence of endometrial hyperplasia, cancer and polyps, (2) determine the
41
42 optimum ET threshold that should be considered abnormal and trigger further investigations. The
43
44 aim was to produce data to help the development of clinical practice guidelines about the
45
46 management work-up.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Materials and methods

The routine demographic and clinical data of the PMB clinic was collected prospectively in a password-protected Microsoft Access database, and then was extracted anonymously and compiled on a Microsoft Excel spreadsheet (Microsoft corporation 2010, Redmond, WA, USA). The database contained the details of 1995 consecutive women referred to Sandwell and West Birmingham Hospitals NHS Trust, West Midlands, UK between 1st January 2011 and 31st January 2015. The study group included 81 (4.1%) postmenopausal women with incidental finding of ET >4mm in the absence of PMB.

The TVS and histopathology reports were accessed using the electronic Clinical Data Archive (CDA, 2007) System that was developed by the IT department of the Trust as a repository of patients' administrative and clinical data. The case notes were reviewed for women undergoing hysteroscopic (diagnostic or therapeutic) procedure to obtain the operation findings.

All postmenopausal women with incidental finding of ET >4mm without PMB were offered Pipelle endometrial biopsy. However, the decision to perform hysteroscopy was individualized after the discussion between the patient and consultant. Women were categorized according to the investigation results into: (i) Group 1: benign endometrium (including benign endometrial polyps), and (ii) Group 2: endometrial atypical hyperplasia and cancer. Endometrial atypical hyperplasia and cancer were combined as a single disease. This is because of the high rate of undercall and progression to cancer when atypical hyperplasia is found (Smith et al., 2014, Gallos et al., 2013).

Statistical analysis

Statistical analyses were carried out using IBM® SPSS® Statistics for windows software version 20 (International Business Machines (IBM) corporation, Armonk, NYC, USA). Continuous variable indices are presented as mean with standard deviation or as median with inter-quartile range (IQR) as distribution demands.

Receiver operator characteristic (ROC) curve was used to identify the ET cut-off point for pathological endometrium. From the ROC curve, we calculated the area under the curve (AUC) with 95% confidence interval (95%CI), *p* value, sensitivity and specificity for the relevant cut-off point. The following equation was used to define the optimum cut-off point (Indrayan, 2013):

$$\sqrt{(1 - \text{sensitivity})^2 + (1 - \text{specificity})^2}$$

Fisher's Exact test was used to find the statistical association between the ET cut-off point and the outcome.

Data were collected as part of the routine investigations and treatment, and the project was considered as "service evaluation"; therefore, ethics approval was not deemed necessary. Service evaluation may not require ethical approval in the UK (NHS Health Research Authority, 2011, UCL Research Ethics Committee 2015).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Results

The prevalence of endometrial atypical hyperplasia and cancer was 4/81 (4.9%) and benign endometrial polyps was 20/81 (24.7%) in postmenopausal women with incidental finding of ET >4mm without PMB. Women's characteristics and investigations findings are shown in table 1 and table 2, respectively. Three women were diagnosed with endometrial cancer. The first was 86-year-old, body mass index (BMI) was not measured as she was wheel-chair dependent, and the ET was 25mm. The second was 75-year-old, BMI was 26, and the ET was 12mm. The third was 83-year-old, BMI was 26, and the ET was 10mm. One woman was diagnosed with endometrial atypical hyperplasia who was 57-year-old, BMI was 33, and the ET was 16mm. None of these four women was a user of HRT or Tamoxifen.

The ROC curve (Figure 1) identified the ET threshold for diagnosing endometrial atypical hyperplasia and cancer as ≥ 10 mm with sensitivity of 100% (95% CI=40-100%), specificity of 60% (95% CI=48-71%), AUC=0.8 (95% CI=0.66-0.93), $p < 0.04$.

The outcome of investigations using ≥ 10 mm endometrial thickness threshold is summarized in table 3. The prevalence of endometrial atypical hyperplasia and cancer was significantly high with ET ≥ 10 mm, $p = 0.03$. For the 35 women with ET ≥ 10 mm, the prevalence of endometrial atypical hyperplasia and cancer was 4/35 (11.4%) and benign endometrial polyps was 16/35 (45.7%).

Discussion

To our knowledge, this is the third prospective study to address the dilemma of management of postmenopausal women referred to secondary care with thickened endometrium without PMB.

The prevalence of endometrial atypical hyperplasia and cancer was 4.9% and benign endometrial polyps was 24.7%. The ROC curve identified the ET cut-off level for endometrial atypical hyperplasia and cancer to be ≥ 10 mm.

In their prospective series, Schmidt *et al* reported similar prevalence of endometrial atypical hyperplasia and cancer (4.9%), but higher prevalence of endometrial polyps (73.3%) (Schmidt *et al.*, 2009), while Giannella *et al* found the prevalence to be 2.1% and 34%, respectively (Giannella *et al.*, 2014). The other three published series in the past decade were retrospective carrying the risk of bias from heterogeneity and missing data. They reported wide variations in the prevalence of endometrial atypical hyperplasia and cancer (1.3%-13.2%) and endometrial polyps (27.2%-59.1%) (Famuyide *et al.*, 2014, Saatli *et al.*, 2014, Yasa *et al.*, 2016). In particular, the high prevalence of endometrial atypical hyperplasia and cancer (70/530, 13.2%) in Saatli *et al* study is hard to explain since it is much higher than the reported prevalence of the disease in women with PMB which varied in the literature between 5-10% (ACOG, 2009, Bachmann *et al.*, 2003).

Ninety percent of women with endometrial cancer present with PMB as previously mentioned. (Bachmann *et al.*, 2003, ACOG, 2009). Nonetheless, there is likely a preclinical phase during which some cancers might be detectable prior to the development of symptoms. In addition,

1
2
3 some cancers do not present with bleeding until they progressed beyond Stage I (Smith et al.,
4 2014). The National Institute for Health and Care Excellence (NICE) sets a 3% positive
5 predictive value to trigger referrals to secondary care for investigations for suspected cancer
6 (NICE, 2015). In this study, the endometrial atypical hyperplasia and cancer prevalence of
7 11.4% indicates that referral of postmenopausal women with incidental finding of thickened
8 endometrium ≥ 10 mm is warranted.
9
10
11
12
13
14
15
16
17
18
19

20
21 Giannella *et al* tested the diagnostic accuracy of various ET cut-off values by comparing
22 histological and hysteroscopic findings in asymptomatic postmenopausal women with ET >4mm.
23 They found that no ET cut-off values had optimal diagnostic accuracy, but an ET cut-off value
24 >10 mm did not miss any cases of endometrial atypical hyperplasia or cancer. At this cut-off
25 value, the atypical hyperplasia and cancer rate was 9.4%. On using ET cut-off value >4 mm,
26 97% of performed hysteroscopies revealed a benign intra-uterine pathology (Giannella et al.,
27 2014). Smith-Bindman *et al* concluded after the decision-analysis study that no cut-off is perfect,
28 and cancer will be missed no matter what cut-off is used. However, using a threshold of 11 mm
29 seems to provide an acceptable trade-off between cancer detection and unnecessary biopsies
30 prompted by an incidental finding (Smith-Bindman et al., 2004). In this study, the ≥ 10 mm
31 threshold identified by the ROC curve had sensitivity of 100%, which would be more reassuring
32 to clinicians than 11mm cut-off.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51
52 Our data strengthens the current body of literature to help development of clinical practice
53 guidelines about the management work-up. The data were collected prospectively, consecutively,
54
55
56
57
58
59
60

1
2
3 and in a standardized fashion, minimizing bias from incomplete data. We included a
4
5 homogeneous cohort of postmenopausal women referred to secondary care with incidental
6
7 finding of ET >4mm without PMB. Nonetheless, given the small sample size, we could not
8
9 include in the analysis women's characteristics and risk factors for endometrial hyperplasia and
10
11 cancer such as age and body mass index.
12
13

14
15
16
17
18 A well-designed multi-center large prospective study is required to reach consensus about the
19
20 optimum ET cut-off to initiate investigation in asymptomatic postmenopausal women with
21
22 incidental finding of thickened endometrium. This study showed that the prevalence of
23
24 pathology is high enough to justify investigation and for the clinical problem not to be trivialized
25
26 by women and doctors alike. The use of ≥ 10 mm ET threshold to prompt investigations may be
27
28 acceptable since it is unlikely to miss endometrial atypical hyperplasia or cancer.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Funding

No fund was obtained for this study but the research fellowship of AG was funded by Aswan University, Egypt, and the Egyptian Cultural Centre and Education Bureau in London, UK.

For Peer Review Only

1
2
3 **Conflict of interest**
4

5 The authors declare no conflicts to interest.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- American College of Obstetricians and Gynecologists (ACOG) 2009. ACOG Committee Opinion No. 440: The Role of Transvaginal Ultrasonography in the Evaluation of Postmenopausal Bleeding. *Obstetrics and Gynecology*, 114, 409-11.
- Bachmann, L. M., ter Riet, G., Clark, T. J., Gupta, J. K. & Khan, K. S. 2003. Probability analysis for diagnosis of endometrial hyperplasia and cancer in postmenopausal bleeding: an approach for a rational diagnostic workup. *Acta Obstetrica et Gynecologica Scandinavica*, 82, 564-9.
- Famuyide, A. O., Breitkopf, D. M., Hopkins, M. R. & Laughlin-Tommaso, S. K. 2014. Asymptomatic Thickened Endometrium in Postmenopausal Women: Malignancy Risk. *Journal of Minimally Invasive Gynecology*, 21, 782-786.
- Farfaras, A., Cumashi, E., Soulos, P., Pavlakis, E., Drakontaidis, A. & Skolarikos, P. 2011. Asymptomatic, postmenopausal women with sonographically thickened endometrium. Pathologic findings following hysteroscopic evaluation. *Climacteric*, 14, 129.
- Gallos, I. D., Krishan, P., Shehmar, M., Ganesan, R. & Gupta, J. K. 2013. Relapse of endometrial hyperplasia after conservative treatment: a cohort study with long-term follow-up. *Human Reproduction*, 28, 1231-6.
- Gambacciani, M., Monteleone, P., Ciaponi, M., Sacco, A. & Genazzani, A. R. 2004. Clinical usefulness of endometrial screening by ultrasound in asymptomatic postmenopausal women. *Maturitas*, 48, 421-4.
- Giannella, L., Mfuta, K., Setti, T., Boselli, F., Bergamini, E. & Cerami, L. B. 2014. Diagnostic accuracy of endometrial thickness for the detection of intra-uterine pathologies and appropriateness of performed hysteroscopies among asymptomatic postmenopausal women. *European Journal of Obstetrics Gynecology and Reproductive Biology*, 177, 29-33.

- 1
2
3 Indrayan, A. 2013. Search for the best threshold of continuous test: ROC curve. *Medical*
4
5 *Biostatistics*. Third ed.: CRC press Taylor & Francis group.
6
7
8 Jade, E., Hoyte, H., Naji, O. & Mamta, P. 2016. Histological outcomes at hysteroscopy in
9
10 postmenopausal women with asymptomatic endometrial thickening: A 6 year view from a DGH.
11
12 *Gynecological Surgery*, 13 (1 Supplement 1), S79.
13
14
15 National Health System (NHS) Health Research Authority (HRA). 2011. *Governance*
16
17 *Arrangements for Research Ethics Committees (GAfREC)* [Online]. UK Health Departments.
18
19 Available: <http://www.hra.nhs.uk> [Accessed 05 December 2017].
20
21
22 National Institutes of Health (NIH) National Cancer Institute (NCI). 2000. *Surveillance,*
23
24 *Epidemiology, and End Results Program (SEER)* [Online]. Bethesda,MD. Available:
25
26 <https://seer.cancer.gov/> [Accessed 05 December 2017].
27
28
29 Royal College of Obstetricians and Gynaecologists (RCOG). 2016. *Management of Endometrial*
30
31 *Hyperplasia Green-Top Guideline No. 67* [Online]. Available: <https://www.rcog.org.uk>
32
33 [Accessed 05 December 2017].
34
35
36 Saatli, B., Yildirim, N., Olgan, S., Koyuncuoglu, M., Emekci, O. & Saygili, U. 2014. The role of
37
38 endometrial thickness for detecting endometrial pathologies in asymptomatic postmenopausal
39
40 women. *Australian and New Zealand Journal of Obstetrics and Gynaecology*, 54, 36-40.
41
42
43 Schmidt, T., Breidenbach, M., Nawroth, F., Mallmann, P., Beyer, I. M., Fleisch, M. C., et al.
44
45 2009. Hysteroscopy for asymptomatic postmenopausal women with sonographically thickened
46
47 endometrium. *Maturitas*, 62, 176-178.
48
49
50 Smith-Bindman, R., Weiss, E. & Feldstein, V. 2004. How thick is too thick? When endometrial
51
52 thickness should prompt biopsy in postmenopausal women without vaginal bleeding. *Ultrasound*
53
54 *in Obstetrics and Gynecology*, 24, 558-65.
55
56
57
58
59

1
2
3 Smith, P. P., O'Connor, S., Gupta, J. & Clark, T. J. 2014. Recurrent postmenopausal bleeding: a
4 prospective cohort study. *Journal of Minimally Invasive Gynecology*, 21, 799-803.
5

6
7 University College of London (UCL) Research Ethics Committee. 2015. *Exemptions* [Online].
8

9 Available: <https://ethics.grad.ucl.ac.uk/exemptions.php> [Accessed 28 August 2017].
10

11
12 Yasa, C., Dural, O., Bastu, E., Ugurlucan, F. G., Nehir, A. & Iyibozkurt, A. C. 2016. Evaluation
13 of the diagnostic role of transvaginal ultrasound measurements of endometrial thickness to detect
14 endometrial malignancy in asymptomatic postmenopausal women. *Archives of Gynecology and*
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Obstetrics, 294, 311-6.

Tables and figures legends

Table 1: Characteristics of postmenopausal women referred with incidental finding of ET >4mm without PMB ($n=81$)

Table 2: Investigations findings of postmenopausal women referred with incidental finding of ET >4mm without PMB ($n=81$)

Table 3: The outcome of investigations using ≥ 10 mm endometrial thickness threshold

Figure 1: Endometrial thickness receiver operator characteristic (ROC) curve for endometrial atypical hyperplasia and cancer. Arrow indicates the point where the 10mm endometrial thickness lies on the curve; Area under the curve (AUC) for the endometrial thickness cut-off ≥ 10 mm is 0.8, 95% confidence interval (CI)=0.66-0.93, $p=0.04$, sensitivity = 100% (95% CI=40-100%), specificity = 60% (95% CI=48-71%)

Table 2: Investigations findings of postmenopausal women referred with incidental finding of ET >4mm without PMB (*n*=81)

Variable	
ET in mm, mean (\pm SD)	10.3 (6.3)
Benign endometrium, <i>n</i> (%)	77 (95.1%)
Endometrium with no hyperplasia or cancer, <i>n</i> (%)	57 (70.4%)
Benign endometrial polyp, <i>n</i> (%), 95%CI)	20 (24.7%, 16-35.7%)
Atypical endometrial hyperplasia and cancer, <i>n</i> (%), 95%CI)	4 (4.9%, 1.6-13%)

ET=endometrial thickness

CI=confidence interval

SD=standard deviation


Table 3: The outcome of investigations using ≥ 10 mm endometrial thickness threshold

Variable	ET ≥ 10mm	ET < 10mm	<i>p</i> value
	<i>n</i> (%)	<i>n</i> (%)	
Benign endometrium			
Endometrium with no hyperplasia or cancer	15 (42.9%)	42 (91.3%)	
Benign endometrial polyp	16 (45.7%)	4 (8.7%)	
Endometrial atypical hyperplasia and cancer	4 (11.4%)	0	0.03*
Total	35	46	

ET=endometrial thickness

CI=confidence interval

**p* is measured by Fisher’s Exact test and considered significant when < 0.05



Endometrial thickness receiver operator characteristic (ROC) curve for endometrial atypical hyperplasia and cancer. Arrow indicates the point where the 10mm endometrial thickness lies on the curve; Area under the curve (AUC) for the endometrial thickness cut-off ≥ 10 mm is 0.8, 95% confidence interval (CI)=0.66-0.93, $p=0.04$, sensitivity = 100% (95% CI=40-100%), specificity = 60% (95% CI=48-71%)

190x275mm (96 x 96 DPI)