

Open Access Maced J Med Sci electronic publication ahead of print,
published on January 25, 2019 as <https://doi.org/10.3889/oamjms.2019.073>

ID Design Press, Skopje, Republic of Macedonia
Open Access Macedonian Journal of Medical Sciences.
<https://doi.org/10.3889/oamjms.2019.073>
eISSN: 1857-9655
Review Article



Hepatitis E Virus Infection in Bulgaria: A Brief Analysis of the Situation in the Country

Magdalena Baymakova^{1*}, Georgi T. Popov¹, Roman Pepovich², Ilija Tsachev³

¹Department of Infectious Diseases, Military Medical Academy, Sofia, Bulgaria; ²Department of Infectious Pathology, Hygiene, Technology and Control of Foods from Animal Origin, Faculty of Veterinary Medicine, University of Forestry, Sofia, Bulgaria; ³Department of Microbiology, Infectious and Parasitic Diseases, Faculty of Veterinary Medicine, Trakia University, Stara Zagora, Bulgaria

Abstract

Citation: Baymakova M, Popov GT, Pepovich R, Tsachev I. Hepatitis E Virus Infection in Bulgaria: A Brief Analysis of the Situation in the Country. Open Access Maced J Med Sci. 2019;10(1):73-78. <https://doi.org/10.3889/oamjms.2019.073>

Keywords: HEV infection; Bulgaria; Public health

***Correspondence:** Magdalena Baymakova, Military Medical Academy, Sofia, Bulgaria. E-mail: dr.baymakova@gmail.com

Received: 16-Oct-2018; **Revised:** 09-Jan-2019; **Accepted:** 10-Jan-2019; **Online first:** 25-Jan-2019

Copyright: © 2019 Magdalena Baymakova, Georgi T. Popov, Roman Pepovich, Ilija Tsachev. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Funding: This study was supported by the National scientific programme "For Women in Science" 2018

Competing Interests: The authors have declared that no competing interests exist

BACKGROUND: Over the past two decades, more thorough investigations for hepatitis E virus (HEV) infection have been done in the world. Reports from Southeast European countries have increased.

AIM: The current article presents a critical analysis of all studies for HEV in Bulgaria.

MATERIAL AND METHODS: A literature search was done using available medical databases. We analysed the literature in PubMed databases and Bulgarian medical databases for English and Bulgarian languages sources. Preference was given to the sources published within the past 24 years (January 1995 – September 2018).

RESULTS: Two thousand two hundred and fifty-seven blood serums were tested for the analysed period (1995 – 2018), and 13.1% of them were positive for acute HEV (Mean \pm SD: 20.38 \pm 25.77%; 95% CI: 1.29 – 39.47%). The following subtypes were established in the country – HEV Subtype 3e, HEV Subtype 3f, HEV Subtype 3c, HEV Subtype 3i, HEV Subtype 3hi and HEV Subtype 1.

CONCLUSION: We hope that the National Health Organizations will take adequate and timely measures to increase the knowledge and research for HEV among Bulgarian citizens.

Introduction

Hepatitis E virus (HEV) infection is an infectious disease which thorough investigation has realised over the past twenty years [1] [2]. First studies of HEV were from the late 1970s and the early 1980s [3], [4]. In Bulgaria, the first human cases were reported by Teoharov et al., in 1995 [5]. The infection has been investigated thoroughly in the country since 2008 [6], [7]. Some Bulgarian authors presented data for HEV infection among Bulgarian patients [8], [9]. However, there are still many unknown questions about this infection in our country.

The aim of this study is a critical analysis of all available articles for HEV infection in Bulgaria.

Materials and Methods

Study Design

A literature search was done using available medical databases. We analysed the literature in PubMed databases and Bulgarian medical databases for English and Bulgarian languages sources using the following keywords "Hepatitis E virus" AND "Bulgaria" AND "HEV" AND "Hepatitis E virus infection" AND "HEV infection". No age restriction was observed. All studies were written in the English or Bulgarian language. The critical analysis included original articles, brief reports and case series. The critical analysis excluded articles with missing data for the number of investigated individuals, case reports

and letters to the editor. Preference was given to the sources published within the past 24 years (January 1995 – September 2018).

Ethics Statement

The study was performed by the principles of the Declaration of Helsinki. Participation in the study was fully voluntary and anonymous and written informed consent was obtained from each person before the medical examination.

Statistical Analysis

The results obtained from the majority of studies do not allow detailed statistical analysis. Therefore, basic statistical indicators such as confidence interval (CI), standard deviation (SD), etc. were applied. Statistical analysis was performed by Excel 2007 (Microsoft, Redmond, Washington, USA) and SPSS Statistics 19.0 (IBM Corp., Armonk, New York, USA). A *P*-value < 0.05 was considered statistically significant.

Results

We found 23 papers (PubMed databases – 5; Bulgarian medical databases – 18). Seven articles fulfilled the inclusion and exclusion criteria and were included in the final data analysis (Table 1) [5], [8], [9], [10], [11], [12], [13]. All articles described hospitalised patients with clinical presentation of acute HEV infection. Two thousand two hundred and fifty-seven blood serums were tested for the analysed period (1995 – 2018), and 13.1% of them were positive for acute HEV (Mean \pm SD: 20.38 \pm 25.77%; 95% CI: 1.29 – 39.47%). Individuals have been tested for anti-HEV IgG and/or anti-HEV IgM. The following subtypes were established in the country – *HEV Subtype 3e* (GenBank accession number 77A 2012-BG, 2588 2011-BG, 2308 2011-BG, et al), *HEV Subtype 3f* (GenBank access. num. 762 2011-BG, 855 2011-BG, 688A 2010-BG, et al.), *HEV Subtype 3c* (GenBank access. num. ISS75 Plov 2014, ISS100 Paz 2014, ISS2 Sof 2013, et al.), *HEV Subtype 3i* (GenBank access. num. 1785 2011-BG, 905 2012-BG), *HEV Subtype 3hi* (GenBank access. num. ISS62 Paz 2014), and *HEV Subtype 1* (GenBank access. num. ISS78 Haskovo – travel Afghanistan 2014) [9], [13]. In Bulgaria HEV infection affected predominately male gender (male/female – 80.0/20.0%). The mean age of HEV-positive people varied between 50.7 \pm 23.0 years (95% CI: 32.26 – 69.06) for Sofia district [8], and 53.6 \pm 14.0 years (95% CI: 45.66 – 61.49) for Plovdiv district [10].

Table 1: Studies on acute hepatitis E virus infection in Bulgaria

First author and reference	Year of Publication	Methodology	Target group	Diagnostics	Investigated, (n)	HEV positive, (n)
Teoharov et al. [5]	1995	Retrospective	Hospitalized	ELISA	53	4
Dikov et al. [8]	2012	Retrospective	Hospitalized	ELISA	117	6
Teoharov et al. [9]	2014	Retrospective	Hospitalized	ELISA HEV RNA	741	67
Petrov et al. [10]	2015	Retrospective	Hospitalized	ELISA	112	12
Baymakova et al. [11]	2016	Retrospective	Hospitalized	ELISA	806	20
Stoykova et al. [12]	2017	Retrospective	Hospitalized	ELISA	325	111
Bruni et al. [13]	2018	Retrospective	Hospitalized	ELISA HEV RNA	103	76

Discussion

HEV infection more frequently affects male than female (61-69%) [14]. Similar data were found in Bulgarian studies for HEV gender distribution (male/female – 80.0/20.0%). But opposite data were reported from Romania (male/female – 40/60%) [15], and Turkey (21.4/78.6%) [16].

The highest incidence of HEV infection is observed in people over fifty years [14]. Similar data are reported from Romania (28% HEV-positive in the age group of 45-65 years) [15], Turkey (50.9 \pm 16.8 years) [16], and Albania (35% HEV-positive in the age group over 50 years) [17]. In Bulgaria, HEV infection occurs mainly in people over 50 years of age.

Up to now, there are only two Bulgarian studies for HEV phylogenetic analysis of human samples [9], [13]. The studies established *HEV Subtype 3e*, *HEV Subtype 3f*, *HEV Subtype 3c*, *HEV Subtype 3i*, *HEV Subtype 3hi* and *HEV Subtype 1* [9], [13]. It is known that *HEV Subtype 3e* occurs mainly in swine, *HEV Subtype 3f* is human subtype, and *HEV Subtype 3i* is mainly wild boar subtype. The established *HEV Subtype 3e* in Bulgaria is close to the reference sequence AB248520 Japan 3e. The established *HEV Subtype 3f* showed high similarity with reference sequence EU495148 France Hu 3f. The third found *HEV Subtype 3i* in Bulgaria is familiar to reference sequence FJ705359 Germany wild boar 3i. These results indicate that the virus might have an autochthonous character in Bulgaria with zoonotic potential and main reservoirs as domestic pigs and wild boars.

In Bulgaria, HEV infection is poorly known and diagnosed. The realised human studies are focused on the clinical manifestations and characteristics of acute clinical form. There are many uncertain things about HEV infection in Bulgaria. First, a basic problem has no established HEV case definition in Bulgaria. Second, has not enough investigations on HEV RNA among Bulgarian patients. Third, there is no licensed diagnostic laboratory for hepatitis E sequencing (HEV RNA analysis). Fourthly, there are no morbidity registers for HEV in Bulgaria.

Therefore, in some areas of the country, it is not known what is the real prevalence of HEV among the population. Fifth, the mortality of HEV is unknown among Bulgarian citizens. Sixthly, HEV seroprevalence is unknown for risk groups (HIV-positive patients, immunocompromised persons, transplanted patients and pregnant women). Seventh, animals' investigations are limited in the country. There is one study for HEV seropositivity among pigs in industrial farms in Bulgaria [18]. The results from this project found 40% HEV-positive pigs [18].

There are some reasons for those missing information and knowledge for HEV in Bulgaria. First, there is a lack of coordination and collaboration between scientific societies of infectious diseases specialists, microbiologists, virologists and gastroenterologists. Second, the researches in the country are poorly financed. Third, the Ministry of Health does not record the incidence and mortality of HEV infection in Bulgaria. Fourthly, National Health Authorities do not conduct a screening programme for blood and blood products for hepatitis E virus. Fifth, the veterinary services in the country do not search for HEV infection routinely in animals (domestic pigs, wild boars, bears, dogs, cats, etc.). Sixthly, lack of efficiency of some hospitals. Seventh, poor diagnostic capacity of some hospital laboratories. Eighth, poorly trained medical staff is seeking, recognising and investigating HEV infection.

In conclusion, the combination of these facts leads to the national position of insufficient evidence of HEV infection in our country. The notified weaknesses could be revised, and the local situations could be improved. We hope that the National Health Organizations will take adequate and timely measures to increase the knowledge and research for HEV among Bulgarian citizens.

Acknowledgement

This study was supported by the National scientific programme "For Women in Science" 2018.

References

1. Khuroo MS, Khuroo MS, Khuroo NS. Hepatitis E: discovery, global impact, control and cure. *World J Gastroenterol*. 2016; 22(31):7030–7045. <https://doi.org/10.3748/wjg.v22.i31.7030> PMID:27610014 PMCID:PMC4988308
2. Pepovich R, Baymakova M, Pishmisheva M, Marutsov P, Pekova L, Tsachev I. Current knowledge on Hepatitis E virus infection. *Vojnosanit Pregl*. 2019; 76(7):[in Press]. <https://doi.org/10.2298/VSP170815159P>
3. Khuroo MS. Study of an epidemic of non-A, non-B hepatitis. Possibility of another human hepatitis virus distinct from post-transfusion non-A, non-B type. *Am J Med*. 1980; 68(6):818–824. [https://doi.org/10.1016/0002-9343\(80\)90200-4](https://doi.org/10.1016/0002-9343(80)90200-4) PMID: 6770682
4. Wong DC, Purcell RH, Sreenivasan MA, Prasad SR, Pavri KM. Epidemic and endemic hepatitis in India: evidence for a non-A, non-B hepatitis virus aetiology. *Lancet*. 1980; 316(8200):876–879. [https://doi.org/10.1016/S0140-6736\(80\)92045-0](https://doi.org/10.1016/S0140-6736(80)92045-0) PMID: 6107544
5. Teoharov P, Tiholova M, Draganov P, Lilyanova V, Ivanova R, Varleva T, et al. First cases of hepatitis E virus infection in Bulgaria. *Infectology (Sofia)*. 1995; 32(3):17–18. [in Bulgarian]
6. Teoharov P, Pishmisheva M, Kovaleva V. Hepatitis E virus infection: a case report. *Infectology (Sofia)*. 2008; 45(1):43–44. [in Bulgarian]
7. Rouseva A, Nikolovska D. Two cases of hepatitis E in Bulgaria. *Pediatr Infect Dis (Sofia)*. 2010; 2(1):26–28. [in Bulgarian]
8. Dikov I, Popov G, Plochev K, Baymakova M. Hepatitis E virus infection in Bulgaria. *Bulg Hepatogastroenterology (Sofia)*. 2012; 14(1):37–40. [in Bulgarian]
9. Teoharov P, Kevorkyan A, Raycheva R, Golkocheva-Markova E, Trandeva-Bankova D, Andonov A. Data on the prevalence of hepatitis E virus in Bulgaria. *Comptes rendus de l'Academie bulgare des Sciences*. 2014; 67(10):1427–1432.
10. Petrov A, Stoycheva M, Vatev N, Boev I. Viral hepatitis E in Plovdiv district: epidemiological, clinical and biochemical characteristics. *Sci Infectol Parasitol (Sofia)*. 2015; 6(1):29–31. [in Bulgarian]
11. Baymakova M, Sakem B, Plochev K, Popov GT, Mihaylova-Garnizova R, Kovaleva V, et al. Epidemiological characteristics and clinical manifestations of hepatitis E virus infection in Bulgaria: a report on 20 patients. *Srp Arh Celok Lek*. 2016; 144(1-2):63–68. <https://doi.org/10.2298/SARH1602063B> PMID:27276860
12. Stoykova Z, Ivanova L, Tsaneva-Damyanova D, Kostadinova T. Hepatitis E virus infection in Northeastern Bulgaria. *Medical Review (Sofia)*. 2017; 53(3):30–34. [in Bulgarian]
13. Bruni R, Villano U, Equestre M, Chionne P, Madonna E, Trandeva-Bankova D, et al. Hepatitis E virus genotypes and subgenotypes causing acute hepatitis, Bulgaria, 2013-2015. *PLoS One*. 2018; 13(6):e0198045. <https://doi.org/10.1371/journal.pone.0198045> PMID: 29879148 PMCID: PMC5991722
14. Aspinall E, Hutchinson S, Goldberg D, Smith-Palmer A. ECDC Surveillance Report: Hepatitis E in the EU/EEA, 2005–2015. Stockholm (Sweden): ECDC, 2017. <https://doi.org/10.2900/059144>
15. Anita A, Gorgan L, Anita D, Oslobanu L, Pavio N, Savuta G. Evidence of hepatitis E infection in swine and humans in the East Region of Romania. *Int J Infect Dis*. 2014; 29:232–237. <https://doi.org/10.1016/j.ijid.2014.10.018> PMID:25447729
16. Eker A, Tansel O, Kunduracilar H, Tokuc B, Yulugkural Z, Yuksel P. Hepatitis E virus epidemiology in adult population in Edirne province, Turkey. *Mikrobiyol Bul*. 2009; 43(2):251–258. [in Turkish] PMID:19621610
17. Kondili LA, Chionne P, Porcaro A, Madonna E, Taffon S, Resuli B, et al. Seroprevalence of hepatitis E virus (HEV) antibody and the possible association with chronic liver disease: a case-control study in Albania. *Epidemiol Infect*. 2006; 134(1):95–101. <https://doi.org/10.1017/S095026880500470X> PMID: 16409655 PMCID: PMC2870364
18. Pishmisheva M, Baymakova M, Golkocheva-Markova E, Kundurzhiev T, Pepovich R, Popov GT, et al. First serological study of hepatitis E virus infection in pigs in Bulgaria. *Comptes rendus de l'Academie bulgare des Sciences*. 2018; 71(7):1001–1008. <https://doi.org/10.7546/CRABS.2018.07.18>