

# THE ASYMPTOTIC NUMBER OF PLANAR, SLIM, SEMIMODULAR LATTICE DIAGRAMS

GÁBOR CZÉDLI

ABSTRACT. A lattice  $L$  is *slim* if it is finite and the set of its join-irreducible elements contains no three-element antichain. We prove that there exists a positive constant  $C$  such that, up to similarity, the number of planar diagrams of slim semimodular lattices of size  $n$  is asymptotically  $C \cdot 2^n$ .

## 1. INTRODUCTION AND THE RESULT

A finite lattice  $L$  is *slim* if  $Ji L$ , the set of join-irreducible elements of  $L$ , contains no three-element antichain. Equivalently,  $L$  is slim if  $Ji L$  is the union of two chains. Slim, semimodular lattices were heavily used while proving a recent generalization of the classical Jordan-Hölder theorem for groups in [4]. These lattices are *planar*, that is, they have planar diagrams, see [4]. Hence it is reasonable to study their planar diagrams, which are called *slim, semimodular (lattice) diagrams* for short. The *size* of a diagram is the number of elements of the lattice it represents. Let  $D_1$  and  $D_2$  be two planar lattice diagrams. A bijection  $\phi: D_1 \rightarrow D_2$  is a *similarity map* if it is a lattice isomorphism preserving the left-right order of (upper) covers and that of lower covers of each element of  $D_1$ . If there is a similarity map  $D_1 \rightarrow D_2$ , then these two diagrams are *similar*, and we will treat them as equal. Let  $N_{\text{ssd}}(n)$  denote the number of slim, semimodular diagrams of size  $n$ , counting them up to similarity. Our target is to prove the following result.

**Theorem 1.1.** *There exists a positive constant  $C < 1$  such that  $N_{\text{ssd}}(n)$  is asymptotically  $C \cdot 2^n$ , that is,  $\lim_{n \rightarrow \infty} (N_{\text{ssd}}(n)/2^n) = C$ .*

Given two composition series in a finite group, the intersections of their members form a slim semimodular lattice with respect to “ $\supseteq$ ”. This follows from Wielandt [13]; see also the proof of Nation [11, Theorem 9.8]. Conversely, [6] proves that every slim semimodular lattice can be represented in this way. Therefore, in a reasonable, order theoretic sense, Theorem 1.1 tells us how many ways the members of two composition series in a group can intersect each other, provided that there are exactly  $n$  intersections and that we make a distinction between the first composition series and the second one.

Note that there are two different methods to deal with  $N_{\text{ssd}}(n)$ . The present one yields the asymptotic statement above, while the method of [1] gives the exact

---

*Date:* June 23, 2015.

*Key words and phrases.* Counting lattices, semimodularity, planar lattice diagram, slim semimodular lattice.

2010 *Mathematics Subject Classification.* 06C10.

This research was supported by the NFSR of Hungary (OTKA), grant numbers K77432 and K83219, and by TÁMOP-4.2.1/B-09/1/KONV-2010-0005.




FIGURE 1. Left and right ranks

values of  $N_{\text{ssd}}(n)$  up to  $n = 50$  (with the help of a usual personal computer). Also, [1] determines the number  $N_{\text{ssl}}(n)$  of slim, semimodular lattices of size  $n$  up to  $n = 50$  while we do not even know  $\lim_{n \rightarrow \infty} (N_{\text{ssl}}(n)/N_{\text{ssl}}(n-1))$ , and it is only a conjecture that this limit exists.

Note also that, besides [1] and [2], there are several papers on counting lattices; see, for example, M. Erné, J. Heitzig, and J. Reinhold [7], M. M. Pawar and B. N. Waphare [12], and J. Heitzig and J. Reinhold [9].

## 2. LATTICE THEORETIC LEMMAS

A minimal non-chain region of a planar lattice diagram  $D$  is called a *cell*. A four-element cell is a *4-cell*; it is also a *covering square*, that is, a cover-preserving four-element Boolean sublattice. We say that  $D$  is a *4-cell diagram* if all of its cells are 4-cells. We shall heavily rely on the following result of G. Grätzer and E. Knapp [8, Lemmas 4 and 5].

**Lemma 2.1.** *Let  $D$  be a finite planar lattice diagram.*

- (i) *If  $D$  is semimodular, then it is a 4-cell diagram. If  $A$  and  $B$  are 4-cells of  $D$  with the same bottom, then these 4-cells have the same top.*
- (ii) *If  $D$  is a 4-cell diagram in which no two 4-cells with the same bottom have distinct tops, then  $D$  is semimodular.*

In what follows, we always assume that  $4 \leq n \in \mathbb{N} = \{1, 2, \dots\}$ , and that  $D$  is a slim, semimodular diagram of size  $n$ . Let  $w_D^\ell$  be the smallest doubly irreducible element of the left boundary chain  $\text{BC}_\ell(D)$  of  $D$ , and let  $\text{rank}_\ell(D)$  be the height of  $w_D^\ell$ . The left-right duals of these concepts are denoted by  $w_D^r$  and  $\text{rank}_r(D)$ . See Figure 1 for an illustration, where  $w_D^\ell$  and  $w_D^r$  are the black-filled elements. By D. Kelly and I. Rival [10, Proposition 2.2], each planar lattice diagram with at least three elements contains a doubly irreducible element  $\neq 0, 1$  on its left boundary. This implies the following statement, on which we will rely implicitly.

**Lemma 2.2.** *Either  $\text{rank}_\ell(D) = \text{rank}_r(D) = 0$  and  $w_D^\ell = w_D^r = 0$ , or  $\text{rank}_\ell(D) > 0$  and  $\text{rank}_r(D) > 0$ .*

For  $a \in D$ , the ideal  $\{x \in D : x \leq a\}$  is denoted by  $\downarrow a$ .

**Lemma 2.3.**  $\text{BC}_\ell(D) \cap \downarrow w_D^\ell \subseteq \text{Ji } D$ .

*Proof.* Suppose, for a contradiction, that the lemma fails, and let  $u$  be the smallest join-reducible element belonging to  $\text{BC}_\ell(D) \cap \downarrow w_D^\ell$ . By D. Kelly and I. Rival [10,

Proposition 2.2], there is a doubly irreducible element  $v$  of the ideal  $\downarrow u = \{x \in D : x \leq u\}$  such that  $v \in \text{BC}_\ell(\downarrow u)$ ; notice that  $v$  also belongs to  $\text{BC}_\ell(D)$ . Clearly,  $v < u$  and  $v$  is join-irreducible in  $D$ . Therefore, since  $v < u \leq w_D^\ell$  and  $w_D^\ell$  is the least doubly irreducible element of  $\text{BC}_\ell(D)$ ,  $v$  is meet-reducible in  $D$ . Hence there exist a  $p \in D$  such that  $v < p$  and  $p \notin \downarrow u$ . Denote by  $u_0$  the unique lower cover of  $u$  in  $\text{BC}_\ell(D)$ . Since  $v < u$ , we have that  $v \leq u_0$ . By semimodularity and  $p \not\leq u_0$ , we obtain that  $u_0 = u_0 \vee v < u_0 \vee p \neq u$ . Hence  $u_0$  has two covers,  $u$  and  $u_0 \vee p$ . Thus  $u_0, u \in \text{BC}_\ell(D)$ ,  $u_0 < u$ ,  $u$  is join-reducible, and  $u_0$  is meet-reducible. This contradicts [5, Lemma 4].  $\square$

Next, we prove the following lemma.

**Lemma 2.4.** *For  $4 \leq n \in \mathbb{N}$ , we have that*

$$(2.1) \quad N_{\text{ssd}}(n-1) + N_{\text{ssd}}(n-3) \leq N_{\text{ssd}}(n),$$

$$(2.2) \quad N_{\text{ssd}}(n) \leq 2 \cdot N_{\text{ssd}}(n-1).$$

*Proof.* The set of slim, semimodular diagrams of size  $n$  is denoted by  $\text{SSD}(n)$ . Let

$$\text{SSD}_{00}(n) = \{D \in \text{SSD}(n) : \text{rank}_\ell(D) = \text{rank}_r(D) = 0\},$$

$$\text{SSD}_{11}(n) = \{D \in \text{SSD}(n) : \text{rank}_\ell(D) = \text{rank}_r(D) = 1\}, \text{ and}$$

$$\text{SSD}_{++}(n) = \text{SSD}(n) - \text{SSD}_{00}(n).$$

Since we can omit the least element and the least three elements, respectively, and the remaining diagram is still slim and semimodular by Lemma 2.1, we conclude that  $|\text{SSD}_{00}(n)| = N_{\text{ssd}}(n-1)$  and  $|\text{SSD}_{11}(n)| = N_{\text{ssd}}(n-3)$ . This implies (2.1). For  $D \in \text{SSD}_{++}(n)$ , we define

$$D^* = D - \{w_D^\ell\}.$$

We know from By D. Kelly and I. Rival [10, Proposition 2.2], mentioned earlier, that

$$(2.3) \quad w_D^\ell \notin \{0, 1\}, \text{ provided } D \in \text{SSD}_{++}(n).$$

This, together with the fact that  $D \in \text{SSD}_{++}(n)$  is not a chain, yields that

$$(2.4) \quad \text{length } D^* = \text{length } D.$$

Let  $w_D^{\ell-}$  denote the unique lower cover of  $w_D^\ell$  in  $D$ . Since each meet-reducible element has exactly two covers by [5, Lemma 2], we conclude from Lemma 2.3 that

$$(2.5) \quad w_{D^*}^\ell = w_D^{\ell-}.$$

It follows from Lemma 2.1 that  $D^* \in \text{SSD}(n-1)$ . From (2.5) we obtain that

$$(2.6) \quad D^* \in \text{SSD}(n-1) \text{ determines } D.$$

Hence  $|\text{SSD}_{++}(n)| \leq |\text{SSD}(n-1)| = N_{\text{ssd}}(n-1)$ . Combining this with  $|\text{SSD}_{00}(n)| = N_{\text{ssd}}(n-1)$  and  $\text{SSD}(n) = \text{SSD}_{00}(n) \dot{\cup} \text{SSD}_{++}(n)$ , where  $\dot{\cup}$  stands for disjoint union, we obtain (2.2).  $\square$

Next, let

$$(2.7) \quad W(n) = \text{SSD}(n-1) - \{D^* : D \in \text{SSD}_{++}(n)\}.$$

Fortunately, this set is relatively small by the following lemma. The upper integer part of a real number  $r$  is denoted by  $\lceil x \rceil$ ; for example,  $\lceil \sqrt{3} \rceil = 2$ .

**Lemma 2.5.** *If  $4 \leq n$ , then  $|W(n)| \leq \sum_{j=2}^{n+1-\lceil\sqrt{n-1}\rceil} N_{\text{ssd}}(j)$ .*

*Proof.* First we show that

$$(2.8) \quad W(n) = \{E \in \text{SSD}(n-1) : w_E^\ell \text{ is a coatom of } E\}.$$

The  $\supseteq$  inclusion is clear from (2.3), (2.4), and (2.5). These facts together with Lemma 2.1 also imply the reverse inclusion since by adding a new cover to  $w_E^\ell$ , to be positioned to the left of  $\text{BC}_\ell(E)$ , we obtain a slim semimodular diagram  $D$  such that  $D^* = E$ .

It follows from Lemma 2.3 that no down-going chain starting at  $w_E^\ell$  can branch out. Thus

$$(2.9) \quad \downarrow w_E^\ell \subseteq \text{BC}_\ell(E) \text{ and } \downarrow w_E^\ell \text{ is a chain.}$$

Since  $w_E^\ell$  is a coatom, we have that

$$(2.10) \quad \text{with the notation } E^\blacktriangleleft = E \setminus \downarrow w_E^\ell, \quad |E^\blacktriangleleft| = |E| - \text{length } E.$$

Clearly,  $E^\blacktriangleleft$  is a join-subsemilattice of  $E$  since it is an order-filter. To prove that

$$(2.11) \quad E^\blacktriangleleft \text{ is a slim, semimodular diagram,}$$

assume that  $x, y \in E^\blacktriangleleft - \{1\}$ . We want to show that  $x \wedge y$ , taken in  $E$ , belongs to  $E^\blacktriangleleft$ . Let  $x_0$  and  $y_0$  be the smallest element of  $\text{BC}_\ell(E) \cap \downarrow x$  and  $\text{BC}_\ell(E) \cap \downarrow y$ , respectively. Since  $x_0, y_0 \in \text{BC}_\ell(E) \cap (\downarrow w_E^\ell - \{w_E^\ell\})$ , the definition of  $w_E^\ell$  implies that  $x_0$  and  $y_0$  are meet-reducible. Hence they have exactly two covers by [5, Lemma 2]. Let  $x_1$  and  $y_1$  denote the cover of  $x_0$  and  $y_0$ , respectively, that do not belong to  $\text{BC}_\ell(E)$ , and let  $x^+$  and  $y^+$  be the respective covers belonging to  $\text{BC}_\ell(E)$ . By the choice of  $x_0$ , we have that  $x^+ \not\leq x$ , whence  $x_1 \leq x$ . Similarly,  $y_1 \leq y$ . Since  $\text{BC}_\ell(E)$  is a chain and the case  $x_0 = y_0$  will turn out to be trivial, we can assume that  $x_0 < y_0$ . We know that  $x_1 \not\leq y_0$  since otherwise  $x_1$  would belong to  $\text{BC}_\ell(E)$  by (2.9). Using semimodularity, we obtain that  $x_1 \vee y_0 \succ y_0$ . Since  $y_0$  has only two covers by [5, Lemma 2] and  $x_1 \leq y^+$  would imply  $x_1 \in \text{BC}_\ell(E)$  by (2.9), it follows that  $x_1 \vee y_0 = y_1$ . Hence  $x_1 \leq y$ ,  $x_1 \leq x$ , and  $x_1 \in E^\blacktriangleleft$  imply that  $x \wedge y$  belongs to (the order filter)  $E^\blacktriangleleft$ . Thus  $E^\blacktriangleleft$  is (to be more precise, determines) a sublattice of (the lattice determined by)  $E$ . The semimodularity of  $E^\blacktriangleleft$  follows from Lemma 2.1. This proves (2.11).

By (2.10) and (2.11), a trivial argument gives that

$$(2.12) \quad E^\blacktriangleleft \in \text{SSD}(n - \text{length } E) \text{ and } E^\blacktriangleleft \text{ determines } E.$$

Next, we have to determine what values  $h = \text{length } E$  can take. Clearly,  $h \leq |E| - 1 = n - 2$ . There are various ways to check that  $|E| \leq (1 + \text{length } E)^2 = (1 + h)^2$ ; this follows from the main theorem of [6], and follows also from the proof of [3, Corollary 2]. Since now  $|E| = n - 1$ , we obtain that  $\lceil\sqrt{n-1}\rceil - 1 \leq h$ . Therefore, combining (2.11) and (2.12), we obtain that

$$W(n) \leq \sum_{h=\lceil\sqrt{n-1}\rceil-1}^{n-2} N_{\text{ssd}}(n-h).$$

Substituting  $j$  for  $n - h$ , we obtain our statement.  $\square$

We conclude this section by the following lemma.

$$\textbf{Lemma 2.6.} \quad 2 \cdot N_{\text{ssd}}(n-1) - \sum_{j=2}^{n+1-\lceil\sqrt{n-1}\rceil} N_{\text{ssd}}(j) \leq N_{\text{ssd}}(n) \leq 2 \cdot N_{\text{ssd}}(n-1).$$

*Proof.* By (2.6) and the definition of  $W(n)$ , we have that

$$\begin{aligned} N_{\text{ssd}}(n) &= |\text{SSD}_{00}(n)| + |\text{SSD}_{++}(n)| = N_{\text{ssd}}(n-1) + |\text{SSD}(n-1) - W(n)| \\ &= N_{\text{ssd}}(n-1) + N_{\text{ssd}}(n-1) - |W(n)|, \end{aligned}$$

and the statement follows from Lemma 2.5 and (2.2).  $\square$

### 3. TOOLS FROM ANALYSIS AT WORK

For  $k \geq 2$ , define  $\kappa_k = N_{\text{ssd}}(k)/N_{\text{ssd}}(k-1)$ . Since  $N_{\text{ssd}}(n-3)/N_{\text{ssd}}(n-1) = 1/(\kappa_{n-1}\kappa_{n-2})$ , dividing the inequalities of Lemma 2.4 by  $N_{\text{ssd}}(n-1)$  we obtain that  $1 + 1/(\kappa_{n-1}\kappa_{n-2}) \leq \kappa_n \leq 2$ , for  $n \geq 4$ . Furthermore, in view of the sentence following (2.7), (2.8) implies easily that  $\kappa_n < 2$  if  $n \geq 7$ . Therefore, since  $\kappa_k \leq 2$  also holds for  $k \in \{2, 3\}$  and  $1 + 1/(2 \cdot 2) = 5/4$ , we conclude that

$$(3.1) \quad 5/4 \leq \kappa_n \leq 2, \quad \text{for } n \geq 4, \quad \text{and } \kappa_n < 2, \quad \text{for } n \geq 7.$$

Clearly,  $N_{\text{ssd}}(k-1) = N_{\text{ssd}}(k)/\kappa_k \leq \frac{4}{5} \cdot N_{\text{ssd}}(k)$  if  $k \geq 4$ . Thus, by iteration, we obtain that

$$(3.2) \quad N_{\text{ssd}}(k-j) \leq (4/5)^j \cdot N_{\text{ssd}}(k), \quad \text{for } j \in \mathbb{N}_0 \text{ and } k \geq j+4.$$

If  $k \geq 5$ , then using  $N_{\text{ssd}}(k) \geq N_{\text{ssd}}(5) \geq 3$  (actually,  $N_{\text{ssd}}(5) = 3$ ), we obtain that

$$(3.3) \quad \begin{aligned} N_{\text{ssd}}(1) + \cdots + N_{\text{ssd}}(k) &= 1 + 1 + 1 + N_{\text{ssd}}(4) + \cdots + N_{\text{ssd}}(k) \\ &\leq 3 + N_{\text{ssd}}(k) \cdot ((4/5)^{k-4} + (4/5)^{k-5} + \cdots + (4/5)^0) \\ &\leq N_{\text{ssd}}(k) + N_{\text{ssd}}(k) \cdot 1/(1-4/5) = 6N_{\text{ssd}}(k). \end{aligned}$$

Combining Lemma 2.6 with (3.3) and (3.2), we obtain that

$$\begin{aligned} 2N_{\text{ssd}}(n-1) - 6 \cdot (4/5)^{\lceil\sqrt{n-1}\rceil-2} \cdot N_{\text{ssd}}(n-1) &\leq \\ 2N_{\text{ssd}}(n-1) - 6N_{\text{ssd}}(n+1-\lceil\sqrt{n-1}\rceil) &\leq \\ \leq N_{\text{ssd}}(n) \leq 2N_{\text{ssd}}(n-1). \end{aligned}$$

Dividing the formula above by  $2N_{\text{ssd}}(n-1)$  and (3.1) by 2, we obtain that

$$(3.4) \quad \max(5/8, 1 - 3 \cdot (4/5)^{\lceil\sqrt{n-1}\rceil-2}) \leq \kappa_n/2 \leq 1, \quad \text{for } n \geq 5.$$

Next, let us choose an integer  $m \geq 5$ , and define

$$z_0 = z_0(m) = \min(3/8, 3 \cdot (4/5)^{\lceil\sqrt{m-1}\rceil-2}).$$

**Lemma 3.1.** *For  $0 \leq z < 1$ , we have  $-\ln(1-z) \leq z/(1-z)$ . If, in addition,  $0 \leq z \leq z_0$ , then  $z/(1-z) \leq z/(1-z_0)$ .*

*Proof.* The second inequality is obvious. The first inequality holds for  $z = 0$  and, for  $0 \leq z < 1$ , the derivative  $1/(1-z)$  of the left side is less than  $1/(1-z)^2$ , that of the right side. This implies the first inequality.  $\square$

With the auxiliary steps made so far, we are ready to start the final argument.

*Proof of Theorem 1.1.* For  $n > m$ , let

$$p_n = \prod_{j=m+1}^n (\kappa_j/2).$$

We obtain from (3.4) that  $\{p_n\}$ , that is,  $\{p_n\}_{n=m+1}^\infty$ , is a decreasing sequence of positive numbers. Clearly,

$$(3.5) \quad N_{\text{ssd}}(n)/2^n = p_n \cdot N_{\text{ssd}}(m)/2^m.$$

Hence it suffices to prove that the sequence  $\{p_n\}$  converges to a positive number, because then its limit is smaller than 1 by (3.1). Let  $s_n = -\ln p_n$ ,  $\mu = 3(1 - z_0)^{-1}$ ,  $\alpha = 4/5$ , and  $\nu = 5\mu/4 = \mu/\alpha$ . Note that  $\{s_n\}$  is an increasing sequence.

Using (3.4) together with Lemma 3.1 at the inequality marked with  $\leq'$  below and (3.4) at the one marked with  $\leq^*$ , and using that the function  $f(x) = \alpha^{\sqrt{x}}$  is decreasing, we obtain that

$$\begin{aligned} 0 < s_n &= \sum_{j=m+1}^n (-\ln(\kappa_j/2)) \leq' \sum_{j=m+1}^n (1 - \kappa_j/2)/(1 - z_0) \\ &\leq^* \mu \cdot \sum_{j=m+1}^n \alpha^{\lceil \sqrt{j-1} \rceil - 2} \leq \mu \cdot \sum_{j=m+1}^n \alpha^{\sqrt{j-1} - 1} = \mu \cdot \sum_{k=m}^{n-1} \alpha^{\sqrt{k} - 1} \\ &= \nu \cdot \sum_{k=m}^{n-1} \alpha^{\sqrt{k}} \leq \nu \cdot \int_{x=m-1}^{n-1} \alpha^{\sqrt{x}} dx \leq \nu \cdot (F(\infty) - F(m-1)), \end{aligned}$$

where  $F(x)$  is a function whose derivative is  $f(x)$ . Let  $\delta = -\ln \alpha = \ln(5/4)$ . It is routine to check (by hand or by computer algebra) that, up to a constant summand,

$$F(x) = -2 \cdot \delta^{-2} \cdot (1 + \delta\sqrt{x}) \cdot \alpha^{\sqrt{x}}.$$

Clearly,  $F(\infty) = \lim_{x \rightarrow \infty} F(x) = 0$ . This proves that the sequence  $\{s_n\}$  converges; so does  $\{p_n\} = \{e^{-s_n}\}$  by the continuity of the exponential function. Therefore, since  $N_{\text{ssd}}(m)/2^m$  in (3.5) does not depend on  $n$ , we conclude Theorem 1.1.  $\square$

**Remark 3.2.** We can approximate the constant in Theorem 1.1 as follows. Since  $e^{-\nu \cdot (F(\infty) - F(m))} \leq e^{-s_n} = p_n \leq 1$  and, by (3.5),  $C = \lim_{n \rightarrow \infty} (p_n N_{\text{ssd}}(m)/2^m)$ , we obtain that

$$(3.6) \quad e^{\nu F(m)} \cdot N_{\text{ssd}}(m)/2^m = e^{-\nu \cdot (F(\infty) - F(m))} \cdot N_{\text{ssd}}(m)/2^m \leq C \leq N_{\text{ssd}}(m)/2^m.$$

Unfortunately, our computing power yields only a very rough estimation. The largest  $m$  such that  $N_{\text{ssd}}(50)$  is known is  $m = 50$ , see [1]. With  $m = 50$  and  $N_{\text{ssd}}(m) = N_{\text{ssd}}(50) = 81\,287\,566\,224\,125$ , it is a routine task to turn (3.6) into

$$0.42 \cdot 10^{-57} \leq C \leq 0.073.$$

We have reasons (but no proof) to believe that  $0.023 \leq C \leq 0.073$ , see the Maple worksheet (version V) available from the authors's home page.

*Acknowledgment.* The author is indebted to Vilmos Totik for helpful discussions.

## REFERENCES

- [1] Czédli, G., Dékány, T., Ozsvárt, L., Szakács, N., Udvari, B.: On the number of slim, semimodular lattices, *Mathematica Slovaca*, to appear
- [2] Czédli, G., Ozsvárt, L., Udvari, B.: How many ways can two composition series intersect? *Discrete Mathematics* **312**, 3523–3536 (2012)
- [3] Czédli, G., Schmidt, E. T.: How to derive finite semimodular lattices from distributive lattices?, *Acta Mathematica Hungarica*, **121**, 277–282 (2008)
- [4] Czédli, G., Schmidt, E. T.: The Jordan-Hölder theorem with uniqueness for groups and semimodular lattices, *Algebra Universalis* **66**, 69–79 (2011)
- [5] Czédli, G., Schmidt, E. T.: Slim semimodular lattices. I. A visual approach, *Order* **29**, 481–497 (2012)
- [6] Czédli, G., Schmidt, E. T.: Composition series in groups and the structure of slim semimodular lattices, *Acta Sci Math. (Szeged)* **79**, 369–390 (2013)
- [7] Erné, M., Heitzig, J., Reinhold, J.: On the number of distributive lattices, *Electron. J. Combin.* **9** (2002), no. 1, Research Paper 24, 23 pp.
- [8] Grätzer, G., Knapp, E.: *Notes on planar semimodular lattices I. Construction*, *Acta Sci. Math. (Szeged)*. **73**, 445–462 (2007)
- [9] Heitzig, J., Reinhold, J.: Counting finite lattices, *Algebra Universalis* **48**, 43–53 (2002)
- [10] Kelly, D., Rival, I.: Planar lattices. *Canad. J. Math.* **27**, 636–665 (1975)
- [11] Nation, J. B.: Notes on Lattice Theory, <http://www.math.hawaii.edu/~jb/books.html>
- [12] Pawar, M. M., Waphare, B. N.: Enumeration of nonisomorphic lattices with equal number of elements and edges, *Indian J. Math.* **45**, 315–323 (2003)
- [13] Wielandt, H: Eine Verallgemeinerung der invarianten Untergruppen, *Math. Zeit.* **45**, 209–244 (1939)

*E-mail address:* [czedli@math.u-szeged.hu](mailto:czedli@math.u-szeged.hu)

*URL:* <http://www.math.u-szeged.hu/~czedli/>

UNIVERSITY OF SZEGED, BOLYAI INSTITUTE, SZEGED, ARADI VÉRTANÚK TERE 1, HUNGARY  
6720