

# On the spectral properties of square matrices that are strictly sign-regular of some order

Rola Al-Seidi (rolanazmi@yahoo.com),

Prof. Jürgen Garloff

Department of Mathematics and Statistics, University of Konstanz, Konstanz, Germany

## Introduction

By "spectral structure of square matrices" we mean properties of the eigenvalues and eigenvectors of matrices. In view of the well-known Perron-Frobenius theory that describes the spectral structure of general entrywise nonnegative matrices, we study the spectral structure of strictly sign-regular matrices of some order. A matrix is called *strictly sign-regular of order  $k$* , denoted by  $SSR_k$ , if all its minors of order  $k$  are non-zero and have the same sign. For example, *totally positive matrices (TP)*, i.e., matrices with all minors positive, are  $SSR_k$  for all  $k$ . Another important class of matrices are those that are  $SSR_k$  for all odd  $k$ . Such matrices have interesting sign variation diminishing properties, and it has been recently shown that they play an important role in the analysis of certain nonlinear cooperative systems.

## Definitions and Notations

We will introduce the following definitions.

- 1 A vector  $x \in \mathbb{R}^n$  is called *totally nonzero* if no entry of  $x$  is zero.
- 2 Let  $v$  be the function from the totally nonzero vectors  $x = [x_1, x_2, \dots, x_n]^T \in \mathbb{R}^n$  into the nonnegative integers defined by

$$v(x) := \#\{i : x_i x_{i+1} < 0\}, \text{ for } i = 1, 2, \dots, n-1,$$

the total sign variation of  $x$ . For general vectors  $x \in \mathbb{R}^n$ ,  $v^-(x)$  is the minimum value of  $v(y)$  among all totally nonzeros  $y$  that agree with  $x$  in its nonzero entries and  $v^+(x)$  is the maximum value of  $v(y)$  among all such vectors. In the case that  $x$  has zero entries,  $v^-(x)$  is also  $v(x')$  in which  $x'$  is simply the result of deleting the zero entries from  $x$ .

- 3 For a vector  $x \in \mathbb{R}^n$ , let

$$v_c^-(x) := \max_i v^-(x_i, x_{i+1}, \dots, x_n, x_1, \dots, x_i),$$

the cyclic number of sign variations.

- 4 Consider the linear time-varying system

$$\dot{x}(t) = A(t)x(t) \quad (1)$$

with  $A(t)$  a continuous matrix function of  $t$ . This system is called *totally positive differential system*, denoted by  $TPDS$ , on a time interval  $(a, b)$  if its transition matrix  $\phi(t, t_0)$  is totally positive for any pair  $(t_0, t)$  with  $a < t_0 < t < b$ . Here the transition matrix is the matrix satisfying  $x(t) = \phi(t, t_0)x(t_0)$ . In particular,  $\phi(t_0, t_0) = I$ . In the special case where  $A(t)$  is a constant matrix, i.e.,  $A(t) \equiv A$ , then  $\phi(t, t_0) = \exp((t - t_0)A)$ . Of course, the transition matrix is real, square, and nonsingular.

- 5 Let  $\mathbb{M}^+ \subset \mathbb{R}^{n \times n}$  denote the subset of  $n \times n$  real matrices that are tridiagonal with positive entries on the super- and sub-diagonals.

- 6 Let  $A \in \mathbb{R}^{n \times n}$ . We say that a set of complex numbers  $c_1, \dots, c_m \in \mathbb{C}^n$  matches  $A$  if  $\sum_{i=1}^m |c_i|^2 > 0$ , and for every  $i$  if the eigenvector  $v^i$  of  $A$  is real then  $c_i$  is real and, if  $v^i, v^{i+1}$  is a conjugate complex pair then  $c_{i+1} = \bar{c}_i$ .

The following example is a simple illustrative example to the previous definitions.

## Example

$$v^+ \begin{bmatrix} 1 \\ -1 \\ 2 \\ -3 \\ 0 \end{bmatrix} = 4,$$

$$v^- \begin{bmatrix} 1 \\ -1 \\ 2 \\ -3 \\ 0 \end{bmatrix} = 3,$$

$$v_c^+ \begin{bmatrix} 1 \\ -1 \\ 2 \\ -3 \\ 0 \end{bmatrix} = 4,$$

$$v_c^- \begin{bmatrix} 1 \\ -1 \\ 2 \\ -3 \\ 0 \end{bmatrix} = 4$$

## Motivation and Previous Works

Applications of strictly sign-regular matrices often come from their interpretation as variation-diminishing transformations. Strictly sign-regular matrices are characterized by the strong variation-diminishing property, i.e.,  $v^+(Ax) \leq v^-(x)$  for all nonzero  $x \in \mathbb{R}^n$ . In particular, multiplying a vector by a  $TP$  matrix can only decrease the number of sign variations in the vector. It was shown that the system (1) is  $TPDS$  if and only if  $A \in \mathbb{M}^+$ . Several studies analysed certain non-linear dynamical systems by showing that the transition matrix in the variational system satisfies a variation-diminishing property with respect to the cyclic number of sign variations in a vector. Motivated by this, Theorem (A) shows that the  $SSR_k$  property is equivalent to a non-standard variation-diminishing property. Theorem (B) provides a simple necessary and sufficient condition for a nonsingular square matrix  $A$  to satisfy a  $CVDP$ .

**Theorem A** Let  $A \in \mathbb{R}^{n \times n}$  be a nonsingular matrix. Pick  $k \in \{1, \dots, n\}$ . Then the following two conditions are equivalent.

- (i) For any vector  $x \in \mathbb{R}^n \setminus \{0\}$  with  $s_c^-(Ax) \leq k - 1$ ,  $v^+(Ax) \leq k - 1$ .
- (ii)  $A$  is  $SSR_k$ .

**Theorem B** Let  $A \in \mathbb{R}^{n \times n}$  be a nonsingular matrix. The following two conditions are equivalent.

- (i) For any vector  $x \in \mathbb{R}^n \setminus \{0\}$   $v_c^+(Ax) \leq v_c^-(x)$ .
- (ii) The matrix  $A$  is  $SSR_r$  for all odd  $r$  in the range  $r \in \{1, \dots, n\}$ .

## Main Results

Our main results extend Theorem (A) to vectors  $x \in \mathbb{C}^n$  and study the properties of the eigenvalues of the matrices in Theorem (B).

- 1 Suppose that  $A \in \mathbb{R}^{n \times n}$  is nonsingular and  $SSR_k$  for some  $k \in \{1, \dots, n-1\}$ . For any  $c_1, \dots, c_{n-1} \in \mathbb{C}$  that match  $A$ , we have

$$v^+ \left( \sum_{i=0}^k c_i v^i \right) \leq k - 1.$$

- 2 Let  $A \in \mathbb{R}^{n \times n}$  be nonsingular and  $SSR_k$  for all odd  $k \in \{1, 3, \dots, n\}$ . Then the following statements are true.
  - 1  $\lambda_1$  is a positive simple eigenvalue of  $A$ .
  - 2 The algebraic multiplicity of any eigenvalue of  $A$  is not greater than 2.
  - 3 The inequalities  $|\lambda_1| > |\lambda_2| \geq |\lambda_3| > |\lambda_4| \geq |\lambda_5| \dots$  hold.
  - 4 For every  $i \in \{3, 5, 7, \dots\}$ ,  $\lambda_{i-1}, \lambda_i$  are either a pair of complex conjugate or both are real and of the same sign.
  - 5 If  $n$  is even, then  $\lambda_n$  is real.

## Acknowledgement

I am gratefully acknowledge *KWIM* for the financial support to attend *EWM – GM 2018*, Graz. This work is done in cooperation with Prof. Michael Margaliot, Tel-Aviv University. We thank him for his support, comments, suggestions, and continuous discussions.

## References

- [1] Ben-Avraham, Tsuff, et al. "Dynamical systems with a cyclic sign variation diminishing property." arXiv preprint :1807.02779 (2018).
- [2] Margaliot, Michael, and Eduardo D. Sontag. "Revisiting totally positive differential systems: a tutorial and new results." arXiv preprint :1802.09590 (2018).
- [3] Karlin, Samuel. Total positivity. Vol. 1. Stanford University Press, 1968.
- [4] Fallat, Shaun M., and Johnson, Charles R. Totally nonnegative matrices. Princeton University Press, 2011.