

Epigenetics

ISSN: 1559-2294 (Print) 1559-2308 (Online) Journal homepage: <http://www.tandfonline.com/loi/kepi20>

SETDB2 and RIOX2 are differentially expressed among renal cell tumor subtypes, associating with prognosis and metastization

Maria João Ferreira, Ana Sílvia Pires-Luís, Márcia Vieira-Coimbra, Pedro Costa-Pinheiro, Luís Antunes, Paula C. Dias, Francisco Lobo, Jorge Oliveira, Céline S. Gonçalves, Bruno M. Costa, Rui Henrique & Carmen Jerónimo

To cite this article: Maria João Ferreira, Ana Sílvia Pires-Luís, Márcia Vieira-Coimbra, Pedro Costa-Pinheiro, Luís Antunes, Paula C. Dias, Francisco Lobo, Jorge Oliveira, Céline S. Gonçalves, Bruno M. Costa, Rui Henrique & Carmen Jerónimo (2017) SETDB2 and RIOX2 are differentially expressed among renal cell tumor subtypes, associating with prognosis and metastization, *Epigenetics*, 12:12, 1057-1064, DOI: [10.1080/15592294.2017.1385685](https://doi.org/10.1080/15592294.2017.1385685)

To link to this article: <https://doi.org/10.1080/15592294.2017.1385685>



Accepted author version posted online: 03 Nov 2017.
Published online: 22 Jan 2018.



Submit your article to this journal [↗](#)



Article views: 116



View related articles [↗](#)



View Crossmark data [↗](#)

RESEARCH PAPER



SETDB2 and RIOX2 are differentially expressed among renal cell tumor subtypes, associating with prognosis and metastization

Maria João Ferreira^{a,#}, Ana Sílvia Pires-Luís^{a,b,#}, Márcia Vieira-Coimbra^a, Pedro Costa-Pinheiro^a, Luís Antunes^c, Paula C. Dias^b, Francisco Lobo^d, Jorge Oliveira^d, Céline S. Gonçalves^{e,f}, Bruno M. Costa^{e,f}, Rui Henrique^{ib a,b,g,*}, and Carmen Jerónimo^{ib a,g,*}

^aCancer Biology and Epigenetics Group – Research Center, Portuguese Oncology Institute of Porto, Porto, Portugal; ^bDepartments of Pathology, Portuguese Oncology Institute of Porto, Porto, Portugal; ^cDepartments of Epidemiology, Portuguese Oncology Institute of Porto, Porto, Portugal; ^dDepartments of Urology, Portuguese Oncology Institute of Porto, Porto, Portugal; ^eLife and Health Sciences Research Institute (ICVS), School of Health Sciences, University of Minho, Braga, Portugal; ^fICVS/3B's – PT Government Associate Laboratory, Braga/Guimarães, Portugal; ^gDepartment of Pathology and Molecular Immunology, Institute of Biomedical Sciences Abel Salazar (ICBAS) – University of Porto, Porto, Portugal

ABSTRACT

Increasing detection of small renal masses by imaging techniques entails the need for accurate discrimination between benign and malignant renal cell tumors (RCTs) as well as among malignant RCTs, owing to differential risk of progression through metastization. Although histone methylation has been implicated in renal tumorigenesis, its potential as biomarker for renal cell carcinoma (RCC) progression remains largely unexplored. Thus, we aimed to characterize the differential expression of histone methyltransferases (HMTs) and histone demethylases (HDMs) in RCTs to assess their potential as metastasis biomarkers. We found that *SETDB2* and *RIOX2* (encoding for an HMT and an HDM, respectively) expression levels was significantly altered in RCTs; these genes were further selected for validation by quantitative RT-PCR in 160 RCTs. Moreover, *SETDB2*, *RIOX2*, and three genes encoding for enzymes involved in histone methylation (*NO66*, *SETD3*, and *SMYD2*), previously reported by our group, were quantified (RT-PCR) in an independent series of 62 clear cell renal cell carcinoma (ccRCC) to assess its potential role in ccRCC metastasis development. Additional validation was performed using TCGA dataset. *SETDB2* and *RIOX2* transcripts were overexpressed in RCTs compared to renal normal tissues (RNTs) and in oncocytomas vs. RCCs, with ccRCC and papillary renal cell carcinoma (pRCC) displaying the lowest levels. Low *SETDB2* expression levels and higher stage independently predicted shorter disease-free survival. In our 62 ccRCC cohort, significantly higher *RIOX2*, but not *SETDB2*, expression levels were depicted in cases that developed metastasis during follow-up. These findings were not apparent in TCGA dataset. We concluded that *SETDB2* and *RIOX2* might be involved in renal tumorigenesis and RCC progression, especially in metastatic spread. Moreover, *SETDB2* expression levels might independently discriminate among RCC subgroups with distinct outcome, whereas higher *RIOX2* transcript levels might identify ccRCC cases with more propensity to endure metastatic dissemination.

ARTICLE HISTORY

Received 12 June 2017
Revised 17 September 2017
Accepted 22 September 2017

KEYWORDS

biomarker; histone methyltransferase; kidney cancer; metastasis; prognosis; renal cell tumor; renal cell carcinoma; *RIOX2*; *SETDB2*

Introduction

Kidney cancer incidence is increasing worldwide, with 62,700 new cases and 14,240 deaths estimated for 2016.¹ Increasing incidence has been attributed to the rising number of incidental small renal tumors diagnosed due to widespread use of imaging techniques, as well as to aging, obesity, and smoking, which are known risk factors for the development of kidney cancer.² Increased detection of small renal masses emphasizes the need for accurate discrimination not only between benign and malignant RCTs, but also among malignant RCTs subtypes. Indeed, renal cell carcinomas (RCCs) that are more likely to behave aggressively and to develop metastases should be clearly distinguished from those that will probably have a more indolent growth and might be managed more conservatively.^{3,4}

Among RCCs, the most frequent are clear cell renal cell carcinoma (ccRCC), papillary renal cell carcinoma (pRCC), and chromophobe renal cell carcinoma (chRCC). Whereas ccRCC is the histotype that more frequently develops metastases, pRCC is more frequently multifocal and chRCC is mostly an indolent cancer that rarely develops metastases, although its differential diagnosis with oncocytoma, a benign tumor, might be challenging.⁵

Metastasis is the foremost cause of cancer-related mortality, despite improvements in diagnosis, surgical techniques, patient care, and adjuvant therapies. Biologic heterogeneity of tumor cells, as well as differences in metastatic tumor microenvironment at different sites may influence response to therapy.⁶

CONTACT Carmen Jerónimo  carmenjeronimo@ipporto.min-saude.pt  Portuguese Oncology Institute of Porto; Research Center-LAB 3, F Bdg., 1st floor; Rua Dr António Bernardino de Almeida, 4200-072 Porto, Portugal.

[#]Joint first authors.

^{*}Joint senior authors.

Thus, understanding pathogenesis of metastases at cellular and molecular level has become a major goal in cancer research.^{6,7} Indeed, management of metastatic renal cell carcinoma (mRCC) remains a major clinical challenge. Although median survival of patients with mRCC has been increasing due to therapeutic advances, specifically in antiangiogenic drugs and tyrosine-kinase inhibitors (from approximately 12 months with cytokine therapy to more than 26 months with VEGF inhibitors therapy),⁸ 5-year survival for advanced kidney cancer was only 11.7% in the period 2007–2013.⁹ Albeit the proportion of patients with mRCC at diagnosis has declined, due to improved imaging techniques as well as more intense screening and incidental case ascertainment, a sizeable number of small RCCs (<4 cm diameter) may present renal capsule invasion, tumor thrombus, or lymphatic and distant metastasis,^{3,4} and their identification constitutes a major challenge.

Altered epigenetic homeostasis has been implicated in tumorigenesis and epigenetic-based biomarkers may assist in diagnosis, prognostication, and prediction of response to targeted therapy.¹⁰ Histone modifications and chromatin modulators, in particular, have been shown to play an important role in cancer progression.¹¹ In RCC, certain histone modifications associate with progression-free survival and correlate with pathological characteristics of tumors.¹² In addition, defects in epigenetic enzymes, involved in chromatin remodeling and packaging, have been implicated in development of RCTs, reflecting the role of these mechanisms in renal tumorigenesis.¹³ Herein, we aimed to investigate the potential of HMTs and HDMs expression as biomarkers of metastatic progression in RCC, using two independent RCT cohorts, complemented with external validation in TCGA dataset. We selected *SETDB2* (an HMT) and *RIOX2* (an HDM), based on an extended characterization of histone methyltransferases in RCTs, previously reported by our group.¹⁴ Additionally, we also tested *SMYD2* and *SETD3* (both HMT), as well as *NO66* (HDM), previously evaluated in our first RCT cohort and TCGA dataset,¹⁴ in the second cohort comprising ccRCC with indolent (non-metastatic) and aggressive (metastatic) disease.

Results

Validation of *RIOX2* and *SETDB2* expression in RCTs

RIOX2 and *SETDB2* expression levels were assessed by quantitative RT-PCR in a series of 160 RCTs and 13 RNTs. The results were fully concordant with those of the TaqMan[®] Array as both genes were significantly overexpressed in RCTs compared to RNTs (P value <0.0001 for *SETDB2* and <0.05 for *RIOX2*; (Fig. 1A and B). Moreover, *RIOX2* and *SETDB2* expression levels differed significantly between benign and malignant RCTs (Fig. 1C and D), and among the four RCT subtypes (Table 1). Oncocytomas displayed the highest *SETDB2* and *RIOX2* expression levels, followed by chRCC (Fig. 1E and F and Table 1). Pairwise comparisons demonstrated that *SETDB2* and *RIOX2* expression levels significantly differed between chRCC and both pRCC and ccRCC, and between pRCC and both chRCC and oncocytoma.

Furthermore, *SETDB2* transcript levels differed significantly between chRCCs and oncocytomas (Fig. 1E and F and Table 1).

SETDB2 and *RIOX2* expression levels and clinicopathological correlates

No significant differences in gender and age were apparent between patients and controls. In RCCs, no statistically significant associations were disclosed between *SETDB2* and *RIOX2* expression levels and Fuhrman or pathological stage categories. In RCTs, expression levels of both genes were significantly higher in females. Moreover, *RIOX2* expression levels significantly associated with patient's age (P value = 0.015). In ccRCCs and pRCCs, *SETDB2* expression levels were significantly lower in patients that developed metastases (Fig. 2A and B).

SETDB2 and *RIOX2* expression levels as prognostic markers

The median follow-up of RCC patients was 175 months (range: 2–375 months). A total of 15 patients died from RCC during this period. In univariable analysis, higher pathological stage (pT3 or higher) associated with shorter survival, whereas gender, age, histological subtype, and Fuhrman grade did not disclose any prognostic value within the available follow-up time. Disease-specific survival (DSS) analysis showed that low *SETDB2* and *RIOX2* levels were significantly associated with worse outcome (P value <0.01 and <0.05, respectively; (Fig. 3A and B). Concerning disease-free survival (DFS) analysis, low *SETDB2* levels significantly associated with shorter time to disease progression (P value <0.0001; Fig. 3C). The same trend was observed for *RIOX2*, but statistical significance was not reached (P value = 0.055; Fig. 3D).

In this series, only one case with local recurrence presented distant metastasis before local recurrence developed, thus DFS is equivalent to metastasis-free survival in this case. In multivariable analysis, a final model including *SETDB2* expression levels and pathological stage was predictive of disease-free survival. Indeed, higher risk of disease progression was depicted for patients with higher pathological stage [HR: 3.03 (1.16–7.80), P value = 0.024] and lower *SETDB2* expression levels [HR: 5.11 (1.72–15.24), P value = 0.003].

RIOX2, *SETDB2*, *SETD3*, *SMYD2*, and *NO66* expression and risk of metastization in ccRCC

No significant differences were apparent for gender (P value = 0.570) and age (P value = 0.402) between ccRCCs patients that developed metastases and those that did not (cohort #2). Furthermore, no statistically significant associations were disclosed between *SETDB2* and *RIOX2* expression levels and Fuhrman grade or pathological stage in this cohort. In this cohort, expression levels of *SETD3*, *SMYD2*, and *NO66*, which we have previously found to associate with shorter disease-specific and disease-free survival,¹⁴ did not significantly differ between the two groups of ccRCC patients (Fig. 4C and E). Concerning




Figure 1. Expression levels of *SETDB2* and *RIOX2* in cohort #1. A: *SETDB2* expression in renal cell tumors (RCTs) and renal normal tissues (RNTs). B: *RIOX2* expression in renal cell tumors (RCTs) and renal normal tissues (RNTs). C: *SETDB2* expression in benign tumors (oncocytoma) and malignant tumors [renal cell carcinoma (RCCs)]. D: *RIOX2* expression in benign tumors (oncocytoma) and malignant tumors (renal cell carcinoma (RCCs)). E: *SETDB2* expression in renal cell tumors subtypes. F: *RIOX2* expression in renal cell tumors subtypes. (*P* values: ****<0.0001; **<0.01; *<0.05).

RIOX2 and *SETDB2* expression levels, only the former differed significantly between metastasized and non-metastasized ccRCCs (Fig. 4B).

***RIOX2* and *SETDB2* expression in RCC Patients from TCGA Dataset**

In TCGA dataset, significantly lower *RIOX2* expression levels were found in RCC compared to RNT, contrarily to our results. Nevertheless, among RCCs, pairwise comparisons showed that *RIOX2* expression levels were significantly higher in chRCCs compared to ccRCCs and pRCCs, paralleling our findings. In ccRCCs from TCGA database, no statistically significant difference was disclosed for *RIOX2* expression levels between the group of patients that developed metastases and those that did not.

Concerning *SETDB2*, lower expression levels were depicted in RCC compared to RNT, as well. In line with our results, however, pairwise comparisons demonstrated that *SETDB2* expression levels were significantly higher in chRCCs compared

to ccRCCs and pRCCs, and expression levels significantly differed among subtypes. Considering only ccRCCs from TCGA database, no statistically significant difference was apparent for *SETDB2* expression levels between the group of patients that developed metastases and those that did not, paralleling our results.

Discussion

Due to the widespread use of imaging tests, the frequency of incidentally detected RCTs has significantly increased, consisting mainly of small and early stage tumors. However, as lymph node and distant metastases may occur even in small RCCs,¹⁵ and the latter constitute the main cause of RCC-related mortality,¹⁶ there is an unmet need for biomarkers capable of accurately discriminate tumors that will metastasize from those that will not, especially among pT1, which are the most amenable to nephron-sparing surgery. Because epigenetic-based biomarkers may assist in diagnosis, prognostication and prediction of response to targeted therapy,¹⁰ we hypothesized that histone modifications and chromatin modulators¹¹ might aid in the identification of RCCs more prone to recur and metastasize. Indeed, previous reports have shown that, in RCC, histone modifications are associated with pathological features and DFS,¹² and defects in chromatin remodelers and chromatin packaging are implicated in RCT development.¹³

In this study, we focused mostly on *RIOX2* and *SETDB2* expression levels as candidate biomarkers for RCC prognostication. Whereas *RIOX2* encodes for an HDM, displaying high transcript or protein levels in RCC and several other cancers, associating with poor prognosis,^{17–24} *SETDB2* encodes for an HMT involved in leukemogenesis,¹⁵ but was not previously associated with solid tumors. These two genes were selected based on previously published array data

Table 1. Comparison of *SETDB2* and *RIOX2* expression among renal normal tissue (RNT), renal cell tumors (RCT), renal cell carcinoma (RCC), and RCT histotypes. For histotype pairwise comparison, the values were statistically significant when *P* value <0.0125 (Bonferroni's correction).

	<i>SETDB2</i> (<i>P</i> value)	<i>RIOX2</i> (<i>P</i> value)
RNT vs. RCT	<0.001	<0.05
RNT vs. RCC	0.001	0.444
Oncocytoma vs. RCC	<0.001	<0.001
ccRCC vs. pRCC	0.392	0.658
ccRCC vs. chRCC	<0.001	<0.001
ccRCC vs. oncocytoma	<0.001	<0.001
pRCC vs. chRCC	<0.001	<0.001
pRCC vs. oncocytoma	<0.001	<0.001
chRCC vs. oncocytoma	<0.001	0.131




Figure 2. Expression levels of *SETDB2* (A) and *RIOX2* (B) in clear cell renal cell carcinomas and papillary renal cell carcinomas (cohort #1) with or without metastasis (*P* value <0.05).

from our team,¹⁴ which were confirmed through analysis of cohort #1 tissue samples. Indeed, both *RIOX2* and *SETDB2* were found overexpressed in RCTs compared to non-paired normal renal tissues, suggesting that their deregulation is associated with neoplastic transformation of renal parenchymal cells. Interestingly, renal oncocytomas displayed the

highest *RIOX2* and *SETDB2* expression levels, significantly differing from RCCs. This result might be of practical value for distinction between oncocytoma and chRCC, especially the eosinophilic variant, as both histotypes display variable degree of morphological overlap, rendering differential diagnosis problematic, particularly in small core biopsies.²⁵




Figure 3. Kaplan-Meier estimated disease-specific survival curves and disease-free survival curves for *SETDB2* (respectively A and C) and *RIOX2* (respectively B and D).




Figure 4. Expression levels of *SETDB2* (A), *RIOX2* (B), *SETD3* (C), *SMYD2* (D), and *NO66* (E) in clear cell renal cell carcinomas (cohort #2) with or without metastasis (P value <0.05).

Unexpectedly, when TCGA dataset was interrogated for *RIOX2* and *SETDB2* expression in renal tissues, RCCs displayed significantly lower expression levels than normal renal tissues, contrarily to our findings in cohort #1. Nonetheless, the expression ranking among RCCs in TCGA paralleled our findings, with chRCC displaying the highest expression levels, compared to ccRCC and pRCC. This discrepancy might be related with the origin of the “normal renal tissue” analyzed. Indeed, whereas in our study RNT samples derived from kidneys not harboring RCC, those of TCGA dataset were collected from macroscopically normal looking areas of organs involved in RCC. As we have previously shown, these “paired” normal renal tissue samples disclose significant epigenetic alterations that may precede neoplastic transformation.²⁶ Thus, the results from cohort #1 and TCGA dataset analysis may not be directly comparable.

A major goal of this study was to ascertain the prognostic value of *RIOX2* and *SETDB2* expression levels in RCCs. Whereas, in univariable analysis, lower *RIOX2* and *SETDB2* expression levels associated with worse DSS, only lower *SETDB2* levels associated with worse DFS. Interestingly, among standard clinicopathological parameters, only stage reached statistical significance. In multivariable analysis, however, only low *SETDB2* expression and stage retained statistical significance, suggesting that assessment of *SETDB2* expression might add relevant prognostic information for the management of RCC patients. Although a previous report has associated higher *RIOX2* immunopositivity with shorter DSS in RCC,¹⁷ these results are not directly comparable with ours as we did not assess protein expression and the proportion of RCC subtypes also differed. Indeed, in our series, survival analysis results were mostly influenced by ccRCC and pRCC, which displayed the lowest expression levels among RCCs. Moreover, *RIOX2* overexpression has been associated with worse prognosis in esophageal cancer²⁴ but with favorable outcome in lung cancer,²⁷ emphasizing that the biological and clinical impact of *RIOX2*

expression is strongly dependent on the primary location and the specific cancer type.

Because DFS was analogous to metastasis-free survival in cohort #1, we looked for differences in *SETDB2* expression levels among tumors with and without metastasis. Interestingly, we found significantly lower *SETDB2* expression in ccRCC and pRCC (the more clinically aggressive histotypes) with metastasis. However, when we attempted to validate these findings in an independent cohort comprising ccRCC with and without metastasis (cohort #2) and in ccRCCs from TCGA dataset, no significant differences were disclosed. Nonetheless, significantly higher *RIOX2* transcript levels were depicted in ccRCC that developed metastasis during follow-up compared to matched ccRCC without metastasis, suggesting that *RIOX2* expression might be a biomarker of progression (metastization) in ccRCC. Interestingly, high *RIOX2* expression levels have been associated with development of lymphatic or distant metastasis in bile duct, gastric and pancreatic carcinomas.^{22,23,28} In TCGA dataset, however, no differences in *RIOX2* expression levels were apparent between ccRCCs that developed metastases and those that did not. These discrepancies might be due to differences in follow-up time and enrolment criteria, as we excluded from analysis cases that presented metastasis at diagnosis and only analyzed cases in which metastases developed after curative-intent surgical treatment. Although we further evaluated three genes encoding for histone modifying enzymes (*SMYD2*, *SETD3*, and *NO66*) previously shown to be associated with worse prognosis in RCC¹⁴ in cohort #2, their expression levels did not significantly differ between ccRCC with and without metastasis.

Although our findings might be limited by the sample size, it should be emphasized that the most frequent histotypes are represented, whereas in many studies only ccRCC cases have been included. Moreover, survival analysis is based on long-term follow-up data, including two patient cohorts and TCGA dataset. Finally, although candidate biomarkers were previously

identified in array-based analysis, validation in independent patient's series was undertaken, whereas several previous studies proposing array-based biomarkers for RCC, have not validated them or have just performed validation in limited series of patients,^{29–32} precluding the assessment of its clinical usefulness.

In conclusion, our results suggest that *SETDB2* and *RIOX2* might be involved in renal tumorigenesis and RCC progression, especially in metastatic spread. Moreover, *SETDB2* expression levels might independently discriminate among RCC patient subgroups with distinct outcome, whereas higher *RIOX2* transcript levels might identify ccRCC cases with more propensity to endure metastatic dissemination.

Materials and Methods

Patients and sample collection

A series of 160 RCTs (cohort #1) comprising 40 cases of each subtype (ccRCC, pRCC, chRCC, and oncocytoma) was prospectively collected from patients consecutively diagnosed and submitted to nephrectomy at the Portuguese Oncology Institute of Porto (IPO Porto). As controls, 13 renal normal tissue (RNT) samples were procured from patients submitted to nephro-ureterectomy due to upper urinary tract urothelial carcinoma, not involving the renal parenchyma. All tissues were immediately frozen and stored at -80°C . Sampling of more than 70% of malignant cells was confirmed using two slides stained with hematoxylin and eosin (H&E) taken before and after frozen section collection for RNA extraction. Relevant clinical data was retrieved from clinical charts.

An independent series of 62 ccRCCs (cohort #2) comprising 31 ccRCCs that have developed metastasis and 31 ccRCCs that did not progress, matched for gender, age, tumor size, grade, and stage at diagnosis were also retrieved from the archives. Tissue samples were prepared as described above.

This study was approved by the ethics committee of IPO Porto (CES-IPOPFG-EPE 518/10).

RNA extraction

For RNA extraction, samples were suspended in TRIzol[®] reagent (Invitrogen[™], Carlsbad, CA, USA; Cat. #15596018) and chloroform (Merck Millipore, Darmstadt, Germany; Cat. #MCX10601) was added after the cells were lysed. RNA concentrations and purity ratios were determined using a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). Samples were then stored at -80°C .

HMT and HDM selection

Selection of candidate HMTs and HDMs was based on the results of previously reported custom made TaqMan[®] Array 96-Well expression plates (Applied Biosystems[®], Foster City, CA, USA; Cat. #4391528).¹⁴ *SMYD2*, *SETD3*, and *NO66* had been previously validated and found to be associated with shorter disease-specific and disease-free survival.¹⁴ Additionally, *SETDB2* and *RIOX2* expression also displayed high fold variation between RCTs and RCCs, as well as between chRCC

and oncocytomas, and were thus selected for further validation. Both presented higher expression in RCT compared to RNT, as well as in oncocytomas than in chRCC.

Validation of selected enzymes

RIOX2 and *SETDB2* mRNA expression levels were firstly evaluated in cohort #1. Subsequently, *RIOX2*, *SETDB2*, *SMYD2*, *NO66*, and *SETD3* expression was assessed in cohort #2.

For validation in cohort #1, 300 ng of mRNA was reverse transcribed and amplified using TransPlex[®] Whole Transcriptome Amplification Kit (Sigma-Aldrich[®], St. Louis, MO, USA) with subsequent purification using QIAquick PCR Purification Kit (QIAGEN, Germany), according to manufacturer's instructions. *RIOX2* and *SETDB2* mRNA levels were evaluated using TaqMan[®] Gene Expression Assays [Applied Biosystems[®], Foster City, CA, USA; Hs99999908 m1 (*GUSβ*), Hs99999901 s1 (18S), Hs01126272 m1 (*SETDB2*), Hs00262155 m1 (*RIOX2*)] according to manufacturer's instructions. For each sample, expression levels were normalized using two internal reference gene, *GUSβ* and *18S*, according to the formula: target gene relative expression = target gene expression level / ((*GUSβ* expression level + *18S* expression level) / 2). Each plate included multiple non-template controls and serial dilutions of a cDNA Human Reference Total RNA (Agilent Technologies, La Jolla, CA, USA; Cat. #750500) to construct a standard curve.

For validation in cohort #2, 1 μg of total RNA was reverse transcribed using the High Capacity cDNA Reverse Transcription kit according to the manufacturer's instructions. *RIOX2*, *SETDB2*, *SMYD2*, *NO66*, and *SETD3* mRNA levels were evaluated using TaqMan[®] Gene Expression Assays [Applied Biosystems[®], Foster City, CA, USA; Cat. #4331182 Hs00220210 m1 (*SMYD2*), Hs00260120 m1 (*SETD3*), Hs02743012 s1 (*NO66*), Hs99999908 m1 (*GUSβ*), Hs99999901 s1 (18S), Hs01126272 m1 (*SETDB2*), Hs00262155 m1 (*RIOX2*)] according to manufacturer's instructions. For each sample, expression levels were normalized using two internal reference gene, *GUSβ* and *18S*, according to the formula: target gene relative expression = target gene expression level / ((*GUSβ* expression level + *18S* expression level) / 2). Each plate included multiple non-template controls and serial dilutions of a cDNA Human Reference Total RNA (Agilent Technologies, La Jolla, CA, USA; Cat. #750500) in order to construct a standard curve.

TCGA dataset analysis in pRCC, chRCC, and ccRCCs patients

The Cancer Genome Atlas (TCGA) dataset was interrogated for data on *RIOX2* and *SETDB2* expression and clinical information, when available, from ccRCCs, pRCCs, and chRCCs patients. All expression data from samples hybridized at the University of North Carolina, Lineberger Comprehensive Cancer Center, using Illumina HiSeq 2000 RNA Sequencing version 2 analysis, were downloaded from TCGA data matrix (<http://tcga-data.nci.nih.gov/tcga/tcgaDownload.jsp>). This dataset included 533 ccRCC, 290 pRCC, and 66 chRCC. The provided value was pre-processed and normalized according to "level 3" specifications of TCGA (see <http://cancergenome.nih.gov/dataportal/> for details). Biospecimen Core Resources (BCRs)

provided the clinical data of each patient. This data is available for download through TCGA data matrix (<http://tcga-data.nci.nih.gov/tcga/dataAccessMatrix.htm>).

Statistical analysis

Non-parametric tests were used to ascertain statistical significance of comparisons among groups. Kruskal-Wallis test (KW) was used for comparisons among multiple groups and Mann-Whitney U test (MW) was used for pairwise comparisons. The prognostic significance of clinicopathological variables (age, gender, histological subtype, pathological stage, Fuhrman grade) and HMTs and HDMs expression levels was assessed by constructing disease-specific and disease-free survival curves using the Kaplan-Meier method with log-rank test (univariable test). The expression levels of *SETDB2* and *RIOX2* were classified as low or high based on the cutoff value of 25th percentile for *SETDB2* expression and 75th percentile for *RIOX2*. A Cox-regression model comprising the different variables (multivariable test) was also constructed. For this analysis, the 120 RCC patients from cohort #1 were included.

Statistical significance was set at *P* value <0.05. Bonferroni correction was applied for pairwise comparisons following multiple groups' analyses. Statistical analysis was performed using SPSS software for Windows, version 22.0 (IBM-SPSS Inc., Chicago, IL, USA), and graphs were built using GraphPad Prism 6.0 software for Windows (GraphPad Software Inc., La Jolla, CA, USA).

Funding


This study was funded by research grants from Research Center of Portuguese Oncology Institute – Porto (CI-IPOP 4–2012 and CI-IPOP 27) and from Associação Portuguesa de Urologia (APU-2010). ASP-L was supported by FCT-Fundação para a Ciência e a Tecnologia fellowship (SFRH/SINTD/94217/2013). CSG is supported by FCT-Fundação para a Ciência e Tecnologia PhD fellowships (SFRH/BD/92786/2013) and BMC is funded by FCT-Fundação para a Ciência e a Tecnologia (IF/00601/2012).

Disclosure statement

None of the authors have any conflict of interest to declare.

ORCID

Rui Henrique  <http://orcid.org/0000-0003-3171-4666>

Carmen Jerónimo  <http://orcid.org/0000-0003-4186-5345>

References

- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin.* 2016;66:7–30. doi:10.3322/caac.21332.
- Znaor A, Lortet-Tieulent J, Laversanne M, Jemal A, Bray F. International variations and trends in renal cell carcinoma incidence and mortality. *Eur Urol.* 2015;67:519–30. doi:10.1016/j.eururo.2014.10.002. PMID:25449206
- Guethmundsson E, Hellborg H, Lundstam S, Erikson S, Ljungberg B, Swedish Kidney Cancer Quality Register G. Metastatic potential in renal cell carcinomas ≤ 7 cm: Swedish Kidney Cancer Quality Register data. *Eur Urol.* 2011;60:975–82. doi:10.1016/j.eururo.2011.06.029. PMID:21741160
- Jewett MA, Mattar K, Basiuk J, Morash CG, Pautler SE, Siemens DR, Tanguay S, Rendon RA, Gleave ME, Drachenberg DE, et al. Active surveillance of small renal masses: progression patterns of early stage kidney cancer. *Eur Urol.* 2011;60:39–44. doi:10.1016/j.eururo.2011.03.030. PMID:21477920
- Moch H, Humphrey PA, Ulbright TM, Reuter VE. WHO Classification of Tumours of the Urinary System and Male Genital Organs. Lyon, France: IARC Press, 2016.
- Sleeman JP. The metastatic niche and stromal progression. *Cancer Metastasis Rev.* 2012;31:429–40. doi:10.1007/s10555-012-9373-9. PMID:22699312
- Gupta GP, Massague J. Cancer metastasis: building a framework. *Cell.* 2006;127:679–95. doi:10.1016/j.cell.2006.11.001. PMID:17110329
- Rodriguez-Vida A, Hutson TE, Bellmunt J, Strijbos MH. New treatment options for metastatic renal cell carcinoma. *ESMO Open.* 2017;2:e000185. doi:10.1136/esmoopen-2017-000185. PMID:28761748
- SEER cancer stat facts: kidney and renal pelvis cancer. Bethesda M, National Cancer Institute.
- Bianco-Miotto T, Chiam K, Buchanan G, Jindal S, Day TK, Thomas M, Pickering MA, O'Loughlin MA, Ryan NK, Raymond WA, et al. Global levels of specific histone modifications and an epigenetic gene signature predict prostate cancer progression and development. *Cancer Epidemiol Biomarkers Prev.* 2010;19:2611–22. doi:10.1158/1055-9965.EPI-10-0555. PMID:20841388
- Chi P, Allis CD, Wang GG. Covalent histone modifications—miswritten, misinterpreted and mis-erased in human cancers. *Nature reviews Cancer.* 2010;10:457–69. doi:10.1038/nrc2876. PMID:20574448
- Ramakrishnan S, Ellis L, Pili R. Histone modifications: implications in renal cell carcinoma. *Epigenomics.* 2013; 5:453–62. doi:10.2217/epi.13.40. PMID:23895657
- Larkin J, Goh XY, Vetter M, Pickering L, Swanton C. Epigenetic regulation in RCC: opportunities for therapeutic intervention? *Nat Rev Urol.* 2012;9:147–55. doi:10.1038/nrurol.2011.236. PMID:22249190
- Pires-Luis AS, Vieira-Coimbra M, Vieira FQ, Costa-Pinheiro P, Silva-Santos R, Dias PC, Antunes L, Lobo F, Oliveira J, Goncalves CS, et al. Expression of histone methyltransferases as novel biomarkers for renal cell tumor diagnosis and prognostication. *Epigenetics.* 2015;10:1033–43. doi:10.1080/15592294.2015.1103578. PMID:26488939
- Mabuchi H, Fujii H, Calin G, Alder H, Negrini M, Rassenti L, Kipps TJ, Bullrich F, Croce CM. Cloning and characterization of CLLD6, CLLD7, and CLLD8, novel candidate genes for leukemogenesis at chromosome 13q14, a region commonly deleted in B-cell chronic lymphocytic leukemia. *Cancer Res.* 2001;61:2870–7. PMID:11306461
- Chin AI, Lam JS, Figlin RA, Belldgrun AS. Surveillance strategies for renal cell carcinoma patients following nephrectomy. *Rev Urol.* 2006;8:1–7. PMID:16985554
- Ishizaki H, Yano H, Tsuneoka M, Ogasawara S, Akiba J, Nishida N, Kojiro S, Fukahori S, Moriya F, Matsuoka K, et al. Overexpression of the myc target gene *Mina53* in advanced renal cell carcinoma. *Pathol Int.* 2007;57:672–80. doi:10.1111/j.1440-1827.2007.02156.x. PMID:17803656
- Teye K, Tsuneoka M, Arima N, Koda Y, Nakamura Y, Ueta Y, Shirouzu K, Kimura H. Increased expression of a Myc target gene *Mina53* in human colon cancer. *Am J Pathol.* 2004;164:205–16. doi:10.1016/S0002-9440(10)63111-2. PMID:14695334
- Ogasawara S, Komuta M, Nakashima O, Akiba J, Tsuneoka M, Yano H. Accelerated expression of a Myc target gene *Mina53* in aggressive hepatocellular carcinoma. *Hepatol Res.* 2010;40:330–6. doi:10.1111/j.1872-034X.2009.00604.x. PMID:20070393
- Komiya K, Sueoka-Aragane N, Sato A, Hisatomi T, Sakuragi T, Mitsuoka M, Sato T, Hayashi S, Izumi H, Tsuneoka M, et al. *Mina53*, a novel c-Myc target gene, is frequently expressed in lung cancers and exerts oncogenic property in NIH/3T3 cells. *J Cancer Res Clin Oncol.* 2010;136:465–73. doi:10.1007/s00432-009-0679-0. PMID:19756735
- Teye K, Arima N, Nakamura Y, Sakamoto K, Sueoka E, Kimura H, Tsuneoka M. Expression of Myc target gene *mina53* in subtypes of human lymphoma. *Oncol Reports.* 2007;18:841–8.

22. Tan XP, Zhang Q, Dong WG, Lei XW, Yang ZR. Upregulated expression of Mina53 in cholangiocarcinoma and its clinical significance. *Oncol Letters*. 2012;3:1037–41.
23. Tan XP, Dong WG, Zhang Q, Yang ZR, Lei XF, Ai MH. Potential effects of Mina53 on tumor growth in human pancreatic cancer. *Cell Biochem Biophys*. 2014;69:619–25. doi:10.1007/s12013-014-9841-7. PMID:24522517
24. Tsuneoka M, Fujita H, Arima N, Teye K, Okamura T, Inutsuka H, Koda Y, Shirouzu K, Kimura H. Mina53 as a potential prognostic factor for esophageal squamous cell carcinoma. *Clin Cancer Res*. 2004;10:7347–56. doi:10.1158/1078-0432.CCR-03-0543. PMID:15534111
25. Volpe A, Finelli A, Gill IS, Jewett MA, Martignoni G, Polascik TJ, Remzi M, Uzzo RG. Rationale for percutaneous biopsy and histologic characterisation of renal tumours. *Eur Urol*. 2012;62:491–504. doi:10.1016/j.eururo.2012.05.009. PMID:22633318
26. Costa VL, Henrique R, Ribeiro FR, Pinto M, Oliveira J, Lobo F, Teixeira MR, Jeronimo C. Quantitative promoter methylation analysis of multiple cancer-related genes in renal cell tumors. *BMC Cancer*. 2007;7:133. doi:10.1186/1471-2407-7-133. PMID:17645803
27. Komiya K, Sueoka-Aragane N, Sato A, Hisatomi T, Sakuragi T, Mitsuoka M, Sato T, Hayashi S, Izumi H, Tsuneoka M, et al. Expression of Mina53, a novel c-Myc target gene, is a favorable prognostic marker in early stage lung cancer. *Lung Cancer*. 2010;69:232–8. doi:10.1016/j.lungcan.2009.10.010. PMID:19914733
28. Xing J, Wang K, Liu PW, Miao Q, Chen XY. Mina53, a novel molecular marker for the diagnosis and prognosis of gastric adenocarcinoma. *Oncol Reports*. 2014;31:634–40. doi:10.3892/or.2013.2918.
29. Jeronimo C, Henrique R. Epigenetic biomarkers in urological tumors: A systematic review. *Cancer Lett*. 2014;342:264–74. doi:10.1016/j.canlet.2011.12.026. PMID:22198482
30. Junker K, Hindermann W, von Eggeling F, Diegmann J, Haessler K, Schubert J. CD70: a new tumor specific biomarker for renal cell carcinoma. *J Urol*. 2005;173:2150–3. doi:10.1097/01.ju.0000158121.49085.ba. PMID:15879877
31. Meyer HA, Tolle A, Jung M, Fritzsche FR, Haendler B, Kristiansen I, Gaspert A, Johannsen M, Jung K, Kristiansen G. Identification of stanniocalcin 2 as prognostic marker in renal cell carcinoma. *Eur Urol*. 2009;55:669–78. doi:10.1016/j.eururo.2008.04.001. PMID:18450365
32. Redova M, Poprach A, Nekvindova J, Iliev R, Radova L, Lakomy R, Svoboda M, Vyzula R, Slaby O. Circulating miR-378 and miR-451 in serum are potential biomarkers for renal cell carcinoma. *J Trans Med*. 2012;10:55. doi:10.1186/1479-5876-10-55.