

# Strategies to improve retention in randomised trials (Review)

Brueton VC, Tierney J, Stenning S, Harding S, Meredith S, Nazareth I, Rait G



**THE COCHRANE  
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2013, Issue 12

<http://www.thecochranelibrary.com>

**WILEY**

---

Strategies to improve retention in randomised trials (Review)

Copyright © 2013 The Authors. The Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

## TABLE OF CONTENTS

HEADER . . . . .	1
ABSTRACT . . . . .	1
PLAIN LANGUAGE SUMMARY . . . . .	2
BACKGROUND . . . . .	3
OBJECTIVES . . . . .	4
METHODS . . . . .	4
RESULTS . . . . .	8
Figure 1. . . . .	9
DISCUSSION . . . . .	15
AUTHORS' CONCLUSIONS . . . . .	18
ACKNOWLEDGEMENTS . . . . .	18
REFERENCES . . . . .	19
CHARACTERISTICS OF STUDIES . . . . .	27
DATA AND ANALYSES . . . . .	88
Analysis 1.1. Comparison 1 Addition of incentive vs none: main analysis, Outcome 1 Retention. . . . .	93
Analysis 2.1. Comparison 2 Addition of incentive: sensitivity analysis: quasi-randomised trials removed, Outcome 1 Retention. . . . .	94
Analysis 3.1. Comparison 3 Addition of incentive: separating research arms of non-factorial trials (three-/four-arm trials), Outcome 1 Retention. . . . .	95
Analysis 4.1. Comparison 4 Addition of telephone follow-up vs incentive, Outcome 1 Retention. . . . .	97
Analysis 5.1. Comparison 5 Addition of monetary incentive to both study arms, Outcome 1 Retention. . . . .	98
Analysis 6.1. Comparison 6 Addition of monetary incentive vs offer of incentive, Outcome 1 Retention. . . . .	99
Analysis 7.1. Comparison 7 Enhanced letter versus standard letter: main analysis, Outcome 1 Retention. . . . .	99
Analysis 8.1. Comparison 8 Communication strategies letter: total design method, Outcome 1 Retention. . . . .	100
Analysis 9.1. Comparison 9 Communication strategies post: main analysis, Outcome 1 Retention. . . . .	101
Analysis 10.1. Comparison 10 Communication strategies: additional reminder vs usual follow-up: main analysis, Outcome 1 Retention. . . . .	102
Analysis 11.1. Comparison 11 Communication strategies additional reminder to trial site vs usual reminder (ICC 0.054), Outcome 1 Retention. . . . .	103
Analysis 12.1. Comparison 12 Communication strategies: questionnaire administered early vs late, Outcome 1 Retention. . . . .	103
Analysis 13.1. Comparison 13 Communication strategies: type of reminder: main analysis, Outcome 1 Retention. . . . .	104
Analysis 14.1. Comparison 14 Questionnaire strategies: new vs standard questionnaire: main analysis, Outcome 1 Retention. . . . .	105
Analysis 15.1. Comparison 15 Questionnaire strategies: new vs standard questionnaire: sensitivity analysis quasi-randomised trial McColl, Outcome 1 Retention. . . . .	106
Analysis 16.1. Comparison 16 Behavioural strategies: main analysis, Outcome 1 Retention. . . . .	107
Analysis 17.1. Comparison 17 Case management, Outcome 1 Retention. . . . .	107
Analysis 18.1. Comparison 18 Methodology strategies, Outcome 1 Retention. . . . .	108
ADDITIONAL TABLES . . . . .	108
APPENDICES . . . . .	113
CONTRIBUTIONS OF AUTHORS . . . . .	125
DECLARATIONS OF INTEREST . . . . .	125
SOURCES OF SUPPORT . . . . .	126
DIFFERENCES BETWEEN PROTOCOL AND REVIEW . . . . .	126
INDEX TERMS . . . . .	126

[Methodology Review]

# Strategies to improve retention in randomised trials

Valerie C Brueton<sup>1</sup>, Jayne Tierney<sup>2</sup>, Sally Stenning<sup>3</sup>, Seeromanie Harding<sup>4</sup>, Sarah Meredith<sup>3</sup>, Irwin Nazareth<sup>5</sup>, Greta Rait<sup>5</sup>

<sup>1</sup>Meta-analysis Group, MRC Clinical Trials Unit, London, UK. <sup>2</sup>Meta-analysis Group, MRC Clinical Trials Unit at UCL, London, UK. <sup>3</sup>MRC Clinical Trials Unit, London, UK. <sup>4</sup>Social and Public Health Sciences Unit, Medical Research Council, Glasgow, UK. <sup>5</sup>Research Department of Primary Care and Population Health, University College London, London, UK

Contact address: Valerie C Brueton, Meta-analysis Group, MRC Clinical Trials Unit, 125 Kingsway, London, WC2B 6NH, UK. [vcb@ctu.mrc.ac.uk](mailto:vcb@ctu.mrc.ac.uk).

**Editorial group:** Cochrane Methodology Review Group.

**Publication status and date:** New, published in Issue 12, 2013.

**Review content assessed as up-to-date:** 13 June 2013.

**Citation:** Brueton VC, Tierney J, Stenning S, Harding S, Meredith S, Nazareth I, Rait G. Strategies to improve retention in randomised trials. *Cochrane Database of Systematic Reviews* 2013, Issue 12. Art. No.: MR000032. DOI: 10.1002/14651858.MR000032.pub2.

Copyright © 2013 The Authors. The Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration. This is an open access article under the terms of the [Creative Commons Attribution](#) Licence, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

### Background

Loss to follow-up from randomised trials can introduce bias and reduce study power, affecting the generalisability, validity and reliability of results. Many strategies are used to reduce loss to follow-up and improve retention but few have been formally evaluated.

### Objectives

To quantify the effect of strategies to improve retention on the proportion of participants retained in randomised trials and to investigate if the effect varied by trial strategy and trial setting.

### Search methods

We searched the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, PreMEDLINE, EMBASE, PsycINFO, DARE, CINAHL, Campbell Collaboration's Social, Psychological, Educational and Criminological Trials Register, and ERIC. We handsearched conference proceedings and publication reference lists for eligible retention trials. We also surveyed all UK Clinical Trials Units to identify further studies.

### Selection criteria

We included eligible retention trials of randomised or quasi-randomised evaluations of strategies to increase retention that were embedded in 'host' randomised trials from all disease areas and healthcare settings. We excluded studies aiming to increase treatment compliance.

### Data collection and analysis

We contacted authors to supplement or confirm data that we had extracted. For retention trials, we recorded data on the method of randomisation, type of strategy evaluated, comparator, primary outcome, planned sample size, numbers randomised and numbers retained. We used risk ratios (RR) to evaluate the effectiveness of the addition of strategies to improve retention. We assessed heterogeneity between trials using the Chi<sup>2</sup> and I<sup>2</sup> statistics. For main trials that hosted retention trials, we extracted data on disease area, intervention, population, healthcare setting, sequence generation and allocation concealment.

---

**Strategies to improve retention in randomised trials (Review)**

Copyright © 2013 The Authors. The Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

## Main results

We identified 38 eligible retention trials. Included trials evaluated six broad types of strategies to improve retention. These were incentives, communication strategies, new questionnaire format, participant case management, behavioural and methodological interventions. For 34 of the included trials, retention was response to postal and electronic questionnaires with or without medical test kits. For four trials, retention was the number of participants remaining in the trial. Included trials were conducted across a spectrum of disease areas, countries, healthcare and community settings. Strategies that improved trial retention were addition of monetary incentives compared with no incentive for return of trial-related postal questionnaires (RR 1.18; 95% CI 1.09 to 1.28, P value < 0.0001), addition of an offer of monetary incentive compared with no offer for return of electronic questionnaires (RR 1.25; 95% CI 1.14 to 1.38, P value < 0.00001) and an offer of a GBP20 voucher compared with GBP10 for return of postal questionnaires and biomedical test kits (RR 1.12; 95% CI 1.04 to 1.22, P value < 0.005). The evidence that shorter questionnaires are better than longer questionnaires was unclear (RR 1.04; 95% CI 1.00 to 1.08, P value = 0.07) and the evidence for questionnaires relevant to the disease/condition was also unclear (RR 1.07; 95% CI 1.01 to 1.14). Although each was based on the results of a single trial, recorded delivery of questionnaires seemed to be more effective than telephone reminders (RR 2.08; 95% CI 1.11 to 3.87, P value = 0.02) and a 'package' of postal communication strategies with reminder letters appeared to be better than standard procedures (RR 1.43; 95% CI 1.22 to 1.67, P value < 0.0001). An open trial design also appeared more effective than a blind trial design for return of questionnaires in one fracture prevention trial (RR 1.37; 95% CI 1.16 to 1.63, P value = 0.0003).

There was no good evidence that the addition of a non-monetary incentive, an offer of a non-monetary incentive, 'enhanced' letters, letters delivered by priority post, additional reminders, or questionnaire question order either increased or decreased trial questionnaire response/retention. There was also no evidence that a telephone survey was either more or less effective than a monetary incentive and a questionnaire. As our analyses are based on single trials, the effect on questionnaire response of using offers of charity donations, sending reminders to trial sites and when a questionnaire is sent, may need further evaluation. Case management and behavioural strategies used for trial retention may also warrant further evaluation.

## Authors' conclusions

Most of the retention trials that we identified evaluated questionnaire response. There were few evaluations of ways to improve participants returning to trial sites for trial follow-up. Monetary incentives and offers of monetary incentives increased postal and electronic questionnaire response. Some other strategies evaluated in single trials looked promising but need further evaluation. Application of the findings of this review would depend on trial setting, population, disease area, data collection and follow-up procedures.

## PLAIN LANGUAGE SUMMARY

### Methods that might help to keep people in randomised trials

#### Background

Most trials follow people up to collect data through personal contact after they have been recruited. Some trials get data from other sources, such as routine collected data or disease registers. There are many ways to collect data from people in trials, and these include using letters, the internet, telephone calls, text messaging, face-to-face meetings or the return of medical test kits. Most trials have missing data, for example, because people are too busy to reply, are unable to attend a clinic, have moved or no longer want to participate. Sometimes data has not been recorded at study sites, or are not sent to the trial co-ordinating centre. Researchers call this 'loss to follow-up', 'drop out' or 'attrition' and it can affect the trial's results. For example, if the people with the most or least severe symptoms do not return questionnaires or attend a follow-up visit, this will bias the findings of the trial. Many methods are used by researchers to keep people in trials. These encourage people to send back data by questionnaire, return to a clinic or hospital for trial-related tests, or be seen by a health or community care worker.

#### Study characteristics

This review identified methods that encouraged people to stay in trials. We searched scientific databases for randomised studies (where people are allocated to one of two or more possible treatments in a random manner) or quasi-randomised studies (where allocation is not really random, e.g. based on date of birth, order in which they attended clinic) that compared methods of increasing retention in trials. We included trials of participants from any age, gender, ethnic, cultural, language and geographic groups.

#### Key results

---

**Strategies to improve retention in randomised trials (Review)**

Copyright © 2013 The Authors. The Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

The methods that appeared to work were offering or giving a small amount of money for return of a completed questionnaire and enclosing a small amount of money with a questionnaire with the promise of a further small amount of money for return of a filled in questionnaire. The effect of other ways to keep people in trials is still not clear and more research is needed to see if these really do work. Such methods are shorter questionnaires, sending questionnaires by recorded delivery, using a trial design where people know which treatment they will receive, sending specially designed letters with a reply self addressed stamped envelope followed by a number of reminders, offering a donation to charity or entry into a prize draw, sending a reminder to the study site about participants to follow-up, sending questionnaires close to the time the patient was last followed-up, managing peoples' follow-up, conducting follow-up by telephone and changing the order of questionnaire questions.

### Quality of evidence

The methods that we identified were tested in trials run in many different disease areas and settings and, in some cases, were tested in only one trial. Therefore, more studies are needed to help decide whether our findings could be used in other research fields.

## BACKGROUND

### Description of the problem or issue

Randomised trials are the gold standard for evaluating the effectiveness and efficacy of interventions. Non-response or loss to follow-up within study groups in randomised trials can compromise study findings by reducing the power of a study to detect a true difference between the control and the intervention group. Differential loss to follow-up may lead to bias through exaggerated effects in favour of one of the groups. This can affect the generalisability and internal validity of the trial and the results (Fewtrell 2008; Schulz 2002).

Missing data from loss to follow-up can be dealt with statistically by various methods including, for example, imputing values based on valid assumptions about the missing data to give a conservative estimate of the treatment effect. However, the risk of bias still remains when trials do not collect adequate data to give accurate estimates (Hollis 1999). Schulz and colleagues suggested that less than 5% loss to follow-up may lead to minimum bias, while 20% loss to follow-up can threaten trial validity, although the pattern of loss to follow-up by treatment may also be an important factor (Schulz 2002). Loss to follow-up from randomised trials can sometimes go unreported and using different, but plausible, assumptions about outcomes for participants lost to follow-up can change the results of randomised trials.

A number of trials have retrospectively examined the predictors of loss to follow-up in different disease areas (Arnow 2007; Snow 2007; Villarruel 2006). In a trial for the treatment of chronic major depression, Arnow examined the predictors of time to, and reason for, dropout of participants (Arnow 2007). Ethnic minorities and participants with comorbid anxiety were more likely to drop out. In a randomised trial of a human immunodeficiency virus (HIV)

prevention intervention for Latino youths, English speakers were more likely to attend follow-up (Villarruel 2006). Snow examined the predictors of clinic attendance and dropout at the 11-year follow-up of the Lung Health study (Snow 2007). Age, gender, number of cigarettes smoked per day, marital status and whether participant's children smoked were predictors of clinic attendance. These analyses showed that attendance for follow-up can be trial and disease specific. An awareness of these factors can help trialists decide which strategies to adopt to improve retention in their randomised trial.

### Description of the methods being investigated

Strategies to improve trial retention include those designed to generate maximum data return or compliance to follow-up procedures. These can include frequency and timing of follow-up visits (follow-up shortly after randomisation versus long-term follow-up), nature of the outcome to be measured (survey based self reported outcomes versus morbidity or mortality reporting), target of the intervention (participants versus providers versus trial sites), and type of intervention (incentives versus communication strategies versus participant case management).

### How these methods might work

These retention strategies are designed to motivate participants (Leathem 2009), or the trial site to continue participating in a trial once they have been recruited and randomised. Some strategies are designed to encourage participants to identify with the trial and to promote a sense of value and belonging, for example, using trial identity cards. Other strategies are designed to keep participants engaged in the trial, for example, by sending participant

newsletters. To encourage a proactive approach to trial retention, strategies can be designed to target participants directly through letters, emails, telephone calls or to target them via the clinicians involved in participant follow-up, for example, through regular communication with trial sites. Strategies have been specifically developed to promote retention in areas of research where it is particularly challenging, such as mental health (Furimsky 2008; Loue 2008), weight loss (Couper 2007; Goldberg 2005), rare diseases (McKinstry 2007), substance abuse (El Khorazaty 2007), research involving minority ethnic groups (Eakin 2007; Loftin 2005; Villacorta 2007), and vulnerable groups such as older people (Burns 2008) or people with HIV (Anastasi 2005).

### Why it is important to do this review

As drop-out or incomplete data causes problems in the conduct, analysis and interpretation of randomised trials, it is important to identify retention strategies that minimise this loss as far as possible.

Davis and colleagues conducted a review of community-based trials published from 1990 to 1999 and described retention strategies and retention outcomes for this area (Davis 2002). Robinson and colleagues conducted a systematic review of strategies for retaining study participants (Robinson 2007). While both reviews identified studies providing data on retention rates from primary studies and strategies used to promote retention, these were not evaluated quantitatively in either review.

A systematic review of strategies to retain participants in population-based cohort studies found that providing incentives was consistently associated with retention in these studies and that response generally increased with increasing incentive value (Booker 2011). Reminder letters, repeat questionnaires and reminder calls also increased response rates. Furthermore, the Edwards et al. Cochrane methodology review on methods to increase response rates to postal and electronic questionnaires found that including monetary incentives, keeping the questionnaire short and contacting people before sending the questionnaire were ways to increase response rates (Edwards 2009). That review was not restricted to research exclusively within randomised trials and covered both healthcare and non-healthcare settings and it is difficult to know which of these strategies would be applicable to randomised trials in health care. Reasons for drop-out in cohort studies and surveys may differ from those in randomised trials. For example, in trials, participants may be randomised to a study group that is not their preferred choice and factors around randomisation and the type of intervention mean that strategies increasing retention in cohort studies and surveys cannot necessarily be extrapolated to randomised trials.

The challenges of boosting recruitment to randomised trials is often described alongside retention in the literature. Some similar strategies may be used in an attempt to both increase recruitment and improve retention, such as giving incentives together with ex-

tra information. Rendell et al. assessed the evidence for the effect of disincentives and incentives on the extent to which clinicians invite eligible people to participate in randomised trials of healthcare interventions (Rendell 2007). No randomised trials of interventions were identified and the authors concluded that some aspects of the conduct of the trial might affect a clinician's willingness to invite people to participate, for example, the way the clinician is invited to take part and the availability of support staff. In another Cochrane methodology review, Treweek et al. assessed strategies to improve recruitment to research studies (Treweek 2010), but recruitment to trials presents different challenges to participant engagement and follow-up. For example, strategies to market a trial and win over participants during the recruitment phase may be different to strategies to keep participants engaged in a trial (Francis 2007).

Many untested strategies are used by researchers to try to improve retention in randomised trials. Therefore, because loss to follow-up can compromise the validity of a trial's findings, delay results and, in some circumstances, increase the costs of the research, a systematic review is needed to assess the effect of strategies to improve retention in randomised trials.

## OBJECTIVES

To quantify the effect of strategies to improve retention in randomised trials.

To investigate if the effect varies by the type of strategy, trial setting and healthcare area.

## METHODS

### Criteria for considering studies for this review

#### Types of studies

We included completed randomised trials that compared strategies to increase retention embedded in host randomised trials (hereafter referred to as retention trials). The retention trials were embedded in real trials (host trials) and not hypothetical trials. The retention trials included at least one randomised comparison of two or more strategies to improve retention, or compared one or more strategies with no strategy. In anticipation of few trials, we included retention trials if they were randomised or quasi-randomised (e.g. had used alternation, date of birth or case record number as a method of allocating participants) (Lefebvre 2008).

Strategies to improve retention were designed for impact after participants were recruited and randomised to either the intervention

or control group of the main and the retention trial. We included trials to increase response to postal and electronic questionnaire. We excluded trials to increase recruitment only. We excluded cohort studies with embedded randomised retention trials, which were the subject of a separate systematic review (Booker 2011).

### Types of data

We included randomised and quasi-randomised retention trials within the context of a host randomised trial with participants from any age, gender, ethnic, cultural, language and geographic groups. We included unpublished and published participant retention data from randomised trials addressing healthcare (including all disciplines and disease areas) and non-healthcare (education, social sciences) topics. We also included trials set in the community that were healthcare related.

### Types of methods

We considered any strategy aimed at increasing retention, directed towards the clinician, researcher or participant. We included strategies compared with each other or with usual study procedures. We also included trials with any combination of strategies to increase retention. Strategies could be participant or trial management focused and include any of the following:

- strategies to motivate participants and clinicians (e.g. incentives or gifts);
- strategies to improve communication with participants or trial sites (e.g. enhanced letters);
- methodology strategies (e.g. shorter length of follow-up or variation in follow-up visit frequency);
- strategies to improve social support for participant retention.

### Types of outcome measures

#### Primary outcomes

We used retention (the proportion of participants retained) at the primary analysis point as defined in each individual retention trial as the primary outcome because it is easier to interpret than attrition/loss to follow-up (i.e. the proportion lost or not retained). In cases where the time point for measurement of the primary outcome was not predefined, we took the first time point reported for analysis. In most cases, this was final response. If retention at a number of time points was reported and no clear time point for the primary outcome for the retention trial was stated, we took data for the nearest time point to the intervention in the retention trial analyses.

#### Secondary outcomes

Retention of participants at secondary analysis points.

### Search methods for identification of studies

We designed a search strategy to identify published and unpublished randomised and quasi-randomised trials that assessed strategies to improve retention in randomised trials in healthcare, education and social science settings. We searched bibliographic databases for published trials and trial registers for trials that had not been fully published, or were unpublished or ongoing. We applied no language restrictions.

#### Electronic searches

Each search comprised an established filter to identify randomised trials plus free-text terms and database subject headings relating to reducing loss to follow-up or increasing retention (Appendix 1). Electronic databases searched included:

- the Cochrane Central Register of Controlled Trials (CENTRAL) (to May 2012);
- PreMEDLINE (to April 2010);
- MEDLINE (1950 to May 2012) (Appendix 2), EMBASE (1980 to May 2012) (Appendix 3) and PsycINFO (1806 to May 2012) (Appendix 4), searched using an Ovid platform;
- Database of Abstracts of Reviews of Effects (DARE, in *The Cochrane Library* May 2012);
- CINAHL (Cumulative Index to Nursing and Allied Health; 1981 to May 2012) (Appendix 5), using the EBSCOHost platform;
- Campbell Collaboration's Social, Psychological, Educational and Criminological Trials Register (C2-SPECTR <http://geb9101.gse.upenn.edu/>; searched May 2009 (website no longer accessible)) (Appendix 6);
- Education Resource Information Centre (ERIC) 1966 to May 2009) (Appendix 7), using Dialog Datarstar.

#### Searching other resources

We handsearched the reference lists of relevant publications and reviews to identify further trial reports (Horsley 2011) (Appendix 8). We also searched the abstracts of Society for Clinical Trials (SCT) meetings from 1980 to 2012, the Current Controlled Trials *meta*Register of Controlled Trials (*m*RCT) ([www.controlled-trials.com/mrct](http://www.controlled-trials.com/mrct)), the Cochrane Methodology Register (in *The Cochrane Library* to April 2012) and the World Health Organization (WHO) trials platform ([apps.who.int/trialsearch](http://apps.who.int/trialsearch)). We conducted a survey of Clinical Trial Units in the UK to identify further eligible trials not identified through other sources and the review was presented at the Society for Clinical Trials 31st Conference in Baltimore, USA in May 2010 and advertised on the Conference

notice board with the aim of identifying potentially eligible trials from outside the UK.

## Data collection and analysis

### Selection of studies

One review author (VB) selected potentially eligible trials from the titles and abstracts retrieved by the searches, using a predesigned study eligibility screening form. We were over inclusive when screening, 0.7% (168/24,304) of records identified were sent for screening to a second review author (GR), which is 23% (168/735) of all potentially eligible records identified. We obtained full-text papers and two review authors (VB, GR) reviewed potentially eligible trials for inclusion. We contacted study authors for electronic copies of papers that we could not access through library sources. We were able to obtain copies of all the potentially eligible papers that we wanted to screen. We resolved disagreements by discussion with a third review author (SS). When necessary, we sought information from the original investigators for potentially eligible trials where we wished to clarify eligibility.

### Data extraction and management

One review author (VB) extracted data from eligible retention trial and associated host trial papers and a second review author (JT) checked the entries. We reached consensus on any disparities by discussion with a third review author (SS). Data extracted for the host trial were aim, setting, disease area, comparators, primary outcome, sample size calculation, inclusion exclusion criteria, sequence generation and allocation concealment, and numbers randomised to each group. For the embedded retention trial, we extracted data for onset in relation to the host trial, source of the sample, aim, primary outcome and type of follow-up. The retention strategy details included type, frequency and timing of administration method of randomisation, numbers randomised, included and retained at primary analysis, and data required for the risk of bias assessment.

### Assessment of risk of bias in included studies

To assess the validity of each retention trial we judged them against the four domains of the Cochrane 'Risk of bias' tool (Higgins 2008a). To assess selection bias, we recorded how the allocation sequence was generated at study level and the methods used to conceal the allocation. We assessed performance bias by recording methods used to blind participants if considered appropriate to do so. For some interventions, participants could not be blinded to the intervention (e.g. where vouchers, cash or gifts were administered). However, in these cases, study personnel could be blinded to the

allocation if administration of the intervention was carried out by someone unaware of the allocation.

As retention is the subject of our review, and retention of participants is the primary outcome, attrition from the trials does not constitute a bias and has not been included in the 'Risk of bias' tables. We assessed each included retention trial for selective outcome reporting by recording the primary outcome for the trial and the outcomes for which results were reported. A judgement was made about each trial for each risk of bias domain assessed. For completed host trials (within which retention trials were embedded), we only assessed sequence generation and allocation concealment, in order to ensure the host trial was randomised.

### Measures of the effect of the methods

We calculated risk ratios (RR) and their 95% confidence intervals (CI) for retention to determine the effect of strategies on this outcome.

### Unit of analysis issues

For retention trials that randomised individuals and clusters, the unit of analysis was the participant. For cluster randomised trials that ignored clustering in the analysis, we inflated the standard errors (SE) to avoid overprecise estimates of effect as follows (Higgins 2008b).

1. We calculated the RR, 95% CI and SE based on participants in the usual way (i.e. ignoring clustering).
2. This standard error was then inflated using the design effect to get an adjusted SE: adjusted SE = SE  $\times$   $\sqrt{\text{design effect}}$ . With the design effect calculated as follows: design effect =  $1 + (M - 1) \text{ICC}$  where M = mean cluster size, ICC = the intracluster correlation coefficient.
3. Where published ICCs were not available, we used the mean ICC from appropriate external estimates for Land 2007. This was the mean of estimates for the return of EuroQol questionnaires (ICC = 0.054) from a source recommended by the *Cochrane Handbook for Systematic Reviews of Interventions* (Section 16.3.4) (Higgins 2008b) and [www.abdn.ac.uk/hsru/documents/iccs-web.xls](http://www.abdn.ac.uk/hsru/documents/iccs-web.xls) (last accessed 27 September 2013).
4. We entered the effect estimate and the new updated SE into Review Manager 5 using the generic inverse variance (RevMan 2012).

Where the number of participants randomised was not clearly stated in the included study report, we contacted the study authors for this information.

### Dealing with missing data

We contacted study authors for data for the risk of bias assessment, numbers randomised to each group and numbers retained in each group at the primary endpoint. We described outcomes with insufficient data qualitatively. For time-to-event outcomes, we used



the time point of the host study primary outcome, taking account of censoring if necessary and if the data were available.

### Assessment of heterogeneity

We measured heterogeneity of the intervention effect using the  $\text{Chi}^2$  statistic at a significance level of 0.10 and the  $I^2$  statistic (Higgins 2003), and explored through subgroup analyses.

### Assessment of reporting biases

We would have investigated reporting bias using tests for funnel plot asymmetry if sufficient data had been available (Egger 1997; Sterne 2008).

### Data synthesis

If there was no substantial heterogeneity, we pooled RRs using the fixed-effect model. If heterogeneity was detected and could not be explained by subgroup or sensitivity analyses, we used the random-effects model or did not pool results.

For factorial trials (Sharp 2006a-h, Kenton 2007a-d), all main effects were included as separate trial comparisons if they addressed different categories of strategies. Where the main effects addressed two or more strategies within the same category (e.g. Bowen 2000abc), we combined the relevant intervention groups and compared them with the control group. We also compared each intervention group with the control group, as separate trial comparisons, in exploratory analyses. For one 2 x 2 x 2 factorial trial (Renfroe 2002a-d), the numbers randomised for each group were not available at the time of analysis so comparison groups were collapsed as far as possible and then treated as separate trial comparisons in the appropriate analyses. For two three-armed trials that compared two similar intervention groups with one control group, we combined the intervention groups and compared it with the control group for the main analyses (Bauer 2004ab, Khadjesari 2011 abc). We also compared each intervention group as separate trial comparisons in exploratory analyses.

These approaches allowed full exploration of the data and also avoided double counting and over-precise pooled estimates of effect in our main analyses. However, this also meant that there were occasionally a greater number of trial comparisons than trials.

Computations for the absolute benefits of effective strategies on questionnaire response and trial retention were based on absolute risk reductions derived from meta-analysis RRs (Cochrane Handbook for Systematic Reviews of Interventions, Section 12.5.4.2: Schünemann 2008).

### Subgroup analysis and investigation of heterogeneity

To explore the effect of different strategies on trial retention, we planned the following subgroup analyses by the type of strategy used in included retention trials.

- Whether the strategy was compared with usual follow-up or other strategies.
- Whether in healthcare or non-healthcare settings.
- Whether assessment of retention was immediate or longer term (e.g. if a response to a questionnaire was expected immediately or at later time points).
- Whether the strategy was participant or management focused.

However, we identified such a diversity of retention trials and interventions that these analyses were inappropriate or not possible. Therefore, different types of strategies were analysed separately and new subgroups were defined within these before we conducted the analyses.

#### (a) Incentives

We subgrouped retention trials or trial comparisons evaluating the addition of an incentive strategy versus none as follows for analysis.

1. Monetary incentives given upfront, defined as money given to the trial participant prior to data collection in cheque, cash or voucher format.
2. Non-monetary incentives, defined as gifts, for example, pens or certificates.
3. Offers of monetary incentives after data collection, defined as a promise of the incentive after return of outcome data through attendance for scheduled follow-up or receipt of follow-up questionnaires.
4. Offers of non-monetary incentives defined as a promise of the non-monetary incentive after return of outcome data through attendance for scheduled follow-up or receipt of follow-up questionnaires.

We subgrouped retention trials or trial comparisons comparing different values of monetary incentives into:

1. those offering incentives;
2. those both giving and offering an incentive for any subsequent data (e.g. sending GBP5 with a questionnaire with an offer of GBP5 if the questionnaire is returned).

We analysed retention trials evaluating the addition of a monetary incentive versus either an offer of a monetary incentive or follow-up by telephone separately.

#### (b) Communication

We grouped retention trials or trial comparisons of the effect of different communication strategies into letter, post and reminder strategies for analysis as follows.

1. Enhanced versus standard cover letter.
2. Total design method versus standard postal communication strategy.
3. Priority versus regular post.
4. Additional reminders versus usual reminders to trial sites.

5. Additional reminders versus usual follow-up to trial participants.
6. Early versus late administration of questionnaire (i.e. sending questionnaires two to three weeks after a follow-up visit versus one to four months after a follow-up visit).
7. Recorded delivery versus telephone reminder.

### (c) Questionnaire structure

We subgrouped trials of questionnaire strategies into length of questionnaire, clarity of meaning, order of questions and layout as follows.

1. Short versus long questionnaire.
2. Long and clear questionnaire versus short and condensed questionnaire.
3. Medical condition questions first versus generic questions first.
4. Relevance of questionnaires: alcohol versus mental health questionnaires.

There were no subgroups for behavioural, case management and methodology retention trials.

Our analyses focused on the primary endpoint of retention. We initially pooled retention trials within subgroups using the fixed-effect model and quantified heterogeneity. We assessed whether these subgroups had a differential impact on retention using the test for interaction. We did not pool trials if results were inconsistent or heterogeneity was excessive.

### Sensitivity analysis

To assess the robustness of the results we planned sensitivity analyses that excluded quasi-randomised retention trials.

## RESULTS


### Description of studies

The studies are described in the [Characteristics of included studies](#), [Characteristics of studies awaiting classification](#), and [Characteristics of excluded studies](#) tables.

### Results of the search

We identified 24,304 abstracts, titles and other records from database searches to May 2012, handsearches of reviews, lists of references in included papers, SCT conference abstracts (to 2012), personal contact with trialists, and the survey of UK Clinical Trials Units ([Figure 1](#)). We screened 735 full-text papers, reports and manuscripts for eligible studies. Of 68 potentially eligible studies, we found 30 to be subsequently ineligible. This left 38 retention trials for inclusion in the review. The retention trials were embedded in real trials (host trials). We identified 11 retention trials from CENTRAL, MEDLINE and CINAHL; 14 from handsearching reviews, conference abstracts, and references lists of eligible papers; and 13 through personal communications or correspondence with clinical trials units. We evaluated six broad types of strategy to improve retention in randomised trials. Most strategies were targeted at increasing questionnaire response. The strategies used for this were incentives, communication, methodology and questionnaire design strategies. There was minimal evidence for the use of behavioural and case management strategies to improve retention.

**Figure 1. Attrition study flow diagram.**



## Included studies

Of the 38 eligible retention trials, 28 were published in full, one as an abstract (Kenton 2007a-d), and one as part of a PhD thesis (Nakash 2007). Four retention trial publications contained two trials each (Khadjesari 2011; McCambridge 2011; McColl 2003; Severi 2011). Eight retention trials are unpublished as of June 2013 (Bailey 1; Bailey 2; Edwards 2001; Land 2007; Letley 2000; MacLennan; Marson 2007; Svoboda 2001).

## Host trials

Twenty-two host trials included a single retention trial (AVID investigators 1997; Boyd 2002; Chaffin 2009; Cooke 2009; Cox 2008; Gail 1992; Dennis 2009; Hughes 1984; International Stroke Trial Group 1997; Kenyon 2001; Lamb 2007; Leigh Brown 2001; Marson 2007 (2); Omenn 2006; Porterhouse 2005; Rothert 2006; Tai 1999; Tilbrook 2011; TOMBOLA 2009a; TOMBOLA 2009b; UK BEAM 2004). Two host trials from this group were unpublished (for the retention trials by Ashby 2011 and Land 2007).

The other host trials included multiple retention trials (one unpublished for the retention trials by Bailey 1 and Bailey 2). Two retention trials (Ford 2006; Subar 2001) were embedded in the US-based Prostate, Lung, Colorectal, Ovarian (PLCO) screening trial of Prorok 2000; two (Avenell 2004; MacLennan) in the RECORD fracture prevention trial (RECORD 2007); two (Edwards 2001; Svoboda 2001) in the CRASH trial (CRASH Trial collaborators 2004); four (Khadjesari 2011 1abc; Khadjesari 2011 2; McCambridge 2011 1; McCambridge 2011 2) in the Down your Drink Trial (Murray 2007); two (Bailey 1; Bailey 2) in a feasibility study for the Sex unzipped website (unpublished); two (Severi 2011 1; Severi 2011 2) in the Text to Stop smoking cessation trial (Free 2011); and two (McColl 2003 1; McColl 2003 2) in the COGENT trial (Eccles 2002).

## Participants and settings

Included retention trials were conducted in a broad spectrum of clinical conditions and geographical settings (see Appendix 9). Eight included retention trials were embedded in trials for the treatment of alcohol and smoking dependency (Bauer 2004ab; Hughes 1989; Khadjesari 2011 1abc; Khadjesari 2011 2; McCambridge 2011 1; McCambridge 2011 2; Severi 2011 1; Severi 2011 2), and four in trials investigating treatments for injuries (Edwards 2001; Gates 2009; Nakash 2007; Svoboda 2001). Six retention trials were set in treatment trials for cancer, cardiovascular disease, epilepsy and back pain (Dorman 1997; Land 2007; Letley 2000; Man 2011; Marson 2007; Renfroe 2002a-d), and

four were embedded in screening trials for cancer, postnatal depression, and elderly diseases (Ford 2006; Kenton 2007a-d; Sharp 2006a-h; Subar 2001). Seven retention trials were embedded in prevention trials, which included two cancer prevention trials for lung and breast cancer (Bowen 2000abc; Sutherland 1996), one migraine prevention trial (Ashby 2011), and three fracture prevention trials (Avenell 2004; Cockayne 2005; MacLennan). Four retention trials were conducted in clinical management trials for orthopaedics, asthma, diabetes and angina (Leigh Brown 1997; McColl 2003 1; McColl 2003 2; Tai 1997). Six retention trials were conducted in other areas: exercise (Cox 2008), parenting (Chaffin 2009), weight management (Couper 2007), neonatal medicine (Kenyon 2005), and sexual health promotion (Bailey 1; Bailey 2).

Twenty-five retention trials were UK based, nine were USA based and two were set in Canada. The remainder were set in Czech Republic and Australia (see Characteristics of included studies table).

Retention trials were embedded in host trials that recruited participants from different settings. Five trials recruited participants directly from the community. Sixteen trials were conducted through secondary care facilities. One trial recruited participants through a combination of state workers compensation programmes, occupational and physician clinic, a surveillance programme and union records. Six UK trials recruited solely through general practitioner (GP) practices and two used a combination of recruitment through GP practices and the media. Seven trials recruited participants via the Internet, six of these were UK based and the other was US based. For one US-based smoking cessation trial, it was unclear how participants were recruited (see Characteristics of included studies table).

## Design of included retention trials

One trial was hosted in a clustered randomised trial and used this design to evaluate a strategy to improve retention (Land 2007). Four retention trials used different factorial designs (Bowen 2000abc; Kenton 2007a-d; Renfroe 2002a-d; Sharp 2006a-h). There was also one three-armed trial (Bauer 2004ab), and three four-armed trials (Khadjesari 2011 1abc; McCambridge 2011 1; McCambridge 2011 2).

Five trials used quasi-randomisation to allocate participants (Bowen 2000abc; Ford 2006; Gates 2009; McColl 2003 1; McColl 2003 2). Two used participant identification numbers (Ford 2006; Gates 2009), and two allocated the first half of a simple random sample of participants to receive one version of a questionnaire, while the remaining half was allocated to a second version (McColl 2003 1; McColl 2003 2). One retention trial used day of clinic visit to allocate participants (Bowen 2000abc).

All trials targeted individual trial participants, except one that targeted trial sites (Land 2007).

We recorded the timing of randomisation in the host trial versus the timing of randomisation in the retention trial. Four trials commenced during a randomised pilot study for the host trial (Khadjesari 2011 1abc; Letley 2000; McCambridge 2011 1; Sutherland 1996). One study started before the host trial (Chaffin 2009). Twenty-nine trials commenced during follow-up for the host trial (Ashby 2011; Avenell 2004; Bailey 1; Bailey 2; Bowen 2000abc; Cockayne 2005; Couper 2007; Cox 2008; Dorman 1997; Edwards 2001; Ford 2006; Gates 2009; Khadjesari 2011 2; Land 2007; Leigh Brown 1997; MacLennan; Man 2011; Marson 2007; McCambridge 2011 2; McColl 2003 1; McColl 2003 2; Nakash 2007; Renfroe 2002a-d; Severi 2011 1; Severi 2011 2; Sharp 2006a-h; Subar 2001; Svoboda 2001; Tai 1997). For one trial, it was unclear when the retention trial started in relation to the host trial (Kenton 2007a-d). Three retention trials started after the host trial had finished (Bauer 2004ab; Hughes 1989; Kenyon 2005): Kenyon 2005 followed-up seven-year-old children of mothers enrolled in the ORACLE trial (Kenyon 2001), Bauer 2004ab followed up participants in the COMMIT smoking cessation trial (Gail 1992), eight years after the original trial was completed and Hughes 1989 followed up participants in a smoking cessation trial six months after that study finished (Hughes 1984).

### Strategies to improve retention

Retention in trials and response to questionnaires were the outcomes measured for all included trials. The included trials evaluated six different types of strategies to improve response or retention. Incentives, communication strategies, variation in questionnaire design, methodology strategies, and combinations of communication and incentive strategies evaluated improving response to postal and electronic questionnaires. Behavioural strategies, case management and some non-monetary incentives were used to encourage participants to return to trial sites for follow-up visits. Each type of strategy is described separately below.

### Outcome measures in the included trials

Thirty-four retention trials measured response to questionnaires. Among these, the questionnaires were by post in 26 trials, electronically in four and one was done by interview. For another three retention trials, response was return of biomedical kits or biomedical kits plus a questionnaire (see [Characteristics of included studies table](#)).

Four included trials measured the number of participants remaining in the trial (Bowen 2000abc; Chaffin 2009; Cox 2008; Ford 2006)

Ten included trials specified that their primary outcome was questionnaire response at a particular time point: McCambridge 2011 1 measured response at one and three months, McCambridge

2011 2 measured response at three and 12 months, and Khadjesari 2011 1abc and Khadjesari 2011 2 measured response within 40 days of the first reminder. For Severi 2011 1, the primary outcome was completed follow-up at 30 weeks from randomisation, Severi 2011 2 used return of specimens one month after a telephone call, Avenell 2004 used retention at one year measured by questionnaire return but also reported retention at four and eight months. Cockayne 2005 and Sharp 2006a-h had final follow-up questionnaire response at any time as their primary outcome.

Two included trials reported questionnaire response at one time point only but without specifying that this was the primary outcome for the trial (Edwards 2001; Svoboda 2001). These trials measured response at three months from the questionnaire being sent. One trial reported trial retention at one time point only (three years) but without specifying that this was the primary outcome for the trial (Ford 2006). This was measured as completing the next cancer screening in a cancer screening trial. In each of these three trials, we used these data for analyses.

Two trials recorded questionnaire response at two time points without stating which was the primary outcome (Dorman 1997; Gates 2009). One trial recorded retention at two time points without stating which was the primary outcome (Cox 2008). We used data for response/retention after the first contact with respondents as the primary outcome for analyses. One trial reported response at three time points (4 weeks, 12 weeks and 9 months), which were all stated as the primary outcome (Nakash 2007). We used the data for week four in our main analysis.

Five trials reported data in survival curves. For these, we used the final analysis point (Ashby 2011; Bowen 2000abc; Chaffin 2009; Land 2007; Sutherland 1996). Authors confirmed data when it had been extracted. Fifteen trials reported the number of questionnaires returned with no time point specified (Bauer 2004ab; Couper 2007; Hughes 1989; Kenton 2007a-d; Kenyon 2005; Leigh Brown 1997; Letley 2000; MacLennan; Man 2011; Marson 2007; McColl 2003 1; McColl 2003 2; Renfroe 2002a-d; Subar 2001; Tai 1997).

### Addition of incentive versus none

There were 14 retention trials of incentives and 19 trial comparisons (Table 1). Thirteen trials were aimed at improving questionnaire response in trials and one trial was aimed at improving return for follow-up at trial site (Bowen 2000abc). The different incentive strategies aimed at improving questionnaire response were vouchers, cash, a charity donation, entry to prize draws, cheques, a certificate of appreciation and offers of study results. Incentive strategies aimed at improving retention were: certificates of appreciation and lapel pins. The value of incentives used in UK evaluations ranged from GBP5 to GBP20 and were in cash, cheque or voucher format. The value of incentives used in US-based studies was USD2 to USD10. For offers of entries into prize draws, the values were higher, ranging from GBP25 to GBP250 for UK prize

draws and USD50 for US-based prize draws. One trial evaluated giving a monetary incentive with a promise of a further incentive for return of trial data (Bailey 2).

### Communication strategies

There were 14 retention trials of communication strategies to improve response to postal questionnaires or return of biomedical test kits, or both, in randomised trials. There were 20 trial comparisons (Table 2). Strategies evaluated were: enhanced letters, additional reminders to participants, priority mailing of questionnaires, time of questionnaire administration, telephone contact and reminders to trial sites of upcoming assessments. One trial used a combination of postal communication strategies known as the total design method (TDM) (Sutherland 1996). This included sending letters in a white envelope with a hospital logo and commemorative stamp, a hand-signed letter on headed notepaper, with a reply self addressed stamped envelope, enclosing the contents. Follow-up was with a postcard sent after seven days followed by two reminder letters. This was compared with a customary method for postal follow-up. One trial evaluated the addition of an electronic SMS (short message service) text reminder on the day participants were due to receive their postal questionnaire (Man 2011).

Five trials evaluated a combination of communication strategies and incentives to improve retention from randomised trials (Couper 2007; Kenton 2007a-d; Renfroe 2002a-d; Sharp 2006a-h). The communication strategies were; first- and second-class outward post (Kenton 2007a-d; Renfroe 2002b; Sharp 2006a-h), stamped and business reply envelopes (Sharp 2006a-h), letters signed by different study personnel (Renfroe 2002c), letters posted at different times (Renfroe 2002d), text messages (Man 2011; Severi 2011 1), and a telephone survey (Couper 2007).

### Questionnaire format

The effect of a change in questionnaire format on response to randomised trial questionnaires was evaluated in eight trials with 10 comparisons (Table 3). Formats evaluated were questionnaire length: short versus long (Dorman 1997; Edwards 2001; McCambridge 2011 1b; McCambridge 2011 2b; Svoboda 2001), long and clear versus short and condensed (Subar 2001), and the order of questions (Letley 2000; McColl 2003 1; McColl 2003 2).

Two further included trials evaluated the effect of the relevance of a questionnaire on response (McCambridge 2011 1a; McCambridge 2011 2a). Relevance was defined as assessing alcohol problems rather than mental health in the context of an Internet-based intervention for hazardous drinkers (McCambridge 2011 1; McCambridge 2011 2).

### Behavioural strategies

There were two trials of behavioural strategies used for retention in randomised trials (Chaffin 2009; Cox 2008). Cox 2008 compared motivational workshops versus information sheets. Chaffin 2009 compared self motivation orientation versus standard information in the context of a parenting programme. In this case, the retention trial was run prior to the host trial with the intention of improving retention in the subsequent parenting programme evaluation trial. The analysis was based on the number eligible for inclusion in the primary analyses for the subsequent parenting programme because we do not know the allocation of those who dropped out between first and second randomisations. Complete time-to-event data were not available for Chaffin 2009, but, as only two participants were censored in the analysis, this is unlikely to have biased the results.

### Case management

One retention trial evaluated the effect of intensive case management procedures on retention of African American male participants in a cancer screening trial (Ford 2006).

### Methodology strategies

One included trial used a trial design where people knew which treatment they received. The trial compared questionnaire response in an open versus blind trial (Avenell 2004).

### Studies excluded from analyses

Two eligible trials could not be included in the analysis (Leigh Brown 1997; Letley 2000). Host trial participants in the retention trial by Leigh Brown 1997 were divided into two groups; one randomised, the other determined by preference of the referring primary care practitioner. The author confirmed that participants in the retention trial were from both randomised and non-randomised groups of the host trial and that these could not be separated.

One recently completed, unpublished trial that is not included in the review examined the effect of newsletters on retention (Mitchell). This trial will be included in the review update.

### Excluded studies

See [Characteristics of excluded studies](#) table.

We excluded trials because they were either part of a non-randomised host study, or they were not a randomised retention trial, or the primary outcome was type of data item missingness. Other excluded trials were aimed at increasing treatment compliance or baseline questionnaire response. We contacted investigators to confirm aspects of eligibility.

## Risk of bias in included studies

See [Characteristics of included studies](#).

### Allocation

All included retention trials reported that participants were randomly allocated to groups for comparison. Twenty-four included trials described adequate sequence generation by a computerised random number generator, block randomisation or use of a table of random numbers table (Avenell 2004; Bailey 1; Bailey 2; Bowen 2000abc; Chaffin 2009; Cockayne 2005; Cox 2008; Hughes 1989; Kenyon 2005; Khadjesari 2011 1abc; Khadjesari 2011 2; Land 2007; Leigh Brown 1997; Letley 2000; MacLennan; Man 2011; Marson 2007; McCambridge 2011 1; McCambridge 2011 2; Nakash 2007; Renfroe 2002a-d; Severi 2011 1; Severi 2011 2; Sutherland 1996). There was insufficient information about the sequence generation for 10 included trials, these were all described as randomised in the retention trial publications (Ashby 2011; Bauer 2004ab; Couper 2007; Dorman 1997; Edwards 2001; Kenton 2007a-d; Sharp 2006a-h; Subar 2001; Svoboda 2001; Tai 1997). Five included trials used quasi-randomisation to allocate participants (Bowen 2000abc; Ford 2006; Gates 2009; McColl 2003 1; McColl 2003 2).

Several methods were used to avoid foreseen allocation of participants; sequence generation by a trial statistician and implemented by a trial manager; sequence generation by an independent researcher, a central randomisation service, or by a nurse using a preprogrammed computer; or allocation by sealed envelopes or sequentially numbered packs. Fifteen trials reported both adequate sequence generation and allocation concealment (Avenell 2004; Bailey 1; Bailey 2; Cockayne 2005; Cox 2008; Hughes 1989; Kenyon 2005; Khadjesari 2011 1abc; Khadjesari 2011 2; Letley 2000; MacLennan; Man 2011; McCambridge 2011 1; McCambridge 2011 2; Nakash 2007).

### Blinding

Blinding of participants was generally not possible in included trials. For example, it is not possible to blind participants to the following strategies to increase trial retention or response to questionnaires: incentive or offer of incentive, behavioural (Cox 2008), or case management strategies (Ford 2006), different types of communication strategies, or questionnaire format strategies. In a number of trials, authors mentioned that participants were aware of the intervention they were getting but were unaware that this was being evaluated (Bowen 2000abc; Chaffin 2009; Kenton 2007a-d; Kenyon 2005; Leigh Brown 1997; MacLennan; Marson 2007; McColl 2003 1; McColl 2003 2). For other trials, blinding of participants or study personnel to the outcome or intervention was not reported. For one trial, a judgement about blinding was not applicable because the study evaluated the effect of blind versus open trials on retention (Avenell 2004).

### Incomplete outcome data

The primary outcome measure for this review was retention, and this was well reported. We contacted authors for clarification of any exclusions after randomisation if this was unclear from retention trial reports.

### Selective reporting

Although retention trial protocols were not available for included trials, the included published and unpublished papers reported all expected outcomes for retention.

### Other potential sources of bias

There were few other potential sources of bias identified from reports of included retention trials. For the behavioural trial by Cox 2008, the authors identified that the “walk and swim sessions were not separated according to the behavioural intervention. Participants were asked not to discuss written materials in the practical sessions”. Therefore, potential contamination between study groups could have led to biased results.

## Effect of methods

### I. Incentive strategies

There were 14 trials of incentives giving 19 trial comparisons with 16,253 participants. There was considerable heterogeneity across incentive subgroups (P value < 0.00001) (Analysis 1.1), so we decided not to pool the results for incentives.

#### Addition of incentive

The three trials (3166 participants) that evaluated the effect of giving monetary incentives to participants showed that the addition of monetary incentives was more effective than no incentive at increasing response to postal questionnaires (RR 1.18; 95% CI 1.09 to 1.28, P value < 0.0001) (Analysis 1.1). A sensitivity analysis excluding the quasi-randomised trial by Gates 2009 still showed that the addition of a monetary incentive remained more effective than none (RR 1.31; 95% CI 1.11 to 1.55, P value = 0.002) (Analysis 2.1).

Based on two Internet-based trials (3613 participants), an offer of a monetary incentive promoted greater return of electronic questionnaires than no offer (RR 1.25; 95% CI 1.14 to 1.38, P value < 0.00001; heterogeneity P value = 0.14) (Analysis 1.1). However, a single trial comparison suggested that an offer of a monetary donation to charity did not increase response to electronic questionnaires (RR 1.02; 95% CI 0.78 to 1.32; P value = 0.90) (Analysis 1.1).

Based on six trials (6322 participants), there was no clear evidence that the addition of non-monetary incentives improved questionnaire response (RR 1.00; 95% CI 0.98 to 1.02, P value = 0.91), but there was some heterogeneity (P value = 0.02) (Analysis 1.1). A sensitivity analysis excluding the quasi-randomised trial (Bowen 2000abc) showed a similar effect (RR 1.00; 95% CI 0.93 to 1.08, P value = 0.99) (Analysis 2.1) and heterogeneity (P value = 0.01). Two trials (1138 participants) evaluating offers of non-monetary incentives suggest that an offer of a non-monetary incentive is neither more nor less effective than no offer (RR 0.99; 95% CI 0.95 to 1.03, P value = 0.60) at improving questionnaire response (Analysis 1.1).

In exploratory analyses, the different incentive arms that were combined for the main analysis did not appear to show differential effects (Analysis 3.1).

#### **Addition of monetary incentive to both study arms**

Two trials (902 participants) show that higher value incentives are better at increasing response to postal questionnaires than lower value incentives (RR 1.12; 95% CI 1.04 to 1.22, P value = 0.005) irrespective of how they are given (Analysis 5.1).

#### **Addition of monetary incentive versus offer of a monetary incentive**

Two trials (297 participants) provided no evidence that giving a monetary incentive is better than an offer of entry into a prize draw for improving response to postal questionnaires (RR 1.04; 95% CI 0.91 to 1.19, P value = 0.56) (Analysis 6.1).

#### **Addition of an offer of entry into a prize draw versus none**

We excluded one trial from the analysis (Leigh Brown 1997). The results showed higher responses in the group offered entry into a prize draw compared with the group not offered entry into the draw (70.5% versus 65.8%).

## **2. Communication strategies**

There were 14 trials of communication strategies and 20 comparisons with 9822 participants.

#### **Addition of telephone survey versus monetary incentive plus questionnaire**

One trial (700 participants) showed no clear evidence that a telephone survey was either more or less effective than a monetary incentive and a questionnaire for improving response (RR 1.08; 95% CI 0.94 to 1.24, P value = 0.27) (Analysis 4.1).

#### **Enhanced versus standard letters**

Results from two trials (2479 participants) showed that an enhanced letter was neither more nor less effective than a standard letter for increasing response to trial postal questionnaires (RR 1.01; 95% CI 0.97 to 1.05, P value = 0.70) (Analysis 7.1).

#### **Total design method versus customary method**

Although based on a single trial (226 participants) the TDM package was more effective than a customary postal communication method at increasing questionnaire return (RR 1.43; 95% CI 1.22 to 1.67, P value < 0.0001) (Analysis 8.1).

#### **Priority versus regular post**

Based on the relevant arms of seven trials (1888 participants), there was no clear evidence that priority post was either more or less effective than regular post at increasing trial questionnaire return (RR 1.02; 95% CI 0.95 to 1.09, P value = 0.55) (Analysis 9.1).

#### **Additional reminder versus usual follow-up practices**

Six trials (3401 participants) evaluated the effect of different additional types of reminders to participants on questionnaire response. There was no evidence that a reminder was either more or less effective than no reminder at improving trial questionnaire response (RR 1.03; 95% CI 0.99 to 1.06, P value = 0.13) (Analysis 10.1).

#### **Additional reminder to trial site versus usual reminder**

Based on one cluster randomised trial (272 participants), a monthly reminder to trial sites of upcoming assessment was neither more nor less effective than the usual follow-up (RR 0.96; 95% CI 0.83 to 1.11, P value = 0.57) (Analysis 11.1).

#### **Early versus late questionnaire administration**

Based on one trial (664 participants), there was no clear evidence that sending questionnaires early either increased or decreased response (RR 1.10; 95% CI 0.96 to 1.26, P value = 0.19) (Analysis 12.1).

#### **Recorded delivery versus telephone reminder**

One small trial (192 participants) found that recorded delivery was more effective than a telephone reminder (RR 2.08; 95% CI 1.11 to 3.87; P value = 0.02) (Analysis 13.1).



### 3. New questionnaire strategies

#### New versus standard questionnaire

Eight trials with 10 comparisons (21,505 participants) evaluated the effect of a new questionnaire format on questionnaire response. Although there was some heterogeneity between the questionnaire subgroups (P value = 0.11) (Analysis 14.1), it did not seem reasonable to pool the results based on such different interventions. Five trials (7277 participants) compared the effect of short questionnaires versus long on postal questionnaire response. There was only a suggestion that short questionnaires may be better (RR 1.04; 95% CI 1.00 to 1.08, P value = 0.07) (Analysis 14.1).

Based on one trial (900 participants; Subar 2001), there is no evidence that long and clear questionnaires were either more or less effective than shorter condensed questionnaires for increasing trial questionnaire response (RR 1.01; 95% CI 0.95 to 1.07, P value = 0.86) (Analysis 14.1).

Two trials (9435 participants; McColl 2003 1; McColl 2003 2) found no evidence that placing disease/condition questions before generic questions is either a more or less effective strategy than a generic questions before disease/condition questions strategy at increasing trial questionnaire response (RR 1.00; 95% CI 0.97 to 1.02, P value = 0.75) (Analysis 14.1). It should be noted that these were quasi-randomised trials (Analysis 15.1).

One trial in this category was not included in the analysis by Letley 2000, outcome data were not available for each study arm when this review was submitted and the overall response rate for this trial was 87%.

In the context of research on reducing alcohol consumption, there was also evidence that more relevant questionnaires (i.e. those relating to alcohol use) increased response rates (RR 1.07; 95% CI 1.01 to 1.14, P value = 0.03).

### 4. Behavioural/motivational strategies

Two community-based trials (273 participants; Chaffin 2009; Cox 2008) showed no evidence that the behavioural/motivational strategies used are either more or less effective than standard information for retaining trial participants (RR 1.08; 95% CI 0.93 to 1.24, P value = 0.31) (Analysis 16.1).

### 5. Case management

One trial (703 participants; Ford 2006) evaluated the effect of intensive case management procedures on retention. There is no evidence that intensive case management was either more or less effective than usual follow-up in the population examined (RR 1.00; 95% CI 0.97 to 1.04, P value = 0.99) (Analysis 17.1).

### 6. Methodology strategies

One fracture prevention trial (538 participants; Avenell 2004) evaluated the effect of participants knowing their treatment allocation (open trial) compared with participants blind/unaware of their allocation on questionnaire response. Using a trial design where people know which treatment they will receive led to higher questionnaire response rates (RR 1.37; 95% CI 1.16 to 1.63, P value = 0.0003) (Analysis 18.1).

#### Reporting bias

Although we planned to investigate potential reporting bias, there were too few studies in most strategies to allow formal testing. However, we were able to obtain considerable data from unpublished trials and those published with limited information, reducing the risk of such biases.

#### Absolute benefits of strategies to improve retention

The absolute benefits of effective strategies on questionnaire response are illustrated in Table 4. The baseline response rates were broadly typical of the response rates seen in trials. The number of questionnaires returned were based on the assumed control arm risk.

Based on a 40% baseline response rate for postal questionnaires, the addition of a monetary incentive was estimated to increase response by 92 questionnaires per 1000 sent (95% CI 50 to 131). With the addition of an offer of a monetary incentive in an Internet-based trial, based on a baseline response rate of 30%, trialists could expect an increase of 140 questionnaires per 1000 (95% CI 86 to 193).

For trials hoping to increase the return of postal questionnaires with chlamydia test kits, the number of kits returned was estimated to increase by 33 per 1000 sent when GBP20 was offered as an incentive, rather than GBP10 (95% CI 11 to 54).

## DISCUSSION

### Summary of main results

Thirty-eight randomised retention trials were included in this review, evaluating six broad types of strategies to increase questionnaire response and retention in randomised trials. In 34 trials, strategies for increasing response to questionnaires were: incentives, communication strategies, new questionnaire format and methodological interventions. Four trials evaluated strategies to improve retention, these were: participant case management, behavioural and non-monetary incentive strategies. Trials were conducted across a spectrum of disease areas, countries, healthcare and community settings.

Strategies with the clearest impact on questionnaire response were: addition of monetary incentives compared with no incentive for return of postal questionnaires, addition of an offer of a monetary incentive when compared with none for return of electronic questionnaires, and an offer of GBP20 vouchers when compared with GBP10 for return of postal questionnaires and biomedical test kits. The evidence was less clear about whether shorter questionnaires rather than longer questionnaires increased response. The evidence was also less clear whether in the context of research on reducing alcohol consumption more relevant questionnaires increased response.

The addition of a non-monetary incentive, an offer of a non-monetary incentive compared with no incentive or, 'enhanced' letters, letters delivered by priority post, or additional reminders compared with standard communication strategies did not increase or decrease trial questionnaire response. Questionnaire structure also did not seem to increase response.

Although each was based on the results of a single trial, recorded delivery (proof of posting and an electronic copy of the signature available online) of questionnaires seemed more effective than telephone reminders, and a 'package' of postal communication strategies with reminder letters appeared better than standard procedures. A trial design where participants knew which treatment they were to receive also appeared more effective than a trial design where they were unaware of the treatment they were about to receive for return of questionnaires in a fracture prevention trial. Further evaluation of these strategies may be needed. Posting questionnaires early, questionnaire order, offers of charity donations or sending reminders to trial sites did not improve response.

Many trial outcome measures were collected using questionnaires, therefore, if response rates can be increased, retention will also be improved. No strategy had a clear impact on increasing the number of participants returning to trial sites for follow-up visits.

## Overall completeness and applicability of evidence

The addition of a GBP5 voucher to usual follow-up procedures was effective for return of postal questionnaires in trials conducted between 2005 and 2009. The more recent unpublished studies by [Bailey 1](#) [Bailey 2](#) found GBP20 vouchers were more effective than GBP10 vouchers for return of postal questionnaires. Splitting the monetary incentive into money given before and after receipt of data could be more effective as a strategy to increase questionnaire follow-up with different population groups and in different trial settings where questionnaire response is low (e.g. with hard to reach groups that may include young male healthy adults, teenagers or residents in areas of high economic deprivation). This could be a cost-effective strategy because if questionnaires are not returned then money is saved. The value of the monetary incentive should not be so high as to be perceived as payment for data but more as an appreciation for efforts made by participants. Offering monetary

incentives may increase the number of questionnaires returned per 1000 participants by at least as much as giving monetary incentives and giving higher valued monetary incentives, but has only been tested in online questionnaires. Offers of monetary incentives were also an effective strategy in the context of an online electronic questionnaire. These could be less costly to increase retention than the addition of a monetary incentive as only those who return the data are reimbursed. This would need further evaluation as the results were based on two Internet-based trials. It would be beneficial for trialists to know which is more effective: an offer of a monetary incentive or an upfront monetary incentive. We did not find any trials that made this direct comparison.

Shorter postal questionnaires have wide applicability to trials and could be considered as a useful strategy to increase trial questionnaire response in online Internet-based trials but there is only a suggestion that these are effective.

Several strategies showed no clear effect. The addition of non-monetary incentives in the form of pens, lapel pins and certificates of appreciation, or offers of non-monetary incentives through offering study results did not increase response or retention. A possible explanation might be how these items are valued by participants, or how they perceive their time is valued. Nevertheless, this result has the potential to reduce trial costs because associated saving could be channelled towards monetary incentives that have been shown to be effective.

The evidence showed that priority post (first-class post or equivalent) did not increase response. It is expensive as a means of communicating with participants and savings can be made by using regular (2nd-class) post instead.

Additional reminders sent to non-responders or as questionnaires were posted; enhanced letters, that is, letters signed by the principal investigator, or letters further explaining the anticipated length of time to complete a questionnaire, were not effective strategies to increase response. Enhanced letters and different types of additional reminders are used by trialists in current research practice. Too many reminders could be counterproductive to improving retention in randomised trials and details of the time expected to undertake specific tasks might be informative but off putting for participants. Nevertheless, letters and reminders are part of the research process and play a role in participant engagement especially if there is little face-to-face contact or in trials with long intervals between data collection time points.

Several strategies to increase questionnaire response need further evaluation to determine their effect but there is only a suggestion that these were effective. If participants are well and engaged with a trial, questionnaire length may not impact on response rates because participants may be happy to feedback on their condition in this way. For other conditions, for example, cancers and terminal illnesses, trial participants might prefer shorter questionnaires if their symptoms are problematic. Telephone follow-up compared with monetary incentive sent with a questionnaire needs further evaluation possibly with a cost-benefit analysis, as both could be

expensive in time and human resources. Although appearing very effective, the total design method for postal questionnaires could be labour intensive to implement, expensive and may no longer be applicable to some participant groups (e.g. young people), or in trials using email, text or the Internet to collect data. Recorded delivery could be useful to ensure trial follow-up supplies reach their intended destination (e.g. biomedical specimen kits and questionnaires). Careful planning of day, date and time of delivery with each participant to avoid inconvenience might be necessary but again this strategy has the potential to be burdensome for trial co-ordinating centres and trial sites to administer. While trialists are assured that follow-up supplies are delivered with this strategy, participants might have the added burden of an extra visit to collect supplies from a sorting post office and this could be costly. The use of open trials to increase questionnaire response can only be applied to trials where blinding is not required and could be counterproductive if a participant or clinician has a treatment preference. Bias associated with loss to follow-up resulting from these preferences could be avoided in blind trials.

Evaluations of strategies that encourage participants to return to trial sites for follow-up visits and monitoring were fewer than strategies to increase response to postal and electronic questionnaires, without further evidence case management and behavioural strategies cannot be recommended for use to encourage participant return.

This review identified no trials from low-income countries. All included studies were conducted in higher-income countries. Therefore, the strategies to increase retention identified by this review may not be generalisable to trials conducted in low-income countries because the interventions identified might not be socially, culturally or economically appropriate for trials run in these regions. The results may also not be applicable to all social groups as we were unable to examine response/retention by social characteristics such as economic disadvantage and social class. Most of the evidence in this review relates to increasing questionnaire follow-up in randomised trials for either the primary or secondary outcome for the host trial. The diversity between strategies and insufficient numbers in each of these categories meant that we could not do subgroup analyses by trial setting and disease area as planned.

## Quality of the evidence

The extent of unpublished trials evaluating retention strategies is unknown; however, this review includes several unpublished trials and we made an effort to capture UK-based unpublished trials through our survey and research contacts. For some comparisons, results were based on one or two trials in a particular context. The inclusion of any further published and unpublished trials in future updates would improve the precision of the results of this review. The six types of strategies that we identified targeted retention of trial participants in randomised trials. We believe response and retention were the relevant dichotomous outcomes to be reported

for this review. Many other strategies used by trialists in practice to reduce attrition/increase response or retention in trials were not identified by this review (e.g. social support strategies; child care, [Loue 2008](#); family support, [De Sousa 2008](#); reduction in the number of visits, [Schulz 2002](#)). Evaluations of trial management strategies are also under-represented in the review (e.g. evaluations of site-specific reports, [El Khorazaty 2007](#); levels of contact by the co-ordinating centre, [Senturia 1998](#); training project staff).

Both published and unpublished included retention trials were fairly well conducted but could be improved. Five of the 39 trials included in the review were quasi-randomised. The motivation for conducting many of the included retention trials was reactive rather than planned upfront (i.e. when loss to follow-up became a problem during trial follow-up, rather than planned prior to host trial commencement).

Most trials used appropriate methods for randomisation or at least stated that they were randomised. For trials that did not describe their methods well or provide further information, there remains a potential risk of selection bias. Sensitivity analyses excluding quasi-randomised trials did not affect the results. In this context, where motivating participants to provide data or attend clinics is often the target of the interventions and so appropriately influences the outcome, lack of blinding is less of a concern. Retention is the outcome and was obtained for all but two trials, so similarly, attrition and selective outcome reporting bias are unimportant. Although the retention trials were fairly well conducted, they could be improved and they were often poorly reported. This may be because they were designed when loss to follow-up became a problem in a trial, rather than preplanned prior to host trial commencement.

## Potential biases in the review process

Many words are used to describe loss to follow-up, for example, attrition, withdrawal and questionnaire non-response. We included these in our search strategy. We attempted to obtain unpublished trials and data by contacting authors and writing to UK clinical trials units and presenting at national and international conferences. We are confident that we have captured most studies and the spectrum of strategies that have been evaluated to date. It is conceivable, however, that less well-reported, ongoing, unpublished trials or trials conducted outside of the UK might have been missed. Most trials used appropriate methods for sequence generation or at least stated that they were randomised and concealed allocation. There is small risk that those that did not describe their methods well or provide further information did not use adequate methods for allocation and concealment and may have biased the results. However, sensitivity analyses excluding quasi-randomised trials did not affect the results. Blinding is hard to achieve in this context, where motivating participants to provide data or attend clinics is often the target of the interventions and so appropriately influences the outcome.

## Agreements and disagreements with other studies or reviews

The strategies that improve retention are, in some cases, the same as or similar to those found to be effective for cohort and cross-sectional study designs. However, prior to our review, it was not clear which of these strategies could be extrapolated to randomised trials. Successful retention strategies used in other study designs may be effective in trials settings and should be tested. Edwards' review on methods to increase response to postal and electronic questionnaires included 513 trials and identified many strategies to increase response to questionnaires (Edwards 2009). Included trials were embedded in surveys, cohort studies and trials, which may explain some of the heterogeneity in effects seen in Edwards' review and reliance on the random-effects model. Unexplained heterogeneity was not a particular problem in this review. Edwards found monetary incentives effective for increasing response to postal questionnaires (Edwards 2009). However, unlike our review, Edwards found that non-monetary incentives were effective for postal and electronic questionnaires. Other strategies found to be effective by Edwards, in agreement with our review, included recorded delivery of questionnaires and shorter questionnaires, although in our review shorter questionnaires need further evaluation. Edwards also found that use of hand-written addresses, stamped return envelopes as opposed to franked return envelopes and first-class outward mailing improved response. Our review found that a 'package' including an enhanced letter incorporating several reminders was effective, but the effectiveness of first-class/priority mail to increase response in randomised trials was unclear. Booker's narrative review of methods to increase retention in population-based cohort studies was based on only 11 randomised trials and no meta-analysis (Booker 2011). The results suggested that incentives were associated with an increase in retention. Nakash's systematic review of ways to increase response to postal questionnaires in health care focused on randomised trials of ways to increase response to postal questionnaires in healthcare research on participant populations (Nakash 2006 (2)). Fifteen trials were included in this meta-analysis, which found that reminder letters, telephone contact and short questionnaires increased response to postal questionnaires in the context of healthcare research. There was no evidence that incentives were effective. Again, this review was not exclusive to evaluations conducted in randomised trials. The Edwards review was broad and focused specifically on methods to enhance response to questionnaires and included studies in non-healthcare settings (Edwards 2009). The reviews by Nakash and Booker focused on retention in specific research areas, health care and cohort studies (Booker 2011; Nakash 2006 (2)). Unlike these reviews, our review focused specifically on a range of strategies evaluated within trials. Therefore, it specifically addressed the question of retention of study participants within randomised trials, which was beyond the scope of the other reviews. Application of these results would depend on trial setting, population, disease area, data collection and follow-up procedures. Moreover, we

identified additional strategies that may improve trial retention, for example, methodological strategies.

This review is the most comprehensive to date on strategies specifically designed to improve retention in randomised trials. We included seven unpublished trials and 18 other trials not included by Edwards (Edwards 2009).

## AUTHORS' CONCLUSIONS

### Implication for methodological research

Trialists may consider including well thought out and adequately powered evaluations of strategies to increase retention in randomised trials. This could include a clear definition of retention strategies and of measures of retention. Trialists conducting future methodology trials can consider incorporating evaluations of strategies to increase retention at the design stage so that power, sample size and funding arrangements are taken into account. Retention trials were often poorly reported without consort diagrams, clear primary outcomes, sample size, sociodemographic composition or power calculations. Considerable time was spent contacting authors for unreported data needed for a robust meta-analysis. Trialists in their reports might consider adhering to the consort guidelines for trial reporting, which would facilitate the synthesis of results in future methodology reviews. There is less research on ways to increase return of participants to trial sites for follow-up and on the effectiveness of strategies to retain trial sites in cluster and individual randomised trials. Research in both areas would be very beneficial to trialists. There is no current system for identifying methodological trials in progress, until a system is set up it may be useful for systematic review authors to incorporate contacting trials units into their search strategy.

## ACKNOWLEDGEMENTS

We would like to thank the following: all authors of included published trials for providing extra unreported data. Principal investigators for data on studies in progress or completed and unpublished: Julia Bailey UCL for data for Bailey 1 and Bailey 2; Graeme MacLennan for data for MacLennan; Stephanie Land for data for Land 2007; the co-ordinators of UK Clinical Trials Units who responded to our survey with information about ongoing or unpublished completed trials; Cara Booker, SPHRU for search strategy information; Angela Young, librarian UCL for assistance with searching databases; and Ian White MRC Biostatistics Unit Cambridge, for helpful comments on the analysis of multiarm trials. Finally, we would like to thank the following: Shaun Treweek, Mike Clarke, Andy Oxman, Karen Robinson and Anne Eisinga for comments on the review protocol and Phil Edwards, Elie Akl,

Lisa Maguire, Jean Suvan, Shaun Treweek, Mike Clarke and Karen Robinson for comments on the full review. This project was funded by the MRC Population Health Sciences Research Network.

## REFERENCES

### References to studies included in this review

#### Ashby 2011 *{published data only}*

\* Ashby R, Turner G, Cross B, Mitchell N, Torgerson D. A randomized trial of electronic reminders showed a reduction in the time to respond to postal questionnaires. *Journal of Clinical Epidemiology* 2011;**64**:208–12.

#### Avenell 2004 *{published data only}*

\* Avenell A, Grant AM, McGee M, McPherson G, Campbell MK, McGee MA, et al. The effects of an open design on trial participant recruitment, compliance and retention - a randomized controlled trial comparison with a blinded, placebo-controlled design. *Clinical Trials* 2004;**1**(6):490–8.

#### Bailey 1 *{unpublished data only}*

Bailey J. Evaluating the effectiveness of incentives in the “Sexunzipped” trial.

#### Bailey 2 *{unpublished data only}*

Bailey J. Evaluating the effectiveness of incentives in the “Sexunzipped” trial.

#### Bauer 2004a *{published data only (unpublished sought but not used)}*

Bauer JE, Rezaishiraz H, Head K, Cowell J, Bepler G, Aiken M, et al. Obtaining DNA from a geographically dispersed cohort of current and former smokers: use of mail-based mouthwash collection and monetary incentives. *Nicotine and Tobacco Research* 2004;**6**(3):439–46.

#### Bauer 2004ab *{published data only (unpublished sought but not used)}*

Bauer JE, Rezaishiraz H, Head K, Cowell J, Bepler G, Aiken M, et al. Obtaining DNA from a geographically dispersed cohort of current and former smokers: use of mail-based mouthwash collection and monetary incentives. *Nicotine and Tobacco Research* 2004;**6**(3):439–46.

#### Bauer 2004b *{published data only (unpublished sought but not used)}*

Bauer JE, Rezaishiraz H, Head K, Cowell J, Bepler G, Aiken M, et al. Obtaining DNA from a geographically dispersed cohort of current and former smokers: use of mail-based mouthwash collection and monetary incentives. *Nicotine and Tobacco Research* 2004;**6**(3):439–46.

#### Bowen 2000a *{published and unpublished data}*

Bowen D, Thornquist M, Anderson K, Barnett M, Valanis B, Goodman G, et al. The use of incentives to increase participation in a chemoprevention trial. *Controlled Clinical Trials* 1995;**16**:41s.

\* Bowen D, Thornquist M, Goodman G, Omenn GS, Anderson K, Barnett M, et al. Effects of incentive items on participation in a randomized chemoprevention trial. *Journal of Health and Psychology* 2000;**5**(1):109–15.

#### Bowen 2000abc *{published and unpublished data}*

\* Bowen D, Thornquist M, Goodman G, Omenn GS, Anderson K, Barnett M, et al. Effects of incentive items on participation in a randomized chemoprevention trial. *Journal of Health and Psychology* 2000;**5**(1):109–15.

#### Bowen 2000b *{published and unpublished data}*

Bowen D, Thornquist M, Goodman G, Omenn GS, Anderson K, Barnett M, et al. Effects of incentive items on participation in a randomized chemoprevention trial. *Journal of Health and Psychology* 2000;**5**(1):109–15.

#### Bowen 2000c *{published and unpublished data}*

Bowen D, Thornquist M, Goodman G, Omenn GS, Anderson K, Barnett M, et al. Effects of Incentive Items on Participation in a Randomized Chemoprevention Trial. *Journal of Health and Psychology* 2000;**5**(1):109–15.

#### Chaffin 2009 *{published and unpublished data}*

Chaffin M, Valle LA, Funderburk B, Gurwitsch R, Silovsky J, Bard D. A motivational intervention can improve retention in PCIT for low-motivation child welfare clients. *Child Maltreatment* 2009;**14**(4):356–68.

#### Cockayne 2005 *{published and unpublished data}*

Cockayne S, Torgerson D. A randomised controlled trial to assess the effectiveness of offering study results as an incentive to increase response rates to postal questionnaires. *BMC Medical Research Methodology* 2005;**5**(1):34.

#### Couper 2007 *{published data only (unpublished sought but not used)}*

Couper PM, Peytchev A, Strecher JV, Rothert K, Anderson J. Following up non respondents to an online weight management intervention: randomized trial comparing mail versus telephone. *Journal of Medical Internet Research* 2007;**9**(2):e16.

#### Cox 2008 *{published and unpublished data}*

Cox KL, Burke V, Beilin LJ, Derbyshire AJ, Grove JR, Blanksby BA, et al. Short and long-term adherence to swimming and walking programs in older women - the Sedentary Women Exercise Adherence Trial (SWEAT 2). *Preventive Medicine* 2008;**46**(6):511–17.

Cox KL, Burke V, Beilin LJ, Grove JR, Blanksby BA, Puddey IB. Blood pressure rise with swimming versus walking in older women: the Sedentary Women Exercise Adherence Trial 2 (SWEAT 2). *Journal of Hypertension* 2006;**24**(2):307–14.

#### Dorman 1997 *{published and unpublished data}*

Dorman P, Slattery J, Farrell B, Dennis MS, Sandercock PA. A randomised comparison of the EuroQol and Short Form-36 after stroke. United Kingdom collaborators in the International Stroke Trial. *BMJ* 1997;**315**(7106):461.

- Edwards 2001** *{published and unpublished data}*  
Edwards P. A comparison of two questionnaires for assessing outcome after head injury. Personal email communication from Edwards P in April 2010.
- Ford 2006** *{published and unpublished data}*  
Ford ME, Havstad S, Vernon SW, Davis SD, Kroll D, Lamerato L, et al. Enhancing adherence among older African American men enrolled in a longitudinal cancer screening trial. *Gerontologist* 2006;**46**(4):545–50.  
Ford ME, Randolph V, Hopkins-Johnson L, Eason SL, Havstad S, Jankowski M, et al. Design of a case management approach to enhance cancer screening trial retention among older African American men. *Journal of Aging and Health* 2004;**16**(5):39s–57s.
- Gates 2009** *{published and unpublished data}*  
Gates S, Williams M, Withers E, Williamson E, Mt-Isa S, Lamb S. Does a monetary incentive improve the response to a postal questionnaire in a randomised controlled trial? The MINT incentive study. *Trials* 2009;**10**(1):44.
- Hughes 1989** *{published and unpublished data}*  
Hughes JFR. Free reprints to increase the return of follow-up questionnaires. Abstracts from the Tenth Meeting of the Society of Clinical Trials, 1989 May 15-18; Minneapolis. 1989; Vol. 10:316–54.
- Kenton 2007a** *{published and unpublished data}*  
Kenton L, Dennis CL, Weston J, Kiss A. The effect of incentives and high priority mailing on postal questionnaire response rates: a mini-RCT. Abstracts from the 28th Meeting of the Society of Clinical Trials; 2007 May 20-23; Montreal. 2007; Vol. 4, issue 4:371–455.
- Kenton 2007a-d** *{published and unpublished data}*  
\* Kenton L, Dennis CL, Watson J, Kiss A. The effect of incentives and high priority mailing on postal questionnaire response rates: a MINI RCT. *Clinical Trials* 2007;**4**(4):371–455.
- Kenton 2007b** *{published and unpublished data}*  
Kenton L, Dennis CL, Weston J, Kiss A. The effect of incentives and high priority mailing on postal questionnaire response rates: a mini-RCT. Abstracts from the 28th Meeting of the Society of Clinical Trials; 2007 May 20-23; Montreal. 2007; Vol. 4, issue 4:371–455.
- Kenton 2007c** *{published and unpublished data}*  
Kenton L, Dennis CL, Weston J, Kiss A. The effect of incentives and high priority mailing on postal questionnaire response rates: a mini-RCT. *Abstracts from the 28th Meeting of the Society of Clinical Trials; 2007 May 20-23; Montreal* 2007;**4**:371–455.
- Kenton 2007d** *{published and unpublished data}*  
Kenton L, Dennis CL, Weston J, Kiss A. The effect of incentives and high priority mailing on postal questionnaire response rates: a mini-RCT. Abstracts from the 28th Meeting of the Society of Clinical Trials; 2007 May 20-23; Montreal. 2007; Vol. 4:371–455.
- Kenyon 2005** *{published and unpublished data}*  
Kenyon S, Pike K, Jones D, Taylor D, Salt A, Marlow N, et al. The effect of a monetary incentive on return of a postal health and development questionnaire: a randomised trial. *BMC Health Services Research* 2005;**5**(1):55.
- Khadjesari 2011** *{published and unpublished data}*  
\* Khadjesari Z, Murray E, Kalaitzaki E, White I, McCambridge J, Thompson S. Impact and costs of incentives to reduce attrition in online trials: two randomised controlled trials. *Journal of Medical Internet Research* 2011;**13**(1):e26.
- Khadjesari 2011 1a** *{published and unpublished data}*  
Khadjesari Z, Murray E, Kalaitzaki E, White IR, McCambridge J, Thompson SG, et al. Impact and costs of incentives to reduce attrition in online trials: two randomized controlled trials. *Journal of Medical Internet Research* 2011;**13**(1):e26.
- Khadjesari 2011 1abc** *{published and unpublished data}*  
Khadjesari Z, Murray E, Kalaitzaki E, White IR, McCambridge J, Thompson SG, et al. Impact and costs of incentives to reduce attrition in online trials: two randomized controlled trials. *Journal of Medical Internet Research* 2011;**13**(1):e26.
- Khadjesari 2011 1ac** *{published and unpublished data}*  
Khadjesari Z, Murray E, Kalaitzaki E, White IR, McCambridge J, Thompson SG, et al. Impact and costs of incentives to reduce attrition in online trials: two randomized controlled trials. *Journal of Medical Internet Research* 2011;**13**(1):e26.
- Khadjesari 2011 1b** *{published and unpublished data}*  
Khadjesari Z, Murray E, Kalaitzaki E, White IR, McCambridge J, Thompson SG, et al. Impact and costs of incentives to reduce attrition in online trials: two randomized controlled trials. *Journal of Medical Internet Research* 2011;**13**(1):e26.
- Khadjesari 2011 1c** *{published and unpublished data}*  
Khadjesari Z, Murray E, Kalaitzaki E, White IR, McCambridge J, Thompson SG, et al. Impact and costs of incentives to reduce attrition in online trials: two randomized controlled trials. *Journal of Medical Internet Research* 2011;**13**(1):e26.
- Khadjesari 2011 2** *{published and unpublished data}*  
Khadjesari Z, Murray E, Kalaitzaki E, White IR, McCambridge J, Thompson SG, et al. Impact and costs of incentives to reduce attrition in online trials: two randomized controlled trials. *Journal of Medical Internet Research* 2011;**13**(1):e26.
- Land 2007** *{published and unpublished data}*  
Land SR, Ritter MW, Costantino JP, Julian TB, Cronin WM, Haile SR, et al. Compliance with patient-reported outcomes in multicenter clinical trials: methodologic and practical approaches. *Journal of Clinical Oncology* 2007;**25**(32):5113–20.
- Leigh Brown 1997** *{published and unpublished data}*  
Leigh Brown AP, Lawrie HER, Kennedy ADM, Webb JAG, Torgerson DJ, Grant AM. Cost effectiveness of a prize draw on response to a postal questionnaire: results of a randomised trial among orthopaedic outpatients in

- Edinburgh. *Journal of Epidemiology and Community Health* 1997;**51**(4):463–4.
- Letley 2000** *{unpublished data only}*  
Letley L, Garratt A. Questionnaire length and internal reliability: a randomised study. Society for Social Medicine 43rd Annual Scientific Meeting; 2000 6-8 Sept; Norwich. 2000.
- MacLennan** *{unpublished data only}*  
MacLennan GS. RECORD Phone Trial.
- Man 2011** *{published data only}*  
Man MS, Tilbrook HE, Jayakody S, Hewitt CE, Cox H, Cross B, et al. Electronic reminders did not improve postal questionnaire response rates or response times: a randomized controlled trial. *Journal of Clinical Epidemiology* 2011;**64**(1001):e1004.
- Marson 2007** *{published and unpublished data}*  
Marson AG. A randomised controlled trial examining the longer-term outcomes of standard versus new antiepileptic drugs: the SANAD trial. *NIHR HTA Report* 2007;**11**(37): 111–3.
- McCambridge 2011** *{published data only}*  
\* McCambridge J, Kalaitzaki E, White RI, Khadjesari Z, Murray E, Linke S. Impact of length or relevance of questionnaires on attrition in online trials: randomized controlled trial. *Journal of Medical Internet Research* 2011; **13**(4):e96.
- McCambridge 2011 1** *{published data only}*  
McCambridge J, Kalaitzaki E, White IR, Khadjesari Z, Murray E, Linke S, et al. Impact of length or relevance of questionnaires on attrition in online trials: randomized controlled trial. *Journal of Medical Internet Research* 2011; **13**(4):e96.
- McCambridge 2011 1a** *{published data only}*  
McCambridge J, Kalaitzaki E, White IR, Khadjesari Z, Murray E, Linke S, et al. Impact of length or relevance of questionnaires on attrition in online trials: randomized controlled trial. *Journal of Medical Internet Research* 2011; **13**(4):e96.
- McCambridge 2011 1b** *{published data only}*  
McCambridge J, Kalaitzaki E, White IR, Khadjesari Z, Murray E, Linke S, et al. Impact of length or relevance of questionnaires on attrition in online trials: randomized controlled trial. *Journal of Medical Internet Research* 2011; **13**(4):e96.
- McCambridge 2011 2** *{published data only}*  
McCambridge J, Kalaitzaki E, White IR, Khadjesari Z, Murray E, Linke S, et al. Impact of length or relevance of questionnaires on attrition in online trials: randomized controlled trial. *Journal of Medical Internet Research* 2011; **13**(4):e96.
- McCambridge 2011 2a** *{published data only}*  
McCambridge J, Kalaitzaki E, White IR, Khadjesari Z, Murray E, Linke S, et al. Impact of length or relevance of questionnaires on attrition in online trials: randomized controlled trial. *Journal of Medical Internet Research* 2011; **13**(4):e96.
- McCambridge 2011 2b** *{published data only}*  
McCambridge J, Kalaitzaki E, White IR, Khadjesari Z, Murray E, Linke S, et al. Impact of length or relevance of questionnaires on attrition in online trials: randomized controlled trial. *Journal of Medical Internet Research* 2011; **13**(4):e96.
- McColl 2003** *{published and unpublished data}*  
\* McColl EM, Eccles MPM, Rousseau NSB, Steen INP, Parkin DWD, Grimshaw JMP. From the generic to the condition-specific? Instrument order effects in quality of life assessment. *Medical Care* 2003;**41**(7):777–90.
- McColl 2003 1** *{published and unpublished data}*  
McColl EM, Eccles MPM, Rousseau NSB, Steen INP, Parkin DWD, Grimshaw JMP. From the generic to the condition-specific? Instrument order effects in quality of life assessment. *Medical Care* 2003;**41**(7):777–90.
- McColl 2003 2** *{published and unpublished data}*  
McColl EM, Eccles MPM, Rousseau NSB, Steen INP, Parkin DWD, Grimshaw JMP. From the generic to the condition-specific? Instrument order effects in quality of life assessment. *Medical Care* 2003;**41**(7):777–90.
- Nakash 2007** *{unpublished data only}*  
\* Nakash RA. *A Study of Response and Non-Response to Postal Questionnaire Follow-up in Clinical Trials: Chapter 6: A Randomised Controlled Trial of a Method of Improving Response to Postal Questionnaire Follow-up in a Clinical Trial [PhD thesis]*. Warwick, UK: University of Warwick, Warwick Medical School, 2007:106–164.
- Renfroe 2002a** *{published and unpublished data}*  
\* Renfroe EG, Heywood G, Foreman L, Schron E, Powell J, Baessler C, et al. The end-of-study patient survey: methods influencing response rate in the AVID trial. *Controlled Clinical Trials* 2002;**23**(5):521–33.  
Renfroe EG, Powell J, Olarte A, Morris M, Hallstrom A. Method of administering patient surveys affects response rate. *Controlled Clinical Trials* 1999;**20**:s62.
- Renfroe 2002a-d** *{published and unpublished data}*  
Renfroe EG, Heywood G, Foreman L, Schron E, Powell J, Baessler C. The end-of-study patient survey: methods influencing response rate in the AVID trial. *Controlled Clinical Trials* 2002;**23**(5):521–33.
- Renfroe 2002b** *{published and unpublished data}*  
Renfroe EG, Heywood G, Foreman L, Schron E, Powell J, Baessler C, et al. The end-of-study patient survey: methods influencing response rate in the AVID trial. *Controlled Clinical Trials* 2002;**23**(5):521–33.
- Renfroe 2002c** *{published and unpublished data}*  
Renfroe EG, Heywood G, Foreman L, Schron E, Powell J, Baessler C, et al. The end-of-study patient survey: methods influencing response rate in the AVID trial. *Controlled Clinical Trials* 2002;**23**(5):521–33.
- Renfroe 2002d** *{published and unpublished data}*  
Renfroe EG, Heywood G, Foreman L, Schron E, Powell J, Baessler C, et al. The end-of-study patient survey: methods influencing response rate in the AVID trial. *Controlled Clinical Trials* 2002;**23**(5):521–33.

**Severi 2011** *{published data only}*

\* Severi E, Free C, Knight R, Robertson S, Edwards P, Hoile E. Two controlled trials to increase participant retention in a randomized controlled trial of mobile phone-based smoking cessation support in the United Kingdom. *Clinical Trials* 2011;**8**(5):654–60.

**Severi 2011 1** *{published data only}*

Severi E, Free C, Knight R, Robertson S, Edwards P, Hoile E. Two controlled trials to increase participant retention in a randomized controlled trial of mobile phone-based smoking cessation support in the United Kingdom. *Clinical Trials* 2011;**8**(5):654–60.

**Severi 2011 2** *{published data only}*

Severi E, Free C, Knight R, Robertson S, Edwards P, Hoile E. Two controlled trials to increase participant retention in a randomized controlled trial of mobile phone-based smoking cessation support in the United Kingdom. *Clinical Trials* 2011;**0**:1–7.

**Sharp 2006a** *{published and unpublished data}*

Cotton SC, Cochran C, Sharp L, Gray N, Gallagher ME. Comparison of three low cost strategies to increase response rates to postal questionnaires. Results from Tombola (Trial of management of borderline and other low-grade abnormal smears). *Clinical Trials* 2008;**5**:388.

\* Sharp L, Cochran C, Cotton SC, Gray NM, Gallagher ME. Enclosing a pen with a postal questionnaire can significantly increase the response rate. *Journal of Clinical Epidemiology* 2006;**59**(7):747–54.

**Sharp 2006a-h** *{published and unpublished data}*

Sharp L, Cochran C, Cotton SC, Gray NM, Gallagher ME. Enclosing a pen with a postal questionnaire can significantly increase the response rate. *Journal of Clinical Epidemiology* 2006;**59**(7):747–54.

**Sharp 2006b** *{published and unpublished data}*

Sharp L, Cochran C, Cotton SC, Gray NM, Gallagher ME. Enclosing a pen with a postal questionnaire can significantly increase the response rate. *Journal of Clinical Epidemiology* 2006;**59**(7):747–54.

**Sharp 2006c** *{published and unpublished data}*

Sharp L, Cochran C, Cotton SC, Gray NM, Gallagher ME. Enclosing a pen with a postal questionnaire can significantly increase the response rate. *Journal of Clinical Epidemiology* 2006;**59**(7):747–54.

**Sharp 2006d** *{published and unpublished data}*

Sharp L, Cochran C, Cotton SC, Gray NM, Gallagher ME. Enclosing a pen with a postal questionnaire can significantly increase the response rate. *Journal of Clinical Epidemiology* 2006;**59**(7):747–54.

**Sharp 2006e** *{published and unpublished data}*

Sharp L, Cochran C, Cotton SC, Gray NM, Gallagher ME. Enclosing a pen with a postal questionnaire can significantly increase the response rate. *Journal of Clinical Epidemiology* 2006;**59**(7):747–54.

**Sharp 2006f** *{published and unpublished data}*

Sharp L, Cochran C, Cotton SC, Gray NM, Gallagher ME. Enclosing a pen with a postal questionnaire can significantly

increase the response rate. *Journal of Clinical Epidemiology* 2006;**59**(7):747–54.

**Sharp 2006g** *{published and unpublished data}*

Sharp L, Cochran C, Cotton SC, Gray NM, Gallagher ME. Enclosing a pen with a postal questionnaire can significantly increase the response rate. *Journal of Clinical Epidemiology* 2006;**59**(7):747–54.

**Sharp 2006h** *{published and unpublished data}*

Sharp L, Cochran C, Cotton SC, Gray NM, Gallagher ME. Enclosing a pen with a postal questionnaire can significantly increase the response rate. *Journal of Clinical Epidemiology* 2006;**59**(7):747–54.

**Subar 2001** *{published and unpublished data}*

Subar AF, Ziegler RG, Thompson FE, Johnson CC, Weissfeld JL, Reding D, et al. Is shorter always better? Relative importance of questionnaire length and cognitive ease on response rates and data quality for two dietary questionnaires. *American Journal of Epidemiology* 2001;**153**(4):404–9.

**Sutherland 1996** *{published and unpublished data}*

Sutherland HJ, Beaton M, Mazer R, Kriukov V, Boyd NF. A randomized trial of the total design method for the postal follow-up of women in a cancer prevention trial. *European Journal of Cancer Prevention* 1996;**5**(3):165–8.

**Svoboda 2001** *{published and unpublished data}*

\* Svoboda P. A comparison of two questionnaires for assessing outcome after head injury. Personal email communication from P Edwards in April 2010.

**Tai 1997** *{published and unpublished data}*

Tai SS, Nazareth I, Haines A, Jowett C. A randomized trial of the impact of telephone and recorded delivery reminders on the response rate to research questionnaires. *Journal of Public Health Medicine* 1997;**19**(2):219–21.

## References to studies excluded from this review

**Arnevik 2009** *{published data only}*

Arnevik E, Wilberg T, Urnes Ø, Johansen M, Monsen J, Karterud S. Psychotherapy for personality disorders: short-term day hospital psychotherapy versus outpatient individual therapy - a randomized controlled study. *European Psychiatry* 2009;**24**:71–8.

**Atherton 2010** *{published data only}*

Atherton H, Oakeshott P, Aghaizu A, Hay P, Kerry S. Use of an online questionnaire for follow-up of young female students recruited to a randomised controlled trial of chlamydia screening. *Journal of Epidemiology and Community Health* 2010;**64**(580):e584.

**Barry 1996** *{published and unpublished data}*

Barry MJ, Walker-Corkery E, Chang Y, Tyll LT, Cherkin DC, Fowler FJ. Measurement of overall and disease-specific health status: does the order of questionnaires make a difference. *Journal of Health Services Research and Policy* 1996;**1**(1):20–7.

**Bednarek 2008** *{published data only}*

Bednarek P, Nichols M, Carlson N, Edelman A, Creinin M, Truitt S, et al. Effect of “observed start” vs. traditional



- “Sunday start” on hormonal contraceptive continuation rates after medical abortion. *Contraception* 2008;**78**:26–30.
- Cox 2003** {published data only}  
\* Cox KL, Burke V, Gorely TJ, Beilin LJ, Puddey IB. Controlled comparison of retention and adherence in home- vs center-initiated exercise interventions in women ages 40-65 years: the S.W.E.A.T. study (Sedentary Women Exercise Adherence Trial). *Preventive Medicine* 2003;**36**:17–29.
- Day 1998** {published data only}  
Day R, Cella D, Ganz P, Constantino J. Electronic monitoring of participant adherence in the NSABP breast cancer prevention trial (BCPT). *Controlled Clinical Trials* 1998;**19**:69s.
- Eaker 2004** {published data only}  
Eaker S, Adami H, Granath F, Wilander E, Sparen P. A large population-based randomized controlled trial to increase attendance at screening for cervical cancer. *Cancer Epidemiology, Biomarkers & Prevention* 2004;**13**(3):346–54.
- Edelstein 2005** {published data only}  
Edelstein S, Minutillo A, Pyle L, Anderson B, Wilfley D, Van Buren D, et al. Rewarding youth behaviour: implementation of incentives in the TODAY trial. *Clinical Trials* 2005;**2**:s23.
- Grabowski 1995** {published data only}  
\* Grabowski JP, Rhoades HP, Elk RP, Schmitz JP, Davis CP, Creson DM, et al. Fluoxetine is ineffective for treatment of cocaine dependence or concurrent opiate and cocaine dependence: two placebo-controlled, double-blind trials. *Journal of Clinical Psychopharmacology* 1995;**15**(3):163–74.
- Hall 1975** {published data only}  
Hall S, Hall R, Borden B, Hanson R. Follow-up strategies in the behavioural treatment of overweight. *Behaviour Research and Therapy* 1975;**13**:167–72.
- Hall 1978** {published data only}  
Hall S, Bass A, Monroe J. Continued contact and monitoring as follow-up strategies: a long-term study of obesity treatment. *Addictive Behaviours* 1978;**3**:139–47.
- Hoffman 1998** {published data only}  
Hoffman S, Burke A, Helzlsouer K, Comstock G. Controlled trial of the effect of length, incentives, and follow-up techniques on response to a mailed questionnaire. *American Journal of Epidemiology* 1998;**148**(10):1007–11.
- Hopkins 1983** {published data only}  
Hopkins K, Podolak J. Class-of-mail and the effects of monetary gratuity on the response rates of mailed questionnaires. *Journal of Experimental Education* 1983;**51**(4):169–70.
- Iglesias 2000** {published data only}  
\* Iglesias C, Torgerson D. Does length of questionnaire matter? A randomised trial of response rates to a mailed questionnaire. *Journal of Health Services Research & Policy* 2000;**5**(4):219–21.
- Iglesias 2001** {published data only}  
Iglesias CP, Birks YF, Torgerson DJ. Improving the measurement of quality of life in older people: the York SF-12. *Quality of Life Research* 2001;**10**:103–11.
- Johnson 2004** {published data only}  
Johnson G, Oxman M, Schmader K, Levin M, Concato J, Williams H, et al. Use of an automated telephone response system for follow-up in a long-term multi-centre clinical trial. *Clinical Trials* 2004;**2**:211.
- Katz 2001** {published data only}  
Katz K, El-Mohandes A, McNeely Johnson D, Jarrett M, Rose A, et al. Retention of low income mothers in a parenting intervention study. *Journal of Community Health* 2001;**26**(3):203–18.
- Leidy 2000** {published and unpublished data}  
Leidy NK, Legro M, Schmier J, Zyczynski T, McHorney C, Coyne K. Does order of administration affect subject response on generic and condition-specific measures of health related quality of life?. *Quality of Life Research* 2000;**9**:337.
- Marsh 1999** {published and unpublished data}  
Marsh P, Kendrick D. Using a diary to record near misses and minor injuries: which method of administration is best. *Injury Prevention* 1999;**5**:305–9.
- McAuley 1994** {published data only}  
McAuley E, Courneya K, Rudolph D, Cox CL. Enhancing exercise in middle aged males and females. *Preventive Medicine* 1994;**23**:498–506.
- McBee 2009** {published data only}  
McBee W, Clemons T, Chew E, SanGiovanni JP. A multi-level approach for retention of participants in a long-term clinical trial. *Clinical Trials* 2009;**6**:493–524.
- Poling 2006** {published data only}  
Poling J, Oliveto A, Petry N, Sofuoglu M, Gonsai K, Gonzalez G, et al. Six-month trial of bupropion with contingency management for cocaine dependence in a methadone-maintained population. *Archives of General Psychiatry* 2006;**63**:219–28.
- Puffer 2004** {published data only}  
Puffer S, Porthouse J, Birks Y, Morton V, Torgerson D. Increasing response rates to postal questionnaires: a randomised trial of variations in design. *Journal of Health Services Research and Policy* 2004;**9**(4):213–7.
- Rhoades 1998** {published data only}  
Rhoades H, Creson D, Elk R, Schmitz J, Grabowski J. Retention, HIV risk, and illicit drug use during treatment: methadone dose and visit frequency. *American Journal of Public Health* 1998;**88**(1):34–9.
- Roberts 2000** {published data only}  
Roberts P, Roberts C, Sibbald B, Torgerson D. The effect of a direct payment or a lottery on questionnaire response rates: a randomised controlled trial. *Journal of Epidemiology and Community Health* 2000;**54**:71–2.
- Schmitz 2005** {published data only}  
Schmitz J, Sayre S, Stotts A, Rothfleisch J, Mooney M. Medication compliance during a smoking cessation clinical trial. *Journal of Clinical Pharmacy and Therapeutics* 2005;**30**:103–11.

trial: a brief intervention using MEMS feedback. *Journal of Behavioral Medicine* 2005;**28**(2):139–47.

**Smeeth 2001ab** {published data only}

Smeeth L, Fletcher AE, Stirling S, Nunes M, Breeze E, Ng E, et al. Randomised comparison of three methods of administering a screening questionnaire to elderly people: findings from the MRC trial of the assessment and management of older people in the community. *BMJ* 2001; **323**(7326):1403.

**Stoner 1998** {published data only}

Stoner T, Dowd B, Carr W, Maldonado G, Church T, Mandel J. Do vouchers improve breast cancer screening rates? Results from a randomized trial. *Health Services Research* 1998;**33**(1):11–28.

**Tassopoulos 2007** {published data only}

Tassopoulos C. The challenges of follow-up in a multi-centre randomised controlled trial. *Clinical Trials* 2007;**4**: 371–455.

**Wu 1997** {published and unpublished data}

Wu AW, Jacobson DL, Berzon RA, Revicki DA, van der Horst C, Fichtenbaum CJ, et al. The effect of mode of administration on medical outcomes study health ratings and EuroQol scores in AIDS. *Quality of Life Research* 1997; **6**(1):3–10.

## References to ongoing studies

**Land 2007 2** {unpublished data only}

Randomised Evaluation of NSABP BAHO Compliance Initiatives. Ongoing study Unclear.

**Mitchell** {unpublished data only}

A Randomised Controlled Trial of Combined Pre-Contact and Participant Update for Increasing Questionnaire Response Rates in Older Women. Ongoing study 2010.

## Additional references

**Anastasi 2005**

Anastasi JK, Capili B, Kim G, Chung A. Clinical trial recruitment and retention of a vulnerable population: HIV patients with chronic diarrhea. *Gastroenterology Nursing* 2005;**28**(6):463–8.

**Arnow 2007**

Arnow BA, Blasey C, Manber R, Constantino MJ, Markowitz JC, Klein DN, et al. Dropouts versus completers among chronically depressed outpatients. *Journal of Affective Disorders* 2007;**97**(1-3):197–202.

**AVID investigators 1997**

The Antiarrhythmics versus Implantable Defibrillators (AVID) Investigators. A comparison of antiarrhythmic-drug therapy with implantable defibrillators in patients resuscitated from near-fatal ventricular arrhythmias. *New England Journal of Medicine* 1997;**337**(22):1576–84.

**Booker 2011**

Booker C, Harding S, Benzeval M. A systematic review of the effect of retention methods on response rates in

population-based cohort studies. *BMC Public Health*. 2011;**11**:249.

**Boyd 2002**

Boyd N, Cousins M, Lockwood G, Trichler D. Dietary fat and breast cancer risk: the feasibility of a clinical trial of breast cancer prevention. *Lipids* 1992;**27**(10):821–6.

**Burns 2008**

Burns D, Soward ACM, Skelly AH, Leeman J, Carlson J. Effective recruitment and retention strategies for older members of rural minorities. *The Diabetes Educator* 2008; **34**(6):1045–52.

**Cooke 2009**

Cooke MW, Marsh JL, Clarke M, Nakash R, Jarvis RM, Hutton JL, et al. Treatment of severe ankle sprain: a pragmatic randomised controlled trial comparing the clinical effectiveness and cost effectiveness of three types of mechanical ankle support with tubular bandage. The CAST trial. NIHR Health Technology Assessment Programme Report 2009; Vol. 13:1–144.

**CRASH Trial collaborators 2004**

CRASH Trial Collaborators. Effect of intravenous corticosteroids on death within 14 days in 10008 adults with clinically significant head injury (MRC CRASH trial): randomised placebo-controlled trial. *Lancet* 2004;**364**: 1321–28.

**Davis 2002**

Davis LL, Broome ME, Cox R. Maximizing retention in community-based clinical trials. *Journal of Nursing Scholarship* 2002;**34**(1):47–53.

**De Sousa 2008**

De Sousa AA, De Sousa J, Kapoor H. An open randomized trial comparing disulfiram and topiramate in the treatment of alcohol dependence. *Journal of Substance Abuse Treatment* 2008;**34**(4):460–3.

**Dennis 2009**

Dennis CL, Hodnett E, Kenton L, Weston J, Zupancic J, Stewart DE, et al. Effect of peer support on prevention of postnatal depression among high risk women: multi site randomised controlled trial. *BMJ* 2009;**338**:a3064.

**Eady 2008**

Eady AM, Wilczynski NL, Haynes RB. PsycINFO search strategies identified methodologically sound therapy studies and review articles for use by clinicians and researchers. *Journal of Clinical Epidemiology* 2008;**61**(1):34–40.

**Eakin 2007**

Eakin EG, Bull SS, Riley K, Reeves MM, Gutierrez S, McLaughlin P. Recruitment and retention of Latinos in a primary care-based physical activity and diet trial: the Resources for Health study. *Health Education Research* 2007;**22**(3):361–71.

**Eccles 2002**

Eccles M, McColl E, Steen N, Rousseau N, Grimshaw J, Parkin D, et al. Effect of computerised evidence based guidelines on management of asthma and angina in adults in primary care: cluster randomised controlled trial. *BMJ* 2002;**325**(7370):941.

**Edwards 2009**

Edwards PJ, Roberts I, Clarke MJ, DiGiuseppi C, Wentz R, Kwan I, et al. Methods to increase response to postal and electronic questionnaires. *Cochrane Database of Systematic Reviews* 2009, Issue 3. [DOI: 10.1002/14651858.MR000008.pub4]

**Egger 1997**

Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**(7109):629–34.

**El Khorazaty 2007**

El Khorazaty MN, Johnson A, Kiely M, El Mohandes A, Subramanian S, Laryea H, et al. Recruitment and retention of low-income minority women in a behavioral intervention to reduce smoking, depression, and intimate partner violence during pregnancy. *BMC Public Health* 2007;**7**(1): 233.

**Fewtrell 2008**

Fewtrell MS, Kennedy K, Singhal A, Martin RM, Ness A, Hadders-Algra M, et al. How much loss to follow-up is acceptable in long-term randomised trials and prospective studies?. *Archives of Disease in Childhood* 2008;**93**(6): 458–61.

**Francis 2007**

Francis D, Roberts I, Elbourne D, Shakur H, Knight R, Garcia J, et al. Marketing and clinical trials: a case study. *Trials* 2007;**8**(1):37.

**Free 2011**

Free C, Knight R, Robertson S, Whittaker R, Edwards P, Zhou W, et al. Smoking cessation support delivered via mobile phone text messaging (txt2stop): a single-blind, randomised trial. *Lancet* 2011;**378**:49–55.

**Furimsky 2008**

Furimsky I, Cheung AH, Dewa CS, Zipursky RB. Strategies to enhance patient recruitment and retention in research involving patients with a first episode of mental illness. *Contemporary Clinical Trials* 2008;**29**(6):862–6.

**Gail 1992**

Gail M, Byar D, Pechacek T, Corle D, for the COMMIT Study Group. Aspects of statistical design for the community intervention trial for smoking cessation (COMMIT). *Controlled Clinical Trials* 1992;**13**(1):6–21.

**Goldberg 2005**

Goldberg JH, Kiernan M. Innovative techniques to address retention in a behavioral weight-loss trial. *Health Education Research* 2005;**20**(4):439–47.

**Higgins 2003**

Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**(7414):557–60.

**Higgins 2008a**

Higgins J, Altman D. Chapter 8: Assessing risk of bias in included studies. In: Higgins JPT, Green S editor(s). *Cochrane Handbook of Systematic Reviews of Interventions*. Chichester: John Wiley & Sons, 2008:188–241.

**Higgins 2008b**

Higgins JPT, Deeks JJ, Altman DG (editors). Chapter 16: Special topics in statistics. In: Higgins JPT, Green S editor(s). *Cochrane Handbook for Systematic Reviews of Interventions*. Chichester: John Wiley & Sons, 2008: 482–529.

**Hollis 1999**

Hollis S, Campbell F. What is meant by intention to treat analysis? Survey of published randomised controlled trials. *BMJ* 1999;**319**(7211):670–4.

**Horsley 2011**

Horsley T, Dingwall O, Sampson M. Checking reference lists to find additional studies for systematic reviews. *Cochrane Database of Systematic Reviews* 2011, Issue 8. [DOI: 10.1002/14651858.MR000026.pub2]

**Hughes 1984**

Hughes JR, Hatsukami DK, Pickensl RW, Krahn D, Malin S, Luknic A. Effect of nicotine on the tobacco withdrawal syndrome. *Psychopharmacology* 1984;**83**(1):82–7.

**International Stroke Trial Group 1997**

International Stroke Trial Collaborative Group. The International Stroke Trial (IST): a randomised trial of aspirin, subcutaneous heparin, both, or neither among 19435 patients with acute ischaemic stroke. *Lancet* 1997;**349**:1569–81.

**Kenyon 2001**

Kenyon SL, Taylor DJ, Tarnow-Mordi W. Broad-spectrum antibiotics for preterm, pre labour rupture of fetal membranes: the ORACLE I randomised trial. *Lancet* 2001;**357**(9261):979–88.

**Lamb 2007**

Lamb S, Gates S, Underwood M, Cooke M, Ashby D, Szczepura A, et al. Managing Injuries of the Neck Trial (MINT): design of a randomised controlled trial of treatments for whiplash associated disorders. *BMC Musculoskeletal Disorders* 2007;**8**(1):7.

**Leathem 2009**

Leathem C, Cupples M, Byrne M, O'Malley M, Houlihan A, Murphy A, et al. Identifying strategies to maximise recruitment and retention of practices and patients in a multi-centre randomised controlled trial of an intervention to optimise secondary prevention for coronary heart disease in primary care. *BMC Medical Research Methodology* 2009;**9**(1):40.

**Lefebvre 2008**

Lefebvre C, Manheimer E, Glanville J. Searching for studies. In: Higgins JPT, Green S editor(s). *Cochrane Handbook for Systematic Reviews of Interventions*. Chichester: John Wiley & Sons, 2008:95–150.

**Leigh Brown 2001**

Leigh Brown A, Lawrie H, Kennedy A, Webb J, Torgerson D, Grant A. The OMENS trial: opportunistic evaluation of musculo-skeletal physician care among orthopaedic outpatients unlikely to require surgery. *Health Bulletin* 2001;**59**(3):198–210.

**Loftin 2005**

Loftin WA, Barnett SK, Bunn PS, Sullivan P. Recruitment and retention of rural African Americans in diabetes research: lessons learned. *The Diabetes Educator* 2005;**31**(2):251–9.

**Loue 2008**

Loue S, Sajatovic M. Research with severely mentally ill Latinas: successful recruitment and retention strategies. *Journal of Immigrant and Minority Health* 2008;**10**(2): 145–53.

**Marson 2007 (2)**

Marson A, Al-Kharusi A, Alwaidh M, Appleton R, Baker G, Chadwick D, et al. The SANAD study of effectiveness of carbamazepine, gabapentin, lamotrigine, oxcarbazepine, or topiramate for treatment of partial epilepsy: an unblinded randomised controlled trial. *Lancet* 2007;**369**(9566): 1000–15.

**McKinstry 2007**

McKinstry B, Hammersley V, Daly F, Sullivan F. Recruitment and retention in a multi-centre randomised controlled trial in Bell's palsy: a case study. *BMC Medical Research Methodology* 2007;**7**(1):15.

**Murray 2007**

Murray E, McCambridge J, Khadjesari Z, White I, Thompson S, Godfrey C, et al. The DYD-RCT protocol: an on-line randomised controlled trial of an interactive computer-based intervention compared with a standard information website to reduce alcohol consumption among hazardous drinkers. *BMC Public Health* 2007;**7**(1):306.

**Nakash 2006 (2)**

Nakash R, Hutton J, Jørstad-Stein E, Gates S, Lamb SE. Maximising response to postal questionnaires - a systematic review of randomised trials in health research. *BMC Medical Research Methodology* 2006;**6**(5):1–9.

**Omenn 2006**

Omenn G, Goodman G, Thornquist M, Almes J, Cullen M, Glass A, et al. Effects of a combination of beta carotene and vitamin A on lung cancer and cardiovascular disease. *New England Journal of Medicine* 1996;**334**(18):1150–5.

**Petrosino 2000**

Petrosino A, Boruch RF, Rounding C, McDonald S, Chalmers I. The Campbell Collaboration Social, Psychological, Educational and Criminological Trials Register (C2-SPECTR) to facilitate the preparation and maintenance of systematic reviews of social and educational interventions. *Evaluation & Research in Education* 2000;**14**(3):206–19.

**Porterhouse 2005**

Porterhouse J, Cockayne S, King C, Saxon L, Steele E, Aspray T, et al. Randomised controlled trial of calcium and supplementation with cholecalciferol (vitamin D3) for prevention of fractures in primary care. *BMJ* 2005;**330**:1–6.

**Prorok 2000**

Prorok PC, Andriole GL, Bresalier RS, Buys SS, Chia D, Crawford D, et al. Design of the Prostate, Lung, Colorectal

and Ovarian (PLCO) cancer screening trial. *Controlled Clinical Trials* 2000;**21**:273s–9s.

**RECORD 2007**

RECORD team. Oral vitamin D3 and calcium for secondary prevention of low-trauma fractures in elderly people (Randomised Evaluation of Calcium Or vitamin D, RECORD): a randomised placebo-controlled trial. *Lancet* 2007;**365**:1621–28.

**Rendell 2007**

Rendell JM, Merritt RK, Geddes J. Incentives and disincentives to participation by clinicians in randomised controlled trials. *Cochrane Database of Systematic Reviews* 2007, Issue 2. [DOI: 10.1002/14651858.MR000021.pub3]

**RevMan 2012**

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). 5.2. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2012.

**Robinson 2007**

Robinson KA, Dennison CR, Wayman DM, Pronovost PJ, Needham DM. Systematic review identifies number of strategies important for retaining study participants. *Journal of Clinical Epidemiology* 2007;**60**(8):757–65.

**Rothert 2006**

Rothert K, Strecher V, Doyle L, Caplan W, Joyce J, Jimison H, et al. Web-based weight management programs in an integrated health care setting: a randomized, controlled trial. *Obesity* 2006;**14**(2):266–72.

**Schulz 2002**

Schulz KF, Grimes DA. Sample size slippages in randomised trials: exclusions and the lost and wayward. *Lancet* 2002;**359**(9308):781–5.

**Schünemann 2008**

Schünemann HJ, Oxman AD, Vist GE, Higgins JPT, Deeks JJ, Glasziou P, et al. Chapter 12: Interpreting results and drawing conclusions. In: Higgins JPT, Green S editor(s). *Cochrane Handbook for Systematic Reviews of Interventions*. Chichester: John Wiley & Sons, 2008:359–87.

**Senturia 1998**

Senturia YD, McNiff Mortimer K, Baker D, Gergen P, Mitchell H, Joseph C, et al. Successful techniques for retention of study participants in an inner-city population. *Controlled Clinical Trials* 1998;**19**(6):544–54.

**Snow 2007**

Snow WM, Connett JE, Sharma S, Murray RP. Predictors of attendance and dropout at the Lung Health Study 11-year follow-up. *Contemporary Clinical Trials* 2007;**28**(1):25–32.

**Sterne 2008**

Sterne J, Egger M, Moher D. Chapter 10: Addressing reporting biases. In: Higgins JPT, Green S editor(s). *Cochrane Handbook for Systematic Reviews of Interventions*. Chichester: John Wiley & Sons, 2008:297–333.

**Tai 1999**

Tai S, Nazareth I, Donegan C, Haines A. Evaluation of general practice computer templates. *Methods of Information in Medicine* 1999;**38**:177–81.

**Tilbrook 2011**

Tilbrook HE, Cox H, Hewitt CE, Kang'ombe AR, Chuang LH, Jayakody S, Aplin JD, et al. Yoga for chronic low back pain. *Annals of Internal Medicine* 2011;**155**:569–78.

**TOMBOLA 2009a**

TOMBOLA Group. Biopsy and selective recall compared with immediate large loop excision in management of women with low grade abnormal cervical cytology referred for colposcopy: multi-centre randomised controlled trial. *BMJ* 2009;**339**:2548.

**TOMBOLA 2009b**

TOMBOLA Group. Cytological surveillance compared with immediate referral for colposcopy in management of women with low grade cervical abnormalities: multi-centre randomised controlled trial. *BMJ* 2009;**339**:2546.

**Treweek 2010**

Treweek S, Mitchell E, Pitkethly M, Cook J, Kjeldstrøm M, Taskila T, et al. Strategies to improve recruitment to randomised controlled trials. *Cochrane Database of Systematic Reviews* 2010, Issue 4. [DOI: 10.1002/14651858.MR000013.pub5]

**UK BEAM 2004**

UKBEAM Trial Team. United Kingdom back pain

exercise and manipulation (UK BEAM) randomised trial: effectiveness of physical treatments for back pain in primary care. *BMJ* 2004;**329**:1377.

**Villacorta 2007**

Villacorta V, Kegeles S, Galea J, Konda KA, Cuba JB, Palacios CFC, et al. Innovative approaches to cohort retention in a community-based HIV/STI prevention trial for socially marginalized Peruvian young adults. *Clinical Trials* 2007;**4**(1):32–41.

**Villarruel 2006**

Villarruel AM, Jemmott JB, III, Jemmott LS. A randomized controlled trial testing an HIV prevention intervention for Latino youth. *Archives of Pediatrics and Adolescent Medicine* 2006;**160**(8):772–7.

**Wong 2006a**

Wong S-L, Wilczynski NL, Haynes RB. Developing optimal search strategies for detecting clinically sound treatment studies in EMBASE. *Journal of the Medical Library Association* 2006;**94**(1):41–7.

**Wong 2006b**

Wong S-L, Wilczynski NL, Haynes RB. Optimal CINAHL search strategies for identifying therapy studies and review articles. *Journal of Nursing Scholarship* 2006;**38**(2):194–9.

\* Indicates the major publication for the study

## CHARACTERISTICS OF STUDIES

### Characteristics of included studies [ordered by study ID]

#### Ashby 2011

Methods	2-arm randomised trial
Data	Adults aged 18-65 years who provided email or mobile telephone contact details for receiving electronic reminders for follow-up in a migraine prevention trial
Comparisons	1. Electronic reminder: either SMS text message, email message, or both sent after the 4-week follow-up study questionnaire sent 2. No electronic reminder sent
Outcomes	Primary: questionnaire response rate defined as proportion of questionnaires returned by participants at final analysis at 40 days Secondary: time to response
Notes	Retention trial embedded in a randomised trial evaluating the effectiveness of food elimination diet based on the ELISA test for food sensitivity for prevention of migraine. Primary outcome for the migraine prevention trial (host trial): change in the number of headache days over 12 weeks using the migraine disability assessment questionnaire (MIDAS). Retention trial identified through mail out to UK clinical trials units

#### *Risk of bias*

Item	Authors' judgement	Description
Allocation concealment?	Yes	Authors response "an independent data manager at the trials unit was responsible for generating the allocation sequence and assigning participants"
Adequate sequence generation?	Unclear	"Randomly generated numbers used to list all participants by ID [identification] number who had provided a mobile phone number and/or and email address. The first half of listed participants were allocated to the intervention group the remaining participants were allocated to the control group"
Blinding?	Unclear	No reference to blinding of either participants or outcome assessors in the study report
Free of selective outcome reporting?	Yes	All defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement

**Avenell 2004**

Methods	Randomised trial
Data	Adults aged $\geq 70$ years with a low trauma osteoporotic fracture in the past 10 years recruited in 1 centre of the RECORD trial
Comparisons	1. An open version of the RECORD trial otherwise identical in design 2. RECORD trial, a randomised double-blind placebo-controlled factorial design of oral calcium 1 g daily and or vitamin D 800 IU/20 $\mu\text{g}$ supplementation
Outcomes	Proportion of eligible participants recruited Proportion remaining in the trial at 1 year Proportion compliant on pill counts at 8 months
Notes	Open version of the Randomised Evaluation of Calcium or Vitamin D (RECORD) Trial treated as the intervention group in the analysis. Proportion retained at 4, 8 and 12 months were reported. Primary outcome for the randomised double-blind placebo-controlled version of the RECORD trial was all new low-energy fractures

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Yes	"Nurse used a pre programmed laptop computer"
Adequate sequence generation?	Yes	"A pre programmed laptop computer to generate random allocation"
Blinding?	Yes	Double-blind randomised trial design compared with an open trial design. For the double-blind randomised trial "allocation remained concealed until the final analyses". "All outcomes were reported or verified by people who were masked to the allocation scheme"
Free of selective outcome reporting?	Yes	All defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement

**Bailey 1**

Methods	Randomised trial embedded in the sex unzipped website feasibility trial
Data	UK English speaking people aged 16-20 years
Comparisons	1. Offer of GBP20 voucher 2. Offer of GBP10 voucher
Outcomes	Retention of participants at 3-month follow-up, i.e. completion of sexual health survey
Notes	Retention trial identified through personal correspondence with the author Sexunzipped website evaluated in an online trial

**Bailey 1** (Continued)

<i>Risk of bias</i>		
Item	Authors' judgement	Description
Allocation concealment?	Yes	Authors response "the trial statisticians generated the randomisation sequence with participants identified by ID [identification] number only and the trial manager implemented it manually"
Adequate sequence generation?	Yes	Authors response "participants were randomised after recruitment but before follow-up to a GBP10 or GBP20 incentive. Randomisation to increased incentive was through simple permutation of the list of remaining recruits"
Blinding?	Unclear	Authors response "allocation sequences were generated without participants' knowledge". "For those allocated to the increased amount of GBP20, this was revealed in a 3 month follow-up email. Those allocated to GBP10 were not aware that others were offered GBP20 (unless friends had enrolled and had discussed the study). Since the trial recruited participants online from all over the UK, this will have reduced the chance of bias due to contamination"
Free of selective outcome reporting?	Yes	Reports the primary outcome

**Bailey 2**

Methods	Randomised trial
Data	UK English speaking people aged 16-20 years
Comparisons	1. Offer of GBP20 voucher: GBP10 in advance and GBP10 on receipt of questionnaire and chlamydia kit 2. Offer of GBP10 voucher: GBP5 in advance and GBP5 on receipt of questionnaire and chlamydia kit
Outcomes	Retention of participants at 3-month follow-up, i.e. completion of sexual health survey and return of chlamydia kit
Notes	Study identified through personal correspondence with author Sex unzipped website evaluated in an online trial



**Bailey 2** (Continued)

<i>Risk of bias</i>		
Item	Authors' judgement	Description
Allocation concealment?	Yes	Authors response "the trial statisticians generated the randomisation sequence (with participants identified by ID [identification] number only), and the trial manager implemented it manually"
Adequate sequence generation?	Yes	Authors response "participants were randomised after recruitment but before follow-up to a GBP10 or GBP20 incentive. Randomisation to increased incentive was through simple permutation of the list of remaining recruits"
Blinding?	Unclear	Authors response "allocation sequences were generated without participants' knowledge". "For those allocated to the increased amount of GBP20, this was revealed in a 3 month follow-up email. Those allocated to GBP10 were not aware that others were offered GBP20 (unless friends had enrolled and had discussed the study). Since the trial recruited participants online from all over the UK, this will have reduced the chance of bias due to contamination"
Free of selective outcome reporting?	Yes	Reports the primary outcome

**Bauer 2004a**

Methods	3-arm randomised trial (first incentive vs. no incentive)	
Data	Pilot study of 15 randomly selected participants from each of 20 communities participating in COMMIT trial	
Comparisons	Enclosed with mouthwash swish collection kits sent to participants "subjects were further randomised to receive" either: 1. USD10 cheque told to keep the cheque whether or not they participated, sent with covering letter and prepaid envelope 2. No incentive, kit sent with covering letter and prepaid envelope All sent 2 weeks after an advance letter with a professionally rendered brochure	
Outcomes	Percentage of mouthwash kits returned reported. No time point given	
Notes	Study embedded in the cluster randomised Community Intervention Trial for Smoking Cessation (COMMIT) trial. Primary outcome: quit rate among heavy smokers. Several attempts to contact authors regarding allocation sequence. Data extracted from Edwards Cochrane review on response to postal questionnaires	
<i>Risk of bias</i>		
Item	Authors' judgement	Description

**Bauer 2004a** (Continued)

Allocation concealment?	Unclear	No reply from author
Adequate sequence generation?	Unclear	“Subjects were further randomised to receive an incentive of....”
Blinding?	Unclear	Insufficient information to make a judgement about this, no reply from author
Free of selective outcome reporting?	Unclear	No reply from author
Other sources of bias?	Unclear	Insufficient information to make a judgement

**Bauer 2004ab**

Methods	3-arm randomised trial (combined incentive vs. no incentive)	
Data	Pilot study of 15 randomly selected participants from each of 20 communities participating in COMMIT trial	
Comparisons	Enclosed with mouthwash swish collection kits sent to participants “subjects were further randomised to receive” either: <ol style="list-style-type: none"> <li>1. USD10 cheque told to keep the cheque whether or not they participated sent with covering letter and prepaid envelope or</li> <li>2. USD2 cheque with covering letter and prepaid envelope or</li> <li>3. No incentive, kit sent with covering letter and prepaid envelope</li> </ol> All sent 2 weeks after an advance letter with a professionally rendered brochure	
Outcomes	Percentage of mouthwash kits returned reported. No time point given	
Notes	Study embedded in the cluster randomised Community Intervention Trial for Smoking Cessation (COMMIT) trial. Primary outcome: quit rate among heavy smokers. Several attempts to contact authors regarding allocation sequence. Data extracted from Edwards Cochrane review on response to postal questionnaires	

**Risk of bias**

Item	Authors' judgement	Description
Allocation concealment?	Unclear	No reply from author
Adequate sequence generation?	Unclear	“Subjects were further randomised to receive an incentive of....”
Blinding?	Unclear	Insufficient information to make a judgement about this no reply from author
Free of selective outcome reporting?	Unclear	No reply from author

**Bauer 2004ab** (Continued)

Other sources of bias?	Unclear	Insufficient information to make a judgement
------------------------	---------	--

**Bauer 2004b**

Methods	3-arm randomised trial (second incentive vs. no incentive)
Data	Pilot study of 15 randomly selected participants from each of 20 communities participating in COMMIT trial
Comparisons	Enclosed with mouthwash swish collection kits sent to participants, “subjects were further randomised to receive” either: <ol style="list-style-type: none"> <li>1. USD2 check with covering letter and prepaid envelope</li> <li>2. No incentive, kit sent with covering letter and prepaid envelope</li> </ol> All sent 2 weeks after an advance letter with a professionally rendered brochure
Outcomes	Percentage of mouthwash kits returned reported. No time point given
Notes	Study embedded in the cluster randomised Community Intervention Trial for Smoking Cessation (COMMIT) trial. Primary outcome: quit rate among heavy smokers. Several attempts to contact authors regarding allocation sequence. Data extracted from Edwards Cochrane review on response to postal questionnaires

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Unclear	No reply from author
Adequate sequence generation?	Unclear	“Subjects were further randomised to receive an incentive of.....”
Blinding?	Unclear	Insufficient information to make a judgement about this
Free of selective outcome reporting?	Unclear	No reply from author
Other sources of bias?	Unclear	Insufficient information to make a judgement

**Bowen 2000a**

Methods	Randomised 2 x 2 factorial trial (first incentive vs. no incentive)
Data	Adults aged 50-69 years recruited from 2 CARET trial sites, participating in the Participant Retention Item Distribution Evaluation (PRIDE) trial

Comparisons	<p>1. Certificate of appreciation preprinted on an 8.5 x 11 inch (21.59 x 27.94 cm) bond off-white paper with gold trim and bold, black lettering. The certificate read “[participant’s name] in recognition of your contribution to an important national study for the prevention of lung cancer, CARET, Cancer Prevention Study, sponsored by The National Cancer Institute”. The participant’s name was computer printed in an attractive font on the certificate. Each certificate had the signatures of the Co-ordinating Center’s principal investigator, study centre investigator and CARET’s project officer from the National Cancer Institute, given during a visit for randomisation or follow-up</p> <p>2. No incentive</p>	
Outcomes	Primary: time of first inactivation (stop taking vitamins or placebos) during the 2-year follow-up period of PRIDE	
Notes	Primary disease outcome for the Carotene and Retinol Efficacy Trial (CARET): lung cancer	
<b>Risk of bias</b>		
Item	Authors’ judgement	Description
Allocation concealment?	Unclear	Author response “the Coordination Center supplied one date-labelled envelope for each day in the enrolment period to the two participating study centres. The envelopes contained the randomisation assignment for the given day. At the end of each day during the randomisation period, study centre staff opened the envelope containing the intervention assignment for the next day. Study centre staff members were otherwise blinded to the allocation sequence”
Adequate sequence generation?	Yes	Author response “we used a block randomisation approach (stratified by study centre) with a 1:1:1:1 intervention arm allocation ratio, where the randomisation unit was the date of enrolment”. Note: treated this as quasi-randomised in the analysis
Blinding?	Yes	Author response “with IRB [Institutional Review Board] approval, the study was conducted without participants’ knowledge of this research. Thus, participants were blinded to their own intervention only in the sense that they were unaware they were randomised to receive particular item(s)”
Free of selective outcome reporting?	Yes	Cumulative incidence of individuals who became inactive during 2-year follow-up re-

**Bowen 2000a** (Continued)

		ported
Other sources of bias?	Unclear	Insufficient information to make a judgement

**Bowen 2000abc**

Methods	Randomised 2 x 2 factorial trial (combined incentive vs. no incentive)
Data	Adults aged 50-69 years recruited from 2 CARET trial sites, participating in the Participant Retention Item Distribution Evaluation (PRIDE) trial
Comparisons	<ol style="list-style-type: none"> <li>1. Certificate of appreciation preprinted on an 8.5 x 11 inch (21.59 x 27.94 cm) bond off-white paper with gold trim and bold, black lettering. The certificate read '[participant's name] in recognition of your contribution to an important national study for the prevention of lung cancer, CARET, Cancer Prevention Study, sponsored by The National Cancer Institute'. The participant's name was computer printed in an attractive font on the certificate. Each certificate had the signatures of the Co-ordinating Center's principal investigator, study centre investigator, and CARET's project officer from the National Cancer Institute, given during a visit for randomisation or follow-up (arm a)</li> <li>2. Lapel pin 1 inch (2.5 cm) in size and designed in cloisonne. Choice between a pin with 6 colours with inscription 'CARET NCI prevention study' and an orange carrot in the middle of the pin or a pin with 5 colours with inscription 'PARTICIPNAT CARET Cancer Prevention Study' and 'Sponsored by NCI' and given during a visit for randomisation or follow-up (arm b)</li> <li>3. Certificate of appreciation (details as before) and lapel pin (details as before) and given during a visit for randomisation or follow-up (arm c)</li> <li>4. No incentive</li> </ol>
Outcomes	Primary: time of first inactivation (stop taking vitamins or placebos) during the 2-year follow-up period of PRIDE
Notes	Primary disease outcome for the Carotene and Retinol Efficacy Trial (CARET): lung cancer

**Risk of bias**

Item	Authors' judgement	Description
Allocation concealment?	Unclear	Author response "the Coordination Center supplied one date-labelled envelope for each day in the enrolment period to the two participating study centres. The envelopes contained the randomisation assignment for the given day. At the end of each day during the randomisation period, study centre staff opened the envelope containing the intervention assignment for the next day. Study centre staff members were otherwise blinded to the allocation sequence"

**Bowen 2000abc** (Continued)

Adequate sequence generation?	Yes	Author response “we used a block randomisation approach (stratified by study centre) with a 1:1:1:1 intervention arm allocation ratio, where the randomisation unit was the date of enrolment”
Blinding?	Yes	Author response “with IRB [Institutional Review Board] approval, the study was conducted without participants’ knowledge of this research. Thus, participants were blinded to their own intervention only in the sense that they were unaware they were randomised to receive particular item(s)”
Free of selective outcome reporting?	Yes	Cumulative incidence of individuals who became inactive during 2 year follow-up reported
Other sources of bias?	Unclear	Insufficient information to make a judgement

**Bowen 2000b**

Methods	Randomised 2 x 2 factorial trial (second incentive vs. no incentive)
Data	Adults aged 50-69 years recruited from 2 CARET trial sites, participating in the Participant Retention Item Distribution Evaluation (PRIDE) trial
Comparisons	<ol style="list-style-type: none"> <li>1. Lapel pin 1 inch (2.5 cm) in size and designed in cloisonne. Choice between a pin with 6 colours with inscription ‘CARET NCI prevention study’ and an orange carrot in the middle of the pin or a pin with 5 colours with inscription ‘PARTICIPNAT CARET Cancer Prevention Study’ and “Sponsored by NCI” and given during a visit for randomisation or follow-up</li> <li>2. No incentive</li> </ol>
Outcomes	Primary: time of first inactivation (stop taking vitamins or placebos) during the 2-year follow-up period of PRIDE
Notes	Primary disease outcome for the Carotene and Retinol Efficacy Trial (CARET): lung cancer

***Risk of bias***

Item	Authors’ judgement	Description
Allocation concealment?	Unclear	Author response “the Coordination Center supplied one date-labelled envelope for each day in the enrolment period to the two participating study centres. The envelopes contained the randomisation assignment for the given

**Bowen 2000b** (Continued)

		day. At the end of each day during the randomisation period, study centre staff opened the envelop containing the intervention assignment for the next day. Study centre staff members were otherwise blinded to the allocation sequence”
Adequate sequence generation?	Yes	Author response “we used a block randomisation approach (stratified by study centre) with a 1:1:1:1 intervention arm allocation ratio, where the randomisation unit was the date of enrolment”
Blinding?	Yes	Author response “with IRB [Institutional Review Board] approval, the study was conducted without participants’ knowledge of this research. Thus, participants were blinded to their own intervention only in the sense that they were unaware they were randomised to receive particular item(s)”
Free of selective outcome reporting?	Yes	Cumulative incidence of individuals who became inactive during 2-year follow-up reported
Other sources of bias?	Unclear	Insufficient information to make a judgement

**Bowen 2000c**

Methods	Randomised 2 x 2 factorial trial (third incentive vs. no incentive)
Data	Adults aged 50-69 years recruited from 2 CARET trial sites, participating in the Participant Retention Item Distribution Evaluation (PRIDE) trial
Comparisons	1. Certificate of appreciation (details as before) <b>and</b> lapel pin (details as before) and given during a visit for randomisation or follow-up 2. No incentive
Outcomes	Primary: time of first inactivation (stop taking vitamins or placebos) during the 2-year follow-up period of PRIDE
Notes	Primary disease outcome for the Carotene and Retinol Efficacy Trial (CARET): lung cancer

**Risk of bias**

Item	Authors’ judgement	Description
------	--------------------	-------------

**Bowen 2000c** (Continued)

Allocation concealment?	Unclear	Author response “the Coordination Center supplied one date-labelled envelope for each day in the enrolment period to the two participating study centres. The envelopes contained the randomisation assignment for the given day. At the end of each day during the randomisation period, study centre staff opened the envelop containing the intervention assignment for the next day. Study centre staff members were otherwise blinded to the allocation sequence”
Adequate sequence generation?	Yes	Author response “we used a block randomisation approach (stratified by study centre) with a 1:1:1:1 intervention arm allocation ratio, where the randomisation unit was the date of enrolment”
Blinding?	Yes	Author response “with IRB [Institutional Review Board] approval, the study was conducted without participants’ knowledge of this research. Thus, participants were blinded to their own intervention only in the sense that they were unaware they were randomised to receive particular item(s)”
Free of selective outcome reporting?	Yes	Cumulative incidence of individuals who became inactive during 2-year follow-up reported
Other sources of bias?	Unclear	Insufficient information to make a judgement

**Chaffin 2009**

Methods	2-arm randomised trial
Data	All parents referred for parenting services at a small inter-city non-profit community agency operating a parenting programme
Comparisons	1. Initial preparenting orientation condition self motivation group 2. Initial preparenting orientation condition standard informational group
Outcomes	Dropout from the parenting group at 12 weeks
Notes	A second randomisation was performed after completion of the orientation programme to parent child interactive therapy vs. standard didactic parenting condition. Dropout recorded at 2-week intervals up to 12 weeks



**Chaffin 2009** (Continued)

<i>Risk of bias</i>		
Item	Authors' judgement	Description
Allocation concealment?	No	"Unblinded randomisation list"
Adequate sequence generation?	Yes	"Computer generated randomisation list"
Blinding?	Yes	Author response "they (parents) were informed only in general terms that we were interested in which types of services helped". Participant interviews were conducted by computer. Blinding of personnel: observational parent-child interaction coding was done by personnel who were not informed about intervention condition assignment
Free of selective outcome reporting?	Yes	All defined outcomes reported
Other sources of bias?	No	"It is possible that therapist effects may have played a role in the outcomes"

**Cockayne 2005**

Methods	2-arm randomised trial
Data	Women aged $\geq 70$ years randomised at 1 centre for a fracture prevention trial due to receive their final follow-up questionnaire in March 2004
Comparisons	1. Final follow-up questionnaire additional question offering results of the trial delivered by post 2. Final follow-up questionnaire no offer of study results delivered by post Both groups received a personalised cover letter showing university sponsorship, along with a business reply envelope. Non-responders within 3 weeks were sent up to 2 reminder letters, questionnaires and business reply envelopes, 3 and 6 weeks after the initial mailing
Outcomes	Return of final follow-up questionnaire by participants. Time point not specified
Notes	Authors contacted to confirm numbers randomised to each arm The Fracture Prevention Trial: calcium 1000 mg plus vitamin D <sub>3</sub> 800 IU plus information sheet on dietary calcium intake and falls prevention vs. information sheet. Primary outcome for the fracture prevention trial: all clinical fractures excluding those of the digits, rib, face and skull

*Risk of bias*

Item	Authors' judgement	Description
------	--------------------	-------------

**Cockayne 2005** (Continued)

Allocation concealment?	Yes	“An independent researcher from the York trials unit randomised eligible women”
Adequate sequence generation?	Yes	“Randomised eligible women in a 3:1 ratio by computer”
Blinding?	Unclear	“Administration of the questionnaire was not blind to the group allocation”
Free of selective outcome reporting?	Yes	All outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement

**Couper 2007**

Methods	2-arm randomised trial (incentive plus postal questionnaire vs. telephone survey)
Data	Adults $\geq$ 18 years, BMI of $\geq$ 25 participating in an Internet-based weight management trial who did not respond to the 12-month questionnaire
Comparisons	1. Telephone call and survey by trained interviewers. Repeated up to 15 times. Attempts made on various days and at various times of the day 2. Postal questionnaire with return address and covering letter signed by directors plus a USD5 bill
Outcomes	Questionnaire response. No time point given
Notes	Internet-based weight management trial compared Internet-based tailored weight management materials with Internet-based non-tailored user navigated weight management materials. Primary outcome for the Internet-based trial: percentage of baseline weight lost

***Risk of bias***

<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Unclear	Not described
Adequate sequence generation?	Unclear	Author response “randomly assigned”
Blinding?	Unclear	Not described
Free of selective outcome reporting?	Unclear	Unclear at the outset what is to be reported
Other sources of bias?	Unclear	Insufficient information to make a judgement

**Cox 2008**

Methods	2-arm randomised trial
Data	All sedentary women aged 50-70 years participating in the SWEAT2 trial
Comparisons	1. 12 work sheets with strategies for goal setting, time management and overcoming barriers to attendance, mini workshops, received worksheets to complete at home - after 6 months received newsletters only. Intervention delivered by a trained facilitator, before an exercise session 2. Information sheets about programme requirements exercise techniques, safety plus 9 newsletters and a report on fitness at 6 months delivered by principle investigator plus 9 newsletters
Outcomes	Programme retention at 6 and 12 months
Notes	Contact with authors to clarify if withdrawals occurred before or after randomisation SWEAT2 compared moderate walking programme vs. swimming programme. Primary outcome adherence to the programme

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Yes	"Computer generated random numbers by a statistician"
Adequate sequence generation?	Yes	"Computer generated random numbers"
Blinding?	Unclear	"Not practical to blind the participants or the research staff to the group assignment" (see other sources of bias below)
Free of selective outcome reporting?	Yes	All defined outcomes reported
Other sources of bias?	Yes	"Walk and swim sessions were not separated according to the behavioural intervention participants asked not to discuss written materials in the practical sessions"

**Dorman 1997**

Methods	2-arm randomised trial
Data	All UK participants entered into the International Stroke Trial between 2 March 1993 and 31 May 1995 who were still alive
Comparisons	1. Short EuroQol posted with personalised letter and reply paid envelope. 1 reminder sent after 2 weeks 2. Long SF36 questionnaire posted with personalised letter and reply paid envelope. 1 reminder sent after 2 weeks

**Dorman 1997** (Continued)

Outcomes	Frequency of response after first and reminder mail out. Data for response to first mail out used	
Notes	International Stroke Trial compared heparin 125,000 IU twice daily + aspirin 300 mg vs. heparin 125,000 IU twice daily; heparin 5000 IU twice daily + aspirin 300 mg daily vs. heparin 5000 IU twice daily vs. aspirin 300 mg daily vs. no heparin or aspirin. Primary outcome: death within 14 days or dependency at 6 months	
<b>Risk of bias</b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Yes	Authors response "generated by computer"
Adequate sequence generation?	Unclear	Author response "generated by computer"
Blinding?	Unclear	Authors report "there was no blinding for either study staff or participants"
Free of selective outcome reporting?	Yes	All defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement

**Edwards 2001**

Methods	2-arm randomised trial	
Data	Adults aged $\geq 16$ years with head injury in the UK CRASH trial	
Comparisons	1. 1-page, 7-question functional dependence questionnaire with covering letter and stamped return envelope. Reminders after 4 and 8 weeks 2. 3-page, 16-question functional dependence questionnaire with covering letter and stamped return envelope. Reminders after 4 and 8 weeks	
Outcomes	No of questionnaires returned within 3 months	
Notes	Authors provided numbers randomised and responded Primary outcome for the CRASH: death from any cause within 2 weeks of injury and death or disability at 6 months	
<b>Risk of bias</b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Yes	"Central computer"
Adequate sequence generation?	Unclear	"Central computer"

**Edwards 2001** (Continued)

Blinding?	Unclear	Author response “questionnaires were packaged and sent to patients by personnel who were independent of the study”
Free of selective outcome reporting?	Yes	Reported non-response
Other sources of bias?	Unclear	Insufficient information to make a judgement

**Ford 2006**

Methods	2-arm quasi-randomised trial
Data	African American men aged 55 years enrolled in the intervention screening arm of the PLCO trial
Comparisons	1. Indepth case management. Case management monthly telephone calls to participants, assisted making medical appointments, helped participants obtain health insurance information, legal aid, transportation services, food programmes, financial support, medication assistance, free medical care, information related to health risks facing African Americans. Provision of PLCO Cancer Screening Trial screening information and the scheduling of annual screening appointments. 2. Regular trial screening procedures. Participants called annually to schedule screening examinations
Outcomes	Number completing the next scheduled PLCO cancer screen at 3 years
Notes	PLCO trial compared digital rectal examination, transvaginal ultrasound and chest x-ray at baseline and 5 years to usual follow-up. PSA and cancer antigen CA125 at baseline, and annually for 5 years. Primary outcome mortality from 4 PLCO cancers

***Risk of bias***

<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Unclear	Not described
Adequate sequence generation?	No	“Randomised by participant id number”
Blinding?	Unclear	Insufficient information to make a judgement about this
Free of selective outcome reporting?	Yes	All defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Gates 2009**

Methods	2-arm quasi-randomised trial
Data	All participants enrolled in the MINT trial due a follow-up questionnaire at 4 or 8 months after whiplash injury
Comparisons	1. GBP5 voucher redeemable at shops <a href="http://www.highstreetvouchers.com">www.highstreetvouchers.com</a> plus questionnaire, cover letter included a sentence explaining that the voucher is to thank participants for their time and effort, delivered by post 2. No voucher and a standard covering letter with the questionnaire
Outcomes	Number of questionnaires returned after first contact with participants
Notes	Number of questionnaires returned in the incentive arm checked with authors Primary outcome for the Managing Injuries of the Neck Trial (MINT) was returned to normal after whiplash injury measured using the Neck Disability Index (NDI)

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Unclear	Author response "lack of concealment of allocations before randomisation was not a major concern because it would have been very difficult for the staff in the study office who were sending out the questionnaires to have selectively allocated systematically different patients to the trial arm"
Adequate sequence generation?	No	Author response "allocation to study arms was according to whether a specific digit of the patients study number was odd or even"
Blinding?	Unclear	Author response "trial office staff were unblinded, they had no influence over any participant's decision to return the questionnaire, and postal and telephone follow-up contacts were performed in a standardised way for all participants, without any reference to whether or not they were participating in the incentive study"
Free of selective outcome reporting?	Yes	All defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Hughes 1989**

Methods	2-arm randomised trial
Data	All smokers for > 1 year in the nicotine gum vs. placebo gum trial

**Hughes 1989** (Continued)

Comparisons	1. A letter promising a free reprint of the study results in return for sending in the questionnaire 2. No offer of results
Outcomes	Questionnaire response. No time point given
Notes	Author contacted regarding numbers randomised Nicotine gum trial was double blind. The primary outcome was alleviation of signs and symptoms of tobacco withdrawal measured using a rating list and POMS (profile of mood states) questionnaire and DSM III criteria for tobacco withdrawal

**Risk of bias**

Item	Authors' judgement	Description
Allocation concealment?	Yes	Author response "by a non involved researcher sequence in sealed envelope, never opened during study"
Adequate sequence generation?	Yes	Author response "using a table of random numbers"
Blinding?	Unclear	Author response "participants fully aware of each condition and which they were in"
Free of selective outcome reporting?	Yes	Response rates reported
Other sources of bias?	Unclear	Insufficient evidence to make a judgement on this

**Kenton 2007a**

Methods	Randomised 2 x 2 factorial trial (incentive vs. offer of incentive)
Data	New mothers with Edinburgh postnatal depression scale score > 9 participating in postpartum depression peer support trial
Comparisons	1. USD2 coin mailed with questionnaire 2. Offer of entry into prize draw for USD50 gift certificate
Outcomes	Questionnaire response. No time point given
Notes	The postpartum depression peer support trial enrolled women < 2 weeks' postpartum at high risk for postnatal depression. New proactive individualised telephone-based peer support with standard postpartum care. The primary outcome was postnatal depression measured using the Edinburgh postnatal depression scale and Structured Clinical Interview Depression (SCID)

**Risk of bias**

Item	Authors' judgement	Description
------	--------------------	-------------

**Kenton 2007a** (Continued)

Allocation concealment?	Yes	Author response “allocation sequence stored in a password protected file only accessed by an external researcher, computer generated”
Adequate sequence generation?	Unclear	Author response “computer generated”
Blinding?	Yes	Author response “participants were not aware of the sub-study and didn’t know other participants were receiving different types of mailings. The trial coordinator was not blinded to the study group”
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Kenton 2007a-d**

Methods	Randomised 2 x 2 factorial trial
Data	New mothers with Edinburgh, postnatal depression scale score > 9 participating in postpartum depression peer support trial
Comparisons	1. USD2 coin mailed with questionnaire 2. Offer of entry into prize draw for USD50 gift certificate 3. USD2 coin with questionnaire sent with high-priority postage stamp 4. Offer of entry into lottery draw for USD50 gift certificate with high-priority postage stamp
Outcomes	Questionnaire response. No time point given
Notes	The postpartum depression peer support trial enrolled women < 2 weeks’ postpartum at high risk for postnatal depression. New proactive individualised telephone-based peer support with standard postpartum care. The primary outcome was postnatal depression measured using the Edinburgh postnatal depression scale and Structured Clinical Interview Depression (SCID)

***Risk of bias***

Item	Authors’ judgement	Description
Allocation concealment?	Yes	Author response “allocation sequence stored in a password protected file only accessed by an external researcher, computer generated”
Adequate sequence generation?	Unclear	Author response “computer generated”



**Kenton 2007a-d** (Continued)

Blinding?	Yes	Author response “participants were not aware of the sub-study and didn’t know other participants were receiving different types of mailings. The trial coordinator was not blinded to the study group”
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Kenton 2007b**

Methods	Randomised 2 x 2 factorial trial (incentive + priority mail vs. offer of incentive + priority mail)
Data	New mothers with Edinburgh postnatal depression scale score > 9 participating in postpartum depression peer support trial
Comparisons	1. USD2 coin with questionnaire sent with high-priority postage stamp 2. Offer of entry into lottery draw for USD50 gift certificate with high-priority postage stamp
Outcomes	Questionnaire response. No time point given
Notes	The postpartum depression peer support trial enrolled women < 2 weeks’ postpartum at high risk for postnatal depression. New proactive individualised telephone-based peer support with standard postpartum care. The primary outcome was postnatal depression measured using the Edinburgh postnatal depression scale and Structured Clinical Interview Depression (SCID)

**Risk of bias**

Item	Authors’ judgement	Description
Allocation concealment?	Yes	Author response “allocation sequence stored in a password protected file only accessed by an external researcher, computer generated”
Adequate sequence generation?	Unclear	Author response “computer generated”
Blinding?	Yes	Author response “participants were not aware of the sub-study and didn’t know other participants were receiving different types of mailings. The trial coordinator was not blinded to the study group”
Free of selective outcome reporting?	Yes	Expected outcomes reported

**Kenton 2007b** (Continued)

Other sources of bias?	Unclear	Insufficient information to make a judgement on this
------------------------	---------	--

**Kenton 2007c**

Methods	Randomised 2 x 2 factorial trial (priority mail vs. no priority mail)
Data	New mothers with Edinburgh postnatal depression scale score > 9 participating in postpartum depression peer support trial
Comparisons	1. USD2 coin sent by high-priority postage stamp plus questionnaire 2. USD2 coin plus questionnaire
Outcomes	Questionnaire response. No time point given
Notes	The postpartum depression peer support trial enrolled women < 2 week' postpartum at high risk for postnatal depression. New proactive individualised telephone-based peer support with standard postpartum care. The primary outcome was postnatal depression measured using the Edinburgh postnatal depression scale and Structured Clinical Interview Depression (SCID)

**Risk of bias**

Item	Authors' judgement	Description
Allocation concealment?	Yes	Author response "allocation sequence stored in a password protected file only accessed by an external researcher, computer generated"
Adequate sequence generation?	Unclear	Author response "computer generated"
Blinding?	Yes	Author response "participants were not aware of the sub-study and didn't know other participants were receiving different types of mailings. The trial coordinator was not blinded to the study group"
Free of selective outcome reporting?	Yes	Expected outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Kenton 2007d**

Methods	Randomised 2 x 2 factorial trial (offer of entry into prize draw plus high-priority postage stamp vs. offer of entry into prize draw)
Data	New mothers with Edinburgh postnatal depression scale score > 9 participating in postpartum depression peer support trial
Comparisons	1. Offer of entry into prize draw for USD50 gift certificate plus high-priority postage stamp 2. Offer of entry into prize draw for USD50 gift certificate
Outcomes	Questionnaire response. No time point given
Notes	The postpartum depression peer support trial enrolled women < 2 weeks' postpartum at high risk for postnatal depression. New proactive individualised telephone-based peer support with standard postpartum care. The primary outcome was postnatal depression measured using the Edinburgh postnatal depression scale and Structured Clinical Interview Depression (SCID)

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Yes	Author response "allocation sequence stored in a password protected file only accessed by an external researcher, computer generated"
Adequate sequence generation?	Unclear	Author response "computer generated"
Blinding?	Yes	Author response "participants were not aware of the sub-study and didn't know other participants were receiving different types of mailings. The trial coordinator was not blinded to the study group"
Free of selective outcome reporting?	Yes	Expected outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Kenyon 2005**

Methods	2-arm randomised trial
Data	Women in ORACLE 1 and 2 participating in the evaluation of health and development study
Comparisons	1. GBP5 voucher mailed with questionnaire redeemable at many high street shops 2. No incentive
Outcomes	Questionnaire response rate. No time point given

**Kenyon 2005** (Continued)

Notes	ORACLE trial: women with preterm prelabour rupture of fetal membranes and women with intact membranes in preterm labour, randomised to erythromycin 250 mg, co-amoxiclav 325 mg, erythromycin 250 mg plus co-amoxiclav 325 mg or placebo x 10 days or until birth. Primary outcome composite of neonatal death, chronic lung disease or major cerebral abnormality before discharge from hospital	
<b>Risk of bias</b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Yes	"Randomly assigned by concealed computer generated allocation"
Adequate sequence generation?	Yes	"Randomly assigned by "computer" "
Blinding?	Yes	Author response "both the participants and the study personnel were blinded to the allocation"
Free of selective outcome reporting?	Yes	Expected outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Khadjesari 2011**

Methods	Khadjesari 1: 4-arm randomised trial Khadjesari 2: 2-arm randomised trial	
Data	See 'Table of characteristics' for <a href="#">Khadjesari 2011 1abc</a> ; <a href="#">Khadjesari 2011 2</a>	
Comparisons	See 'Table of characteristics' for <a href="#">Khadjesari 2011 1abc</a> ; <a href="#">Khadjesari 2011 2</a>	
Outcomes	See 'Table of characteristics' for <a href="#">Khadjesari 2011 1abc</a> ; <a href="#">Khadjesari 2011 2</a>	
Notes	See 'Table of characteristics' for <a href="#">Khadjesari 2011 1abc</a> ; <a href="#">Khadjesari 2011 2</a>	
<b>Risk of bias</b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Yes	"Allocation was concealed", "randomisation could not be subverted by the study team"
Adequate sequence generation?	Yes	Computer-generated randomisation sequence
Blinding?	Unclear	Unclear
Free of selective outcome reporting?	Yes	All outcomes reported

**Khadjesari 2011** (Continued)

Other sources of bias?	Unclear	Insufficient information to make a judgement about this
------------------------	---------	---

**Khadjesari 2011 1a**

Methods	4-arm randomised trial (offer of incentives vs. no offer)
Data	Non-responders at 1 week to the pilot DYD trial 3-month online follow-up questionnaire. Adults aged $\geq 18$ scoring 5 or more on the AUDIT-C test
Comparisons	1. Offer of GBP5 Amazon.co.uk voucher. Emailed voucher code on receipt of response (arm a) 2. Email with no offer of incentive (control)
Outcomes	Proportion completing questionnaire after 40 days
Notes	Primary outcome for DYD: total past week alcohol consumption measured using the TOTAL measure of total alcohol consumption. Trial identified through personal communication

**Risk of bias**

Item	Authors' judgement	Description
Allocation concealment?	Yes	"Allocation was concealed randomisation could not be subverted by the study team"
Adequate sequence generation?	Yes	Computer-generated randomisation sequence
Blinding?	Unclear	Unclear
Free of selective outcome reporting?	Yes	Expected outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**Khadjesari 2011 1abc**

Methods	4-arm randomised trial
Data	Non-responders at 1 week to the pilot DYD trial 3-month online follow-up questionnaire. Adults aged $\geq 18$ scoring 5 or more on the AUDIT-C test
Comparisons	1. Offer of GBP5 Amazon.co.uk voucher. Emailed voucher code on receipt of response (arm a) 2. Offer of GBP5 donation to Cancer Research UK. Emailed hyperlink to charity's website showing donation when response received (arm b) 3. Offer of entry into GBP250 prize draw emailed confirmation of entry when response

**Khadjesari 2011 1abc** (Continued)

	received (arm c) 4. Email prompt for completion of questionnaires with no offer of incentive (control arm)
Outcomes	Proportion completing questionnaire after 40 days
Notes	Primary outcome for DYD: total past week alcohol consumption measured using the TOTAL measure of total alcohol consumption. Trial identified through personal communication
<b>Risk of bias</b>	
<b>Item</b>	<b>Authors' judgement</b> <b>Description</b>
Allocation concealment?	Yes      "Allocation was concealed randomisation could not be subverted by the study team"
Adequate sequence generation?	Yes      Computer-generated randomisation sequence
Blinding?	Unclear      Unclear
Free of selective outcome reporting?	Yes      Expected outcomes reported
Other sources of bias?	Unclear      Insufficient information to make a judgement about this

**Khadjesari 2011 1ac**

Methods	4-arm randomised trial (incentives combined vs. no incentive)
Data	Non-responders at 1 week to the pilot DYD trial 3-month online follow-up questionnaire. Adults aged ≥ 18 scoring 5 or more on the AUDIT-C test
Comparisons	1. Offer of GBP5 Amazon.co.uk voucher. Emailed voucher code on receipt of response (arm a) 2. Offer of entry into GBP250 prize draw emailed confirmation of entry when response received (arm c) 3. Email with no offer of incentive (control arm)
Outcomes	Proportion completing questionnaire after 40 days
Notes	Primary outcome for DYD: total past week alcohol consumption measured using the TOTAL measure of total alcohol consumption. Trial identified through personal communication Separate comparison for arm b of attrition trial see <a href="#">Khadjesari 2011 1b</a>
<b>Risk of bias</b>	
<b>Item</b>	<b>Authors' judgement</b> <b>Description</b>

**Khadjesari 2011 1ac** (Continued)

Allocation concealment?	Yes	“Allocation was concealed randomisation could not be subverted by the study team”
Adequate sequence generation?	Yes	Computer-generated randomisation sequence
Blinding?	Unclear	Unclear
Free of selective outcome reporting?	Yes	Expected outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**Khadjesari 2011 1b**

Methods	4-arm randomised trial (offer of donation to charity vs. no offer)
Data	Non-responders at 1 week to the pilot DYD trial 3-month online follow-up questionnaire. Adults aged $\geq 18$ scoring 5 or more on the AUDIT-C test
Comparisons	1. Offer of GBP5 donation to Cancer Research UK. Emailed hyperlink to charity’s website showing donation when response received (arm b) 2. Email with no offer of incentive (control)
Outcomes	Proportion completing questionnaire after 40 days
Notes	Primary outcome for DYD: total past week alcohol consumption measured using the TOTAL measure of total alcohol consumption. Trial identified through personal communication

***Risk of bias***

<b>Item</b>	<b>Authors’ judgement</b>	<b>Description</b>
Allocation concealment?	Yes	“Allocation was concealed randomisation could not be subverted by the study team”
Adequate sequence generation?	Yes	Computer-generated randomisation sequence
Blinding?	Unclear	Unclear
Free of selective outcome reporting?	Yes	Expected outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**Khadjesari 2011 1c**

Methods	4-arm randomised trial (offer of entry into prize draw vs. no offer)
Data	Non-responders at 1 week to the pilot DYD trial 3-month online follow-up questionnaire. Adults aged $\geq 18$ scoring 5 or more on the AUDIT-C test
Comparisons	1. Offer of entry into GBP250 prize draw emailed confirmation of entry when response received (arm c) 2. Email with no offer of incentive (control arm)
Outcomes	Proportion completing questionnaire after 40 days
Notes	Primary outcome for DYD: total past week alcohol consumption measured using the TOTAL measure of total alcohol consumption. Trial identified through personal communication

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Yes	"Allocation was concealed randomisation could not be subverted by the study team"
Adequate sequence generation?	Yes	Computer-generated randomisation sequence
Blinding?	Unclear	Unclear
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**Khadjesari 2011 2**

Methods	2-arm randomised trial
Data	All DYD trial study participants
Comparisons	1. Offer of a GBP10 Amazon.co.uk voucher. Email with voucher code sent on completion of questionnaire 2. Reminder email with no voucher offer
Outcomes	The proportion of participants that completed the questionnaire after 40 days
Notes	Primary outcome for DYD: total past week alcohol consumption measured using the TOTAL measure of total alcohol consumption. Trial identified through personal communication

***Risk of bias***



Item	Authors' judgement	Description
Allocation concealment?	Yes	"Allocation was concealed", "randomisation could not be subverted by the study team"
Adequate sequence generation?	Yes	Computer-generated randomisation sequence
Blinding?	Unclear	Unclear
Free of selective outcome reporting?	Yes	All outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**Land 2007**

Methods	2-arm randomised trial
Data	Research sites participating in the NSABP B35 trial
Comparisons	1. Automated prospective monthly reminder to trial sites of upcoming participant-reported outcome assessments. Reminder listed participants expected to complete Behavioral and Health Outcomes forms in upcoming 3 months 2. No monthly assessment reminder
Outcomes	Receipt of questionnaire at any time
Notes	B35 Anastrozole vs. tamoxifen for the treatment of ductal carcinoma in situ women aged 55 years plus. Primary outcome: time to first breast cancer reoccurrence

**Risk of bias**

Item	Authors' judgement	Description
Allocation concealment?	Unclear	Author response to question "describe the methods if any, used to conceal the allocation sequence for the prospective reminder study in B35", "not applicable. Institutions were all randomly assigned before trial initiation, so there was no sequence"
Adequate sequence generation?	Yes	Author response "prospective reminder in B-35, three strata were generated, and half in each were randomly assigned to receive the reminder"
Blinding?	No	Author response "there was no blinding the prospective reminder was received by clinical staff"
Free of selective outcome reporting?	Yes	Primary outcome reported

**Land 2007** (Continued)

Other sources of bias?	Unclear	Insufficient information to make a judgement
------------------------	---------	--

**Leigh Brown 1997**

Methods	2-arm randomised trial
Data	Participants in OMENS due to receive follow-up questionnaires between March and December 1995
Comparisons	1. Offer of GBP25 gift voucher monthly prize draw. Postcard reminder after 10 days and 3 weeks with reference to offer 2. No offer. Post card reminders after 10 days and 3 weeks with no reference to offer
Outcomes	Questionnaire response. No time point given
Notes	OMENS effectiveness of musculoskeletal medicine vs. care by orthopaedic surgeon-led services for the treatment of non-surgical orthopaedic outpatients Primary outcomes: change in participant reported health (SF-36 and EuroQol) and marginal health

**Risk of bias**

Item	Authors' judgement	Description
Allocation concealment?	No	Author response "table of random numbers"
Adequate sequence generation?	Yes	Author response "table of random numbers"
Blinding?	Unclear	Author response "those randomised to take part in the prize draw were aware of the intervention and those randomised to be excluded from the draw were unaware. The trial team were aware of the allocation so that they could arrange the monthly draw"
Free of selective outcome reporting?	Yes	Defined outcomes reported outcome reported
Other sources of bias?	Unclear	Insufficient evidence to make a judgement about this

**Letley 2000**

Methods	2-arm randomised trial
Data	Participants in the UKBEAM feasibility trial, aged 18-65 years with low back pain and a score of 4 or more on the RDQ
Comparisons	Review author (VB) checked which was control and which was intervention 1. RDQ before SF-36 in a 26-page questionnaire 2. SF-36 before RDQ in a 26-page questionnaire

**Letley 2000** (Continued)

Outcomes	Questionnaire response at 3 months
Notes	Study complete, no data available at 8 September 2011. UKBEAM: compared the effectiveness of exercise, manipulation, exercise and manipulation. Primary outcome for UKBEAM scores on the RDQ at 3 and 12 months

**Risk of bias**

Item	Authors' judgement	Description
Allocation concealment?	Yes	Author reported "sequence generation was via remote service ensuring allocation concealment"
Adequate sequence generation?	Yes	Author reported "sequence generation was random using randomised permuted blocks"
Blinding?	Unclear	No data
Free of selective outcome reporting?	Yes	Overall response rates reported
Other sources of bias?	Unclear	No data

**MacLennan**

Methods	2-arm randomised trial
Data	Participants who had not returned questionnaires at 12, 24 and 36 months in the RECORD trial
Comparisons	1. Telephone call from RECORD office before first reminder questionnaire sent. Participant asked to complete questionnaire and to try to fill in all questions. Further telephone call from study nurse or RECORD office if not returned after 3 weeks 2. Repeat mailing of usual follow-up letter and questionnaire. Telephone call by study nurse or RECORD office if not returned after 3 weeks (routine follow-up)
Outcomes	Proportion of first reminder questionnaires returned Proportion of questionnaires returned at 4 months Completeness of data
Notes	Primary outcome all new low-energy fractures (self reported) EQ-5D Short form-12. Trial identified through mail out to UK clinical trial units

**Risk of bias**

Item	Authors' judgement	Description
Allocation concealment?	Yes	Author response "central randomisation service"

Adequate sequence generation?	Yes	Author response “randomised to receive the ”intervention“ or not using the central randomisation service”
Blinding?	Yes	Author response “blinding was not possible in the ”intervention“ arm. The control were blinded. Trial staff phoning were not blinded Outcome assessment in the sub-study was objective: questionnaire returned yes/no. Trial staff and trial participants were blinded to the RECORD trial allocation”
Free of selective outcome reporting?	Yes	All defined outcomes reported

**Man 2011**

Methods	2-arm randomised trial
Data	Adults 18-65 years sent a 6-month follow-up questionnaire in a trial of yoga for chronic back pain
Comparisons	<ol style="list-style-type: none"> <li>1. SMS text message: “Yoga trial: You should receive a questionnaire any day now. The data is important to us so please return it as soon as you can. Many thanks”</li> <li>2. Email message: “Thank you for taking part in the Yoga for Low Back Pain Trial. This is an automatic reminder. You should receive your six month questionnaire any day now. The answers you give in the questionnaire are very important to the trial. Therefore, we should be most grateful if you would complete and return the questionnaire (and any other documents) as soon as possible please. Thank you”</li> <li>3. SMS text message plus email message</li> <li>4. No SMS test message or email message</li> </ol>
Outcomes	Postal questionnaire response
Notes	Yoga for chronic back pain trial: primary outcome functional limitations and disability measured by the RDQ. Setting 13 non-national health service settings in the UK. Participants were recruited through general practices

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Yes	Computer-generated sequence to randomly allocate participants
Adequate sequence generation?	Yes	Independent data manager generated a computer sequence
Blinding?	Unclear	No data
Free of selective outcome reporting?	Yes	Reported response to questionnaires
Other sources of bias?	Unclear	No data

**Marson 2007**

Methods	2-arm randomised trial
Data	Participants in SANAD due a QoL questionnaire at 1 year
Comparisons	1. Cover letter with estimate of the length of time required to complete questionnaire 2. Standard cover letter without estimate of the length of time required to complete questionnaire
Outcomes	Response rate. No time point given
Notes	SANAD compared the long-term effects of antiepileptic drugs in people with epilepsy. Participants randomised to arm a received carbamazepine, gabapentin, lamotrigine, oxcarbazepine, or topiramate and valproate, lamotrigine, or arm B received topiramate. Primary outcome time to treatment failure and time to achieve a 12-month remission of seizures

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	No	Author response "the list was given to the study researcher who worked sequentially through the list from left to right from top to bottom"
Adequate sequence generation?	Yes	Author response "list of allocations was generated electronically using random permuted blocks of length 20"
Blinding?	Yes	Author response "this was a postal study, blinding was not necessary for participants; they received either a time-framed letter or non time-framed letter. Clinical personnel did not see the letters, as these were sent postally directly to participants from the study research office"
Free of selective outcome reporting?	Yes	Attrition reported
Other sources of bias?	Unclear	Insufficient evidence to make a judgement about this

**McCambridge 2011**

Methods	McCambridge 1: 4-arm randomised trial in the pilot phase of the DYD trial; McCambridge 2: 4-arm randomised trial in the host DYD trial
Data	See 'Table of characteristics' for <a href="#">McCambridge 2011 1</a> ; <a href="#">McCambridge 2011 1a</a> ; <a href="#">McCambridge 2011 1b</a> ; <a href="#">McCambridge 2011 2</a> ; <a href="#">McCambridge 2011 2a</a> ; <a href="#">McCambridge 2011 2b</a>
Comparisons	See 'Table of characteristics' for <a href="#">McCambridge 2011 1</a> ; <a href="#">McCambridge 2011 1a</a> ; <a href="#">McCambridge 2011 1b</a> ; <a href="#">McCambridge 2011 2</a> ; <a href="#">McCambridge 2011 2a</a> ; <a href="#">McCambridge 2011 2b</a>

**McCambridge 2011** (Continued)

Outcomes	See 'Table of characteristics' for <a href="#">McCambridge 2011 1</a> ; <a href="#">McCambridge 2011 1a</a> ; <a href="#">McCambridge 2011 1b</a> ; <a href="#">McCambridge 2011 2</a> ; <a href="#">McCambridge 2011 2a</a> ; <a href="#">McCambridge 2011 2b</a>	
Notes	See 'Table of characteristics' for <a href="#">McCambridge 2011 1</a> ; <a href="#">McCambridge 2011 1a</a> ; <a href="#">McCambridge 2011 1b</a> ; <a href="#">McCambridge 2011 2</a> ; <a href="#">McCambridge 2011 2a</a> ; <a href="#">McCambridge 2011 2b</a>	
<b><i>Risk of bias</i></b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Yes	See 'Risk of bias' table <a href="#">McCambridge 2011 1</a> ; <a href="#">McCambridge 2011 1a</a> ; <a href="#">McCambridge 2011 1b</a> ; <a href="#">McCambridge 2011 2</a> ; <a href="#">McCambridge 2011 2a</a> ; <a href="#">McCambridge 2011 2b</a>
Adequate sequence generation?	Yes	See 'Risk of bias' table <a href="#">McCambridge 2011 1</a> ; <a href="#">McCambridge 2011 1a</a> ; <a href="#">McCambridge 2011 1b</a> ; <a href="#">McCambridge 2011 2</a> ; <a href="#">McCambridge 2011 2a</a> ; <a href="#">McCambridge 2011 2b</a>
Blinding?	Unclear	See 'Risk of bias' table <a href="#">McCambridge 2011 1</a> ; <a href="#">McCambridge 2011 1a</a> ; <a href="#">McCambridge 2011 1b</a> ; <a href="#">McCambridge 2011 2</a> ; <a href="#">McCambridge 2011 2a</a> ; <a href="#">McCambridge 2011 2b</a>
Free of selective outcome reporting?	Yes	See 'Risk of bias' table <a href="#">McCambridge 2011 1</a> ; <a href="#">McCambridge 2011 1a</a> ; <a href="#">McCambridge 2011 1b</a> ; <a href="#">McCambridge 2011 2</a> ; <a href="#">McCambridge 2011 2a</a> ; <a href="#">McCambridge 2011 2b</a>
Other sources of bias?	Unclear	See 'Risk of bias' table <a href="#">McCambridge 2011 1</a> ; <a href="#">McCambridge 2011 1a</a> ; <a href="#">McCambridge 2011 1b</a> ; <a href="#">McCambridge 2011 2</a> ; <a href="#">McCambridge 2011 2a</a> ; <a href="#">McCambridge 2011 2b</a>

**McCambridge 2011 1**

Methods	4-arm randomised trial in the pilot phase of the DYD trial
Data	Adults aged $\geq 18$ years scoring $\geq 5$ on the AUDIT-C test participating in the pilot phase of the DYD trial 1-month data used
Comparisons	<ol style="list-style-type: none"> <li>1. APQ 23 items</li> <li>2. AUDIT 10</li> <li>3. LDQ 10</li> <li>4. CORE-OM 23/34 items (mental health assessment)</li> </ol>

Outcomes	Response to electronic questionnaires	
Notes	Primary outcome for DYD: total past week alcohol consumption measured using the TOTAL measure of total alcohol consumption. Trial identified through personal communication	
<b>Risk of bias</b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Yes	"Allocation was concealed randomisation could not be subverted by the study team"
Adequate sequence generation?	Yes	Computer-generated randomisation sequence
Blinding?	Unclear	Unclear
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**McCambridge 2011 1a**

Methods	4-arm randomised trial ('relevance' of questionnaire: alcohol vs. mental health data)	
Data	Adults aged $\geq 18$ years scoring 5 or more on the AUDIT-C test participating in the pilot phase of the DYD trial 1-month data used	
Comparisons	1. APQ 23 items 2. CORE-OM 23/34 items (Mental health assessment)	
Outcomes	Response to electronic questionnaires	
Notes	Primary outcome for DYD: total past week alcohol consumption measured using the TOTAL measure of total alcohol consumption. Trial identified through personal communication	
<b>Risk of bias</b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Yes	"Allocation was concealed randomisation could not be subverted by the study team"
Adequate sequence generation?	Yes	Computer-generated randomisation sequence
Blinding?	Unclear	Unclear

**McCambridge 2011 1a** (Continued)

Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**McCambridge 2011 1b**

Methods	4-arm randomised trial (short vs. long questionnaire comparison)
Data	Adults aged $\geq 18$ years scoring 5 or more on the AUDIT-C test participating in the DYD trial. For this comparison 1-month follow-up data from <a href="#">McCambridge 2011 1</a> used
Comparisons	1. AUDIT 10 items + LDQ 10 items 2. APQ 23 items
Outcomes	Response to electronic questionnaires
Notes	Primary outcome for DYD: total past week alcohol consumption measured using the TOTAL measure of total alcohol consumption. Trial identified through personal communication

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Yes	"Allocation was concealed randomisation could not be subverted by the study team"
Adequate sequence generation?	Yes	Computer-generated randomisation sequence
Blinding?	Unclear	Unclear
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**McCambridge 2011 2**

Methods	4-arm randomised trial in the host DYD trial
Data	Adults aged $\geq 18$ years scoring 5 or more on the AUDIT-C test participating in the pilot phase of the DYD trial 3-month data used
Comparisons	1. APQ 23 items 2. AUDIT 10 3. LDQ 10



	4. CORE-OM 23/34 items (Mental health assessment)	
Outcomes	Response to electronic questionnaires	
Notes	Primary outcome for DYD: total past week alcohol consumption measured using the TOTAL measure of total alcohol consumption. Trial identified through personal communication	
<b><i>Risk of bias</i></b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Yes	"Allocation was concealed randomisation could not be subverted by the study team"
Adequate sequence generation?	Yes	Computer-generated randomisation sequence
Blinding?	Unclear	Unclear
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**McCambridge 2011 2a**

Methods	4-arm randomised trial ('relevance' of questionnaire alcohol vs. mental health)	
Data	Adults aged $\geq 18$ years scoring 5 or more on the AUDIT-C test participating in the DYD trial. For this comparison 3-month follow-up data from <a href="#">McCambridge 2011 2</a> used	
Comparisons	1. AUDIT 10 items + LDQ 10 items 2. CORE-10	
Outcomes	Response to electronic questionnaires	
Notes	Primary outcome for DYD: total past week alcohol consumption measured using the TOTAL measure of total alcohol consumption. Trial identified through personal communication	
<b><i>Risk of bias</i></b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Yes	"Allocation was concealed randomisation could not be subverted by the study team"
Adequate sequence generation?	Yes	Computer-generated randomisation sequence

**McCambridge 2011 2a** (Continued)

Blinding?	Unclear	Unclear
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**McCambridge 2011 2b**

Methods	4-arm randomised trial (short vs. long questionnaire)
Data	Adults aged $\geq 18$ years scoring 5 or more on the AUDIT-C test participating in the DYD trial. For this comparison, 3-month follow-up data from <a href="#">McCambridge 2011 2</a> used
Comparisons	1. AUDIT 10 items + LDQ 10 items 2. APQ 23 items
Outcomes	Response to electronic questionnaires
Notes	Primary outcome for DYD: total past week alcohol consumption measured using the TOTAL measure of total alcohol consumption. Trial identified through personal communication

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Yes	"Allocation was concealed randomisation could not be subverted by the study team"
Adequate sequence generation?	Yes	Computer-generated randomisation sequence
Blinding?	Unclear	Unclear
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**McColl 2003**

Methods	2-arm randomised trial
Data	40 adults aged $\geq 18$ years with asthma from 62 general practices participating in the COGENT trial
Comparisons	See 'Table of characteristics' for <a href="#">McColl 2003 1</a> ; <a href="#">McColl 2003 2</a>

**McColl 2003** (Continued)

Outcomes	See 'Table of characteristics' for <a href="#">McColl 2003 1</a> ; <a href="#">McColl 2003 2</a>	
Notes	See 'Table of characteristics' for <a href="#">McColl 2003 1</a> ; <a href="#">McColl 2003 2</a>	
<b><i>Risk of bias</i></b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	No	See 'Risk of bias' table <a href="#">McColl 2003 1</a> ; <a href="#">McColl 2003 2</a>
Adequate sequence generation?	Unclear	See 'Risk of bias' table <a href="#">McColl 2003 1</a> ; <a href="#">McColl 2003 2</a>
Blinding?	Yes	See 'Risk of bias' table <a href="#">McColl 2003 1</a> ; <a href="#">McColl 2003 2</a>
Free of selective outcome reporting?	Yes	See 'Risk of bias' table <a href="#">McColl 2003 1</a> ; <a href="#">McColl 2003 2</a>
Other sources of bias?	Unclear	See 'Risk of bias' table <a href="#">McColl 2003 1</a> ; <a href="#">McColl 2003 2</a>

**McColl 2003 1**

Methods	2-arm randomised trial	
Data	40 adults aged $\geq 18$ years with asthma from 62 general practices participating in the COGENT trial	
Comparisons	1. Asthma specific instruments (NASQ and AQLQ) followed by generic questions in medical outcomes SF-36 version 1 and EQ-5D 2. Generic questions followed by condition specific	
Outcomes	Questionnaire response rates, speed and patterns. No time point given	
Notes	COGENT before and after cluster randomised, 2 x 2 factorial trial. Evaluated use of computerised support system for decision making for implementing evidenced-based clinical guidelines for the management of asthma (and angina see <a href="#">McColl 2003 2</a> ). General practices randomised to computerised guidelines for the management of angina or asthma. Primary outcome adherence to guidelines	
<b><i>Risk of bias</i></b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	No	Author response "practices provided us with a computerised list of all patients fulfilling eligibility criteria. We drew a simple random sample of patients from that list first 40 from list received version 1, second 40 version 2"

**McColl 2003 1** (Continued)

Adequate sequence generation?	Unclear	Author response “practices provided us with a computerised list of all patients fulfilling eligibility criteria. We drew a simple random sample of patients from that list. First 40 received version 1, second 40 version 2. Logistically, truly random allocation of the 80 patients to versions 1 and 2 would have been impractical to implement”
Blinding?	Yes	Author response “respondents were not alerted to the fact that we were experimenting with the order of instruments”
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient evidence to make a judgement about this

**McColl 2003 2**

Methods	2-arm randomised trial
Data	40 adults aged $\geq 18$ years with asthma from 62 general practices participating in the COGENT trial
Comparisons	1. Angina-specific instruments (SAQ) followed by generic questions (SF-36 version 1 and EQ-5D) 2. Generic questions followed by condition-specific questions
Outcomes	Questionnaire response rates, speed and patterns. No time point given
Notes	COGENT before and after cluster randomised, 2 x 2 factorial trial. Evaluated use of computerised support system for decision making for implementing evidenced-based clinical guidelines for the management of angina (and asthma see <a href="#">McColl 2003 1</a> ). General practices randomised to computerised guidelines for the management of angina or asthma. Primary outcome adherence to guidelines

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	No	Author response “practices provided us with a computerised list of all patients fulfilling eligibility criteria. We drew a simple random sample of patients from that list. First 40 from list received version 1, second 40 version 2”
Adequate sequence generation?	Unclear	Author response “practices provided us with a computerised list of all patients fulfilling eligibility criteria. We drew a simple random sample of patients from that list. First 40 received version 1, second 40 version 2. Logistically, truly random allocation of the 80 patients to versions 1 and 2 would have been impractical to implement”

**McColl 2003 2** (Continued)

Blinding?	Yes	Author response “respondents were not alerted to the fact that we were experimenting with the order of instruments”
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient evidence to make a judgement about this

**Nakash 2007**

Methods	2-arm randomised trial
Data	Participants aged $\geq 16$ years from 5 trial centres randomised to CAST between November 2003 and July 2005
Comparisons	1. Trial calendar: monthly customised calendar, included prenotification caption on the months the participant is due to receive the questionnaire with reminder caption the following month 2. No trial calendar
Outcomes	Questionnaire response at 4, 12 weeks and 9 months. Response at 4 weeks used for analysis Amount of prompting required to return questionnaires at each time point Percentage of missing data of the core outcome
Notes	CAST tubular bandage, below knee cast, Aircast® ankle brace and Bledsoe® boot compared in people with acute severe ankle sprain. Primary outcome FAOS, FLP, SF-12 and EuroQol 5

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Yes	“Baseline packs compiled in advance and stored at trial sites. On randomisation the next consecutively numbered pack was taken. Allocation concealed until participant recruited into CAST and pack opened”
Adequate sequence generation?	Yes	“Computer generated random sequence”
Blinding?	Yes	“No blinding of participants or clinic staff”, “data inputting was blind to allocation”
Free of selective outcome reporting?	Yes	All defined outcomes reported
Other sources of bias?	Unclear	Insufficient evidence to make a judgement about this

**Renfroe 2002a**

Methods	2 x 2 x 2 x 2 factorial randomised trial (certificate of appreciation vs. no certificate)
Data	All surviving participants at AVID study termination
Comparisons	1. Certificate of appreciation with cover letter signed by physician or co-ordinator sent either by express or standard post, 2-3 weeks after last AVID follow-up visit or 1-4 months after last AVID follow-up visit 2. No certificate with cover letter signed by physician or co-ordinator, sent either by express or standard post, 2-3 weeks after last AVID follow-up visit or 1-4 months after last AVID follow-up visit
Outcomes	Questionnaire response. No time point given
Notes	AVID conducted in people resuscitated from ventricular fibrillation or cardioverted for ventricular tachycardia. Compared implanted cardioverter defibrillator vs. antiarrhythmic drugs. Primary outcome overall mortality

***Risk of bias***

<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Unclear	The AVID CTC prepared the resulting 16 sets of distinct participant survey packets. These participant-specific packets were then mailed to the study co-ordinators, who distributed them to participants as instructed
Adequate sequence generation?	Yes	Author response "all patients in the study were randomised. Details of the randomisation scheme are no longer extant, but given the factorial design I think it's safe to assume that the randomization for each factor was by permuted blocks of size 2"
Blinding?	Yes	Participants were instructed to mail the completed surveys directly to the CTC in postage-paid, self-addressed return envelopes. Only their AVID study number identified participants, assuring confidentiality of their survey responses
Free of selective outcome reporting?	Yes	All defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Renfro 2002a-d**

Methods	2 x 2 x 2 x 2 factorial randomised trial
Data	All surviving participants at AVID study termination
Comparisons	<ol style="list-style-type: none"> <li>1. Certificate of appreciation</li> <li>2. No certificate of appreciation</li> <li>3. Overnight express delivery</li> <li>4. Regular post</li> <li>5. Cover letter signed by physician</li> <li>6. Cover letter signed by study co-ordinator</li> <li>7. Early administration</li> <li>8. Late administration</li> </ol>
Outcomes	Questionnaire response. No time point given
Notes	AVID conducted in people resuscitated from ventricular fibrillation or cardioverted for ventricular tachycardia. Compared implanted cardioverter defibrillator vs. antiarrhythmic drugs. Primary outcome overall mortality

***Risk of bias***

<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Unclear	The AVID CTC prepared the resulting 16 sets of distinct participant survey packets. These participant-specific packets were then mailed to the study co-ordinators, who distributed them to participants as instructed
Adequate sequence generation?	Yes	Author response "all patients in the study were randomised. Details of the randomisation scheme are no longer extant, but given the factorial design I think it's safe to assume that the randomization for each factor was by permuted blocks of size 2"
Blinding?	Yes	Participants were instructed to mail the completed surveys directly to the CTC in postage-paid, self-addressed return envelopes. Only their AVID study number identified participants, assuring confidentiality of their survey responses
Free of selective outcome reporting?	Yes	All defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Renfroe 2002b**

Methods	2 x 2 x 2 x 2 factorial randomised trial (express vs. regular post)
Data	All surviving participants at AVID study termination
Comparisons	1. Overnight express delivery with cover letter signed by physician or co-ordinator with or without a certificate of appreciation 2-3 weeks after last AVID follow-up visit or 1-4 months after last AVID follow-up visit 2. Regular post with cover letter signed by physician or co-ordinator with or without a certificate of appreciation 2-3 weeks after last AVID follow-up visit or 1-4 months after last AVID follow-up visit
Outcomes	Questionnaire response. No time point given
Notes	AVID conducted in people resuscitated from ventricular fibrillation or cardioverted for ventricular tachycardia. Compared implanted cardioverter defibrillator with antiarrhythmic drugs. Primary outcome overall mortality

***Risk of bias***

<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Unclear	The AVID CTC prepared the resulting 16 sets of distinct participant survey packets. These participant-specific packets were then mailed to the study co-ordinators, who distributed them to participants as instructed
Adequate sequence generation?	Yes	Author response "all patients in the study were randomised. Details of the randomisation scheme are no longer extant, but given the factorial design I think it's safe to assume that the randomisation for each factor was by permuted blocks of size 2"
Blinding?	Yes	Participants were instructed to mail the completed surveys directly to the CTC in postage-paid, self-addressed return envelopes. Only their AVID study number identified participants, assuring confidentiality of their survey responses
Free of selective outcome reporting?	Yes	All defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this



**Renfroe 2002c**

Methods	2 x 2 x 2 x 2 factorial randomised trial (physician vs. study co-ordinator signed cover letter)
Data	All surviving participants at AVID study termination
Comparisons	1. Cover letter signed by physician sent either by express or standard post, with or without a certificate of appreciation, 2-3 weeks after last AVID follow-up visit or 1-4 months after last AVID follow-up visit 2. Cover letter signed by study co-ordinator sent either by express or standard post, with or without a certificate of appreciation 2-3 weeks after last AVID follow-up visit or 1-4 months after last AVID follow-up visit
Outcomes	Questionnaire response. No time point given
Notes	AVID conducted in participants resuscitated from ventricular fibrillation or cardioverted for ventricular tachycardia. Compared implanted cardioverter defibrillator vs. antiarrhythmic drugs. Primary outcome overall mortality

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Unclear	The AVID CTC prepared the resulting 16 sets of distinct participant survey packets. These participant-specific packets were then mailed to the study co-ordinators, who distributed them to participants as instructed
Adequate sequence generation?	Yes	Author response "all patients in the study were randomised. Details of the randomisation scheme are no longer extant, but given the factorial design I think it's safe to assume that the randomisation for each factor was by permuted blocks of size 2"
Blinding?	Yes	Participants were instructed to mail the completed surveys directly to the CTC in postage-paid, self-addressed return envelopes. Only their AVID study number identified participants, assuring confidentiality of their survey responses
Free of selective outcome reporting?	Yes	Report all outcomes
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Renfroe 2002d**

Methods	2 x 2 x 2 x 2 factorial randomised trial (early vs. late administration of questionnaire)
Data	All surviving participants at AVID study termination
Comparisons	1. Questionnaire sent 2-3 weeks after last AVID follow-up visit by express or standard post with cover letter signed by physician or co-ordinator with or without a certificate of appreciation 2. Questionnaire sent 1-4 months after last AVID follow-up visit, by express or standard post with cover letter signed by physician or co-ordinator with or without a certificate of appreciation
Outcomes	Questionnaire response. No time point given
Notes	AVID conducted in people resuscitated from ventricular fibrillation or cardioverted for ventricular tachycardia. Compared implanted cardioverter defibrillator vs. antiarrhythmic drugs. Primary outcome overall mortality

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Unclear	The AVID CTC prepared the resulting 16 sets of distinct participant survey packets. These participant-specific packets were then mailed to the study co-ordinators, who distributed them to participants as instructed
Adequate sequence generation?	Yes	Author response "all patients in the study were randomised. Details of the randomisation scheme are no longer extant, but given the factorial design I think it's safe to assume that the randomisation for each factor was by permuted blocks of size 2"
Blinding?	Yes	Participants were instructed to mail the completed surveys directly to the CTC in postage-paid, self-addressed return envelopes. Only their AVID study number identified participants, assuring confidentiality of their survey responses
Free of selective outcome reporting?	Yes	All defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Severi 2011**

Methods	Two 2-arm randomised trials
Data	UK smokers aged $\geq 16$ years participating in the Txt2stop trial
Comparisons	See 'Table of characteristics' for <a href="#">Severi 2011 1</a> and <a href="#">Severi 2011 2</a>
Outcomes	See 'Table of characteristics' for <a href="#">Severi 2011 1</a> and <a href="#">Severi 2011 2</a>
Notes	See 'Table of characteristics' for <a href="#">Severi 2011 1</a> and <a href="#">Severi 2011 2</a>

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Unclear	"Concealed from the investigators"
Adequate sequence generation?	Yes	"Allocated through minimisation using minim software"
Blinding?	Yes	"Single blind controlled trial, with those recording and assessing outcomes blind to the intervention"
Free of selective outcome reporting?	Yes	All defined outcomes reported

**Severi 2011 1**

Methods	2-arm randomised trial
Data	UK smokers aged $\geq 16$ participating in the Txt2stop trial
Comparisons	<p>1. Text message "Be proud of yourself for helping medical research! Thank you for filling in the txt 2 stop questionnaire" plus a fridge magnet: the message on the fridge magnet was placed in a sealed envelope, this said: "medical research is important to society" and pointed out that by taking part in Txt2stop the participants are benefiting society. Fridge magnet sent by post 16-20 weeks after randomisation. Text message sent 3 days after Txt2stop postal follow-up questionnaire</p> <p>2. Text message reminding the participant the follow-up questionnaire was due 3 days after the Txt2stop questionnaire had been sent. The text message said "Thank you for filling in the Txt2stop questionnaire". Sent 3 days after the text to stop postal follow-up questionnaire</p>
Outcomes	<p>Primary outcome: completed follow-up questionnaires at 30 weeks from randomisation</p> <p>Secondary outcome: completed follow-up questionnaires at 26 weeks from randomisation</p>
Notes	Txt2stop UK-based smoking cessation trial evaluating the effectiveness of the Txt2stop mobile phone text messaging smoking cessation programme on biochemically verified continuous smoking abstinence at 6 months. Compared Txt2stop motivational messages vs. behaviour change support to text messages unrelated to quitting

***Risk of bias***

**Severi 2011 1** (Continued)

Item	Authors' judgement	Description
Allocation concealment?	Unclear	"Concealed from the investigators"
Adequate sequence generation?	Yes	"Allocated through minimisation using minim software"
Blinding?	Yes	"Single blind controlled trial, with those recording and assessing outcomes blind to the intervention"
Free of selective outcome reporting?	Yes	All defined outcomes reported

**Severi 2011 2**

Methods	2-arm randomised trial
Data	UK smokers aged $\geq 16$ years participating in the Txt2stop trial
Comparisons	1. Telephone call from female principal investigator (senior clinician and researcher) to participants 6 weeks overdue returning specimen to increase participant follow-up plus standard Txt2stop follow-up procedures 2. Standard Txt2stop follow-up procedures
Outcomes	Completed cotinine sample follow-up at the end of May 2009 for Txt2stop (1 month after a telephone call)
Notes	Txt2stop UK-based smoking cessation trial evaluating the effectiveness of the Txt2stop mobile phone text messaging smoking cessation program on biochemically verified continuous smoking abstinence at 6 months. Compared Txt2stop motivational messages vs. behaviour change support to text messages unrelated to quitting

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Unclear	"Concealed from the investigators"
Adequate sequence generation?	Yes	"Allocated through minimisation using minim software"
Blinding?	Yes	"Single blind controlled trial, with those recording and assessing outcomes blind to the intervention"
Free of selective outcome reporting?	Yes	All defined outcomes reported

**Sharp 2006a**

Methods	2 x 2 x 2 factorial randomised trial (trial-branded pen vs. no pen: comparison 1)
Data	Women due to receive a TOMBOLA psychosocial questionnaire during June-August 2003 for 12, 18, 24, 30, 34, 36 months' follow-up
Comparisons	1. TOMBOLA-branded pen + despatch first-class post + second-class stamped preaddressed return envelope 2. No TOMBOLA-branded pen + despatch first-class post + second-class stamped preaddressed return envelope
Outcomes	Response rates at any time
Notes	TOMBOLA compared 2 management policies (colposcopy vs. 6-monthly smears) in women aged 20-59 years with low-grade abnormal cervical smear. Primary outcome: cumulative incidence of cervical intraepithelial neoplasia grade $\geq 2$

***Risk of bias***

<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Unclear	2 authors did randomisation
Adequate sequence generation?	Unclear	"Computer randomised"
Blinding?	Unclear	Outcome not influenced by lack of blinding
Free of selective outcome reporting?	Yes	Expected outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Sharp 2006a-h**

Methods	2 x 2 x 2 factorial randomised trial
Data	Women due to receive a TOMBOLA psychosocial questionnaire during June-August 2003 for 12, 18, 24, 30, 34, 36 months' follow-up
Comparisons	1. TOMBOLA-branded pen + despatch first-class post + second-class stamped preaddressed return envelope 2. No TOMBOLA-branded pen + despatch first-class post + second-class stamped preaddressed return envelope 3. TOMBOLA-branded pen + despatch first-class post + free post business-reply envelope 4. No TOMBOLA-branded pen + despatch first-class post + free post business-reply envelope 5. TOMBOLA-branded pen + despatch second-class post + second-class stamped preaddressed return envelope 6. No TOMBOLA-branded pen + despatch second-class post + second-class stamped preaddressed return envelope

**Sharp 2006a-h** (Continued)

	7. TOMBOLA-branded pen + despatch second-class post + free post business-reply envelope 8. No TOMBOLA-branded pen + despatch second-class post + free post business-reply envelope	
Outcomes	Response rates at any time	
Notes	TOMBOLA compared 2 management policies (colposcopy vs. 6-monthly smears) in women aged 20-59 years with low-grade abnormal cervical smear. Primary outcome: cumulative incidence of cervical intraepithelial neoplasia grade $\geq 2$	
<b><i>Risk of bias</i></b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Unclear	2 authors did randomisation
Adequate sequence generation?	Unclear	"Computer randomised"
Blinding?	Unclear	Outcome not influenced by lack of blinding
Free of selective outcome reporting?	Yes	Expected outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Sharp 2006b**

Methods	2 x 2 x 2 factorial randomised trial (trial-branded pen vs. no pen: comparison 2)	
Data	Women due to receive a TOMBOLA psychosocial questionnaire during June-August 2003 for 12, 18, 24, 30, 34, 36 months' follow-up	
Comparisons	1. TOMBOLA-branded pen + despatch first-class post + free post business-reply envelope 2. No TOMBOLA-branded pen + despatch first-class post + free post business-reply envelope	
Outcomes	Response rates at any time	
Notes	TOMBOLA compared 2 management policies (colposcopy vs. 6-monthly smears) in women aged 20-59 years with low-grade abnormal cervical smear. Primary outcome: cumulative incidence of cervical intraepithelial neoplasia grade $\geq 2$	
<b><i>Risk of bias</i></b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Unclear	2 authors did randomisation
Adequate sequence generation?	Unclear	"Computer randomised"

**Sharp 2006b** (Continued)

Blinding?	Unclear	Outcome not influenced by lack of blinding
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Sharp 2006c**

Methods	2 x 2 x 2 factorial randomised trial (trial branded pen vs. no pen: comparison 3)
Data	Women due to receive a TOMBOLA psychosocial questionnaire during June-August 2003 for 12, 18, 24, 30, 34, 36 months' follow-up
Comparisons	1. TOMBOLA-branded pen + despatch second-class post + second-class stamped preaddressed return envelope 2. No TOMBOLA-branded pen + despatch second-class post + second-class stamped preaddressed return envelope
Outcomes	Defined outcomes reported
Notes	TOMBOLA compared 2 management policies (colposcopy vs. 6-monthly smears) in women aged 20-59 years with low-grade abnormal cervical smear. Primary outcome: cumulative incidence of cervical intraepithelial neoplasia grade $\geq 2$

***Risk of bias***

<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Unclear	2 authors did randomisation
Adequate sequence generation?	Unclear	"Computer randomised"
Blinding?	Unclear	Outcome not influenced by lack of blinding
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Sharp 2006d**

Methods	2 x 2 x 2 factorial randomised trial (trial-branded pen vs. no pen: comparison 4)
Data	Women due to receive a TOMBOLA psychosocial questionnaire during June-August 2003 for 12, 18, 24, 30, 34, 36 month follow-up
Comparisons	1. TOMBOLA-branded pen + despatch second-class post + free post business-reply envelope 2. No TOMBOLA-branded pen + despatch second-class post + free post business-reply envelope
Outcomes	Response rates at any time
Notes	TOMBOLA compared 2 management policies (colposcopy vs. 6-monthly smears) in women aged 20-59 years with low-grade abnormal cervical smear. Primary outcome: cumulative incidence of cervical intraepithelial neoplasia grade $\geq 2$

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Unclear	2 authors did randomisation
Adequate sequence generation?	Unclear	"Computer randomised"
Blinding?	Unclear	Outcome not influenced by lack of blinding
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Sharp 2006e**

Methods	2 x 2 x 2 factorial randomised trial (first vs. second-class post comparison 1)
Data	Women due to receive a TOMBOLA psychosocial questionnaire during June-August 2003 for 12, 18, 24, 30, 34, 36 months' follow-up
Comparisons	1. TOMBOLA-branded pen + despatch first-class post + second-class stamped preaddressed return envelope 2. TOMBOLA-branded pen + despatch second-class post + second-class stamped preaddressed return envelope
Outcomes	Defined outcomes reported
Notes	TOMBOLA compared 2 management policies (colposcopy vs. 6-monthly smears) in women 20-59 years with low-grade abnormal cervical smear. Primary outcome: cumulative incidence of cervical intraepithelial neoplasia grade $\geq 2$



Sharp 2006e (Continued)

<i>Risk of bias</i>		
Item	Authors' judgement	Description
Allocation concealment?	Unclear	2 authors did randomisation
Adequate sequence generation?	Unclear	"Computer randomised"
Blinding?	Unclear	Outcome not influenced by lack of blinding
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

Sharp 2006f

Methods	2 x 2 x 2 factorial randomised trial (first vs. second-class post comparison 2)
Data	Women due to receive a TOMBOLA psychosocial questionnaire during June-August 2003 for 12, 18, 24, 30, 34, 36 months' follow-up
Comparisons	1. No TOMBOLA-branded pen + despatch first-class post + second-class stamped preaddressed return envelope 2. No TOMBOLA-branded pen + despatch second-class post + second-class stamped preaddressed return envelope
Outcomes	Response rates at any time
Notes	TOMBOLA compared 2 management policies (colposcopy vs. 6-monthly smears) in women 20-59 years with low-grade abnormal cervical smear. Primary outcome: cumulative incidence of cervical intraepithelial neoplasia grade $\geq 2$

<i>Risk of bias</i>		
Item	Authors' judgement	Description
Allocation concealment?	Unclear	2 authors did randomisation
Adequate sequence generation?	Unclear	"Computer randomised"
Blinding?	Unclear	Outcome not influenced by lack of blinding
Free of selective outcome reporting?	Yes	Defined outcomes reported

**Sharp 2006f** (Continued)

Other sources of bias?	Unclear	Insufficient information to make a judgement on this
------------------------	---------	--

**Sharp 2006g**

Methods	2 x 2 x 2 factorial randomised trial (first vs. second-class comparison 3)	
Data	Women due to receive a TOMBOLA psychosocial questionnaire during June-August 2003 for 12, 18, 24, 30, 34, 36 months' follow-up	
Comparisons	1. TOMBOLA-branded pen + despatch first-class post + second-class stamped preaddressed return envelope 2. TOMBOLA-branded pen + despatch first-class post + free post business-reply envelope	
Outcomes	Response rates at any time	
Notes	TOMBOLA compared 2 management policies (colposcopy vs. 6-monthly smears) in women aged 20-59 years with low-grade abnormal cervical smear. Primary outcome: cumulative incidence of cervical intraepithelial neoplasia grade $\geq 2$	

**Risk of bias**

Item	Authors' judgement	Description
Allocation concealment?	Unclear	2 authors did randomisation
Adequate sequence generation?	Unclear	"Computer randomised"
Blinding?	Unclear	Outcome not influenced by lack of blinding
Free of selective outcome reporting?	Yes	Defined outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Sharp 2006h**

Methods	2 x 2 x 2 factorial randomised trial (first vs. second-class comparison 4)	
Data	Women due to receive a TOMBOLA psychosocial questionnaire during June-August 2003 for 12, 18, 24, 30, 34, 36 month follow-up	
Comparisons	1. No TOMBOLA-branded pen + despatch first-class post + second-class stamped preaddressed return envelope 2. No TOMBOLA-branded pen + despatch first-class post + free post business-reply envelope	

**Sharp 2006h** (Continued)

Outcomes	Response rates at any time	
Notes	TOMBOLA compared 2 management policies (colposcopy vs. 6-monthly smears) in women 20-59 years with low-grade abnormal cervical smear. Primary outcome: cumulative incidence of cervical intraepithelial neoplasia grade $\geq 2$	
<b>Risk of bias</b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	Unclear	2 authors did randomisation
Adequate sequence generation?	Unclear	"Computer randomised"
Blinding?	Unclear	Outcome not influenced by lack of blinding
Free of selective outcome reporting?	Yes	Expected outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement on this

**Subar 2001**

Methods	2-arm randomised trial	
Data	Participants aged 55-74 years from 3 centres participating in PLCO trial (control arm)	
Comparisons	1. Diet history questionnaire DHQ (36-page food frequency questionnaire) plus 1-page questionnaire on time taken to complete questionnaire 2. PLCO food frequency questionnaire (16-page food frequency questionnaire) plus 1-page questionnaire on time taken to complete questionnaire	
Outcomes	Response rate. No time point given	
Notes	PLCO trial compared digital rectal examination, transvaginal ultrasound and chest x-ray at baseline and at 5 years vs. usual follow-up. PSA and cancer antigen CA125 at baseline, and annually for 5 years. Primary outcome: mortality from PLCO cancers	
<b>Risk of bias</b>		
<b>Item</b>	<b>Authors' judgement</b>	<b>Description</b>
Allocation concealment?	No	Author response "none known"
Adequate sequence generation?	Unclear	Author response "at each centre half were randomised to received". Authors contacted "we would have likely used some computer generated randomisation scheme"

**Subar 2001** (Continued)

Blinding?	Unclear	Author response “the respondents were not blinded. Not known if personnel were 'blinded”
Free of selective outcome reporting?	Yes	Expected outcome reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**Sutherland 1996**

Methods	2-arm randomised trial
Data	226 women taking part in the feasibility study for the Canadian diet and breast cancer prevention feasibility trial
Comparisons	1. Total design method for postal follow-up: white envelope with hospital logo and commemorative stamp; headed notepaper; reply self addressed stamped envelope enclosing contents, hand signature on letters. Postcard sent after 7 days, reminders sent twice 2. Customary method for postal follow-up: brown envelope with return address stamped on, computer-printed labels, no signature on letter, reply self addressed stamped envelope folded and inserted behind forms, no reminder
Outcomes	Time to return of questionnaire at 70 days
Notes	Canadian diet and breast cancer prevention feasibility trial compared teaching women aged over 30 years of age with at least 50% of breast volume occupied by radiological changes of dysplasia how to reduce dietary fat to a level of 15% of calories vs. no teaching. Primary outcome recruitment and retention

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Unclear	Not clear
Adequate sequence generation?	Yes	Author response “computer generated random numbers”
Blinding?	Unclear	Author response “personel knew the allocation a subject had received but their only contact with subjects was the follow-up phone call in some allocated to the ”customary“ method”
Free of selective outcome reporting?	Yes	Expected outcome reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**Svoboda 2001**

Methods	2-arm randomised trial
Data	Adults $\geq$ 16 years with head injury in the CRASH trial (Czech)
Comparisons	1. 1-page, 7-question functional dependence questionnaire sent with a covering letter and a stamped return envelope. Reminders sent after 4 and 8 weeks 2. 3-page, 16-question functional dependence questionnaire sent with a covering letter and a stamped return envelope. Reminders sent after 4 and 8 weeks
Outcomes	Number of questionnaires returned within 3 months
Notes	Numbers randomised and responded provided by authors. Primary outcome for the CRASH trial death from any cause within 2 weeks of injury and death or disability at 6 months

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Yes	Author response "central computer"
Adequate sequence generation?	Yes	Author response "random allocation central computer"
Blinding?	Unclear	Author response "the questionnaires were packaged and sent to patients by personnel who were independent of the study"
Free of selective outcome reporting?	Yes	Expected outcome reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

**Tai 1997**

Methods	2-arm randomised trial
Data	All host study participants lost to follow-up in the evaluation of general practice computer templates trial
Comparisons	1. Recorded delivery reminder letter with QoL questionnaire included, sent once 2. Telephone calls repeated up to 3 times at 10-12 a.m. or 2-5.30 p.m. Message left on answering machine after third call
Outcomes	Number of questionnaires returned. No time point given
Notes	Evaluation of general practice computer templates cluster randomised trial compared computerised templates for asthma and diabetes management in general practice. Primary outcome frequency of use of computer templates assessed by examining computerised records of those who responded to QoL questionnaires

***Risk of bias***

Item	Authors' judgement	Description
Allocation concealment?	Yes	Author response "computer generated by statistician independent to trial manager"
Adequate sequence generation?	Unclear	Author response "computer generated by statistician"
Blinding?	Unclear	Study personnel were not blind to the intervention the participants received
Free of selective outcome reporting?	Yes	Expected outcomes reported
Other sources of bias?	Unclear	Insufficient information to make a judgement about this

APQ-23: Alcohol Problems Questionnaire; AQLQ: Asthma Quality of Life Questionnaire; AUDIT-C: Alcohol Use Disorders Identification Test; AVID: Antiarrhythmics Versus Implantable Defibrillators; CTC: Clinical Trial Centre; COGENT: COlorectal cancer GENEtics; CORE OM: CORE Outcome Measure; CRASH: Corticosteroid Randomisation after Significant Head injury trial; DYD: Down Your Drink; ELISA: enzyme-linked immunosorbent assay; FAOS: Foot and Ankle Outcome Score; FLP: Functional Limitations Profile; LDQ: Leeds Dependency Questionnaire; MIDAS: MIgraine Disability Assessment Questionnaire; NASQ Newcastle Asthma Symptoms Questionnaire; NSABP: National Surgical Adjuvant Breast and Bowel project; PLCO: Prostate, Lung, Colorectal, Ovarian; PSA: prostate-specific antigen; QoL: quality of life; RDQ: Roland Disability Questionnaire; RECORD: Randomised Evaluation of Calcium Or Vitamin D; SANAD: Standard And New Antiepileptic Drugs; SF-36: Short form-36-item health survey; TOMBOLA: Trial Of Management of Borderline and Other Low-grade Abnormal smears; UKBEAM: UK Back pain Exercise And Manipulation.

### Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Arnevik 2009	This retention trial was not embedded in a randomised trial
Atherton 2010	Comparison of Internet vs. postal questionnaires not randomised
Barry 1996	Retention trial compared distribution of scores for subjects completing different questionnaire versions. Author confirmed retention/questionnaire return was not an outcome measure
Bednarek 2008	Retention trial outcome is continuation of treatment
Cox 2003	Retention trial outcome treatment compliance
Day 1998	Retention trial measured adherence to treatment. Authors do not have retention data
Eaker 2004	Retention trial embedded in a cohort

(Continued)

Edelstein 2005	Retention study is not a randomised trial. Incentives not randomised. Author confirmed these were not instituted to help with retention but with adherence to pill taking and life style modification requirements
Grabowski 1995	Substudy aim is retention in treatment comparing different follow-up schedules for addiction treatment trial
Hall 1975	Not a randomised/quasi-randomised retention trial
Hall 1978	Not a randomised/quasi-randomised retention trial
Hoffman 1998	Retention trial embedded in a blood bank cohort
Hopkins 1983	Retention trial embedded in a survey
Iglesias 2000	Retention trial embedded in a cohort of general practitioner practice participants
Iglesias 2001	Retention trial embedded in the recruitment phase of the host trial
Johnson 2004	Retention study not embedded in a randomised trial
Katz 2001	Retention study is not a randomised trial. Authors confirmed the effectiveness of gift incentives was not evaluated in a substudy for the Pride in Parenting trial
Leidy 2000	Retention study appears to be a randomised trial but no response from authors to establish if retention was an outcome. For the substudy, trial sites randomised to 1 of 2 orders of administration of quality of life questionnaires. Response rates not reported. Missing data, internal consistency reliability, mean score values, relationship between the 2 measures evaluated
Marsh 1999	Host study was not a randomised trial. "Practices were randomly allocated to the intervention group using random number tables. Each intervention practice matched with one control practice"
McAuley 1994	Retention study is not a randomised trial. There is a single randomisation stratified by classes in the morning and early evening. No response from authors regarding randomisation to class times
McBee 2009	Retention study not a randomised trial. Authors confirm strategies to improve retention were not evaluated in an Age-Related Eye Disease Study 2 (AREDS2) substudy
Poling 2006	Substudy aim is about diagnostic compliance. 4-arm trial comparing contingency management with or without active bupropion and voucher control with or without active bupropion. Here contingency management and voucher control are aimed at getting information on the disease condition/response to treatment for the primary outcome of the host trial i.e. negative urine sample for cocaine and opioids. Contingency management and voucher control are not related to retention in the host trial but related to diagnostic compliance
Puffer 2004	Retention RCT was embedded in a survey. Authors confirmed that the 2 x 2 factorial study testing four different questionnaire designs was embedded in a survey
Rhoades 1998	Substudy retention in treatment. 2 x 2 trial of dose and visit frequency of attending a clinic either 2 or 5 days per week. Primary outcome was retention in treatment for all randomizations. Similar to <a href="#">Grabowski 1995</a> trial

(Continued)

Roberts 2000	Retention trial embedded in a survey about menopause services
Schmitz 2005	Substudy about compliance to treatment and pill taking behaviour rather than trial retention
Smeeth 2001ab	Substudy about response to baseline assessment
Stoner 1998	Retention study was not a randomised trial. Host study was a cluster randomised trial. Effectiveness of vouchers not evaluated in a substudy
Tassopoulos 2007	Not a retention randomised trial
Wu 1997	Substudy designed to evaluate whether scores are different using 3 modes of questionnaire administration, rather than retention

### Characteristics of ongoing studies [ordered by study ID]

#### Land 2007 2

Trial name or title	Randomised Evaluation of NSABP BAHO Compliance Initiatives
Methods	Randomised trial embedded in Protocol B36
Data	Participants or trial sites/institutions participating in the B36 trial. Pre- and postmenopausal women aged > 18 years, with histologically confirmed invasive breast adenocarcinoma
Comparisons	Menstrual history calendar Usual procedures
Outcomes	To improve form submission compliance in Behavioural and Health Outcomes (BAHO) protocols
Starting date	Unclear
Contact information	Land@nsabp.pitt.edu
Notes	Study identified through personal correspondence with author B36: US-based comparison of 2 combination chemotherapy regimens to treat women with breast cancer

#### Mitchell

Trial name or title	A Randomised Controlled Trial of Combined Pre-Contact and Participant Update for Increasing Questionnaire Response Rates in Older Women
Methods	Randomised trial embedded in the SCOOP screening trial
Data	Women aged 70-85 years not receiving treatment for osteoporosis



**Mitchell** (Continued)

Comparisons	1. Prenotification to complete the 2-year follow-up questionnaire plus a study update 2. No study update
Outcomes	Response to postal questionnaires at 2-year follow-up
Starting date	2010
Contact information	natasha.mitchell@york.ac.uk
Notes	Study identified through mail out to UK clinical trials units SCOOP: UK-based pragmatic randomised controlled trial of the effectiveness of screening for osteoporosis in older women for the prevention of fractures. 10-year absolute risk of fracture calculated from a World Health Organization algorithm based on screening questionnaire data, x-ray and dual-energy x-ray absorptiometry scan results compared with usual care

## DATA AND ANALYSES

### Comparison 1. Addition of incentive vs none: main analysis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	14		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Addition of monetary incentive	3	3166	Risk Ratio (M-H, Fixed, 95% CI)	1.18 [1.09, 1.28]
1.2 Addition of offer of monetary incentive/prize draw	2	3613	Risk Ratio (M-H, Fixed, 95% CI)	1.25 [1.14, 1.38]
1.3 Addition of non-monetary incentive	6	6322	Risk Ratio (M-H, Fixed, 95% CI)	1.00 [0.98, 1.02]
1.4 Addition of offer of non-monetary incentive	2	1138	Risk Ratio (M-H, Fixed, 95% CI)	0.99 [0.95, 1.03]
1.5 Addition of offer of monetary donation to charity	1	815	Risk Ratio (M-H, Fixed, 95% CI)	1.02 [0.78, 1.32]

### Comparison 2. Addition of incentive: sensitivity analysis: quasi-randomised trials removed

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	7		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Addition of monetary incentive	2	1022	Risk Ratio (M-H, Fixed, 95% CI)	1.31 [1.11, 1.55]
1.2 Addition of non-monetary incentive	5	1594	Risk Ratio (M-H, Fixed, 95% CI)	1.00 [0.93, 1.08]

### Comparison 3. Addition of incentive: separating research arms of non-factorial trials (three-/four-arm trials)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	14		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Addition of monetary incentive	3	3066	Risk Ratio (M-H, Fixed, 95% CI)	1.17 [1.09, 1.27]
1.2 Offer of monetary incentive	3	4224	Risk Ratio (M-H, Fixed, 95% CI)	1.24 [1.13, 1.37]
1.3 Addition of non-monetary incentive	8	10793	Risk Ratio (M-H, Fixed, 95% CI)	1.00 [0.98, 1.01]

#### Comparison 4. Addition of telephone follow-up vs incentive

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Telephone survey vs. monetary incentive and questionnaire	1	700	Risk Ratio (M-H, Fixed, 95% CI)	1.08 [0.94, 1.24]

#### Comparison 5. Addition of monetary incentive to both study arms

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	2	902	Risk Ratio (M-H, Fixed, 95% CI)	1.12 [1.04, 1.22]
1.1 Addition of GBP10 plus offer of GBP10 vs. addition of GBP5 plus offer of GBP5	1	485	Risk Ratio (M-H, Fixed, 95% CI)	1.16 [1.04, 1.30]
1.2 Addition of GBP20 voucher offer vs. addition of GBP10 voucher offer	1	417	Risk Ratio (M-H, Fixed, 95% CI)	1.08 [0.97, 1.21]

#### Comparison 6. Addition of monetary incentive vs offer of incentive

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	2	297	Risk Ratio (M-H, Fixed, 95% CI)	1.04 [0.91, 1.19]
1.1 Addition of monetary incentive vs. offer of entry into prize draw	2	297	Risk Ratio (M-H, Fixed, 95% CI)	1.04 [0.91, 1.19]

#### Comparison 7. Enhanced letter versus standard letter: main analysis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	2		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Enhanced letter vs. standard letter	2	2479	Risk Ratio (M-H, Fixed, 95% CI)	1.01 [0.97, 1.05]

**Comparison 8. Communication strategies letter: total design method**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Total design method for postal questionnaires vs. customary method	1	226	Risk Ratio (M-H, Fixed, 95% CI)	1.43 [1.22, 1.67]

**Comparison 9. Communication strategies post: main analysis**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	7		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Priority vs. regular post	7	1888	Risk Ratio (M-H, Fixed, 95% CI)	1.02 [0.95, 1.09]

**Comparison 10. Communication strategies: additional reminder vs usual follow-up: main analysis**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	6		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Additional reminder vs. usual follow-up procedures	6	3401	Risk Ratio (M-H, Fixed, 95% CI)	1.03 [0.99, 1.06]

**Comparison 11. Communication strategies additional reminder to trial site vs usual reminder (ICC 0.054)**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	1		Risk Ratio (Fixed, 95% CI)	Subtotals only
1.1 Monthly reminder of upcoming assessments to trial site vs. usual reminders	1	272	Risk Ratio (Fixed, 95% CI)	0.96 [0.83, 1.11]

### Comparison 12. Communication strategies: questionnaire administered early vs late

---

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Early vs. late administration	1	664	Risk Ratio (M-H, Fixed, 95% CI)	1.10 [0.96, 1.26]

---

### Comparison 13. Communication strategies: type of reminder: main analysis

---

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Recorded delivery vs. telephone reminder	1	192	Risk Ratio (M-H, Fixed, 95% CI)	2.08 [1.11, 3.87]

---

### Comparison 14. Questionnaire strategies: new vs standard questionnaire: main analysis

---

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	10		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Short vs. long questionnaire	5	7277	Risk Ratio (M-H, Fixed, 95% CI)	1.04 [1.00, 1.08]
1.2 Long and clear vs. short and condensed questionnaires	1	900	Risk Ratio (M-H, Fixed, 95% CI)	1.01 [0.95, 1.07]
1.3 Question order: condition first vs. generic first questions	2	9435	Risk Ratio (M-H, Fixed, 95% CI)	1.00 [0.97, 1.02]
1.4 Questionnaire: relevant vs. less relevant to condition	2	3893	Risk Ratio (M-H, Fixed, 95% CI)	1.07 [1.01, 1.14]

---

**Comparison 15. Questionnaire strategies: new vs standard questionnaire: sensitivity analysis quasi-randomised trial McColl**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	8		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Short vs. long questionnaire	5	7277	Risk Ratio (M-H, Fixed, 95% CI)	1.04 [1.00, 1.08]
1.2 Long and clear vs. short and condensed questionnaires	1	900	Risk Ratio (M-H, Fixed, 95% CI)	1.01 [0.95, 1.07]
1.3 Questionnaire: relevant vs. less relevant to condition	2	3893	Risk Ratio (M-H, Fixed, 95% CI)	1.07 [1.01, 1.14]

**Comparison 16. Behavioural strategies: main analysis**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	2		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Motivation vs. information	2	273	Risk Ratio (M-H, Fixed, 95% CI)	1.08 [0.93, 1.24]

**Comparison 17. Case management**

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Case management vs. usual follow-up	1	703	Risk Ratio (M-H, Fixed, 95% CI)	1.00 [0.97, 1.04]

**Comparison 18. Methodology strategies**


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Retention	1		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only
1.1 Open vs. blind trial design	1	538	Risk Ratio (M-H, Fixed, 95% CI)	1.37 [1.16, 1.63]

## Analysis 1.1. Comparison 1 Addition of incentive vs none: main analysis, Outcome 1 Retention.


Review: Strategies to improve retention in randomised trials

Comparison: 1 Addition of incentive vs none: main analysis

Outcome: 1 Retention



(Continued ...)




(1) data for overall response used for ockayne

### Analysis 2.1. Comparison 2 Addition of incentive: sensitivity analysis: quasi-randomised trials removed, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials


Comparison: 2 Addition of incentive: sensitivity analysis: quasi-randomised trials removed

Outcome: 1 Retention



(Continued ...)






### Analysis 3.1. Comparison 3 Addition of incentive: separating research arms of non-factorial trials (three-/four-arm trials), Outcome 1 Retention.


Review: Strategies to improve retention in randomised trials

Comparison: 3 Addition of incentive: separating research arms of non-factorial trials (three-/four-arm trials)

Outcome: 1 Retention



(Continued . . .)




### Analysis 4.1. Comparison 4 Addition of telephone follow-up vs incentive, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 4 Addition of telephone follow-up vs incentive

Outcome: 1 Retention




### Analysis 5.1. Comparison 5 Addition of monetary incentive to both study arms, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 5 Addition of monetary incentive to both study arms

Outcome: 1 Retention




### Analysis 6.1. Comparison 6 Addition of monetary incentive vs offer of incentive, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 6 Addition of monetary incentive vs offer of incentive

Outcome: 1 Retention




### Analysis 7.1. Comparison 7 Enhanced letter versus standard letter: main analysis, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 7 Enhanced letter versus standard letter: main analysis

Outcome: 1 Retention




**Analysis 8.1. Comparison 8 Communication strategies letter: total design method, Outcome 1 Retention.**

Review: Strategies to improve retention in randomised trials

Comparison: 8 Communication strategies letter: total design method

Outcome: 1 Retention




### Analysis 9.1. Comparison 9 Communication strategies post: main analysis, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 9 Communication strategies post: main analysis

Outcome: 1 Retention




### Analysis 10.1. Comparison 10 Communication strategies: additional reminder vs usual follow-up: main analysis, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 10 Communication strategies: additional reminder vs usual follow-up: main analysis

Outcome: 1 Retention






### Analysis 11.1. Comparison 11 Communication strategies additional reminder to trial site vs usual reminder (ICC 0.054), Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 11 Communication strategies additional reminder to trial site vs usual reminder (ICC 0.054)

Outcome: 1 Retention




### Analysis 12.1. Comparison 12 Communication strategies: questionnaire administered early vs late, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 12 Communication strategies: questionnaire administered early vs late

Outcome: 1 Retention




**Analysis 13.1. Comparison 13 Communication strategies: type of reminder: main analysis, Outcome 1 Retention.**

Review: Strategies to improve retention in randomised trials

Comparison: 13 Communication strategies: type of reminder: main analysis

Outcome: 1 Retention




## Analysis 14.1. Comparison 14 Questionnaire strategies: new vs standard questionnaire: main analysis, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 14 Questionnaire strategies: new vs standard questionnaire: main analysis

Outcome: 1 Retention




## Analysis 15.1. Comparison 15 Questionnaire strategies: new vs standard questionnaire: sensitivity analysis quasi-randomised trial McColl, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 15 Questionnaire strategies: new vs standard questionnaire: sensitivity analysis quasi-randomised trial McColl

Outcome: 1 Retention




### Analysis 16.1. Comparison 16 Behavioural strategies: main analysis, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 16 Behavioural strategies: main analysis

Outcome: 1 Retention




### Analysis 17.1. Comparison 17 Case management, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 17 Case management

Outcome: 1 Retention




### Analysis 18.1. Comparison 18 Methodology strategies, Outcome 1 Retention.

Review: Strategies to improve retention in randomised trials

Comparison: 18 Methodology strategies

Outcome: 1 Retention



## ADDITIONAL TABLES

Table 1. Trials evaluating incentive strategies

Trial or trial comparison	Incentive groups	Control group	Outcome type
<b>Addition of incentive vs. none</b>			
<a href="#">Bauer 2004ab</a>	a) USD10 cheque b) USD2 cheque Arms combined for main analysis	No incentive	DNA specimen kit return plus postal questionnaire response
<a href="#">Gates 2009</a>	GBP5 voucher	No incentive	Postal questionnaire response
<a href="#">Kenyon 2005</a>	GBP5 voucher	No incentive	Postal questionnaire response
<a href="#">Khadjesari 2011 1ac</a>	a) Offer GBP5 voucher c) Offer of entry into GBP250 prize draw, groups combined for	No incentive	Internet-based questionnaire response

**Table 1. Trials evaluating incentive strategies** (Continued)

	main analysis		
<a href="#">Khadjesari 2011 2</a>	Offer of GBP10 Amazon.co.uk voucher	No incentive	Internet-based questionnaire response
<a href="#">Bowen 2000abc</a>	a) Certificate b) Pin c) Pin and certificate groups combined for main analysis	No incentive	Participants retention
<a href="#">Renfroe 2002a</a>	Certificate of appreciation	No certificate of appreciation	Postal questionnaire response
<a href="#">Sharp 2006a</a>	Pen	No pen	Postal questionnaire response
<a href="#">Sharp 2006b</a>	Pen	No pen	Postal questionnaire response
<a href="#">Sharp 2006c</a>	Pen	No pen	Postal questionnaire response
<a href="#">Sharp 2006d</a>	Pen	No pen	Postal questionnaire response
<a href="#">Cockayne 2005</a>	Offer of study results	No offer	Postal questionnaire response
<a href="#">Hughes 1989</a>	Offer of free reprint of results	No offer	Postal questionnaire response
<a href="#">Khadjesari 2011 1b</a>	Offer of GBP5 charity donation	No offer	Internet-based questionnaire response
<b>Addition of monetary incentive to both groups</b>			
<a href="#">Bailey 1 unpublished</a>	Offer of GBP20 shopping voucher	Offer of GBP10 shopping voucher	Postal questionnaire response
<a href="#">Bailey 2 unpublished</a>	Shopping voucher: GBP10 in advance and GBP10 on data return	Shopping voucher: GBP5 in advance and GBP5 on data return	Questionnaire response and chlamydia kit return
<b>Addition of monetary incentive vs. offer of incentive</b>			
<a href="#">Kenton 2007a</a>	USD2 coin	Draw for USD50 gift voucher	Postal questionnaire response
<a href="#">Kenton 2007b</a>	USD2 coin	Draw for USD50 gift voucher	Postal questionnaire response
<b>Offer of prize draw vs. no offer</b>			
<a href="#">Leigh Brown 1997</a>	Aware of monthly prize draw of GBP25 gift voucher	No offer of draw	Postal questionnaire response

**Table 2. Trials evaluating communication strategies**

Trial or trial comparison	Communication strategy	Control arm	Outcome type
<b>Enhanced letter vs. standard letter</b>			
Renfroe 2002c	Cover letter signed by physician	Cover letter signed by co-ordinator	Postal questionnaire response
Marson 2007	Letter explaining the approximate length of time to complete questionnaire	Standard letter	Postal questionnaire response
<b>Total design method vs. customary method</b>			
Sutherland 1996	Total design method for postal follow-up	Standard method for postal follow-up	Postal questionnaire response
<b>Priority vs. regular post</b>			
Renfroe 2002b	Express delivery	Standard delivery	Postal questionnaire response
Sharp 2006e	Despatch first-class stamp	Despatch second-class stamp	Postal questionnaire response
Sharp 2006f	Despatch first-class stamp	Despatch second-class stamp	Postal questionnaire response
Sharp 2006g	Second-class return envelope	Free post return envelope	Postal questionnaire response
Sharp 2006h	Second-class return envelope	Free post return envelope	Postal questionnaire response
Kenton 2007c	Priority mail	Standard mail	Postal questionnaire response
Kenton 2007d	Priority mail	Standard mail	Postal questionnaire response
<b>Additional reminder vs. usual follow-up</b>			
Ashby 2011	Electronic reminder	No electronic reminder	Postal questionnaire response
MacLennan unpublished	Telephone reminder	No telephone reminder	Postal questionnaire response
Nakash 2007	Trial calendar given at recruitment with questionnaire due dates	No calendar	Postal questionnaire response
Severi 2011 1	Text message and fridge magnet both emphasising social benefits of study participation	Text message reminder sent 3 days after questionnaire	Postal questionnaire response
Severi 2011 2	Telephone reminder from principle investigator	Standard procedures.	Return of cotinine samples



**Table 2. Trials evaluating communication strategies** (Continued)

Man 2011	SMS text message as follow-up questionnaire sent out	No SMS text message	Postal questionnaire response
<b>Additional trial site reminder vs. usual reminder</b>			
Land 2007	Prospective monthly reminder of upcoming assessments to trial sites	No extra reminder to trial sites	Postal questionnaire response
<b>Early vs. late administration of questionnaire</b>			
Renfro 2002d	Questionnaire sent 2-3 weeks after last AVID follow-up visit	Questionnaire sent 1-4 months after last AVID follow-up visit	Postal questionnaire response
<b>Recorded delivery vs. telephone reminder</b>			
Tai 1997	Recorded delivery reminder	Telephone reminder	Postal questionnaire response
<b>Addition telephone follow-up vs. incentive</b>			
Couper 2007	Telephone survey by trained interviewer	Postal questionnaire and USD5 bill	Post and questionnaire response

AVID: Antiarrhythmics Versus Implantable Defibrillators; SMS: short message service.

**Table 3. Trials evaluating new questionnaire strategies**

Trial or trial comparison	Questionnaire strategy	Control arm	Outcome type
<b>Short vs. long</b>			
Dorman 1997	Short EuroQol	Long SF-36 questionnaire.	Postal questionnaire response
Edwards 2001 unpublished	1-page, 7-question functional dependence questionnaire	3-page, 16-question functional dependence questionnaire	Postal questionnaire response
Svoboda 2001 unpublished	1-page, 7-question functional dependence questionnaire	3-page, 16-question functional dependence questionnaire	Postal questionnaire response
McCambridge 2011 1b	AUDIT Short + LDQ	APQ	Internet-based questionnaire response
McCambridge 2011 2b	AUDIT Short + LDQ	APQ	Internet-based questionnaire response

**Table 3. Trials evaluating new questionnaire strategies** (Continued)

<b>Long and clear vs. short and condensed</b>			
Subar 2001	DHQ (36-page food frequency questionnaire)	PLCO (16-page food frequency questionnaire)	Postal questionnaire response and onsite completion
<b>Question order</b>			
McColl 2003 1	Asthma condition-specific questions first followed by generic	Generic questions followed by condition specific	Postal questionnaire response
McColl 2003 2	Angina condition-specific questions followed by generic	Generic questions followed by condition specific	Postal questionnaire response
Letley 2000 unpublished	RDQ at front and SF-36 at back	SF-36 at front RDQ at back	Postal questionnaire response
<b>Relevance of questionnaire</b>			
McCambridge 2011 1a	APQ23 items	CORE-OM Mental health assessment 23/34 items	Internet-based questionnaire response
McCambridge 2011 2a	AUDIT Short + LDQ	CORE-OM Mental health assessment 10 items	Internet-based questionnaire response

APQ: Alcohol Problems Questionnaire; AUDIT: Alcohol Use Disorders Identification Test; LDQ Leeds Dependency Questionnaire; PLCO: Prostate, Lung, Colorectal, Ovarian; SF-36: Short Form 36 item.

**Table 4. Gain in number of questionnaires returned per 1000 questionnaires sent**

Example of proportion of questionnaires returned in control arm	Strat- egy to im- prove re- tention	RR	1	RR	30%	40%	50%	60%	70%	80%	90%

**Table 4. Gain in number of questionnaires returned per 1000 questionnaires sent** (Continued)

Addition of monetary incentive versus none	1.18	0.847	107	92	76	61	5	3	2
Addition of offer of monetary incentive/prize draw versus none	1.25	0.800	140	120	100	80	60	40	20
Addition of higher value monetary incentive versus addition of lower amount	1.12	0.890	77	66	55	44	33	22	11

RR: risk ratio.

## APPENDICES

### Appendix I. Attrition terms

Cochrane Central Register of Controlled Trials (CENTRAL) and Database of Abstracts of Reviews of Effects (DARE in *The Cochrane Library*) were searched using attrition terms shown below.

(minimi\$ adj2 attrition).ab,ti.

(prevent\$ adj2 attrition).ab,ti.

(lessen\$ adj2 attrition).ab,ti.

(decreas\$ adj2 attrition).ab,ti.

(reduc\$ adj2 attrition).ab,ti.

(minimi\$ adj2 drop-out).ab,ti.

(prevent\$ adj2 drop-out).ab,ti.

(lessen\$ adj2 drop-out).ab,ti.

(decreas\$ adj2 drop-out).ab,ti.

(reduc\$ adj2 drop-out).ab,ti.

(minimi\$ adj2 drop-out\$).ab,ti.

(prevent\$ adj2 drop-out\$).ab,ti.

(lessen\$ adj2 drop-out\$).ab,ti.  
 (decreas\$ adj2 drop-out\$).ab,ti.  
 (reduc\$ adj2 drop-out\$).ab,ti.  
 (minimi\$ adj2 drop-out\$).ab,ti.  
 (prevent\$ adj2 drop-out\$).ab,ti.  
 (lessen\$ adj2 drop-out\$).ab,ti.  
 (decreas\$ adj2 drop-out\$).ab,ti.  
 (reduc\$ adj2 drop-out\$).ab,ti.  
 (minimi\$ adj2 dropout\$).ab,ti.  
 (prevent\$ adj2 dropout\$).ab,ti.  
 (lessen\$ adj2 dropout\$).ab,ti.  
 (decreas\$ adj2 dropout\$).ab,ti.  
 (reduc\$ adj2 dropout\$).ab,ti.  
 (strateg\$ adj2 drop-out\$).ab,ti.  
 (strateg\$ adj2 dropout\$).ab,ti.  
 (loss adj2 follow-up).ab,ti.  
 (lost adj2 follow-up).ab,ti.  
 (loss adj2 followup).ab,ti.  
 (lost adj2 followup).ab,ti.  
 (minimi\$ adj2 withdrawal).ab,ti.  
 (prevent\$ adj2 withdrawal).ab,ti.  
 (lessen\$ adj2 withdrawal).ab,ti.  
 (decreas\$ adj2 withdrawal).ab,ti.  
 (reduc\$ adj2 withdrawal).ab,ti.  
 (minimi\$ adj2 withdrawal\$).ab,ti.  
 (prevent\$ adj2 withdrawal\$).ab,ti.  
 (lessen\$ adj2 withdrawal\$).ab,ti.  
 (decreas\$ adj2 withdrawal\$).ab,ti.  
 (reduc\$ adj2 withdrawal\$).ab,ti.  
 (strateg\$ adj2 attrition).ab,ti.  
 (strateg\$ adj2 drop-out).ab,ti.  
 (strateg\$ adj2 dropout).ab,ti.  
 (strateg\$ adj2 follow-up).ab,ti.  
 (strateg\$ adj2 followup).ab,ti.  
 (increas\$ adj2 retention).ab,ti.  
 (encourag\$ adj2 retention).ab,ti.  
 (maximi\$ adj2 retention).ab,ti.  
 (promot\$ adj2 retention).ab,ti.  
 (improv\$ adj2 retention).ab,ti.  
 (strateg\$ adj2 response\$).ab,ti.  
 (strateg\$ adj2 (questionnaire\$ adj3 response\$)).ab,ti.  
 (increas\$ adj2 (questionnaire\$ adj3 response\$)).ab,ti.  
 (encourag\$ adj2 (questionnaire\$ adj3 response\$)).ab,ti.  
 (maximi\$ adj2 (questionnaire\$ adj3 response\$)).ab,ti.  
 (promot\$ adj2 (questionnaire\$ adj3 response\$)).ab,ti.  
 (improv\$ adj2 (questionnaire\$ adj3 response\$)).ab,ti.  
 (increas\$ adj2 response\$).ab,ti.  
 (encourag\$ adj2 response\$).ab,ti.  
 (maximi\$ adj2 response\$).ab,ti.  
 (promot\$ adj2 response\$).ab,ti.  
 (improv\$ adj2 response\$).ab,ti.  
 (retention adj2 strateg\$).ab,ti.  
 retention rate\$.ab,ti.

(retention adj2 method\$).ab,ti.  
(retention adj2 technique\$).ab,ti.  
attrition rate\$.ab,ti.  
(questionnaire\$ adj3 (response\$ adj2 method\$)).ab,ti.  
(questionnaire\$ adj3 (response adj2 technique\$)).ab,ti.  
(questionnaire adj response rate\$).ab,ti. (1145)  
(difficult\$ adj2 (retain\$ or retention)).ab,ti.

Participant Dropouts/

The search syntax was adapted as follows for different search interfaces

**MEDLINE; EMBASE; PsycINFO via Ovid,**

**pt-** Publication type.

**adj2-** words within 2 words of each other.

**ab-** word in abstract.

**sh-** sub heading.

**ti** word in title.

**/** - Subject heading Medline.

**\$** - Truncation symbol.

Codes used to de duplicate in Ovid were:

use mesz

use emez

use psyh

**Cochrane Central Register of Controlled Trials (CENTRAL) and Database of Abstracts of Reviews of Effects (DARE) via *The Cochrane Library***

\* Truncation symbol

NEAR/2 - words within 2 words of each other.

:kw- keyword

Codes used to de duplicate in CENTRAL were:

“accession number ” near pubmed

“accession number ” near2 embase

**CINAHL (Cumulative Index to Nursing and Allied Health; 1981 to present) via EBSCOHost**

**MH** Major heading (CINAHL via EBSCOHost -)

**+ -** (e.g. Treatment Outcomes+) (CINAHL via EBSCOHost -)

**N2** - words within 2 words of each other.

\* Truncation symbol.

**Campbell Collaboration’s Social, Psychological, Educational and Criminological Trials Register (C2-SPECTR <http://geb9101.gse.upenn.edu/>)**

\* - Truncation symbol.

**Education Resource Information Centre (ERIC) via Dialog datastar.**

**\$** - Truncation symbol

**ab-** word in abstract.

**ti** word in title.

### **MeSH Headings**

**exp Participant Dropouts/:** This was used in MEDLINE, only as a subject heading.

In PsycINFO **Experimental attrition** was used

In CINAHL plus **Research subject retention** was used (“research dropouts” - term scope = mechanisms used to keep study participants willing and able to contribute to participate in the study for its duration).

## Appendix 2. MEDLINE search strategy

Search strategy for **MEDLINE** Cochrane Highly Sensitive Search Strategy for identifying randomized trials in MEDLINE: Sensitivity and precision maximising version, 2008 revision [Lefebvre 2008](#); Ovid format.

- #1 randomized controlled trial.pt.
- #2 controlled clinical trial.pt.
- #3 randomized.ab.
- #4 placebo.ab.
- #5 clinical trials as topic.sh.
- #6 randomly.ab.
- #7 trial.ti.
- #8 1 or 2 or 3 or 4 or 5 or 6 or 7
- #9 exp animals/ not humans. sh.
- #10 8 not 9
- #11 10 AND Attrition terms ([Appendix 1](#))

## Appendix 3. EMBASE search strategy

**EMBASE** search using a sensitivity and specificity maximising search strategy for identifying clinically sound treatment studies [Wong 2006a](#).

- #1 random\$.tw.
- #2 placebo\$.ti,ab,sh.
- #3 double-blind\$.tw.
- #4 1 or 2 or 3
- #5 4 AND Attrition terms ([Appendix 1](#))

## Appendix 4. PsycINFO search strategy

**PsycINFO** search strategy for identifying high quality studies on treatment: Sensitivity and specificity maximising version. [Eady 2008](#)

- #1 double-blind.ab,ti.
- #2 "random\$ assigned.".ab,ti.
- #3 control.ab,ti.
- #4 1 or 2 or 3
- #5 4 AND Attrition terms ([Appendix 1](#))

## Appendix 5. CINAHL search strategy

**CINAHL** search strategy for identifying therapy studies and review articles: Sensitivity and specificity maximising version [Wong 2006b](#)

- #1 PT Clinical trial
- #2 (MH "Treatment Outcomes+")
- #3 randomi?ed
- #4 1 or 2 or 3
- #5 4 AND Attrition terms ([Appendix 1](#))

## Appendix 6. C2-SPECTR search strategy

Campbell Collaboration's Social, Psychological, Educational and Criminological Trials Register C2-SPECTR advanced search Non-indexed fields and indexed fields.

Terms used: {retention} OR {attrition} OR {dropout} OR {drop-out} OR {withdrawal} OR {response}.

## Appendix 7. ERIC search strategy

ERIC search strategy [Petrosino 2000](#)

#1 RANDOM\$.TI,AB.

#2 RANDOM\$.TI,AB.

#3 (ALLOCAT\$ OR ALLOT\$ OR ASSIGN\$ OR BASIS OR DIVID\$ OR ORDER\$).TI,AB.

#4 (2 NEAR 3).TI,AB.

#5 RANDOM\$.TI,AB. NOT (4 ADJ or1).TI,AB.

#6 ((SINGL\$ OR DOUBL\$ OR TREBL\$ OR TRIPL\$) NEAR (BLIND\$ OR MASK\$)).TI,AB.

#7 ((COMPAR\$ OR CONTROL\$ OR EXPERIMENT\$ OR INTERVENT\$ OR THERAP\$ OR TREATMENT\$) NEAR (GROUP\$ OR CLASS\$)).TI,AB.

#8 (ALLOCAT\$ OR ALLOT\$ OR ASSIGN\$ OR DIVID\$ OR ORDER\$).TI,AB.

#9 (7 NEAR 8).TI,AB.

#10 crossover.TI,AB.

#11 (LATIN NEAR SQUARE).TI,AB.

#12 ((CLINIC\$ OR CONTROL\$) NEAR (TRIAL\$ OR STUDY\$ OR STUDIES\$)).TI,AB.

#13 PLACEBO\$

#14 (1 OR 4 OR 5 OR 6 OR 9 OR 10 OR 11 OR 12 OR 13).TI,AB.

#15 Attrition

#16 (attrition ADJ research ADJ studies). TI,AB.

#17 14 AND 16

#18 17 AND Attrition terms ([Appendix 1](#))

## Appendix 8. Reference lists of reviews and other publications searched

1. Torgerson G. Should we pay the patient? Review of financial incentives to enhance patient compliance. *BMJ* 1997;315.
2. Jeffery M, Hickey BE, Hider PN. Follow-up strategies for patients treated for non-metastatic colorectal cancer. *Cochrane Database of Systematic Reviews* 2007 Art No : CD002200 DOI: 10.1002/14651858.CD002200.pub2 2011.
3. Rudy EB, Estok PJ, Kerr ME, Menzel L. Research incentives: money versus gifts. *Nursing Research* 1994;43:253.
4. Bruzzese JM, Gallagher R, McCann-Doyle S, Reiss PT, Wijetunga NA. Effective methods to improve recruitment and retention in school-based substance use prevention studies. *Journal of School Health* 2009;79:400-7.
5. Rojas MP, Telaro E, Russo A, Moschetti I, Coe L, Fossati R et al. Follow-up strategies for women treated for early breast cancer. *Cochrane Database of Systematic Reviews*, 2000, Art. No.: CD001768. DOI:10.1002/14651858.CD001768.pub2.
6. Froelicher ES, Miller NH, Buzaitis A, Pfenninger P, Misuraco A, Jordan S et al. The Enhancing Recovery in Coronary Heart Disease Trial (ENRICH): strategies and techniques for enhancing retention of patients with acute myocardial infarction and depression or social isolation. *Journal of Cardiopulmonary Rehabilitation* 2003;23:269-80.
7. Sarre G, Dyas J, Clarke A, Keen C. *Planning Recruitment Options: Strategies for Primary Care Research*. 2009.
8. Heo M, Papademetriou E, Meyers BS. Design characteristics that influence attrition in geriatric antidepressant trials: meta-analysis. *International Journal of Geriatric Psychiatry* 2009;24:990-1001.
9. Fabricatore AN, Wadden TA, Moore RH, Butryn ML, Gravalles EA, Erondy NE et al. Attrition from randomized controlled trials of pharmacological weight loss agents: a systematic review and analysis. *Obesity Reviews* 2009;10:333-41.
10. Northouse LL, Rosset T, Phillips L, Mood D, Schafenacker A, Kershaw T. Research with families facing cancer: the challenges of accrual and retention. *Research in Nursing & Health* 2006;29:199-211.
11. Mc Daid C, Hodges Z, Fayter D, Stirk L, Eastwood A. Increasing participation of cancer patients in randomised controlled trials: a systematic review. *Trials* 2006;7:16.

12. McFarlane J: Strategies for successful recruitment and retention of abused women for longitudinal studies. *Issues in Mental Health Nursing* 2007;28:883-97.
13. Burns D, Soward ACM, Skelly AH, Leeman J, Carlson J. Effective recruitment and retention strategies for older members of rural minorities. *The Diabetes Educator* 2008;34:1045-52.
14. Davis LL, Broome ME, Cox R. Maximizing retention in community-based clinical trials. *Journal of Nursing Scholarship* 2002; 34:47-53.
15. Edwards P, Roberts I, Sandercock P, Frost C. Follow-up by mail in clinical trials: does questionnaire length matter? *Control Clinical Trials* 2004;25:31-52.
16. Edwards P, Cooper R, Roberts I, Frost C. Meta-analysis of randomised trials of monetary incentives and response to mailed questionnaires. *Journal of Epidemiology and Community Health* 2005;59:987-99.
17. Edwards PJ, Roberts IG, Clarke MJ, DiGiuseppi C, Wentz R, Kwan I et al. Methods to increase response rates to postal and electronic questionnaires. *Cochrane Database of Systematic Reviews* 2009, Issue 3 Art No: MR000008 2009.
18. Nakash R, Hutton J, Jorstad-Stein E, Gates S, Lamb S. Maximising response to postal questionnaires - a systematic review of randomised trials in health research. *BMC Medical Research Methodology* 2006;6:5.
19. Robinson KA, Dennison CR, Wayman DM, Pronovost PJ, Needham DM. Systematic review identifies number of strategies important for retaining study participants. *Journal of Clinical Epidemiology* 2007;60:757.
20. Schellings R, Kessels AG, ter Riet G, Knottnerus JA, Sturmans F. Randomized consent designs in randomized controlled trials: systematic literature search. *Contemporary Clinical Trials* 2006;27:320-32.
21. Warner KE, Murt HA. Economic incentives for health. *Annual Review of Public Health* 1984;5:107-33.
22. Wiemann CM, Chacko MR, Tucker JD, Velasquez MM, Smith PB, DiClemente RJ et al. Enhancing recruitment and retention of minority young women in community-based clinical research. *Journal of Pediatric and Adolescent Gynecology* 2005;18:403-7.
23. Yabroff KR, Kerner JF, Mandelblatt JS. Effectiveness of interventions to improve follow-up after abnormal cervical cancer screening. *Preventive Medicine* 2000;31:429-39.
24. van der Bij AK, Laurant MGH, Wensing M. Effectiveness of physical activity interventions for older adults: a review. *American Journal of Preventive Medicine* 2002;22:120-33.
25. Yancey AK, Ortega AN, Kumanyika SK. Effective recruitment and retention of minority research participants. *Annual Review of Public Health* 2006;27:1-28.
26. Scott P, Edwards P. Personally addressed hand-signed letters increase questionnaire response: a meta-analysis of randomised controlled trials. *BMC Health Services Research* 2006;6:111.

## Appendix 9. Characteristics of host trials

Clinical area main trial	Condition	Participants	Setting	Attrition Study
Treatment dependence	of Alcohol	Adults scoring +5 on AUDIT-C, mean age 37 years in an online trial comparing interactive computer intervention plus website information vs. website information for modifying alcohol intake <a href="#">Murray 2007</a>	Community: online	<a href="#">Khadjesari 2011 1abc</a>
	Alcohol	Adults scoring +5 on AUDIT-C mean age 37 years in an online trial comparing in-	Community: online	<a href="#">Khadjesari 2011 2</a>



(Continued)

		teractive computer inter- vention plus website in- formation vs. website in- formation for modifying alcohol intake <a href="#">Murray 2007</a>		
	Alcohol	Adults scoring +5 on AUDIT-C mean age 37 years in an on- line trial comparing in- teractive computer inter- vention plus website in- formation vs. website in- formation for modifying alcohol intake <a href="#">Murray 2007</a>	Community: online	<a href="#">McCambridge 2011 1</a>
	Alcohol	Adults scoring +5 on AUDIT-C mean age 37 years in an on- line trial comparing in- teractive computer inter- vention plus website in- formation vs. website in- formation for modifying alcohol intake <a href="#">Murray 2007</a>	Community: online	<a href="#">McCambridge 2011 2</a>
	Smoking	Adult smokers aged 38- 77 years in a smoking cessation trial of pub- lic education through media and community wide events, healthcare providers work sites and other organisations vs. no intervention <a href="#">Gail 1992</a>	USA community	<a href="#">Bauer 2004ab</a>
	Smoking	Adult smokers mean age 36.7 years in a trial of nicotine gum vs. placebo gum. Smokers for > 1 year <a href="#">Hughes 1984</a>	USA community	<a href="#">Hughes 1989</a>
	Smoking	Adult smokers willing to quit aged > 16 years in a trial compar- ing Txt2stop motiva-	UK community	<a href="#">Severi 2011 1</a>

(Continued)

		tional messages and behaviour change support vs. text messages unrelated to quitting <a href="#">Free 2011</a>		
	Smoking	Adult smokers willing to quit aged > 16 years in a trial comparing Txt2stop motivational messages and behaviour change support vs. text messages unrelated to quitting <a href="#">Free 2011</a>	UK community	<a href="#">Severi 2011 2</a>
Treatment of injury	Neck	MINT trial: adults with whiplash injury 18-87 years in a 2 x 2 cluster randomised trial comparing whiplash book vs. usual advice. Individuals randomised to physiotherapy vs. single advice session reinforcing advice given <a href="#">Lamb 2007</a>	UK hospital trusts	<a href="#">Gates 2009</a>
	Ankle	Cast trial: adults aged 16-57 years with acute severe ankle sprain in a trial comparing tubular bandage vs. below knee cast vs. Aircast® ankle brace vs. Bledsoe® boot <a href="#">Cooke 2009</a>	UK accident and emergency departments	<a href="#">Nakash 2007*</a>
	Head	Adults with head injury aged > 16 years in trial of 48-hour infusion of methylprednisolone vs. placebo <a href="#">CRASH Trial collaborators 2004</a>	UK hospital intensive care units	<a href="#">Edwards 2001*</a>
	Head	Adults with head injury > 16 years CRASH trial: 48-hour infusion of methylprednisolone vs. placebo <a href="#">CRASH Trial collaborators 2004</a>	Czech republic hospital intensive care units	<a href="#">Svoboda 2001*</a>

(Continued)

Treatment of disease	Breast cancer	Women with ductal carcinoma in situ aged > 49 years in a trial comparing anastrozole vs. tamoxifen (unpublished)	USA, Canada, Puerto Rico hospitals	<a href="#">Land 2007</a>
	Stroke	Acute stroke patients 50-80 years in an international stroke trial of heparin 125,000 IU twice daily + aspirin 300 mg daily vs. heparin 125,000 IU twice vs. heparin 5000 IU twice daily + aspirin 300 mg daily, heparin 5000 IU twice daily vs. aspirin 300 mg daily vs. no heparin or aspirin <a href="#">International Stroke Trial Group 1997</a>	UK hospital	<a href="#">Dorman 1997</a>
	Ventricular fibrillation, ventricular tachycardia	Adults cardioverted from ventricular tachycardia or resuscitated from ventricular fibrillation aged 54-76 years participating in the AVID trial comparing an implanted cardioverter defibrillator vs. antiarrhythmic drugs <a href="#">AVID investigators 1997</a>	USA hospital	<a href="#">Renfroe 2002a-d</a>
	Epilepsy	Adults with epilepsy mean 38.3 years in the SANAD trial. Arm a: carbamazepine vs. gabapentin vs. lamotrigine vs. oxcarbazepine vs. topiramate. Arm b: valproate vs. lamotrigine (LTG) vs. vs topiramate (TPM) <a href="#">Marson 2007 (2)</a>	UK hospital outpatient departments	<a href="#">Marson 2007</a> †
	Back pain	Adults with low back pain aged 18-65 years in a trial comparing exercise manipulation vs. exercise plus manipulation <a href="#">UK BEAM 2004</a>	UK primary care	<a href="#">Letley 2000</a> ‡

(Continued)

	Back pain	Adults aged 18-65 years in a trial comparing yoga vs. usual care <a href="#">Tilbrook 2011</a>	UK primary care	<a href="#">Man 2011</a>
Screening	Prostate, lung, ovarian, colorectal cancer	Adults aged 55-74 years in PLCO trial comparing PSA and CA125 at baseline, and annually for 5 years. Digital rectal examination, transvaginal ultrasound and chest x-ray at baseline and 5 years vs. usual follow-up <a href="#">Prorok 2000</a>	USA trial sites	<a href="#">Subar 2001</a> , <a href="#">Ford 2006</a> (African American men aged $\geq 55$ years only from PLCO)
	Cervical	Women with low-grade abnormal cervical smear aged 20-59 years in the TOMBOLA trial: Colposcopy vs. 6-monthly smears <a href="#">TOMBOLA 2009a</a> <a href="#">TOMBOLA 2009b</a>	UK	<a href="#">Sharp 2006a-h</a>
	Postnatal depression	Women child-bearing aged > 18 years < 2 weeks' postpartum at high risk of postnatal depression in a trial comparing proactive individualised telephone-based peer support vs. standard postpartum care <a href="#">Dennis 2009</a>	Canada community	<a href="#">Kenton 2007a-d</a>
Prevention	Fracture	Adults with history of osteoporotic fracture > 70 years in the RECORD trial: oral calcium + vitamin D vs. oral calcium vs. vitamin D vs. placebo <a href="#">RECORD 2007</a>	UK hospital	<a href="#">MacLennan*</a>
	Fracture	Adults with history of osteoporotic fracture aged > 70 years in the RECORD trial: oral calcium + vitamin D vs.	UK hospital	<a href="#">Avenell 2004</a>

(Continued)

		oral calcium vs. vitamin D vs. placebo <a href="#">RECORD 2007</a>		
	Fracture	Women with hip fracture risk factors aged > 70 years in a fracture prevention trial of calcium 1000 mg plus vitamin D <sub>3</sub> 800 IU plus information sheet on dietary calcium intake and falls prevention vs. information sheet <a href="#">Porterhouse 2005</a>	UK primary care	<a href="#">Cockayne 2005</a>
	Migraine	Adults history of 2 migraine attacks aged 18-65 years with migraine randomised to true diet vs. sham diet (unpublished)	UK community	<a href="#">Ashby 2011</a>
	Lung cancer	Adults exposed to smoking and asbestos aged > 45 years in the CARET trial 2 x 2: beta-carotene + retinol daily vs. beta-carotene vs. retinol vs. placebo <a href="#">Omenn 2006</a>	USA trial sites	<a href="#">Bowen 2000abc</a>
	Breast cancer	Women with 50% of breast volume dysplasia > 30 years in Canadian diet and cancer prevention trial. Counselling and individualised dietary prescription vs. taught principals of a healthy diet not counselled to change fat content <a href="#">Boyd 2002</a>	Canada Hospital clinic	<a href="#">Sutherland 1996</a>
Clinical management	Asthma	Adult with asthma > 70 years in COGENT trial: computerised decision support guidelines for asthma vs. angina care <a href="#">Eccles 2002</a>	UK primary care	<a href="#">McCull 2003</a>

(Continued)

	Asthma and diabetes	Adults with asthma mean age 47 years. Study template for diabetes vs. study template for asthma <a href="#">Tai 1999</a>	UK primary care	<a href="#">Tai 1997</a>
	Angina	Adult with asthma aged > 70 years in the COGENT trial: computerised decision support guidelines for asthma vs angina care <a href="#">Eccles 2002</a>	UK primary care	<a href="#">McCull 2003</a>
	Orthopaedics	Adults with non-surgical musculoskeletal condition $\geq 18$ years in OMENS trial: orthopaedic patients management from single musculoskeletal medicine physician vs. orthopaedic surgeon-led management <a href="#">Leigh Brown 1997</a>	UK Hospital orthopaedic outpatient department	<a href="#">Leigh Brown 1997</a>
Other areas	Exercise	Women sedentary 50-70 years. SWEAT2 trial: moderate walking programme vs. swimming programme <a href="#">Cox 2008</a>	Australia community	<a href="#">Cox 2008</a>
	Parenting	Adults referred for parenting mean age 29 years. Parent-child interactive therapy vs. standard didactic parenting condition <a href="#">Chaffin 2009</a>	USA community	<a href="#">Chaffin 2009</a>
	Weight management	Adults with BMI > 25 aged > 18 years. Internet-based tailored weight management materials vs. Internet-based non-tailored user navigated weight management materials <a href="#">Rothert 2006</a>	USA community	<a href="#">Couper 2007</a>

(Continued)

	Effect of antibiotics on neonatal outcomes	Women < 37 weeks' gestation in ORACLE 1 + 2 trial: 2 x 2 factorial: co-amoxiclav + erythromycin vs. co-amoxiclav vs. erythromycin vs. placebo; 4 times daily x 10 days or until birth <a href="#">Kenyon 2005</a>	UK secondary care/community	<a href="#">Kenyon 2005</a>
	Health promotion	People aged 16-20 years in the sex unzipped pilot feasibility trial: interactive intervention website vs. information only website (main trial unpublished)	UK on line	<a href="#">Bailey 1*</a>
	Health promotion	People aged 16-20 years in the sex unzipped pilot feasibility trial: interactive intervention website vs. information only website (main trial unpublished)	UK on line	<a href="#">Bailey 2*</a>

\* unpublished; ‡ unpublished abstract; † report; published abstract; appendix

AUDIT: Alcohol Use Disorders Identification Test; AVID: Antiarrhythmics Versus Implantable Defibrillators; BMI: body mass index; CARET: Carotene and Retinol Efficacy Trial; COGENT: COlorectal cancer GENeTics; CRASH: Clinical Randomisation of an Antifibrinolytic in Significant Haemorrhage; IU: international unit; OMENS: ; ORACLE: ; PLCO: Prostate, Lung, Colorectal, Ovarian; PSA: prostate-specific antigen; RECORD: Randomised Evaluation of Calcium Or vitamin D; SANAD: Standard And New Antiepileptic Drugs; SWEAT: Sedentary Women Exercise Adherence Trial; TOMBOLA: Trial Of Management of Borderline and Other Low-grade Abnormal smears

## CONTRIBUTIONS OF AUTHORS

VB wrote the protocol for the review with comments from JT, GR, SS, SM, SH and IN. JT and VB designed the searches with comments from SH. VB conducted the searches and screened all abstracts and full papers of potentially eligible trials. VB and GR screened potentially eligible trial papers. SS acted as a third reviewer. Data extraction was conducted by VB and checked by JT. JT designed the analysis plan with VB. VB conducted the analysis with advice on interpretation of results from JT, SS, IN and GR. VB wrote the first draft of the review with comments from all authors.

## DECLARATIONS OF INTEREST

None known.

## SOURCES OF SUPPORT

### Internal sources

- None, Not specified.

### External sources

- None, Not specified.

## DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We grouped trials according to: the type of strategy used, whether the strategy was compared with usual follow-up or other strategies. The diversity of trials and interventions identified in the review were not anticipated. Therefore, the prespecified analyses were not appropriate. Instead, new subgroups were defined prior to analyses. We had planned to include trials that were targeted at treatment or follow-up compliance. We have only included trials that targeted follow-up compliance as the strategies used are transferable to other trials. We had planned to assess whether retention was immediate or longer term (e.g. if a response to a questionnaire was expected immediately or at time points in the future). Time points were poorly reported, where these were reported we used data for this analysis for the primary outcome time point, there was insufficient reporting of this variable to group other time points for further analysis. We had also planned to group participant or management-focused strategies. Only one unpublished trial (Land) evaluated a management-focused strategy to reduce attrition.

As treatment compliance was not a focus of this review, search strategies with the terms 'compliance' were removed for the 2009 to 2012 updates. Most untruncated 'response' search terms were removed because hits relating to 'response' were captured by the search term 'response\*'. To avoid references to treatment response, the search term '*questionnaire*' was added to 'response' in all remaining search terms with 'response' or 'response\*' to make the search specific to questionnaire response. Search updates (2009 to 2012) for EMBASE, MEDLINE and PsycINFO were de duplicated in Ovid. MEDLINE and EMBASE records were excluded for search updates in CENTRAL. C2 Spectre and ERIC searches were not updated from 1 May 2009, as C2-SPECTR [geb9101.gse.upenn.edu/](http://geb9101.gse.upenn.edu/): was not accessible and the search platform for ERIC changed from Datastarweb to Proquest in December 2011. The latter limits searches to 10 lines of text.

## INDEX TERMS

### Medical Subject Headings (MeSH)

Case Management; Correspondence as Topic; Patient Compliance [psychology; \*statistics & numerical data]; Patient Dropouts [statistics & numerical data]; Questionnaires; Randomized Controlled Trials as Topic [\*statistics & numerical data]; Reward



## MeSH check words

Humans