



Citation for published version:

McKimm-Breschkin, JL, Barrett, S, Pilling, PA, Hader, S, Watts, A & Streltsov, V 2018, 'Structural and functional analysis of anti-influenza activity of 4-, 7-, 8- and 9-deoxygenated 2,3-difluoro-N-acetylneuraminic acid derivatives', *Journal of Medicinal Chemistry*, vol. 61, no. 5, pp. 1921–1933.
<https://doi.org/10.1021/acs.jmedchem.7b01467>

DOI:

[10.1021/acs.jmedchem.7b01467](https://doi.org/10.1021/acs.jmedchem.7b01467)

Publication date:

2018

Document Version

Peer reviewed version

[Link to publication](#)

This document is the Accepted Manuscript Version of a Published Work that appeared in final form in *Journal of Medicinal Chemistry*, copyright (C) American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see: <http://dx.doi.org/10.1021/acs.jmedchem.7b01467>

University of Bath

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Article

Structural and functional analysis of anti-influenza activity of 4-, 7-, 8- and 9-deoxygenated 2,3-difluoro-N-acetylneuraminic acid derivatives

Jennifer Lois McKimm-Breschkin, Susan Barrett, Patricia A. Pilling, Stefan Hader, Andrew G Watts, and Victor A Streltsov

J. Med. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.jmedchem.7b01467 • Publication Date (Web): 03 Feb 2018

Downloaded from <http://pubs.acs.org> on February 13, 2018

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1
2
3 **Structural and functional analysis of anti-influenza activity of 4-, 7-, 8- and 9-deoxygenated**
4
5 **2,3-difluoro-N-acetylneuraminic acid derivatives**
6
7
8
9

10 **Authors**

11 Jennifer L. McKimm-Breschkin^{a,*}, Susan Barrett^a, Patricia A. Pilling^a, Stefan Hader^b, Andrew G.
12
13 Watts^b and Victor A. Streltsov^c
14
15
16
17
18

19 **Affiliations**

20
21 ^a CSIRO Manufacturing, 343 Royal Parade, Parkville, 3052 Australia
22
23

24 ^b Department of Pharmacy and Pharmacology, University of Bath, Claverton Down, Bath BA2
25
26 7AY, United Kingdom
27

28 ^c The Florey Institute of Neuroscience and Mental Health, 30 Royal Parade, Parkville, VIC 3052,
29
30 Australia
31
32
33
34

35 ***Corresponding author**

36 Jennifer L McKimm-Breschkin

37
38 jmbvirology@gmail.com
39
40
41
42
43

44 Phone +613 8344 1106
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Competitive inhibitors of the influenza neuraminidase (NA) were discovered almost 20 years ago, with zanamivir and oseltamivir licensed globally. These compounds are based on a transition state analogue of the sialic acid substrate. We recently showed that 5-(Acetylamino)-2,3,5-trideoxy-2,3-difluoro-D-erythro- β -L-manno-2-nonulopyranosonic acid (DFSA) and its derivatives are also potent inhibitors of the influenza NA. They are mechanism based inhibitors, forming a covalent bond between the C2 of the sugar ring and Y406 in the NA active site, thus inactivating the enzyme. We have now synthesized a series of deoxygenated DFSA derivatives in order to understand the contribution of each hydroxyl in DFSA to binding and inhibition of the influenza NA. We have investigated their relative efficacy in enzyme assays *in vitro*, in cell culture and by X-ray crystallography. We found loss of the 8- and 9-OH had the biggest impact on the affinity of binding and antiviral potency.

1. Introduction

There are two types of influenza which cause significant morbidity and mortality in humans, influenza A and B. Both influenza A and B viruses have two surface glycoproteins, the hemagglutinin (HA) responsible for binding to sialic acid containing receptors on the cell surface, and the neuraminidase (NA) enzyme which is responsible for cleavage of sialic acids from the cell surface, to facilitate the spread of progeny virions. Influenza A is further divided into subtypes depending on the HA (18 subtypes) and NA (11 subtypes). Two influenza A subtypes are endemic in humans, (H1N1) and (H3N2). Zoonotic transmission may occasionally occur from birds and swine, including the highly pathogenic avian (H5N1) and (H7N9) viruses and the A(H1N1)pdm09 pandemic swine virus.

There are 9 residues in the NA active site which are highly conserved across all influenza A and B and which interact directly with substrate. These in turn are stabilized by a second shell of 10-11 highly conserved residues, primarily by an extensive network of hydrogen bonds.¹⁻⁴ These are defined as framework as they make no direct contact with the substrate but hold the functional residues in place for binding and catalysis.

There are now four drugs which target the NA, neuraminidase inhibitors (NAIs), which due to the high conservation of the NA active site are effective against all strains of influenza. Four NAIs are licensed in various parts of the world for the treatment and prevention of influenza, oseltamivir, zanamivir (**1**) (Figure 1), peramivir and laninamivir,⁵ a long acting derivative of zanamivir. By inhibiting the influenza NA they prevent virus release, resulting in aggregation of viruses at the cell surface.⁶ Because of differences in the chemistry of the NAIs and subtle differences in NA structures they can bind with a different affinity to the same subtype. e.g. oseltamivir has lower IC₅₀s than zanamivir for N2 NAs, whereas zanamivir has lower IC₅₀s than

1
2
3 oseltamivir for N1 and influenza B NAs^{5,7}. Additionally, due to these differences, resistance can
4 be both NAI and subtype specific.^{5,7} While mutations conferring resistance are primarily in
5 active site residues, mutations remote from the active site can also confer resistance.⁵ Thus, in
6 developing novel NAIs they need to be evaluated against different influenza types and subtypes
7 and against different known resistant mutants of influenza.

8
9
10 We recently demonstrated that the difluorosialic acid (DFSA) mechanism based inhibitor 5-
11 (Acetylamino)-2,3,5-trideoxy-2,3-difluoro-D-erythro- β -L-manno-2-nonulopyranosonic acid (**2**)
12 (Figure 1) and its C-4 amino- and guanidino- derivatives, could inhibit the NA enzyme activity
13 and replication of influenza viruses both *in vitro* and *in vivo*.⁸ We showed by X-ray
14 crystallography that **2** leads to inactivation of enzyme activity by the formation of a covalent
15 bond between the C-2 of the sugar ring and the hydroxyl group of Y406 in the influenza enzyme
16 active site. This covalently linked sialosyl-enzyme intermediate was trapped along with a
17 transition-state analogue resembling an oxocarbenium ion.⁹ Also, the secondary non-catalytic,
18 receptor binding site was occupied by DFSA in the influenza N9 NA–DFSA complex.

19
20
21 In addition to the 4-amino and 4-guanidino derivatives of **2**, we have now synthesized a series of
22 deoxygenated DFSA derivatives **3–6** (Figure 1) in order to understand the contribution of each
23 hydroxyl in binding and inhibition of these sialic acid based inhibitors to the influenza NA.¹⁰

24
25
26 The influenza strains selected for evaluating the impact of removal of the OH groups were those
27 of clinical importance to humans, including the endemic human strains seasonal A(H1N1),
28 pandemic (H1N1)pdm09, A(H3N2) and B, and avian A(H5N1) and the A(H1N9) reassortant, as
29 a model for the avian A(H7N9) NA. NAI resistant mutants of some of these viruses, with
30 different resistance profiles, were also used in the enzyme assays. Some of these were previously
31 tested against other DFSA inhibitors⁸ and interestingly not all mutants were resistant to the
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 DFSA inhibitors. Among the resistant viruses tested, the impacts of the amino acid substitutions
4 on sensitivity to the known NAIs are as follows: D197E in influenza B NA affects binding of all
5 sialic acid derived NAIs, due to altered interactions of the N-acetyl group on the ring with
6
7 R152,¹¹ H274Y in N1 NAAs affects rotation of E276 necessary for binding the pentyl ether side
8 chain of oseltamivir¹² with only a small impact on binding of zanamivir, E119V in N2 NAAs
9 affects interactions of the 4-amino group on oseltamivir with the active site, E119G in N9 NAAs
10 specifically confers resistance to zanamivir, due to changes in the solvent structure and altered
11 interactions of G119 with the 4-guanidino group.¹³ An H252Y substitution between avian
12 (H5N1) clade 1 and 2 viruses leads to reduced oseltamivir sensitivity.¹⁴

13
14 The NAIs oseltamivir and **1** are competitive inhibitors and a fluorescence based enzyme
15 inhibition assay is routinely used for evaluating the sensitivity of influenza virus NAAs *in vitro*.⁷
16
17 ¹⁵⁻¹⁷ However, normally only a single IC₅₀ value is calculated after 30-60 min.^{7, 17} While a single
18 IC₅₀ value provides details on the relative affinity of these NAAs it does not provide any
19 information on the relative rates of drug binding. As compound **1** is a competitive inhibitor, its
20 efficacy is strongly reflected by its affinity (IC₅₀) for the NA. However, since compounds **2-6** are
21 mechanism based inhibitors,⁸ their efficacy is dependent not only on the affinity, but upon both
22 the rate of inactivation, (dependent on both the Michaelis complex formation as well as the
23 subsequent covalent bond formation), and the rate of reactivation (breakdown of the covalent
24 intermediate followed by dissociation of the product). Kinetic analyses of inhibitor binding and
25 dissociation requires large amounts of purified enzyme/virus and are labour intensive,^{8, 18}
26 limiting the numbers of virus/inhibitor combinations that can be readily assayed.

27
28 We have developed two microtiter fluorescence based enzyme inhibition assays which use
29 unpurified virus, and instead of taking a single reading, monitor the reactions in real time. The

1
2
3 IC₅₀ kinetics assay monitors the changes in IC₅₀ values for each 10 min over 60 min, in two
4
5 separate reactions with and without pre-incubation. This provides information on not only
6
7 relative affinity (IC₅₀), but the relative rates of binding and dissociation of each of the
8
9 compounds^{16, 19} A solid phase reactivation assay monitors the time to recover activity after
10
11 removal of inhibitor over several hours.¹⁹ We have previously used these assays to demonstrate
12
13 slow binding kinetics and rates of dissociation of the NAIs consistent with those observed using
14
15 purified NA in the classic assays.^{16, 19-22} Additionally, we have shown that while wild type
16
17 viruses generally demonstrate slow binding of the known NAIs, the NAs of resistant mutants
18
19 demonstrate rapid binding and dissociation of the NAIs.^{11, 15, 16, 19, 23, 24}
20
21
22
23 We have used these assays here to compare relative inhibition of the NAs of a large panel of wild
24
25 type and drug resistant mutants by each of the deoxy derivatives **3-6**. For the sake of simplicity,
26
27 we will refer to inactivation as binding and reactivation as dissociation as we are also comparing
28
29 them to the binding and dissociation of competitive NAIs.
30
31
32
33 In addition, we have evaluated the efficacy of **3-6** to inhibit replication of wild type viruses in
34
35 cell culture in a plaque reduction assay. We show that loss of the 8- and 9-OH have the biggest
36
37 impact on the affinity of binding and antiviral potency both *in vitro* and in cell culture.
38
39
40 We have also used structural analysis by X-ray crystallography of the N9 NA in complex with
41
42 each of the inhibitors, to help explain the biological impact of the removal of the OH groups on
43
44 binding. While we have used the G70C (H1N9) virus as a model virus and its NA for structural
45
46 studies for many years, the N9 subtype has become more relevant due to the emergence of a
47
48 highly lethal (H7N9) virus in China.^{25, 26}
49
50
51
52
53

54 **2. Results**

55
56
57
58
59
60

2.1 IC₅₀ Kinetics

Compounds **1-6** were all evaluated in the microtiter based IC₅₀ kinetics assays against each of the wild type and mutant viruses in the panel. IC₅₀ values for each time point were calculated as the concentration of inhibitor which reduced the NA enzyme activity by 50% compared to the control uninhibited enzyme activity. We used zanamivir as a reference for a slow binding inhibitor with high affinity.¹⁶ The greater the difference in final (60 min) IC₅₀ values between the reactions without and with preincubation (-/+ ratio), the slower the binding (Figures 2 and 3, Table 1). We have previously shown for the NAIs that fast binding inhibitors have a -/+ ratio of around 1 for the 60 min IC₅₀ values, and slower binding inhibitors have ratios of >1.5-2, indicating that preincubation enhances binding. Additionally, the rate of increase of the IC₅₀ values after preincubation with inhibitor upon addition of MUNANA substrate, gives a qualitative indication of the dissociation rate. The faster the increase, i.e. the greater the difference between the 10 min and 60 min IC₅₀ values, the faster the dissociation. A lower ratio indicates slower dissociation (Figures 2 and 3, Supporting information Table S1). All assays were carried out in duplicate as described in the Experimental section. The IC₅₀ value for each 10 min time point is the mean of the individual IC₅₀ values of the duplicates for that virus-inhibitor reaction. The profiles of the bar graphs (Figures 2 and 3) provide an easy overview of changes in IC₅₀ values for the large number of virus and compound combinations, showing whether the IC₅₀ values increase or decrease with and without preincubation over the 60 min reaction time, indicative of fast or slow binding/dissociation. Comparisons of the 60 min IC₅₀ values without and with preincubation, as an indication of relative slow or fast binding are all presented in Table 1. Comparisons of the change in IC₅₀ values between 10 and 60 min, as an indication of the relative dissociation rate, are all presented in Supporting information, Table S1.

2.1.1 N1 NAs

The compounds were tested against the seasonal (H1N1) and H274Y oseltamivir resistant viruses, swine and human isolates of (H1N1)pdm09 viruses, and two avian (H5N1) viruses, which differ in oseltamivir sensitivity. None of the inhibitors showed the classic slow binding profile of the bar graphs seen with zanamivir (**1**), where the IC_{50} values continue to decrease without preincubation from 10 to 60 min, and are significantly lower after preincubation (Figure 2, Table 1).

Despite large sequence differences between the (H1N1)pdm09 and (H5N1) NAs their kinetics profiles were very similar for each of the inhibitors (Figures 2B, 2C), but both were different to the seasonal (H1N1) NA and its H274Y mutant (Figure 2A). Consistent with previous findings, clade 1 (H5N1) NA showed faster dissociation of **1** than clade 2 NA^{16, 19} (Figure 2C and Supporting information Table S1).

IC_{50} kinetics profiles for compounds **2** and **3** were similar to those previously seen with NAI resistant mutant NAs.¹⁶ They showed fast binding/dissociation to both the (H1N1)pdm09 and (H5N1) NAs, with similar IC_{50} values with or without preincubation. In contrast, **2** demonstrated slow binding to the seasonal (H1N1) NA, while the loss of 4-OH increased both the affinity and rate of binding for compound **3** (Figure 2 and Table 1).

Removal of the 7-OH resulted in slightly higher affinity of **4** for the wild type seasonal (H1N1) NA and slow binding. As seen with **2**, loss of the 7-OH resulted in reduced affinity to the H274Y mutant NA. However, unlike previous mutant NAs, it had different binding kinetics. The IC_{50} values were similar with or without preincubation, but continued to decrease in both reactions. Rather than fast binding this indicated slower binding/dissociation (Figure 2A, Table 1, Supporting information Table S2).

1
2
3 Loss of the 8-OH resulted in compound **5** binding with a much lower affinity to all the N1 NAs
4 compared to compounds **2**, **3** or **4**. Kinetics showed an unusual pattern of binding to the
5
6 (H1N1)pdm09 and clade 2 (H5N1) NAs, where the IC₅₀ values were not significantly different
7
8 with or without preincubation, however they continued to decrease after the addition of substrate
9
10 (Figure 2B, C). We have not previously observed this with other NAIs binding to wild type NAs,
11
12 but it suggested there was slower turnover, possibly because of both slow binding and slow
13
14 dissociation. In contrast, there was an increase in IC₅₀ values for the clade 1 (H5N1) NA after
15
16 preincubation with **5**, indicating a faster dissociation (Figure 2C, Supporting information Table
17
18 S1).
19
20
21
22

23
24 Loss of the 9-OH also had a significant impact on the affinity of the compounds, with a 1-2 log₁₀
25
26 increase in IC₅₀ values for all the N1 NAs. There were larger differences in the IC₅₀ values with
27
28 and without incubation for the seasonal (H1N1) and the H274Y mutant NAs with both **5** and **6**,
29
30 indicating slower binding compared to some of the other compounds. Except for the clade 1
31
32 (H5N1) NA the IC₅₀ values continued to decrease with or without preincubation for these two
33
34 compounds. This suggested both **5** and **6** had a slow turnover.
35
36

37 **2.1.2 N2 NA**

38
39 IC₅₀ kinetics for the H3N2 and oseltamivir resistant E119V NAs showed loss of the 4-OH
40
41 improved affinity (lower IC₅₀ values) of **3** binding to the wild type H3N2 NA, compared to **2**
42
43 although it was still fast, with no significant effect of preincubation on IC₅₀ values (Figure 3A,
44
45 Table 1). While **2** binds with higher affinity to the E119V mutant than to wild type, in contrast **3**
46
47 binds with lower affinity to the mutant NA.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Loss of the 7-OH improved affinity, with lower IC₅₀ values for **4** than for **2** for both the wild type
4 and E119V mutant NAs. As seen for **2**, compound **4** bound with higher affinity to the mutant
5 NA, and it also showed slower binding than to the wild type NA.
6
7

8
9
10 While loss of either the 8-OH or 9-OH resulted in significantly reduced affinity, IC₅₀ values were
11 similar for each of the compounds for both the wild type and E119V mutant. Compound **5**
12 showed the classic slow binding profile for both wild type and E119V NAs, with about a 10-fold
13 decrease in IC₅₀ values after preincubation, and slow dissociation, with a small increase in IC₅₀
14 values with time after addition of substrate. Compound **6** showed the unusual profile, similar to
15 the seasonal (H1N1), with the IC₅₀ values continuing to decrease with or without preincubation
16 (Figure 3, Table 1, Supporting information Table S1).
17
18
19
20
21
22
23
24

25 26 **2.1.3 N9 NA**

27
28 The G70C H1N9 reassortant was used to evaluate inhibition of the N9 NA, as a model for the N9
29 NA of the pathogenic (H7N9) viruses recently spreading in China.^{25, 26} Compound **2** was fast
30 binding to the wild type N9 NA. Although the IC₅₀ values were comparable between the E119G
31 mutant and wild type NAs (Figure 3B, Table 1 and Supporting information Table S2) the bar
32 graph profiles showed the IC₅₀ values decreased with or without preincubation for the E119G
33 mutant NA, suggesting slower turnover. In contrast to **2**, compound **3** demonstrated higher
34 affinity, but slower binding to the wild type N9 NA (Figure 3B, Table 1). While the E119G
35 mutant showed the classic resistance profile of higher IC₅₀ values, fast binding and dissociation,
36 with IC₅₀ values increasing with or without preincubation, as seen for this mutant with
37 zanamivir¹⁶ (Table 1, Supporting information Tables S1, S2), the loss of the 4-OH did not affect
38 the affinity of binding to this mutant NA compared to **2**. It did however decrease binding of **3**
39 compared to the wild type N9 NA (Supporting information Table S2).
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Compared to **2**, loss of the 7-OH led to a higher affinity of **4** for the wild type N9 NA. In
4
5 contrast, while **4** had reduced binding to the E119G NA compared to the wild type (Supporting
6
7 information Table S2), loss of the 7-OH had no impact on binding to this mutant NA compared
8
9 to **2**. (Supporting information Table S1).

10
11
12 Both **5** and **6** demonstrated a further decrease in IC₅₀ values with or without preincubation over
13
14 the 60 min reaction time, indicating slow turnover. As seen for all the other viruses the loss of
15
16 either the 8-OH or the 9-OH reduced their affinity. The E119G substitution also reduced binding
17
18 of both inhibitors, compared to the wild type NA and compared to **2** (Supporting information,
19
20 Table S2). This reinforces the importance of both the 8- and 9-OH groups for high affinity
21
22 binding. It further highlights the complex interactions involved in ligand binding, since the 8-
23
24 and 9-OH groups are on the opposite side of the active site, compared to E/G119.

28 29 **2.1.4 B NA**

30
31 The B/Perth viruses were used to evaluate efficacy against wild type and broadly NAI resistant
32
33 influenza B NAs. The D197E mutation results in reduced affinity and faster dissociation of **1**
34
35 (Figure 3C, Table 1, Supporting information Table S1), and because the D197E substitution
36
37 affects interaction of R152 with the N-acetyl group on the sugar ring, it would be expected to
38
39 affect binding of all sialic acid based inhibitors.

40
41
42 Compared to **2**, loss of the 4-OH, as with the influenza A NAs, led to higher affinity of **3** (Figure
43
44 3C). Unexpectedly **3** bound with similar affinity to both the wild type and E197 mutant NA,
45
46 somehow compensating for the reduced binding due to the altered E197/R152 interactions with
47
48 the N-acetyl group (Figure 3C, Supporting information Table S2). Both showed the classic slow
49
50 binding profile, with IC₅₀ values after preincubation being lower than without preincubation
51
52 (Figure 3C, Table 1).

1
2
3 Compounds **4**, **5** and **6** all had reduced affinity (higher IC_{50} values) for the mutant NA compared
4
5 to the wild type NA, (Supporting information Table S2). However, unlike **1**, compounds **2-6** still
6
7 demonstrated slow binding to the B/Perth wild type and mutant NA (Table 1 and Supporting
8
9 information Table S1).

10
11
12 Loss of the 7-OH resulted in a marginally higher affinity compared to **2** (Table 1), but loss of the
13
14 8- or 9-OH groups again reduced affinity of **5** and **6** for both wild type and the D197E NAs,
15
16 while maintaining the slow binding, and slow dissociation seen with other viruses (Table 1,
17
18 Supporting information Table S1).

19
20
21 With the NAIs the ratio of -/+ preincubation has been a good indicator of slow and fast binding.
22
23 However, for compounds **5** and **6** the -/+ ratios in Table 1 clearly do not fully reflect what
24
25 appears to be slow turnover, since the IC_{50} values continue to decrease after preincubation
26
27 indicating additional compound is still binding. This demonstrates the importance of looking at
28
29 the profiles of the IC_{50} kinetics graphs, demonstrating the changes in IC_{50} values over time, rather
30
31 than a single IC_{50} value or -/+ ratio.
32
33

34 35 **2.2. Reactivation**

36
37
38 While our IC_{50} kinetics assays give an indication of the relative rate of dissociation after the
39
40 addition of substrate in the preincubation reaction (Supporting information Table S1), this is still
41
42 in the presence of inhibitor. The classic reactivation experiments react purified NAs or viruses
43
44 with excess inhibitor, then remove the unbound inhibitor by spin columns and monitor recovery
45
46 of enzyme activity after addition of substrate.^{8, 18, 27-29} However this requires significant
47
48 quantities of virus or NA and is labor intensive. We had also seen a different pattern for the IC_{50}
49
50 kinetics especially for compounds **5** and **6** than previously seen with other NAIs binding to wild
51
52 type virus NAs, where although preincubation did not lead to obvious enhanced binding, the IC_{50}
53
54
55
56
57
58
59
60

1
2
3 values continued to decrease either with or without preincubation. We therefore used our 96-well
4 solid phase reactivation assay^{19, 24} to test dissociation of **2-6** from the wild type virus NAs as a
5 further confirmatory assay to determine whether the novel profiles for **5** and **6** did reflect very
6 slow dissociation compared to **2-4**. The relative rate of reactivation is expressed as a $T_{1/2}$ which is
7 the time required for reaching half the maximum average rate of the uninhibited control. Results
8 are shown in Table 2. The $T_{1/2}$ results for each virus and inhibitor were compared to the $T_{1/2}$ for
9 the virus controls without inhibitor, by the Student's *t* test. Consistent with the IC_{50} kinetics,
10 reactivation was significantly slower for compound **1** for all of the NAs ($P < 0.001$).¹⁹

11
12 For the (H1N1) A/Mississippi seasonal strain, reactivation was significantly slower for **2-6**, with
13 **6** being the slowest.

14
15 For the (H1N1)pdm09 and both (H5N1) strains, reactivation after removal of **2** and **3** was very
16 rapid, with $T_{1/2}$ values barely more than 10 min. In fact, some ultimately showed an enhanced
17 rate of MUNANA cleavage compared to the control with no inhibitor. This is consistent with the
18 IC_{50} kinetics showing similar values with or without preincubation. Reactivation after removal of
19 **3** was marginally faster than reactivation after removal of **4** for each of these viruses, but was not
20 significantly faster. Turnover of **5** and **6** were slower, as also seen in the IC_{50} kinetics, although
21 only significantly slower for the (H1N1)pdm09 and (H5N1) clade 2 viruses.

22
23 For the H3N2 NA, in contrast to compound **1**, reactivation after removal of compounds **3** and **4**
24 was very fast, comparable to the control. Reactivation after removal of compound **2** was just
25 significantly slower than the control. Turnover of **5** and **6** were significantly slower, again
26 consistent with the IC_{50} kinetics showing decreases in IC_{50} values in both reactions.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 For the NWS/G70C N9 NA reactivation was very rapid after removal of compounds **2**, **3** and **4**
4 while reactivation after removal of **5** and **6** was significantly slower reflecting the results in the
5
6 IC₅₀ kinetics where the IC₅₀s continued to decrease in the preincubation reaction.
7

8
9
10 Reactivation of the B/Perth NA activity after removal of the inhibitors was slowest of all of the
11
12 inhibitors **2-6**, even slower than for **1**. Results were consistent with the IC₅₀ kinetics, with **3**
13
14 having the fastest turnover of all the deoxy inhibitors for B/Perth, and **5** and **6** having the slowest
15
16 turnover of any virus NA-DFSA combination.
17

18 19 **2.3. Plaque reduction assay (PRA)**

20
21
22 We have demonstrated in enzyme assays that while the loss of the 4-OH generally leads to lower
23
24 affinity, it results in faster turnover. In contrast loss of the 8- and 9-OH groups result in lower
25
26 affinity, but slower turnover. Since the ability of the covalent DFSA inhibitors to inhibit virus
27
28 replication will be a combination of the relative affinity as well as their turnover we tested their
29
30 efficacy against a selection of the wild type viruses in the PRA to see whether there was greater
31
32 correlation with the slow turnover of affinity. The EC₅₀ is the effective concentration of inhibitor
33
34 to reduce plaque sizes by 50% compared to the uninhibited control. We tested the
35
36
37 A/Mississippi/03/01 seasonal (H1N1) virus, the G70C H1N9 reassortant, the B/Perth/211/2001
38
39 virus and two different human H3N2 viruses, A/Fukui/45/04 and A/Victoria/503/06, as some
40
41 H3N2 are known to have lower sensitivity to the other NAIs in cell culture, due to lower affinity
42
43 of their HA. This lowers their dependence on the NA, making them appear less sensitive in a
44
45 PRA, due to the altered HA and NA balance.
46
47

48
49 Inhibition of the (H1N1) A/Mississippi virus was comparable for compounds **2**, **3** and **4** in the
50
51 PRA (Table 3, Supporting information Figure S1). Despite very slow dissociation of **5** its
52
53
54
55
56
57
58
59
60

1
2
3 efficacy was less than that of the other inhibitors, and **6** was the least potent, reflecting the
4
5 relative IC₅₀ values in the enzyme inhibition assays.
6

7
8 All inhibitors, including **1** showed lower potency against the H3N2 A/Fukui/45/04 (Table 3)
9
10 virus in the PRA compared to the other viruses. This is likely due to the HA having a lower
11
12 affinity for the sialic acid receptor, hence less NA activity is needed for virus to elute and
13
14 spread.³⁰ Although the IC₅₀ values in the enzyme assay were lower for compounds **3** and **4**
15
16 compared to compound **2**, there was no difference detectable in their potency in the PRA.
17

18
19 Inhibition by **5** was even poorer, reflecting its lower efficacy in the enzyme assay, as we saw no
20
21 inhibition even at 100 μM in the PRA.
22

23
24 Since the A/Victoria/503/06 H3N2 virus was more sensitive to zanamivir in the PRA, we also
25
26 tested this virus with the DFSA panel. Both **3** and **4** showed higher potency compared to DFSA
27
28 **2**, with lower efficacy of **5** and **6** (Table 3). This is consistent with the enzyme assay results seen
29
30 with the A/Fukui/45/04 N2 NA.
31

32
33 All the inhibitors were mostly more potent against the B/Perth virus in the PRA than against the
34
35 other influenza A viruses tested (Table 3), which was also observed in the IC₅₀ kinetics assays.
36
37 Once again, although **2**, **3** and **4** had faster turnover compared to **5** and **6**, they were more potent
38
39 than **5** and **6** with their slower turnover.
40

41
42 Thus, it appears that efficacy of inhibition of virus replication in cell culture primarily reflects
43
44 the affinity of the compound for the NA, rather than the turnover rate.
45
46
47
48

49 **2.4. X-ray crystal structures**

50 **2.4.1 Trapped intermediates in the active site**

51
52
53
54
55
56
57
58
59
60

1
2
3 As seen previously with DFSA, **2**⁹ binding of the deoxy compounds **3-6** resulted in the trapping
4 of intermediates in the N9 NA. There is a clear rotation of the COOH group and loss of the
5 fluorine (F2) at the C2 position of the ligand. The ligands were trapped in the form of the two n-
6 deoxy (n=4,7,8,9) 3-fluorosialyl-enzyme intermediates bound in the active site (Figure 4). A
7 covalent bond of $\sim 1.4 \text{ \AA}$ was observed in the electron density map between C2 of n-deoxy
8 (n=4,7,8,9) 3-fluoro sialic acid and the phenolic oxygen of Y406 (Figure 4) consistent with
9 previous results recently reported.⁸ A covalent bond has also been observed previously in the
10 structures of the bacterial GH33 sialidase NanI³¹, N2 NA with a DFSA³² and N9 NA with a
11 DFSA derivative 3-equatorial-2,3-difluoro-4-guanidino sialic acid (FeqGuDFSA).⁸
12 Again, similar to the **2**-NA N9 complex⁹ in addition to the covalent intermediate species
13 detected, we also identified non-bonded species which we assumed to be unsaturated forms of n-
14 deoxy 3-fluoro sialic acid intermediates with fractional occupancies: 20% / 80% (**3**), 45% / 55%
15 (**4**), 60% / 40% (**5**), and 25% / 75% (**6**), respectively for covalently bonded and non-bonded
16 species, (Figure 4A, C, E, G).

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
The distances between the anomeric C2 of non-bonded forms and the phenolic oxygen of Y406
are $\sim 2.5 \text{ \AA}$, much longer than the corresponding C-O covalent bond lengths observed for the
covalent intermediates. This non-bonded form of the n-deoxy fluorosialic acid intermediates is
consistent with an oxocarbenium ion like transition state in which the carboxylate group is
coplanar with the oxocarbenium ion plane with considerable sp^2 hybridization at the anomeric
carbon. The oxocarbenium ion positive charge at the ring oxygen O6 is stabilized by hydrogen
bonds with waters at $2.7\text{-}3.4 \text{ \AA}$ in all N9 complexes. The oxocarbenium ion is thus sandwiched
between the active site negatively charged residues, in particular the deprotonated tyrosine and
waters. Further, the oxocarbenium ion is also stabilized by electrostatic interactions with the

1
2
3 negatively charged carboxylate group adjacent to the anomeric carbon C2 and the fluorine
4 attached to the C3 carbon. These anionic charges assist with the departure of the negatively
5 charged fluorine leaving group and the formation of a covalent intermediate bound to the
6 phenolic oxygen of tyrosine. Because the oxocarbenium ion has double-bond character between
7 the anomeric carbon C2 and the ring oxygen O6, this part of the structure must be planar. Thus,
8 for a 6-membered ring, C2, O6, C1 and C3 must lie in a plane. This arrangement can be
9 accommodated in a 6-membered ring only in the half chair conformation similar to a glycal
10 DANA generated at a low rate from sialic acid soaked in crystals of the influenza B virus NA at
11 4°C for an extended time.³³ Similarly, the n-deoxy-DFSA fluorinated substrate analogues
12 facilitated trapping enzyme bound species resembling intermediates or “transition states”. The
13 structures of assumed bonded and non-bonded ligands were refined from the observed
14 experimental electron density with stereochemical constraints. The ligands are stabilized by an
15 extended hydrogen bond network involving residues and structured water molecules, most of
16 them are well documented in the NA structures with sialic acid¹ and FeqGuDFSA.⁸ In addition,
17 the fluorines make contacts in the range of 3.0-3.9 Å with the side chain nitrogen atom of R118
18 as well as short contacts in the range of 2.5-3.5 Å with the carboxyl oxygen of D151. Interactions
19 between fluorine and a protonated nitrogen are well known and have been computed at up to
20 13.5 kcal/mol.⁸

2.4.2 Ligands in additional receptor binding sites

21 The omit³⁴ electron density (Figure 4B, D, F, H) revealed one additional n-deoxy-DFSA
22 (n=4,7,8,9) site formed by three serines: **S367**, **S370** and **S372** (N2 numbering used throughout).
23 This is the known second site in the N9 NA, previously reported to bind sialic acid, which is
24 discrete from the catalytic site.³⁵ The n-deoxy-DFSA position is overlapping well with that for **2**⁹
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 and sialic acid in this second binding site³⁵ confirming that the secondary ligand sites in N9 NA
4
5 are not specific to the sialic acid complex only.
6

7 8 **2.4.3 Compound 3**

9
10 The secondary receptor binding site confirms the original structure of **3** (**Figure 4B**), prior to
11
12 binding and cleavage. According to kinetic assays above **3** binds with higher affinity than **2** to
13
14 the wild type N9 (Figure 3, Table 1). While the N9 E119G mutant NA has reduced binding of **3**
15
16 compared to the wild type, binding is similar to **2**, but shows a different kinetics profile
17
18 suggesting a faster turnover (Figure 3). E119 must provide important interactions, which are lost
19
20 with G119. This is most likely contacts of ~3 Å between the carboxyl groups of E119 and D151
21
22 and the F3 fluorines (Figure 4A). In the case of the wild type N9, the negatively charged 4-OH
23
24 group next to the electronegative F3 seems to destabilize the F3 contacts with E119 and D151
25
26 and the loss of 4-OH improves binding of **3**.
27
28
29

30 31 **2.4.4 Compound 4**

32
33 The crystal structure shows the second site (Figure 4D), again confirming the original **4** structure
34
35 prior to cleavage. The 7-OH group was mainly involved in contacts of ~2.5 Å with two waters as
36
37 seen in **2**⁹ and **3** complexes (Figure 4A). Loss of the 7-OH with those water contacts (Figure 4C)
38
39 would be expected to lead to minor changes in affinity to both the wild N9 and its E119G
40
41 mutant. In fact, **4** binds with slightly higher affinity to the wild type than **2**, and loss of the 7-OH
42
43 has no impact on binding to the mutant G119 NA compared to **2**.
44
45
46

47 48 **2.4.5 Compound 5**

49
50 In contrast to **3** and **4** N9 complexes, which were obtained by a short soak in ligands, we could
51
52 only obtain complexes for **5** and **6** by co-crystallization. The secondary ligand site in the co-
53
54 crystallized **5**-N9 complex showed fragmented electron density (Figure 4F), suggesting
55
56
57
58
59
60

fractional population of this site and reduced binding affinity. Also, in the active site (Figure 4E), 8-OH forms important contacts of 2.5-3.5 Å with COOH groups of E276 and E277, of 3.6 Å with R292 and of ~3Å with a water. Due to loss of the 8-OH there is an upwards displacement of the sugar ring of the ligand to the Y406, but there is also some rotation of both the E276 and E277 compared to other complexes.

2.4.6 Compound 6

Similar to **5**, the complex with **6** was only obtained by co-crystallization and showed high atomic displacement parameters and fragmented electron density for **6** in the secondary ligand binding site (Figure 4H). This again suggests slower binding and reduced affinity as seen for **5**. The active site shows (Figure 4G) loss of contacts between 9-OH and E276 (2.5 Å), R224 (3.3 Å), and a water (2.8 Å). Loss of these important contacts would correlate with the reduced binding affinity of **5** and **6** to not only the N9 but to all other NAs tested.

3. Discussion

Zanamivir (**1**) is reported to be a competitive, slow binding inhibitor of the influenza NAs. It has been suggested that the 4-guanidino side chain in **1** needs to displace water for high affinity binding, hence leading to slow binding.^{20, 36} Without a guanidino substitution at the 4-position, as in 2-deoxy-2,3-didehydro-N-acetylneuraminic acid (DANA) and the 4-amino derivatives of sialic acid, binding to the NA is rapid.¹⁶ Binding of **2** to influenza NAs would thus be expected to be rapid. The aim with a covalent inhibitor is to have high affinity binding, but additionally a slower turnover, since unlike competitive inhibitors, once a covalent inhibitor is hydrolyzed it cannot inhibit another NA.⁸ We were interested in testing the impact of removal of one of each of the 4-, 7-, 8- or 9-OH from **2** on binding affinity (inactivation), and dissociation/turnover

1
2
3 (reactivation) with a variety of different wild type and mutant N1, N2, N9 and influenza B NAs.
4
5 Insight into the roles of each of the OH groups could provide important information on potential
6
7 sites for substitution to enhance potency of the DFSA inhibitors, as well as understanding how
8
9 ligand modifications affect binding to known NAI resistant mutants.
10
11
12
13

14
15 Our previous development of microtiter based assays has enabled us to test a large number of
16
17 combinations of inhibitors and wild type and mutant viruses for their relative potency and
18
19 kinetics of binding and dissociation in enzyme assays.^{11, 15, 16, 19, 23, 24, 37} We observed much more
20
21 variation in affinity and binding kinetics of **2** and **3-6** among different wild type viruses
22
23 compared to **1**, oseltamivir and peramivir.¹⁶ We showed here that while binding of **2** and some of
24
25 the deoxy derivatives did not show the classic profile of slow binding inhibitors in the IC₅₀
26
27 kinetics reaction to all the tested influenza A NAs, some also did demonstrate slow binding.
28
29 Unexpectedly they all showed slow binding to not only the influenza B wild type NA, but also to
30
31 the mutant influenza B NA. Additionally, in contrast to other NAIs where potency against
32
33 influenza B is generally lower than to influenza A NAs, **2** and its deoxygenated derivatives **3-6**
34
35 generally bound with higher affinity to the B/Perth NA, compared to influenza A NAs. This is
36
37 consistent with our previous observations that **2** and its 4-amino and 4-guanidino derivatives
38
39 bound with high affinity to influenza B NA.⁸
40
41
42
43

44
45 We have previously shown that mutations which confer reduced susceptibility to the NAIs ,
46
47 result in loss of slow binding, so that preincubation makes no difference,^{11, 15, 16, 37} and inhibitors
48
49 dissociate rapidly.¹⁹ However while some of the deoxy DFSA inhibitors did show slow binding,
50
51 and they had reduced affinity for mutant NAs compared to their wild type parent, many still
52
53 showed either slow binding or slow turnover in the mutant NAs. Some of the mutant NAs even
54
55
56
57
58
59
60

1
2
3 bound the compound with higher affinity compared to the wild type NA. Thus, the mutations had
4 minimal impact on binding **3-6**, compared to **2**.

7
8 In general, the loss of the 4-OH resulted in higher affinity binding (lower IC₅₀ values) compared
9 to **2** for wild type NAs, while the loss of the 8- and 9-OH resulted in lower affinity, but slower
10 binding and turnover. We have previously determined that if the IC₅₀ after preincubation (+) is
11 lower than without preincubation (-), then the inhibitor is slow binding (-/+ ratio >1.5-2).
12

13
14 However, here we observed a novel profile in the IC₅₀ kinetics particularly for **5** and **6**, where
15 with or without preincubation the IC₅₀ values continued to decrease during the 60 min reaction
16 time. This suggested a nett slower binding and dissociation, hence a slower turnover, which was
17 not necessarily reflected by the -/+ ratio. This emphasizes the importance of looking at the
18 graphs for the whole reaction time and not just a single IC₅₀. This interpretation was supported
19 by the results of our reactivation assays, which demonstrated a slower reactivation than for the
20 other compounds. We have only observed this unusual pattern of binding with **1** binding to an
21 NAI resistant seasonal (H1N1) with a novel Y155H mutation.¹⁵
22
23

24
25 Additionally, we evaluated the inhibitors in a plaque reduction assay in cell culture, providing
26 information on the nett effect of affinity and turnover on inhibition of virus replication and the
27 importance of each OH group. While theoretically the inhibitor with the slowest turnover might
28 be expected to have the highest nett potency in terms of inhibition of virus replication *in vivo*,
29 this was not the case. Potency was more a reflection of the relative affinity, rather than the
30 turnover. Both **5** and **6** which had the slowest turnover of all the compounds, had both the lowest
31 affinity and the lowest potency. Conversely, **3** which had a rapid turnover, but higher affinity,
32 was more potent than **5** and **6**. Thus, our studies have importantly demonstrated that slowing
33 down turnover of these mechanism based inhibitors will not necessarily result in higher potency.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Structural analysis also supported the observed variation in ligand binding, showing the
4 importance of water molecules in stabilizing ligands in the active site as well as the interactions
5 of the 8-OH and 9-OH with E276/E277/R224/R292. X-ray crystallography demonstrated that
6 like **2**⁹ compounds **3-6** could also bind in the second sialic acid binding site.³⁵ This is a non-
7 catalytic site and the biological role of this site is yet not understood. The NAIs do not bind this
8 site, since they have a DANA boat-like conformation. Only ligands with the sialic acid chair
9 conformation can bind, in a similar way to binding to the HA.⁹ One study has shown that the
10 ability of NA to bind red blood cells via this second site correlated with the cleavage efficiencies
11 of other multivalent substrates.³⁸ Others have more recently shown that substrate binding via the
12 second sialic acid-binding site in (H7N9) viruses enhances NA catalytic efficiency against the
13 same substrate.³⁹ Hence binding in this site may affect affinity of the DFSA inhibitors, but not
14 the kinetics. Thus an inhibitor which can bind both in the secondary sialic acid binding site as
15 well as the NA active site, might have advantages in reducing virus binding and spread,
16 particularly for avian viruses which continue to pose a pandemic threat.

37 38 **4. Conclusions**

39
40 Although the efficacy of covalent inhibitors is dependent on both the affinity and turnover, we
41 have shown here that with the deoxy-DFSA inhibitors **3-6**, the advantages of a slower turnover
42 can be counteracted by lower affinity. Hence the design of covalent inhibitors needs to
43 incorporate modifications leading to both high affinity and slow turnover. Loss of the 8-OH and
44 9-OH had the greatest impact on binding and affinity of the covalent inhibitors tested here.

Experimental Section

Materials

Zanamivir **1** was kindly provided by GSK (Stevenage, UK) and compound **2** was kindly provided by the group of Prof. Steve Withers (Vancouver, BC). Compounds **3-6** were prepared as previously described.¹⁰ The purity of final compounds was $\geq 95\%$. Purity was determined by elemental analysis and is available in the Supporting Information.

Viruses and cells

The following wild type and drug resistant viruses were grown in cell culture as previously described.^{8, 16}: B/Perth/211/2001 wild type and D197E cross-resistant mutant,⁴⁰ A/Mississippi/03/01 seasonal (H1N1) and H274Y oseltamivir and peramivir resistant mutant,⁷ A/Fukui/45/04 H3N2 wild type and E119V oseltamivir resistant mutant,¹⁷ A/Victoria/503/06 H3N2 grown in MDCK cells,⁴¹ NWS/G70C H1N9 and E119G zanamivir and peramivir resistant mutant.^{13, 42} The following irradiated viruses were provided by the CSIRO Australian Animal Health Laboratory for enzyme assays only (H1N1)pdm09 A/California/7/09 and A/Swine/Shepparton/6/2009, and A/Chicken//Vietnam/08/2004 (H5N1) clade 1 and A/Chicken/Bangli/BBVD-563/2007 clade 2 (H5N1).³⁷

Plaque reduction assays (PRAs) in Madin Darby Canine Kidney cells (MDCK) were carried out as previously described^{8, 23} using an overlay of DMEM/ F12 in 0.5% immunodiffusion-grade agarose (MP Biomedicals, Australia), containing 1 mg/mL L-1-tosylamido-2-phenylethyl chloromethyl ketone ('TPCK')-treated trypsin (Worthington, USA). Serial ten-fold dilutions of inhibitors were incorporated into the overlay²³ and cells were fixed at 3 days with 1% formalin, and stained with 0.05% neutral red and scanned. The EC₅₀ is the effective inhibitor concentration

1
2
3 causing a 50% decrease in plaque size. Where there was greater than 50% reduction in plaque
4 size between two drug concentrations a range was used.
5
6
7
8
9

10 **Enzyme assays**

11
12 NA enzyme activity was measured using the fluorescent substrate 4-Methylumbelliferyl N-
13 acetyl- α -D-neuraminic acid (MUNANA, Carbosynth, Berkshire, UK). MUNANA was diluted to
14 a final concentration of 100 μ M in 50 mM sodium acetate pH 5.5 and 5 mM CaCl₂. We used a
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

11
12 All viruses were initially titrated to determine a dilution which gave similar NA enzyme activity.
13
14 Inhibition assays were carried out using the constant amount of each virus, incubated with seven
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

11
12 While the IC₅₀ for NAIs is routinely calculated for a single time point after the addition of
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 information on the relative rates of inhibitor binding, and the relative rates of dissociation of
4 many different wild type and mutant viruses and NAIs.^{11, 15, 16, 23, 24, 37} For IC₅₀ kinetics two
5
6 separate inhibition assays were carried out. In the first reaction virus was preincubated with
7
8 serial 10-fold dilutions of inhibitor for 30 min at room temperature, prior to the addition of
9
10 substrate, and in the second reaction virus, serial 10-fold dilutions of inhibitor and MUNANA
11
12 substrate were all co-incubated. Fluorescence was read every minute for 60 min and the percent
13
14 inhibition of enzyme activity for each 10 min time point compared to the uninhibited control was
15
16 plotted against the concentration of inhibitor. The IC₅₀ values were determined for each 10 min,
17
18 hence 6 inhibition curves and 6 IC₅₀ values were calculated for each reaction. Each virus-
19
20 inhibitor combination was tested in duplicate for each of the pre incubation and no preincubation
21
22 reactions. IC₅₀ values were calculated independently for each 10 min time point for each of the
23
24 duplicates and the mean IC₅₀ values were then plotted as bar graphs to demonstrate the changes
25
26 in IC₅₀ with time i.e. the IC₅₀ kinetics.
27
28
29
30
31

32
33 Solid phase reactivation assays were carried out as recently described.^{19, 24} Briefly, after binding
34
35 virus to multiple wells in a black ELISA plate (Greiner Fluorotrack 600), inhibitor was added at
36
37 >10 times the IC₅₀ and incubated for 30 min. Inhibitor was then washed off, MUNANA substrate
38
39 was added, and the reactivation of enzyme activity was monitored after each 10 min for 4 hours.
40
41 The relative rates of activity were calculated compared to the uninhibited virus control for each
42
43 30 min period, and if reactivation was rapid, rates for the initial 10 and 20 min were calculated.
44
45 The T_{1/2} was calculated as the time taken for the rate to reach half the maximum rate of the
46
47 control.¹⁹ Two or three wells for each drug were used per assay in two independent assays.
48
49
50
51
52
53

54 **Statistics**

55
56
57
58
59
60

1
2
3 Data for reactivation assays were analyzed for statistical significance using a Student's *t* test,
4 where a *P* value of <0.05 was considered statistically significant.
5
6
7
8
9

10 **Crystallization**

11
12 NA from influenza virus A/NWS/Tern/Australia/G70C/75 was purified as described previously¹³
13 and crystallized in potassium phosphate buffer (1.7 M, pH 6.7) as previously described.³⁵ The
14 N9-NA **3** and **4** complexes were prepared by soaking crystals in cryo-protectant solution
15 containing crystallization well solution plus 20% glycerol and 2 mM concentration of inhibitor
16 for 30 min at ambient temperature and 15 min at 4°C, respectively. The N9-NA **5** and **6**
17 complexes were prepared by co-crystallization.
18
19
20
21
22
23
24
25
26
27

28 **X-ray data collection, structure determination, and refinement**

29
30 X-ray diffraction data sets for single crystals of N9 NA with inhibitor complexes in well solution
31 plus 15% (v/v) glycerol as cryo-protectant were collected at -173°C with 360° rotation, 1σ
32 oscillation and 1s exposure with wavelength of ~0.95 Å using the MX1/MX2 beamlines at the
33 Australian synchrotron. The data sets were processed with HKL2000.⁴³ Further data collection
34 statistics are given in Table 4. The positions of one N9 NA molecule were identified in the
35 asymmetric units by PHASER⁴⁴ molecular replacement using the structure of N9 (PDB entry:
36 1NNC)³⁵ without ligands. The structures with protein molecules alone were refined and then the
37 inhibitors were built into the observed residual electron densities.
38
39
40
41
42
43
44
45
46
47
48

49 Initial refinement of the inhibitor position did not account for all the residual density observed in
50 all N9 NA protein structures. In particular, there was slight negative residual density around the
51 F3 atom and positive residual density in the region between the inhibitor and the Y406 (N9 NA)
52
53
54
55
56
57
58
59
60

1
2
3 residue. The continuous bridge of residual electron density between the C2 atom of the inhibitor
4 and hydroxyl oxygen of the aromatic side chain of Y406 residue suggested the C-O covalent
5 bond, however the refined bond distance was always longer than the expected covalent bond of
6 $\sim 1.4 \text{ \AA}$. Next, the covalent intermediate species accompanied by unsaturated forms of fluorosialic
7 acid as oxocarbenium ion transition-state analogues were refined in N9 NA protein structures.

8
9
10 The geometry of unsaturated forms of fluorosialic acid species is similar to that of DANA in
11 which the carboxylate group is coplanar with the ring plane with considerable sp^2 hybridization
12 at the anomeric carbon. Since a fluorine atom is refined at the chiral position, the double bond is
13 between the ring oxygen and the anomeric carbon, with positive charge at the oxygen resembling
14 an oxocarbenium ion transition state. Adjusting the occupancies of two bonded and non-bonded
15 conformations the length of the covalent linkage for bonded species was refined to a chemically
16 sensible value of $\sim 1.4 \text{ \AA}$. After that the corresponding LINK record was added to the PDB files
17 to define this bond. The occupancies of two conformations were refined to 20% / 80% (**3**), 45% /
18 55% (**4**), 60% / 40% (**5**), and 25% / 75% (**6**) respectively for covalently bonded and non-bonded
19 species. The refinement of occupancy was carried iteratively alternating fixed B-factors or
20 occupancies until physically meaningful and similar for both species values of atomic B-factors
21 were obtained.

22
23
24 In addition, the secondary binding site for molecules of all deoxy-DFSA were identified in the
25 N9 NA complexes, by difference Fourier methods during the course of the refinement. It should
26 be noted that both soaking and co-crystallization were sufficient to observe the initial binding
27 events of **2** including secondary sites and a variable mixed occupancy of the active site. Electron
28 densities for secondary sites of co-crystallized complexes of **5** and **6** were less defined and
29 somewhat fragmented.

1
2
3 Water molecules in the active site and elsewhere were then added. Iterative refinement and
4 model building were conducted using REFMAC⁴⁵ and XFIT/MIFit.^{46, 47} N-linked glycans were
5 observed at all predicted sites (N86, N146 and N201) and are included in the refined atomic
6 model. Refinement yielded a model for 388 residues for the N9 with 11 attached glycans, 1
7 calcium ion, two alternating conformations of the inhibitor, 1 n-deoxy DFSA molecule at
8 additional site and 430 (3), 423 (4), 483 (5), and 354 (6) water molecules (Table 4). Progress of
9 the refinement was monitored using the Rfree statistics based on a test set encompassing 5% of
10 the observed diffraction amplitudes.⁴⁸ The coordinates of the complexes have been deposited in
11 the PDB. Deposition codes and further experimental and data processing details are given in
12 Table 4. Final omit electron densities associated with the ligands in active site and secondary
13 binding sites are displayed in Figure 4 for all N9 NA – inhibitor complexes.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI:

Comparison of IC₅₀ values for 10 and 60 min in the preincubation reaction (**Table S1**),

Fold difference between wild type and mutant 60 min IC₅₀ values in the preincubation reaction (**Table S2**).

Plaque reduction assay for seasonal influenza (H1N1) A/Mississippi/03/01 (**Figure S1**),

Molecular formula strings (**PDF**)

Molecular formula strings (**CSV**)

Accession Codes

Crystal structures of influenza neuraminidase N9 in complex with compounds **3** (PDB 5W26), **4** (PDB 5W2U), **5** (PDB 5W2W), and **6** (PDB 5W2Y). Authors will release the atomic coordinates and experimental data upon article publication.

Corresponding author Jennifer L McKimm-Breschkin

+61 3 8344 1106

jmbvirology@gmail.com

Current addresses

Jennifer L McKimm-Breschkin

Department of Microbiology and Immunology, University of Melbourne, Peter Doherty Institute, Melbourne, 3000 Australia

Stefan Hader

1
2
3 Kings College London, Strand, London, WC2R 2LS, UK Stefan.hader@kcl.ac.uk
4
5
6

7 **Author contributions**

8
9
10 JMB analysed data, prepared NA for crystallisation, wrote the manuscript, SB carried out IC₅₀
11
12 kinetics, reactivation assays and plaque assays, SH synthesized the compounds, AW wrote the
13
14 manuscript, supervised synthesis of the compounds, PP crystallized the NA, VS did the X-ray
15
16 crystallography and structural analysis and wrote the manuscript.
17
18
19

20 **Acknowledgments**

21
22 MRC Grant G0600514.
23

24
25
26 Pandemic Influenza grant NHMRC 595625
27

28
29 SH was funded by an MRC (UK) PhD studentship.
30

31
32 We acknowledge the use of the Australian Synchrotron protein crystallography MX beamlines,
33
34 Victoria, Australia
35

36 **Abbreviations:** NA neuraminidase, NAI neuraminidase inhibitor, DFSA difluoro sialic acid,
37
38
39

40 **Conflict of Interest**

41
42
43 The authors declare no competing financial interest
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- (1) Varghese, J. N.; McKimm-Breschkin, J. L.; Caldwell, J. B.; Kortt, A. A.; Colman, P. M. The structure of the complex between influenza virus neuraminidase and sialic acid, the viral receptor. *Proteins* **1992**, *14*, 327-332.
- (2) Burmeister, W. P.; Ruigrok, R. W.; Cusack, S. The 2.2 Å resolution crystal structure of influenza B neuraminidase and its complex with sialic acid. *EMBO J.* **1992**, *11*, 49-56.
- (3) Varghese, J. N.; Colman, P. M. Three-dimensional structure of the neuraminidase of influenza virus A/Tokyo/3/67 at 2.2 Å resolution. *J. Mol. Biol.* **1991**, *221*, 473-486.
- (4) Colman, P. M.; Hoyne, P. A.; Lawrence, M. C. Sequence and structure alignment of paramyxovirus hemagglutinin-neuraminidase with influenza virus neuraminidase. *J. Virol.* **1993**, *67*, 2972-2980.
- (5) McKimm-Breschkin, J. L. Influenza neuraminidase inhibitors: antiviral action and mechanisms of resistance. *Influenza Other Respir Viruses* **2013**, *7 Suppl 1*, 25-36.
- (6) Tarbet, E. B.; Hamilton, S.; Vollmer, A. H.; Luttick, A.; Ng, W. C.; Pryor, M.; Hurst, B. L.; Crawford, S.; Smee, D. F.; Tucker, S. P. A zanamivir dimer with prophylactic and enhanced therapeutic activity against influenza viruses. *J. Antimicrob. Chemother.* **2014**, *69*, 2164-2174.
- (7) Monto, A. S.; McKimm-Breschkin, J. L.; Macken, C.; Hampson, A. W.; Hay, A.; Klimov, A.; Tashiro, M.; Webster, R. G.; Aymard, M.; Hayden, F. G.; Zambon, M. Detection of influenza viruses resistant to neuraminidase inhibitors in global surveillance during the first 3 years of their use. *Antimicrob. Agents Chemother.* **2006**, *50*, 2395-2402.
- (8) Kim, J. H.; Resende, R.; Wennekes, T.; Chen, H. M.; Bance, N.; Buchini, S.; Watts, A. G.; Pilling, P.; Streltsov, V. A.; Petric, M.; Liggins, R.; Barrett, S.; McKimm-Breschkin, J. L.;

1
2
3 Niikura, M.; Withers, S. G. Mechanism-based covalent neuraminidase inhibitors with broad-
4 spectrum influenza antiviral activity. *Science* **2013**, *340*, 71-75.
5
6

7 (9) Streltsov, V. A.; Pilling, P.; Barrett, S.; McKimm-Breschkin, J. L. Catalytic mechanism
8 and novel receptor binding sites of human parainfluenza virus type 3 hemagglutinin-
9 neuraminidase (hPIV3 HN). *Antiviral Res.* **2015**, *123*, 216-223.
10
11
12

13 (10) Hader, S.; Watts, A. G. The synthesis of a series of deoxygenated 2,3-difluoro-N-
14 acetylneuraminic acid derivatives as potential sialidase inhibitors. *Carbohydr. Res.* **2013**, *374*,
15 23-28.
16
17
18

19 (11) Oakley, A. J.; Barrett, S.; Peat, T. S.; Newman, J.; Streltsov, V. A.; Waddington, L.;
20 Saito, T.; Tashiro, M.; McKimm-Breschkin, J. L. Structural and functional basis of resistance to
21 neuraminidase inhibitors of influenza B viruses. *J. Med. Chem.* **2010**, *53*, 6421-6431.
22
23
24
25

26 (12) Collins, P. J.; Haire, L. F.; Lin, Y. P.; Liu, J.; Russell, R. J.; Walker, P. A.; Skehel, J. J.;
27 Martin, S. R.; Hay, A. J.; Gamblin, S. J. Crystal structures of oseltamivir-resistant influenza virus
28 neuraminidase mutants. *Nature* **2008**, *453*, 1258-1261.
29
30
31
32

33 (13) Blick, T. J.; Tiong, T.; Sahasrabudhe, A.; Varghese, J. N.; Colman, P. M.; Hart, G. J.;
34 Bethell, R. C.; McKimm-Breschkin, J. L. Generation and characterization of an influenza virus
35 neuraminidase variant with decreased sensitivity to the neuraminidase-specific inhibitor 4-
36 guanidino-Neu5Ac2en. *Virology* **1995**, *214*, 475-484.
37
38
39
40
41
42
43

44 (14) McKimm-Breschkin, J. L.; Selleck, P. W.; Usman, T. B.; Johnson, M. A. Reduced
45 sensitivity of influenza A (H5N1) to oseltamivir. *Emerg. Infect. Dis.* **2007**, *13*, 1354-1357.
46
47
48

49 (15) McKimm-Breschkin, J. L.; Williams, J.; Barrett, S.; Jachno, K.; McDonald, M.; Mohr, P.
50 G.; Saito, T.; Tashiro, M. Reduced susceptibility to all neuraminidase inhibitors of influenza
51
52
53
54
55
56
57
58
59
60

1
2
3 H1N1 viruses with haemagglutinin mutations and mutations in non-conserved residues of the
4 neuraminidase. *J. Antimicrob. Chemother.* **2013**, *68*, 2210-2221.

5
6
7 (16) Barrett, S.; Mohr, P. G.; Schmidt, P. M.; McKimm-Breschkin, J. L. Real time enzyme
8 inhibition assays provide insights into differences in binding of neuraminidase inhibitors to wild
9 type and mutant influenza viruses. *PLoS One* **2011**, *6*, e23627.

10
11
12 (17) Tashiro, M.; McKimm-Breschkin, J. L.; Saito, T.; Klimov, A.; Macken, C.; Zambon, M.;
13 Hayden, F. G. Surveillance for neuraminidase-inhibitor-resistant influenza viruses in Japan,
14 1996-2007. *Antivir. Ther.* **2009**, *14*, 751-761.

15
16
17 (18) Watts, A. G.; Opezzo, P.; Withers, S. G.; Alzari, P. M.; Buschiazzo, A. Structural and
18 kinetic analysis of two covalent sialosyl-enzyme intermediates on *Trypanosoma rangeli*
19 sialidase. *J. Biol. Chem.* **2006**, *281*, 4149-4155.

20
21
22 (19) Barrett, S.; McKimm-Breschkin, J. L. Solid phase assay for comparing reactivation rates
23 of neuraminidases of influenza wild type and resistant mutants after inhibitor removal. *Antiviral*
24 *Res.* **2014**, *108*, 30-35.

25
26
27 (20) Pegg, M. S.; von Itzstein, M. Slow-binding inhibition of sialidase from influenza virus.
28 *Biochem. Mol. Biol. Int.* **1994**, *32*, 851-858.

29
30
31 (21) Kati, W. M.; Saldivar, A. S.; Mohamadi, F.; Sham, H. L.; Laver, W. G.; Kohlbrenner, W.
32 E. GS4071 is a slow-binding inhibitor of influenza neuraminidase from both A and B strains.
33 *Biochem. Biophys. Res. Commun.* **1998**, *244*, 408-413.

34
35
36 (22) Baum, E. Z.; Wagaman, P. C.; Ly, L.; Turchi, I.; Le, J.; Bucher, D.; Bush, K. A point
37 mutation in influenza B neuraminidase confers resistance to peramivir and loss of slow binding.
38 *Antiviral Res.* **2003**, *59*, 13-22.

- 1
2
3 (23) McKimm-Breschkin, J. L.; Rootes, C.; Mohr, P. G.; Barrett, S.; Streltsov, V. A. In vitro
4 passaging of a pandemic H1N1/09 virus selects for viruses with neuraminidase mutations
5 conferring high-level resistance to oseltamivir and peramivir, but not to zanamivir. *J.*
6
7
8
9
10 *Antimicrob. Chemother.* **2012**, *67*, 1874-1883.
- 11
12 (24) McKimm-Breschkin, J. L.; Barrett, S. Neuraminidase mutations conferring resistance to
13 laninamivir lead to faster drug binding and dissociation. *Antiviral Res.* **2015**, *114*, 62-66.
- 14
15 (25) Zhou, J.; Wang, D.; Gao, R.; Zhao, B.; Song, J.; Qi, X.; Zhang, Y.; Shi, Y.; Yang, L.;
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- (25) Zhou, J.; Wang, D.; Gao, R.; Zhao, B.; Song, J.; Qi, X.; Zhang, Y.; Shi, Y.; Yang, L.;
Zhu, W.; Bai, T.; Qin, K.; Lan, Y.; Zou, S.; Guo, J.; Dong, J.; Dong, L.; Wei, H.; Li, X.; Lu, J.;
Liu, L.; Zhao, X.; Huang, W.; Wen, L.; Bo, H.; Xin, L.; Chen, Y.; Xu, C.; Pei, Y.; Yang, Y.;
Zhang, X.; Wang, S.; Feng, Z.; Han, J.; Yang, W.; Gao, G. F.; Wu, G.; Li, D.; Wang, Y.; Shu, Y.
Biological features of novel avian influenza A (H7N9) virus. *Nature* **2013**, *499*, 500-503.
- (26) Qi, W.; Jia, W.; Liu, D.; Li, J.; Bi, Y.; Xie, S.; Li, B.; Hu, T.; Du, Y.; Xing, L.; Zhang, J.;
Zhang, F.; Wei, X.; Eden, J. S.; Li, H.; Tian, H.; Li, W.; Su, G.; Lao, G.; Xu, C.; Xu, B.; Liu, W.;
Zhang, G.; Ren, T.; Holmes, E. C.; Cui, J.; Shi, W.; Gao, G. F.; Liao, M. Emergence and
adaptation of a novel highly pathogenic H7N9 influenza virus in birds and humans from a 2013
human-infecting low-pathogenic ancestor. *J. Virol.* **2018**, *92*(2) e00921-17
- (27) Kiso, M.; Kubo, S.; Ozawa, M.; Le, Q. M.; Nidom, C. A.; Yamashita, M.; Kawaoka, Y.
Efficacy of the new neuraminidase inhibitor CS-8958 against H5N1 influenza viruses. *PLoS*
Pathog. **2010**, *6*, e1000786.
- (28) Bantia, S.; Arnold, C. S.; Parker, C. D.; Upshaw, R.; Chand, P. Anti-influenza virus
activity of peramivir in mice with single intramuscular injection. *Antiviral Res.* **2006**, *69*, 39-45.

- 1
2
3 (29) Bantia, S.; Upshaw, R.; Babu, Y. S. Characterization of the binding affinities of
4
5 peramivir and oseltamivir carboxylate to the neuraminidase enzyme. *Antiviral Res.* **2011**, *91*,
6
7 288-291.
8
9
10 (30) Asaoka, N.; Tanaka, Y.; Sakai, T.; Fujii, Y.; Ohuchi, R.; Ohuchi, M. Low growth ability
11
12 of recent influenza clinical isolates in MDCK cells is due to their low receptor binding affinities.
13
14 *Microbes Infect* **2006**, *8*, 511-519.
15
16
17 (31) Newstead, S. L.; Potter, J. A.; Wilson, J. C.; Xu, G.; Chien, C.-H.; Watts, A. G.; Withers,
18
19 S. G.; Taylor, G. L. The structure of *Clostridium perfringens* NanI sialidase and its catalytic
20
21 intermediates. *J. Biol. Chem.* **2008**, *283*, 9080-9088.
22
23
24 (32) Vavricka, C. J.; Liu, Y.; Kiyota, H.; Sriwilaijaroen, N.; Qi, J.; Tanaka, K.; Wu, Y.; Li, Q.;
25
26 Li, Y.; Yan, J.; Suzuki, Y.; Gao, G. F. Influenza neuraminidase operates via a nucleophilic
27
28 mechanism and can be targeted by covalent inhibitors. *Nat Comm* **2013**, *4*, 1491.
29
30
31 (33) Burmeister, W. P.; Henrissat, B.; Bosso, C.; Cusack, S.; Ruigrok, R. W. Influenza B virus
32
33 neuraminidase can synthesize its own inhibitor. *Structure* **1993**, *1*, 19-26.
34
35
36 (34) Winn, M. D.; Ballard, C. C.; Cowtan, K. D.; Dodson, E. J.; Emsley, P.; Evans, P. R.;
37
38 Keegan, R. M.; Krissinel, E. B.; Leslie, A. G. W.; McCoy, A.; McNicholas, S. J.; Murshudov, G.
39
40 N.; Pannu, N. S.; Potterton, E. A.; Powell, H. R.; Read, R. J.; Vagin, A.; Wilson, K. S. Overview
41
42 of the CCP4 suite and current developments. *Acta Crystallogr D* **2011**, *67*, 235-242.
43
44
45 (35) Varghese, J. N.; Colman, P. M.; van Donkelaar, A.; Blick, T. J.; Sahasrabudhe, A.;
46
47 McKimm-Breschkin, J. L. Structural evidence for a second sialic acid binding site in avian
48
49 influenza virus neuraminidases. *Proc. Natl. Acad. Sci. U. S. A.* **1997**, *94*, 11808-11812.
50
51
52 (36) von Itzstein, M.; Wu, W. Y.; Kok, G. B.; Pegg, M. S.; Dyason, J. C.; Jin, B.; Van Phan,
53
54 T.; Smythe, M. L.; White, H. F.; Oliver, S. W.; Colman, P. M.; Varghese, J. N.; Ryan, D. M.;

1
2
3 Woods, J. M.; Bethell, R. C.; Hotham, V. J.; Cameron, J. M.; Penn, C. R. Rational design of
4 potent sialidase-based inhibitors of influenza virus replication. *Nature* **1993**, *363*, 418-423.

5
6
7 (37) McKimm-Breschkin, J. L.; Barrett, S.; Pudjiatmoko; Azhar, M.; Wong, F. Y.; Selleck, P.;
8 Mohr, P. G.; McGrane, J.; Kim, M. I222 neuraminidase mutations further reduce oseltamivir
9 susceptibility of Indonesian Clade 2.1 highly pathogenic avian influenza A(H5N1) viruses. *PLoS*
10 *One* **2013**, *8*, e66105.

11
12
13 (38) Uhlenendorff, J.; Matrosovich, T.; Klenk, H. D.; Matrosovich, M. Functional significance
14 of the hemadsorption activity of influenza virus neuraminidase and its alteration in pandemic
15 viruses. *Arch. Virol.* **2009**, *154*, 945-957.

16
17
18 (39) Dai, M.; McBride, R.; Dortmans, J. C.; Peng, W.; Bakkers, M. J.; de Groot, R. J.; van
19 Kuppeveld, F. J.; Paulson, J. C.; de Vries, E.; de Haan, C. A. Mutation of the second sialic acid-
20 binding site, resulting in reduced neuraminidase activity, preceded the emergence of H7N9
21 influenza A virus. *J. Virol.* **2017**, *91*: e00049-17.

22
23
24 (40) Hurt, A. C.; Iannello, P.; Jachno, K.; Komadina, N.; Hampson, A. W.; Barr, I. G.;
25 McKimm-Breschkin, J. L. Neuraminidase inhibitor-resistant and -sensitive influenza B viruses
26 isolated from an untreated human patient. *Antimicrob. Agents Chemother.* **2006**, *50*, 1872-1874.

27
28
29 (41) Mohr, P. G.; Deng, Y. M.; McKimm-Breschkin, J. L. The neuraminidases of MDCK
30 grown human influenza A(H3N2) viruses isolated since 1994 can demonstrate receptor binding.
31 *Virol J.* **2015**, *12*, 67.

32
33
34 (42) Smith, B. J.; McKimm-Breshkin, J. L.; McDonald, M.; Fernley, R. T.; Varghese, J. N.;
35 Colman, P. M. Structural studies of the resistance of influenza virus neuraminidase to inhibitors.
36 *J. Med. Chem.* **2002**, *45*, 2207-2212.

- 1
2
3 (43) Otwinowski, Z.; Minor, W. Processing of X-ray diffraction data collected in oscillation
4 mode. *Methods Enzymol.* **1997**, 307-326.
5
6
7 (44) McCoy, A. J.; Grosse-Kunstleve, R. W.; Adams, P. D.; Winn, M. D.; Storoni, L. C.;
8 Read, R. J. Phaser crystallographic software. *J Appl Crystallogr* **2007**, 40, 658-674.
9
10
11 (45) Murshudov, G. N.; Vagin, A. A.; Dodson, E. J. Refinement of macromolecular structures
12 by the maximum-likelihood method. *Acta Crystallogr.* **1997**, D53, 240-255.
13
14
15 (46) McRee, D. E.; Badger, J. MIFit Manual. Version 3.0 for Windows and LINUX. © *Rigaku*
16
17 **2003-6.**
18
19
20 (47) McRee, D. E. XtalView/Xfit-A versatile program for manipulating atomic coordinates
21 and electron density. *J Struct Biol* **1999**, 125, 156-165.
22
23
24 (48) Brünger, A. T. The Free R value: a novel statistical quantity for assessing the accuracy of
25 crystal structures. *Nature* **1992**, 355, 472-474.
26
27
28 (49) Schrodinger, L. *The PyMOL Molecular Graphics System, Version 1.3r1.*, **2010.**
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 Comparison of 60 min IC₅₀ values with (+) and without (-) preincubation

	sH1N1	sH1N1	pdmH1N1	pdmH1N1	H5N1	H5N1	H3N2	H1N9	H1N9		B	
μM	wt	H274Y	Cal	Shep	Cl 1	Cl 2	H3N2 wt	E119V	wt	E119G	B wt	D197E
1-	0.0078	0.0060	0.0120	0.0058	0.0120	0.0112	0.0324	0.0519	0.0057	0.8159	0.1674	0.4338
1+	0.0019	0.0022	0.0011	0.0014	0.0025	0.0013	0.0038	0.0034	0.0027	0.6784	0.0089	0.2575
-/+	4.09	2.72	10.85	4.23	4.71	8.62	8.53	15.26	2.10	1.20	18.81	1.68
2-	0.116	0.139	0.791	0.791	2.168	0.889	1.665	0.465	1.103	1.201	0.414	1.257
2+	0.080	0.116	0.601	0.470	1.662	0.863	1.377	0.238	1.192	1.148	0.065	0.174
-/+	1.45	1.19	1.32	1.68	1.30	1.03	1.21	1.95	0.93	1.05	6.37	7.23
3-	0.048	0.024	0.428	0.474	0.985	0.535	0.313	3.705	0.119	1.278	0.076	0.250
3+	0.069	0.133	0.490	0.490	1.396	0.583	0.318	3.675	0.011	1.050	0.027	0.032
-/+	0.69	0.18	0.87	0.97	0.71	0.92	0.98	1.01	11.23	1.22	2.79	7.75
4-	0.061	0.098	0.457	0.418	0.460	0.281	0.472	0.191	0.438	0.985	0.228	1.550
4+	0.037	0.104 ^d	0.330	0.316	0.710	0.316	0.345	0.102	0.289	0.874 ^d	0.040	0.212
-/+	1.65	0.95	1.39	1.32	0.65	0.89	1.37	1.88	1.51	1.13	5.76	7.30

5-	8.29	1.15	25.81	25.77	21.66	28.20	28.70	25.14	23.02	132.55	7.62	40.05
5+	2.61	0.28	22.35 ^d	21.96 ^d	17.05	24.52 ^d	5.01	4.40	10.91 ^d	107.87 ^d	1.15	14.48 ^d
-/+	3.17	4.04	1.15	1.17	1.27	1.15	5.73	5.71	2.11	1.23	6.60	2.77
6-	10.76	6.42	31.71	29.64	20.68	23.77	44.57	36.26	25.38	234.81	9.19	79.03
6+	4.50 ^d	4.69 ^d	20.96	20.68	25.40	26.10	29.51 ^d	25.99 ^d	22.24 ^d	210.19 ^d	4.48 ^d	32.23 ^d
-/+	2.39	1.37	1.51	1.43	0.81	0.91	1.51	1.40	1.14	1.12	2.05	2.45

The IC₅₀ values are the means of two individual IC₅₀ values for each virus-inhibitor combination.

^a Virus abbreviations: sH1N1=A/Mississippi/03/01 seasonal (H1N1), pdmH1N1 Cal =pandemic H1N1 A/California/7/09 H1N1, pdmH1N1

Shep =pandemic H1N1 A/Swine/Shepparton/6/2009, H5N1 Cl 1=A/Chicken/Vietnam/08/2004 (H5N1) clade 1, H5N1 Cl 2=

A/Chicken/Bangli/BBVD-563/2007 (H5N1)clade 2, H3N2=A/Fukui/45/04 (H3N2) , H1N9=NWS/G70C H1N9, B=B/Perth/211/2001

^b - = no preincubation, + = preincubated for 30 min prior to addition of MUNANA substrate,

^c The greater the difference between no preincubation and preincubation the slower the binding, -/+ ~1 indicates fast, >1.5-2 indicates slow binding.

^d IC₅₀s continue to decrease both with and without preincubation (Figures 1 and 2), so the ratio may not properly reflect the extent of slow binding.

Table 2 Time (T_{1/2} min) for reaction rates to reach 50% of the maximum rate of the uninhibited control after removal of inhibitor

Virus	Control		1 ¹⁹	2	3	4	5	6
	Ave (SD)	Ave (SD)						
H1N1 A/Mississippi	15.6 (6.3)	41.3 (7.7) ^{b***}	24.8 (2.3) [*]	26.7 (0.4) [*]	29.2 (1.6) ^{**}	30.4 (9.0) ^{**}	37.5 (4.4) ^{***}	
pdmH1N1 A/Shep	12.5 (3.1)	ND	11.8 (1.6)	11.6 (1.8)	15.8 (0.4)	17.8 (3.7) ^{**}	24.5 (3.5) ^{***}	
H5N1 Clade 1	15.6 (3.7)	27.7 (2.7) ^{***}	12.0 (1.1)	13.7 (1.0)	15.0 (2.7)	19.4 (5.5)	17.2 (3.5)	
H5N1 Clade 2	12.1 (1.0)	72.8 (5.1) ^{***}	8.5 (0.5)	10.6 (2.5)	12.3 (2.0)	23.6 (7.7) ^{***}	16.9 (3.3) ^{**}	
H3N2 A/Fukui	12.8 (3.5)	108.7 (18.8) ^{***}	17.6 (2.1) [*]	13.4 (4.0)	13.9 (3.8)	25.1 (2.9) ^{***}	29.7 (7.2) ^{***}	
H1N9 G70C	10.9 (2.1)	32.1 (1.6) ^{***}	11.1 (0.8)	12.0 (1.1)	13.5 (0.8)	32.2 (1.0) ^{****}	30.6 (0.6) ^{***}	
B/Perth	10.5 (2.2)	24.6 (3.5) ^{***}	43.6 (4.2) ^{***}	29.2 (4.8) ^{***}	68.7 (14.5) ^{***}	185.5 (31.9) ^{***}	>240 ^{***}	

Results are from at least two independent assays, with ≥ 2 replicates per assay.

^a Virus abbreviations: A/Mississippi =A/Mississippi/03/01 seasonal (H1N1), pdmH1N1 A/Shep =pandemic H1N1 A/Swine/Shepparton/6/2009, H5N1 Clade 1=A/Chicken/Vietnam/08/2004, H5N1 Clade 2= A/Chicken/Bangli/BBVD-563/2007, H3N2 A/Fukui =A/Fukui/45/04 (H3N2), H1N9 G70C=NWS/G70C H1N9, B/Perth=B/Perth/211/2001.

1
2
3
4
5 **b** The $T_{1/2}$ **value** for each inhibitor was compared to the control values by the Student's *t* test, significantly slower compared to control ^{*} $P < 0.05$,

6
7 ^{**} $P < 0.01$, ^{***} $P < 0.001$
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 3 EC₅₀s of compounds causing a 50% reduction in plaque size

Virus	1	2	3	4	5	6
	nM	μM	μM	μM	μM	μM
H1N1 A/Mississippi	≤1	1	1	0.1-1	10	10-100
H3N2 A/Fukui	100	100	100	10-100	≥100	ND
H3N2 MDCK A/Victoria	0.1-1	100	10	10	≥100	≥100
H1N9 G70C	1-10	1-10	1-10	10	100	100
B/Perth	10	1	1	1-10	10-100	10-100

A/Mississippi =A/Mississippi/03/01 seasonal (H1N1), H3N2 A/Fukui =A/Fukui/45/04 (H3N2),
A/Victoria =A/Victoria/503/06, H1N9 G70C=NWS/G70C H1N9, B/Perth=B/Perth/211/2001.

Results are from two independent assays.

Table 4.**X-ray data collection and refinement statistics.**

	N9 NA -3	N9 NA -4	N9 NA -5	N9 NA -6
PDB entry	5W26	5W2U	5W2W	5W2Y
Wavelength (Å)	0.95324	0.95369	0.95324	0.95373
Resolution range (Å)	42.00 – 1.90 (1.95 – 1.90) ^a	48.38 – 2.00 (2.05 – 2.00)	48.30 – 1.85 (1.90 – 1.85)	48.48 – 2.39 (2.45 – 2.39)
Space group	I 4 3 2	I 4 3 2	I 4 3 2	I 4 3 2
Unit cell	$a=b=c=180.94$ Å, $\alpha=\beta=\gamma=90^\circ$	$a=b=c=180.03$ Å, $\alpha=\beta=\gamma=90^\circ$	$a=b=c=180.71$ Å, $\alpha=\beta=\gamma=90^\circ$	$a=b=c=181.38$ Å, $\alpha=\beta=\gamma=90^\circ$
Unique reflections	37832 (2735)	32364 (1723)	40605 (2158)	19147 (1035)
Multiplicity	72.0 (25.0)	78.4 (32.0)	75.0 (29.0)	35.4 (16.6)
Completeness (%)	99.9 (99.9)	99.6 (99.8)	99.8 (99.0)	98.5 (98.2)
Mean $I/\sigma(I)$	39.0 (1.5)	10.0 (1.3)	10.2 (1.2)	11.4 (1.8)
Wilson B-factor	22.5	23.1	21.8	22.2
Matthews coefficient, V_m	2.83	2.83	2.83	2.83

	(Å ³ /Da)				
	Bulk solvent $k_{\text{sol}}(\text{e}/\text{Å}^3)$, $B_{\text{sol}}(\text{Å}^2)$	0.40, 47.9	0.41, 61.5	0.39, 52.3	0.41, 42.9
	$R_{\text{merge}}^{\text{b}}$	0.170 (0.750)	0.308 (0.760)	0.297 (0.820)	0.394 (0.762)
	$R_{\text{work}}^{\text{d}}$	0.136 (0.289)	0.142 (0.273)	0.140 (0.231)	0.141 (0.257)
	$R_{\text{free}}^{\text{e}}$	0.177 (0.294)	0.187 (0.308)	0.171 (0.262)	0.250 (0.274)
	Number of non-hydrogen atoms	3718	3715	3793	3655
	macromolecules	3090	3094	3087	3094
	ligands	197	197	222	208
	water	430	423	483	352
	metals	1	1	1	1
	Protein residues	388	388	388	388
	RMS(bonds) (Å)	0.020	0.021	0.019	0.017
	RMS(angles) (°)	2.017	2.078	1.997	1.937
	Average B-factor (Å ²)	24.0	25.0	24.0	23.0
	Ramachandran favoured (%)	96	96	95	95
	Ramachandran allowed (%)	4	4	5	5

1
2
3
4
5
6
7 ^aStatistics for the highest-resolution shell are shown in parentheses. ^b $R_{\text{merge}} = \frac{\sum_{hkl} \sum_j |I_j - \langle I_j \rangle|}{\sum_{hkl} \sum_j I_j}$, where hkl specifies unique indices, j
8 indicates equivalent observations of hkl , I_j and σ_j^2 are the observed intensities and their errors, and $\langle I_j \rangle$ and $\langle \sigma_j \rangle$ are the mean values. ^d $R = \frac{\sum_{hkl} ||$
9 $F_o | - F_c |}{\sum_{hkl} | F_o |}$, where $| F_o |$ and $| F_c |$ are the observed and calculated structure factor amplitudes, respectively. ^eRepresents 5% of the data.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Figure Legends

Figure 1

Chemical structures of Zanamivir (**1**), 2,3-difluoro sialic acid (DFSA) (**2**), 4-deoxy (**3**), 7-deoxy (**4**), 8-deoxy (**5**) and 9-deoxy (**6**) DFSA.

Figure 2

IC₅₀ Kinetics showing changes in IC₅₀ values during the 60 min reaction period with (+) or without preincubation (-) for N1 NAs. The IC₅₀ value for each 10 min time point is the mean of the individual IC₅₀ values of the duplicates for that virus-inhibitor reaction. The greater the difference in final (60 min) IC₅₀ values between the reactions without and with preincubation (-/+ ratio), the slower the binding. Additionally, the rate of increase of the IC₅₀ values after preincubation with inhibitor upon addition of MUNANA substrate, gives a qualitative indication of the dissociation rate. The faster the increase, i.e. the greater the difference between the 10 min and 60 min IC₅₀ values, the faster the dissociation. IC₅₀ values for compounds **5** and **6** continue to decrease for some viruses even after preincubation. This novel profile suggests compounds are very slow binding. (A) Seasonal (H1N10 and H274Y mutant, which has reduced sensitivity to oseltamivir and peramivir. (B) pdmH1N1 Cal = human isolate A/California/7/09, Sh=swine isolate A/Swine/Shepparton/6/2009. (C) H5N1 cl 1= A/Chicken//Vietnam/08/2004 clade1, cl 2= A/Chicken/Bangli/BBVD-563/2007 clade 2

Figure 3

IC₅₀ Kinetics showing changes in IC₅₀ values during the 60 min reaction period with (+) or without (-) preincubation for N2, N9 and B NAs. The IC₅₀ value for each 10 min time point is the mean of the individual IC₅₀ values of the duplicates for that virus-inhibitor reaction. The

1
2
3 greater the difference in final (60 min) IC_{50} values between the reactions without and with
4 preincubation (-/+ ratio), the slower the binding. Additionally, the rate of increase of the IC_{50}
5 values after preincubation with inhibitor upon addition of MUNANA substrate, gives a
6 qualitative indication of the dissociation rate. The faster the increase, i.e. the greater the
7 difference between the 10 min and 60 min IC_{50} values, the faster the dissociation. IC_{50} values
8 for compounds **5** and **6** continue to decrease for some viruses even after preincubation. This
9 novel profile suggests compounds are very slow binding. (A) H3N2 wild type and E119V
10 mutant, which has reduced sensitivity to oseltamivir. (B) N9 wild type and E119G mutant,
11 which has reduced sensitivity to zanamivir. (C) B/Perth wild type and D197E mutant which
12 has reduced sensitivity to all NAIs.
13
14
15
16
17
18
19
20
21
22
23
24
25

26 **Figure 4**

27
28 The N9 NA active site complexes with the trapped covalently bound 3-fluoro-sialyl-enzyme
29 intermediate in yellow and non-bonded 3-fluoro-oxocarbenium “transition-state” analogue in
30 green are shown in the left side panels. The n-deoxy-DFSA (**n=4,7,8,9**) secondary sites are
31 shown in purple in the right side panels. Red spheres represent water molecules. (A) and (B)
32 compound **3**; (C) and (D) compound **4**; (E) and (F) compound **5**; and (G) and (H) compound
33 **6**, respectively. All sites are overlaid with the omit³⁴ electron density map shown as a grey
34 mesh contoured at 1σ within 1.6 Å of ligands. The figure was produced using PyMol.⁴⁹
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1




Figure 2




Figure 3




Figure 4




Table of Contents graphic

