

The Annals of Probability
2019, Vol. 47, No. 5, 2801–2829
<https://doi.org/10.1214/18-AOP1322>
© Institute of Mathematical Statistics, 2019

SELF-AVOIDING WALK ON NONUNIMODULAR TRANSITIVE GRAPHS

BY TOM HUTCHCROFT

University of Cambridge

We study self-avoiding walk on graphs whose automorphism group has a transitive nonunimodular subgroup. We prove that self-avoiding walk is ballistic, that the bubble diagram converges at criticality, and that the critical two-point function decays exponentially in the distance from the origin. This implies that the critical exponent governing the susceptibility takes its mean-field value, and hence that the number of self-avoiding walks of length n is comparable to the n th power of the connective constant. We also prove that the same results hold for a large class of repulsive walk models with a self-intersection based interaction, including the weakly self-avoiding walk. All of these results apply in particular to the product $T_k \times \mathbb{Z}^d$ of a k -regular tree ($k \geq 3$) with \mathbb{Z}^d , for which these results were previously only known for large k .

1. Introduction. A *self-avoiding walk* (SAW) on a graph G is a path in G that visits each vertex at most once. In the probabilistic study of self-avoiding walk, one fixes a graph (often the hypercubic lattice \mathbb{Z}^d), and is interested in both enumerating the number of n -step SAWs and studying the asymptotic behaviour of a uniformly random SAW of length n . This leads to two particularly important questions.

QUESTION 1.1. *What is the asymptotic rate of growth of the number of SAWs of length n ?*

QUESTION 1.2. *How far from the origin is the endpoint of a typical SAW of length n ?*

These questions are simple to state but are often very difficult to answer. Substantial progress has been and continues to be made for SAW on Euclidean lattices. In particular, a very thorough understanding of SAW on \mathbb{Z}^d for $d \geq 5$ has been established in the seminal work of Hara and Slade [23, 24]. The low-dimensional cases $d = 2, 3, 4$ continue to present serious challenges. For a comprehensive introduction to and overview of this literature, we refer the reader to [2, 31].

Received September 2017; revised June 2018.

MSC2010 subject classifications. Primary 60K35; secondary 05A99, 60B99.

Key words and phrases. Self-avoiding walk, nonunimodular, transitive graph, mean-field, nonamenable, bubble diagram.

Recently, the study of SAW on more general graphs has gathered momentum. In particular, a systematic study of SAW on transitive graphs has been initiated in a series of papers by Grimmett and Li [11, 13–17, 19], which is primarily concerned with properties of the connective constant. Other works on SAW on non-Euclidean transitive graphs include [3, 10, 12, 29, 32, 33]; see [18] for a survey of these results.

In this paper, we give complete answers to Question 1.1 and Question 1.2 for self-avoiding walk on graphs whose automorphism group admits a *nonunimodular transitive subgroup* (defined in the next subsection). Although graphs whose entire automorphism group is nonunimodular are generally considered to be rather contrived and unnatural, the class of graphs with a nonunimodular transitive *subgroup* of automorphisms is much larger. Indeed, it includes natural examples such as the product $T_k \times \mathbb{Z}^d$ of a k -regular tree with \mathbb{Z}^d for every $k \geq 3$ (or indeed $T_k \times H$ where H is an arbitrary transitive graph), for which the results of this paper were only previously known for sufficiently large values of k (see the discussion at the end of this subsection).

Our proofs are inspired by the analysis we carried out for percolation on the same class of graphs in our paper [26], which relies on similar methodology. It should be remarked that although every graph possessing a transitive nonunimodular subgroup of automorphisms is necessarily nonamenable [36], we never use this fact in our analysis.

Our first theorem answers Question 1.1 in the nonunimodular context. Let G be a transitive graph, let 0 be a fixed root vertex of G and let $Z(n)$ be the number of length n SAWs in G starting at 0 . Hammersley and Morton [20] observed that $Z(n)$ satisfies the submultiplicative inequality $Z(n+m) \leq Z(n)Z(m)$, from which it follows by Fekete’s lemma that there exists a constant $\mu_c = \mu_c(G)$, known as the *connective constant* of G , such that

$$\mu_c = \lim_{n \rightarrow \infty} Z(n)^{1/n} = \inf_{n \geq 1} Z(n)^{1/n},$$

so that in particular

$$\mu_c^n \leq Z(n) \leq \mu_c^{n+o(n)}$$

for every $n \geq 0$. We also define the *susceptibility* $\chi(z)$ to be the generating function

$$\chi(z) = \sum_{n \geq 0} z^n Z(n),$$

which has radius of convergence $z_c := \mu_c^{-1}$. The connective constant is not typically expected to have a nice or interesting value (a notable exception is the hexagonal lattice [9, 34]), and it is usually much more interesting to estimate the subexponential correction to $Z(n)$ than it is to estimate μ_c . We stress that submultiplicativity arguments alone do not yield any control of this subexponential correction whatsoever.

THEOREM 1.3 (Counting walks). *Let G be a connected, locally finite graph, and suppose that $\text{Aut}(G)$ has a transitive nonunimodular subgroup. Then there exists a constant C such that*

$$\frac{z_c}{z_c - z} \leq \chi(z) \leq \frac{Cz_c}{z_c - z} \quad \text{and} \quad \mu_c^n \leq Z(n) \leq C\mu_c^n$$

for every $0 \leq z < z_c$ and $n \geq 0$.

In fact, we are able to obtain explicit estimates on the constants that appear in this theorem; see Remark 3.5. The lower bounds of the theorem are trivial consequences of submultiplicativity. The upper bounds on $\chi(z)$ and $Z(n)$ in this theorem are equivalent up to the choice of constant; see Lemma 3.4. Probabilistically, the upper bound on $Z(n)$ means that the concatenation of two uniformly chosen n -step SAWs has probability at least $1/C > 0$ to be self-avoiding for every $n \geq 0$.

Our next theorem answers Question 1.2 in the nonunimodular context. We define \mathbb{P}_n to be the uniform measure on self-avoiding walks of length n in G starting at 0, and denote the random self-avoiding walk sampled from \mathbb{P}_n by $X = (X_i)_{i=0}^n$. For each $z \geq 0$ and $x \in V$, we define the *two-point function*:

$$G(z; x) = \sum_{\omega \in \Omega} z^{|\omega|} \mathbb{1}(\omega : 0 \rightarrow x \text{ self-avoiding}) = \sum_{n \geq 0} z^n Z(n) \mathbb{P}_n(X_n = x).$$

In the following theorem, $d(0, x)$ denotes the graph distance between 0 and x .

THEOREM 1.4 (Speed and two-point function decay). *Let $G = (V, E)$ be a connected, locally finite graph, and suppose that $\text{Aut}(G)$ has a transitive nonunimodular subgroup. Then there exists a positive constant c such that*

$$G(z_c; x) \leq \exp[-cd(0, x)]$$

for every $x \in V$ and

$$\mathbb{P}_n(d(0, X_n) \geq cn) \geq 1 - \exp[-cn]$$

for every $n \geq 0$.

Let us briefly survey related theorems in the literature. It is reasonable to conjecture that the conclusions of Theorems 1.3 and 1.4 hold for every transitive non-amenable graph. Indeed, it is plausible that the conclusion of Theorem 1.3 holds for every transitive graph with at least quintic volume growth. The conjectures are trivial when the graph is a tree. Li [29] has shown that SAW is ballistic on a certain class of infinitely ended transitive graphs, and Madras and Wu [32] and Benjamini [3] have shown that SAW on certain specific hyperbolic lattices has linear mean displacement; the latter paper also establishes that the conclusions of Theorem 1.3 hold for the lattices they consider. Gilch and Müller [10] have proven that the conclusions of Theorem 1.3 hold for free products of quasi-transitive graphs (which are always infinitely ended).

Nachmias and Peres [33] proved that the conclusions of Theorems 1.3 and 1.4 hold for every transitive nonamenable graph satisfying

$$(1) \quad (d - 1)\rho < \mu_c,$$

where d is the degree of the graph, ρ is its spectral radius and μ_c is its connective constant. In particular, this holds whenever $\rho \leq 1/2$ [35], as well as for nonamenable transitive graphs of large girth (where what constitutes “large” depends on the spectral radius and the degree). However, we do *not* expect this criterion to apply to $T_k \times \mathbb{Z}^d$ for small k . [Deciding whether or not a graph satisfies (1) can be difficult as to do so one must estimate the connective constant. It can be shown that the criterion (1) does not hold for $T_k \times \mathbb{Z}$ when $k = 3$ by upper bounding μ_c by the growth rate of the number of walks that avoid their last six locations: the latter quantity is exactly computable with computer assistance. On the other hand, using the methods of Lyons and Peres ([30], Theorem 7.37), one can show that (1) does hold for $T_k \times \mathbb{Z}$ when $k \geq 6$. Similarly, using the methods of Yamamoto [38] it should be possible to give a perturbative proof of Theorems 1.3 and 1.4 for $T_k \times \mathbb{Z}$ for every $k \geq 4$.] Furthermore, our nonperturbative approach also lets us handle *anisotropic* SAW on $T_k \times \mathbb{Z}^d$, in which the walk is weighted to prefer \mathbb{Z}^d -edges to T_k -edges, and this bias may be arbitrarily strong (see Section 1.2). It seems very unlikely that such a result could be established using perturbative techniques.

For the hypercubic lattice, Hara and Slade [23, 24] proved that $Z(n)$ grows like μ_c^n as $n \rightarrow \infty$ whenever $d \geq 5$. In the same setting, they also proved that the distance from the origin to the endpoint of an n -step SAW is typically of order $n^{1/2}$. Hara [22] later proved that the critical two-point function decays like $\|x\|^{-d+2}$. (Both behaviours are the same as for simple random walk.) For $d = 4$, it is conjectured that similar asymptotics hold up to logarithmic corrections. See [1] and references therein for an account of substantial recent progress on four-dimensional *weakly* self-avoiding walk. For $d = 2, 3$, the gap between what is known and what is conjectured is very large; important results include those of [7, 8, 21, 28]; see [2, 31] and references therein for more details.

1.1. *Tilted walks and the modular function.* We now define unimodularity and nonunimodularity. Let $G = (V, E)$ be a connected, locally finite graph, and let $\text{Aut}(G)$ be the group of automorphisms of G . Recall that a subgroup $\Gamma \subseteq \text{Aut}(G)$ is said to be *transitive* if it acts transitively on G , that is, if for any two vertices $u, v \in V$ there exists $\gamma \in \Gamma$ such that $\gamma u = v$. The *modular function* $\Delta = \Delta_\Gamma : V^2 \rightarrow (0, \infty)$ of a transitive subgroup $\Gamma \subseteq \text{Aut}(G)$ is defined to be

$$\Delta(x, y) = \frac{|\text{Stab}_y x|}{|\text{Stab}_x y|},$$

where $\text{Stab}_x y$ is the orbit of y under the stabilizer of x in Γ . The group Γ is said to be *unimodular* if $\Delta \equiv 1$, and *nonunimodular* otherwise. The most important properties of the modular function are the *co-cycle identity*, which states that

$$\Delta(x, y)\Delta(y, z) = \Delta(x, z)$$

for every $x, y, z \in V$ and the *tilted mass-transport principle*, which states that if $F : V^2 \rightarrow [0, \infty]$ is invariant under the diagonal action of Γ , meaning that $F(\gamma x, \gamma y) = F(x, y)$ for every $\gamma \in \Gamma$ and $x, y \in V$, then

$$\sum_{y \in V} F(x, y) = \sum_{y \in V} F(y, x) \Delta(x, y).$$

See [30], Chapter 8, for proofs of these properties and further background, and [4], Section 4, for a probabilistic interpretation of the modular function. Note that Δ is itself invariant under the diagonal action of Γ .

The prototypical example of a pair (G, Γ) of a graph together with a nonunimodular transitive subgroup $\Gamma \subseteq \text{Aut}(G)$ is given by the k -regular tree T_k with $k \geq 3$ together with the group Γ_ξ of automorphisms fixing some specified end ξ of T_k . (An *end* of a tree is an equivalence class of infinite simple paths, where paths may start from any vertex and two simple paths are considered equivalent if the sets of vertices they visit have finite symmetric difference.) Let us briefly give an explicit description of the modular function in this example. Every vertex v of T_k has exactly one neighbour that is closer to the end ξ than it is. We call this vertex the *parent* of v . (In other words, the parent of v is the unique neighbour of v that lies in the unique simple path that starts at v and is in the equivalence class ξ .) All other neighbours of v are said to be *children* of v . This leads to a partition of T into levels $(L_n)_{n \in \mathbb{Z}}$, unique up to choice of index, such that if v is in L_n then its parents are in L_{n+1} and its children are in L_{n-1} . The modular function in this example is given explicitly by

$$\Delta(u, v) = (k - 1)^n \iff u \in L_m \text{ and } v \in L_{m+n} \text{ for some } m \in \mathbb{Z}.$$

From this example, many further examples can be built. In particular, if G is an arbitrary transitive graph and $T_k \times G$ is the product of T_k with G , then $\text{Aut}(T_k \times G)$ has a nonunimodular transitive subgroup of automorphisms isomorphic Γ to $\Gamma_\xi \times \text{Aut}(G)$ and with modular function $\Delta_\Gamma((u, x), (v, y)) = \Delta_{\Gamma_\xi}(u, v) \Delta_{\text{Aut}(G)}(x, y)$; see, for example, [5, 26, 37] for further examples.

As in [26], the key to our analysis is to define *tilted* versions of classical quantities such as the susceptibility. These quantities will be similar to their classical analogues, but will have an additional parameter, λ , and will be weighted in some sense by the modular function to the power λ . We will show that these tilted quantities behave in similar ways to their classical analogues (corresponding to $\lambda = 0$) but, crucially, will have different critical values associated to them.

For each $\lambda \in \mathbb{R}$ and $n \geq 0$, we define

$$Z(\lambda; n) = \sum_{x \in V} \sum_{\omega \in \Omega} \mathbb{1}[\omega : 0 \rightarrow x \text{ a length } n \text{ SAW}] \Delta^\lambda(0, x),$$

and define the *tilted susceptibility* to be

$$\chi_w(z, \lambda) = \sum_{n \geq 0} z^n Z_w(\lambda; n).$$

Since every self-avoiding walk ω of length $n + m$ is the concatenation of two self-avoiding walks ω_1 and ω_2 of lengths n and m , respectively, the co-cycle identity implies that

$$Z(\lambda; n + m) \leq Z(\lambda; n)Z(\lambda; m)$$

for every $n, m \geq 0$. It follows by Fekete’s lemma, as before, that

$$(2) \quad \mu_{c,\lambda} = \mu_{c,\lambda}(G, \Gamma) := \lim_{n \rightarrow \infty} Z(\lambda; n)^{1/n} = \inf_{n \geq 1} Z(\lambda; n)^{1/n}$$

and that $z_{c,\lambda} = z_{c,\lambda}(G, \Gamma) := \mu_{c,\lambda}^{-1}$ is the radius of convergence of $\chi(z; \lambda)$.

The tilted mass-transport principle leads to a symmetry between λ and $1 - \lambda$. Indeed, it implies that

$$\begin{aligned} Z(\lambda; n) &= \sum_{x \in V} \sum_{\omega \in \Omega} \mathbb{1}[\omega : 0 \rightarrow x \text{ a length } n \text{ SAW}] \Delta^\lambda(0, x) \\ &= \sum_{x \in V} \sum_{\omega \in \Omega} \mathbb{1}[\omega : x \rightarrow 0 \text{ a length } n \text{ SAW}] \Delta^\lambda(x, 0) \Delta(0, x) \\ (3) \quad &= \sum_{x \in V} \sum_{\omega \in \Omega} \mathbb{1}[\omega : 0 \rightarrow x \text{ a length } n \text{ SAW}] \Delta^{1-\lambda}(0, x) = Z(1 - \lambda; n) \end{aligned}$$

for every $\lambda \in \mathbb{R}$ and $n \geq 0$, and hence that

$$\chi(z, \lambda) = \chi(z, 1 - \lambda)$$

for every $\lambda \in \mathbb{R}$, and $z \geq 0$. In particular, it follows that $z_{c,\lambda} = z_{c,1-\lambda}$ for every $\lambda \in \mathbb{R}$. Moreover, it is easy to see that $Z(\lambda, n)$ is a convex function of λ for each fixed n , and combined with (3) this implies that both $Z(\lambda, n)$ and $\chi_w(z, \lambda)$ are decreasing on $(-\infty, 1/2]$ and increasing on $[1/2, \infty)$, while $z_{c,\lambda}(w)$ is increasing on $(-\infty, 1/2]$ and decreasing on $[1/2, \infty)$. This leads to a special role for $\lambda = 1/2$, which we call the *critical tilt*. We call $z_t = z_{c,1/2}$ the *tiltability threshold* and call $[0, z_t)$ the *tiltable phase*.

The main technical result of this paper is the following. We will show in Section 3 that it easily implies Theorems 1.3 and 1.4.

THEOREM 1.5. *Let G be a connected, locally finite graph, and let Γ be a transitive nonunimodular subgroup of $\text{Aut}(G)$. Then the function*

$$\mathbb{R} \rightarrow (0, z_t], \quad \lambda \mapsto z_{c,\lambda}$$

is continuous, and is strictly increasing on $(-\infty, 1/2]$.

Besides implying Theorems 1.3 and 1.4, Theorem 1.5 also immediately yields *tilted* versions of those theorems, at least when λ does not take its critical value of $1/2$. We define a probability measure on the set of self-avoiding walks of length n starting at 0 by

$$\mathbb{P}_{\lambda,n}(\{\omega\}) = Z(\lambda; n)^{-1} \Delta^\lambda(0, \omega^+),$$

where ω^+ denotes the endpoint of ω .

THEOREM 1.6. *Let G be a connected, locally finite graph, let Γ be a transitive nonunimodular subgroup of $\text{Aut}(G)$. Then the following hold:*

1. *For every $\lambda \neq 1/2$, there exists a constant C_λ such that*

$$\frac{z_{c,\lambda}}{z_{c,\lambda} - z} \leq \chi(z, \lambda) \leq \frac{C_\lambda z_{c,\lambda}}{z_{c,\lambda} - z} \quad \text{and} \quad \mu_c^n \leq Z_w(\lambda; n) \leq C_\lambda \mu_{c,\lambda}^n$$

for every $0 < z < z_{c,\lambda}$ and $n \geq 0$.

2. *For every $z \in [0, z_t)$, there exists a constant c_z such that*

$$G(z; x) \leq e^{-c_z d(0,x)}$$

for every $x \in V$.

3. *For every $\lambda \neq 1/2$, there exist positive constants c_λ and c'_λ such that*

$$(4) \quad \mathbb{P}_{\lambda,n}[\text{sgn}(\lambda - 1/2) \log \Delta(0, X_n) \geq c_\lambda n] \geq 1 - e^{-c_\lambda n}$$

for every $n \geq 0$, and hence

$$(5) \quad \mathbb{P}_{\lambda,n}[d(0, X_n) \geq c'_\lambda n] \geq 1 - e^{-c'_\lambda n},$$

for every $n \geq 0$.

REMARK 1.7. The estimate (4) also yields additional information concerning the untilted case $\lambda = 0$. Indeed, it shows that for all $\lambda \neq 1/2$ the height $\log \Delta(0, X_n)$ of the walk behaves ballistically. Moreover, it shows that the walk is typically displaced in the downward direction when $\lambda < 1/2$ and the upward direction when $\lambda > 1/2$.

Theorem 1.6 naturally leads to questions concerning the critically tilted case $\lambda = 1/2$. We present some such questions along with some partial results in Section 4.

1.2. Other repulsive walk models. All our results apply more generally to a large family of repulsive walk models, including the self-avoiding walk as a special case. This generalization does not add substantial complications to the proof. In this section, we define the family of models that we will consider and state the generalized theorem.

Let G be a connected, locally finite graph, let Γ be a transitive subgroup of $\text{Aut}(G)$ and let E^\rightarrow be the set of oriented edges of G . An oriented edge e of G is oriented from its tail e^- to its head e^+ and has reversal e^\leftarrow . Let $n \geq 0$. A path of length n in G is a pair of functions $\omega\{0, \dots, n\} \rightarrow V$ and $\omega\{(i, j) : 0 \leq i, j \leq n : |i - j| = 1\} \rightarrow E^\rightarrow$ such that $\omega(i + 1, i) = \omega(i, i + 1)^\leftarrow$ for every $0 \leq i < n$ and $\omega(i, i + 1)^\leftarrow = \omega(i)$ and $\omega(i, i + 1)^\rightarrow = \omega(i + 1)$ for every $0 \leq i < n$. (Note that, by a slight abuse of notation, we denote both the path and each of these functions by the same letter, usually ω .) In other words, a path of length n is a multi-graph

homomorphism from the line graph with n edges into G . A *path* in G is a path of some length n , and we write $|\omega|$ for the length of the path ω . Note that length one paths are just oriented edges, while length zero (a.k.a. *trivial*) paths are just vertices. We write ω^- and ω^+ for the first and last vertices of ω , and write ω^\leftarrow for the reversal of ω . We say that an ordered pair of paths (ω_1, ω_2) in G are *contiguous* if $\omega_1^+ = \omega_2^-$. We will usually write simply that ω_1 and ω_2 are contiguous, with the ordering being implicit. Given a pair of contiguous paths (ω_1, ω_2) , we define their concatenation $\omega_1 \circ \omega_2$ in the natural way. By slight abuse of terminology, we say that a contiguous pair of paths (ω_1, ω_2) are *disjoint* if $\omega_1(i) \neq \omega_2(j)$ for every $i < |\omega_1|$ and $j > 0$.

Let G be a graph, let Γ be a transitive group of automorphisms of G and let Ω be the set of paths of finite length in G . Consider a *weight function* $w : \Omega \rightarrow [0, \infty)$, which is always taken to give weight one to every trivial path. We say that w is Γ -*invariant* if $w(\gamma\omega) = w(\omega)$ for every $\gamma \in \Gamma$ and $\omega \in \Omega$. We say that w is *reversible* if $w(\omega) = w(\omega^\leftarrow)$ for every $\omega \in \Omega$. We say that w is *repulsive* if

$$w(\omega \circ \eta) \leq w(\omega)w(\eta)$$

for every contiguous pair $(\omega, \eta) \in \Omega^2$. We say that a weight function $w : \Omega \rightarrow [0, \infty)$ is *nondegenerate* if $w(\omega) = 1$ for every ω with $|\omega| = 0$, and $w(\omega) \geq 1$ for every ω with $|\omega| = 1$. We say that the weight function w is *zero-range* if $w(\omega \circ \eta) = w(\omega)w(\eta)$ whenever ω and η are contiguous and disjoint.

For brevity, we will call a weight function *good* if it is Γ -invariant, reversible, nondegenerate, zero-range and repulsive. Important examples of good weight functions include

$$w(\omega) = \mathbb{1}[\omega \text{ is self-avoiding}]$$

which is the weight function for self-avoiding walk, and

$$w(\omega) = \exp\left[-g \sum_{0 \leq i < j \leq |\omega|} \mathbb{1}(\omega(i) = \omega(j))\right]$$

for $g \geq 0$, which is the weight function for weakly self-avoiding walk (a.k.a. the Domb–Joyce model [6]). Another very natural example is the *anisotropic* self-avoiding walk on $T \times \mathbb{Z}^d$:

$$w(\omega) = \mathbb{1}[\omega \text{ self-avoiding}] \\ \times \exp[a\#\{\mathbb{Z}^d \text{ edges used by } \omega\} + b\#\{\text{tree edges used by } \omega\}]$$

for $a, b \geq 0$. This walk prefers to use T edges if $a < b$ and prefers to use \mathbb{Z}^d edges if $a > b$. Finally, to illustrate the flexibility of the definition, let us also give examples of some more exotic good weight functions:

$$w(\omega) = \mathbb{1}[\omega \text{ does not visit any vertex more than twice}],$$

$$w(\omega) = \mathbb{1}[\omega(i) \neq \omega(j) \text{ if } |i - j| \text{ is not prime}],$$

$$w(\omega) = \mathbb{1}[\text{the subgraph of } G \text{ spanned by the edges traversed by } \omega \text{ is a tree}].$$

(Tyler Helmuth has informed us that the first of these examples is related to ϕ^6 field theory.)

For each Γ -invariant weight function w , we define

$$Z_w(n) = \sum_{\omega \in \Omega} w(\omega) \mathbb{1}[\omega^- = 0, |\omega| = n]$$

to be the total weight of all walks of length n . For every n with $Z_w(n) > 0$, we define a probability measure on the set of paths of length n starting at 0 by

$$\mathbb{P}_{w,n}(\{\omega\}) = Z_w(n)^{-1} w(\omega).$$

The two-point function $G_w(z; x)$, the susceptibility $\chi_w(z)$, and the critical parameter $z_c(w)$ are defined analogously to the case of self-avoiding walk, as are the tilted variants $\chi_w(z, \lambda)$, $z_{c,\lambda}(w)$ and $\mathbb{P}_{w,\lambda,n}$. Note that the tilted quantities all depend implicitly on the choice of Γ . Similar to the case of SAW, if w is Γ -invariant and reversible then the tilted mass-transport principle implies that

$$Z_w(\lambda; n) = Z_w(1 - \lambda; n),$$

for every $\lambda \in \mathbb{R}$ and $n \geq 0$, and hence that $\chi_w(\lambda; z) = \chi_w(1 - \lambda; z)$ and $z_{c,\lambda}(w) = z_{c,1-\lambda}(w)$ for every $\lambda \in \mathbb{R}$. The same statements concerning monotonicity of these quantities on $(-\infty, 1/2]$ and $[1/2, \infty)$ hold for arbitrary good weight functions as they did for SAW, and for the same reasons.

If w is Γ -invariant and repulsive, then $Z_w(\lambda; n)$ is submultiplicative for every λ by the co-cycle identity, so that

$$\mu_{c,\lambda}(w) := z_{c,\lambda}(w)^{-1} = \lim_{n \rightarrow \infty} Z_w(\lambda; n)^{1/n} = \inf_{n \geq 1} Z_w(\lambda; n)^{1/n} \in [-\infty, \infty)$$

is well defined by Fekete’s lemma. If furthermore w is good then the nondegeneracy and zero-range properties imply the lower bound $\mu_{c,\lambda} \geq 1$ for every $\lambda \in \mathbb{R}$.

We will prove the following generalization of Theorem 1.5 to arbitrary good weight functions. Again, we stress that this applies in particular to any graph of the form $T \times G$, where T is a regular tree of degree at least 3 and G is an arbitrary transitive graph.

THEOREM 1.8. *Let G be a connected, locally finite graph, let Γ be a transitive nonunimodular subgroup of $\text{Aut}(G)$, and let $w : \Omega \rightarrow [0, \infty)$ be a good weight function. Then the function*

$$\mathbb{R} \rightarrow (0, z_t(w)], \quad \lambda \mapsto z_{c,\lambda}(w)$$

is continuous, and is strictly increasing on $(-\infty, 1/2]$.

Theorem 1.8 has the following straightforward consequences, which generalize Theorem 1.6. In particular, analogues of Theorems 1.3 and 1.4 for general good weight functions w follow from the untilted case $\lambda = 0$.

THEOREM 1.9. *Let G be a connected, locally finite graph, let Γ be a transitive nonunimodular subgroup of $\text{Aut}(G)$, and let $w : \Omega \rightarrow [0, \infty)$ be a good weight function. Then the following hold:*

1. *For every $\lambda \neq 1/2$, there exists a positive constant C_λ such that*

$$\frac{z_{c,\lambda}}{z_{c,\lambda} - z} \leq \chi_w(z, \lambda) \leq \frac{C_\lambda z_{c,\lambda}}{z_{c,\lambda} - z} \quad \text{and} \quad \mu_c^n \leq Z_w(\lambda; n) \leq C_\lambda \mu_{c,\lambda}^n,$$

for every $0 < z < z_{c,\lambda} = z_{c,\lambda}(w)$ and $n \geq 0$.

2. *For every $z \in [0, z_t)$, there exists a positive constant c_z such that*

$$G_w(z; x) \leq e^{-c_z d(0,x)}$$

for every $x \in V$.

3. *For every $\lambda \neq 1/2$, there exist positive constants c_λ, c'_λ such that*

$$\mathbb{P}_{w,\lambda,n}[\text{sgn}(\lambda - 1/2) \log \Delta(0, X_n) \geq c_\lambda n] \geq 1 - e^{-c_\lambda n}$$

for every $n \geq 0$, and hence

$$\mathbb{P}_{w,\lambda,n}[d(0, X_n) \geq c'_\lambda n] \geq 1 - e^{-c'_\lambda n},$$

for every $n \geq 0$.

REMARK 1.10. The reader may find it an enlightening (and easy) exercise to prove the third item of Theorem 1.9 for simple random walk (i.e., in the case $w(\omega) \equiv 1$) and observe what happens in the case $\lambda = 1/2$.

1.3. About the proofs and organization. The proof of Theorem 1.5 and Theorem 1.8 starts by using Fekete’s lemma to get bounds on *bridges* at $z_{c,\lambda}$. The fact that Fekete’s lemma can be used to obtain surprisingly strong bounds for critical models on graphs of exponential growth was first exploited in [25], and is also central to our work on percolation in the nonunimodular setting [26]. We then convert this control of bridges into a control of walks: this conversion centres around a tilted version of a generating function inequality of Madras and Slade [31]. In Section 3, we use Theorem 1.8 to prove Theorem 1.9. In Section 4, we examine the critically tilted case $\lambda = 1/2$, posing several open problems and giving some small partial results.

2. Proof of Theorems 1.5 and 1.8. Let G be a connected, locally finite graph, let $\Gamma \subseteq \text{Aut}(G)$ be transitive and nonunimodular, let 0 be a fixed root vertex and let w be a good weight function. We define the *height* of a vertex v to be $\log \Delta(0, v)$, and define

$$\mathbb{H} = \{\log \Delta(u, v) : u, v \in V\}$$

to be the set of height differences that appear in G . We say that v is *higher* than u if $\Delta(u, v) > 1$ and that v is *lower* than u if $\Delta(u, v) < 1$. We also define $\mathbb{H}_+ =$

$\mathbb{H} \cap (0, \infty)$ and $\mathbb{H}_{\geq 0}$ to be the sets of positive and nonnegative heights that appear in G , respectively.

We define the *level* L_t to be the set of vertices at height t , that is,

$$L_t = \{v \in V : \log \Delta(0, v) = t\}.$$

We also define

$$t_0 = \sup\{\log \Delta(u, v) : u \sim v\}$$

to be the maximum height difference between adjacent vertices, and define the *slab* S_t to be

$$S_t = \{v \in V : t \leq \log \Delta(0, v) < t + t_0\}.$$

We say that $\omega \in \Omega$ is an *up-bridge* if its endpoint ω^+ is not lower than any of its other points, and its starting point ω^- is not higher any of its other points. Similarly, we say that $\omega \in \Omega$ is a *down-bridge* if its endpoint ω^+ is not higher than any of its other points, and its starting point ω^- is not lower any of its other points. Thus, ω is an up-bridge if and only if its reversal ω^\leftarrow is a down-bridge.

We will specify that a walk ω is an up-bridge by writing a superscript of “u.b.”, and a down-bridge by writing a superscript of “d.b.”. For each $t \in \mathbb{R}$, we define

$$a_w(z; t) = \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\omega : 0 \xrightarrow{\text{u.b.}} L_t)$$

and

$$d_w(z; t) = \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\omega : 0 \xrightarrow{\text{d.b.}} L_{-t}),$$

both of which are equal to zero when $t \notin \mathbb{H}_{\geq 0}$. The Γ -invariance of w and the tilted mass-transport principle implies that

$$\begin{aligned} a_w(z; t) &= \sum_{v \in V} \mathbb{1}(\Delta(0, v) = e^t) \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\omega : 0 \xrightarrow{\text{u.b.}} v) \\ &= \sum_{v \in V} \Delta(0, v) \mathbb{1}(\Delta(v, 0) = e^t) \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\omega : v \xrightarrow{\text{u.b.}} 0), \end{aligned}$$

and applying the reversibility of w and the identity $\Delta(0, v) = \Delta(v, 0)^{-1}$ we obtain that

$$\begin{aligned} a_w(z; t) &= \sum_{v \in V} \Delta(0, v) \mathbb{1}(\Delta(0, v) = e^{-t}) \sum_{\omega \in \Omega} w(\omega^\leftarrow) z^{|\omega^\leftarrow|} \mathbb{1}(\omega^\leftarrow : v \xrightarrow{\text{u.b.}} 0) \\ &= \sum_{v \in V} \Delta(0, v) \mathbb{1}(\Delta(0, v) = e^{-t}) \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\omega : 0 \xrightarrow{\text{d.b.}} v) \\ (6) \quad &= e^{-t} d_w(z; t) \end{aligned}$$

for every $t \in \mathbb{H}_{\geq 0}$. Finally, define the sequence $(A_w(z; n))_{n \geq 0}$ by

$$A_w(z; n) = \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}[\omega : 0 \xrightarrow{\text{u.b.}} S_{nt_0}] = \sum_{nt_0 \leq t < (n+1)t_0} a_w(z; t).$$

LEMMA 2.1. *The sequence $A_w(z; n)$ satisfies the generalized supermultiplicative estimate*

$$A_w(z; n + m + 2) \geq z^2 A_w(z; n) A_w(z; m)$$

for every $n, m \geq 0$ and $0 \leq z \leq 1$.

PROOF. It suffices to construct for each $n, m \geq 0$ an injective function

$$\begin{aligned} & \{\omega_1 \in \Omega : \omega_1 : 0 \xrightarrow{\text{u.b.}} S_{nt_0}\} \times \{\omega_2 \in \Omega : \omega_2 : 0 \xrightarrow{\text{u.b.}} S_{mt_0}\} \\ & \rightarrow \{\omega \in \Omega : \omega : 0 \xrightarrow{\text{u.b.}} S_{(n+m+2)t_0}\} \end{aligned}$$

such that

$$(7) \quad w(\omega) \geq w(\omega_1)w(\omega_2)$$

and

$$(8) \quad |\omega| \leq |\omega_1| + |\omega_2| + 2$$

for every pair of up-bridges $\omega_1 : 0 \xrightarrow{\text{u.b.}} S_{nt_0}$ and $\omega_2 : 0 \xrightarrow{\text{u.b.}} S_{mt_0}$.

For each $v \in V$, we fix an automorphism $\gamma_v \in \Gamma$ such that $\gamma_v 0 = v$. We let η_1 and η_2 be paths of length one and two respectively that start at 0 and whose endpoints have height t_0 and $2t_0$, respectively. Let $\omega_1 : 0 \rightarrow S_{nt_0}$ and $\omega_2 : 0 \rightarrow S_{mt_0}$ be up-bridges. Then we have that

$$(n + m)t_0 \leq \log \Delta(0, \omega_1^+) + \log \Delta(0, \omega_2^+) < (n + m + 2)t_0.$$

We define ω as follows:

- If the sum of the heights of ω_1^+ and ω_2^+ is greater than or equal to $(n + m + 1)t_0$, we let ω be the composition $\omega_1 \circ \gamma_1 \eta_1 \circ \gamma_2 \omega_2$, where $\gamma_1 = \gamma_{\omega_1^+}$ sends 0 to the final vertex of ω_1 , and $\gamma_2 = \gamma_{(\gamma_1 \eta_1)^+}$ sends 0 to the final vertex of $\gamma_1 \eta_1$.
- If the sum of the heights of ω_1^+ and ω_2^+ is strictly less than $(n + m + 1)t_0$, we let ω be the composition $\omega_1 \circ \gamma_1 \eta_2 \circ \gamma_2 \omega_2$, where $\gamma_1 = \gamma_{\omega_1^+}$ sends 0 to the final vertex of ω_1 , and $\gamma_2 = \gamma_{(\gamma_1 \eta_2)^+}$ sends 0 to the final vertex of $\gamma_1 \eta_2$.

The path ω is clearly an up-bridge $0 \rightarrow S_{(n+m+2)t_0}$, and clearly satisfies the length bound (8), while the weight bound (7) follows from the assumption that w is non-degenerate and zero-range. To see that the function $(\omega_1, \omega_2) \mapsto \omega$ is injective, observe that ω_1 is necessarily equal to the longest initial segment of ω that has height strictly less than $(n + 1)t_0$ at its endpoint, while ω_2 is necessarily equal to the

longest final segment of ω that has height difference strictly less than $(m + 1)t_0$ between its starting point and endpoint. \square

Next, we define H_n^+ to be the *upper half-space* $H_n^+ := \bigcup_{s \geq nt_0} L_s$, and define

$$b_w(z; n) = \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}[\omega : 0 \rightarrow H_n^+].$$

The following lemma complements Lemma 2.2 and will be used to prove continuity of $\lambda \mapsto z_{c, \lambda}(w)$.

LEMMA 2.2. *The sequence $b_w(z; n)$ satisfies the generalized submultiplicative estimate*

$$b_w(z; n + m + 1) \leq b_w(z; n) b_w(z; m)$$

for every $n, m \geq 0$ and $z \geq 0$.

PROOF. For each $v \in V$, let $\gamma_v \in \Gamma$ be such that $\gamma_v 0 = v$, and for each $n \geq 0$ let

$$H_n^+(v) = \gamma_v H_n^+ = \{u \in V : \log \Delta(v, u) \geq nt_0\}.$$

Let $\omega : 0 \rightarrow H_{n+m+1}^+$. Write $\omega = \omega_1 \circ \omega_2$, where ω_1 is the portion of ω up until it enters H_n^+ for the first time and ω_2 is the remaining portion of ω . Then $\omega_1^+ \in S_{nt_0}$ by definition of t_0 , and it follows that $H_{n+m+1}^+ \subseteq H_m^+(\omega_1^+)$. Thus, using repulsivity of w to bound $w(\omega) \leq w(\omega_1)w(\omega_2)$, we have that

$$\begin{aligned} b_w(z; n + m + 1) &\leq \sum_{\omega_1 : 0 \rightarrow H_n^+} w(\omega_1) z^{|\omega_1|} \sum_{\omega_2 : \omega_1^+ \rightarrow H_m^+(\omega_1^+)} w(\omega_2) z^{|\omega_2|} \\ &= b_w(z; n) b_w(z; m), \end{aligned}$$

where transitivity of Γ and Γ -invariance of w are used in the second line. \square

We now recall Fekete’s lemma, one form of which states that if $(c_n)_{n \geq 0}$ is a sequence taking values in $[-\infty, \infty]$ that satisfies the generalized subadditive inequality

$$c_{n+m+n_0} \leq c_n + c_m + C$$

for some constants n_0 and C , then

$$\lim_{n \rightarrow \infty} \frac{1}{n} c_n = \inf_{n \geq 0} \frac{c_n + C}{n + n_0} \in [-\infty, \infty],$$

so that in particular the limit on the left-hand side exists.

Applying Fekete’s lemma in light of Lemmas 2.1 and 2.2, we obtain that the quantities

$$(9) \quad \begin{aligned} \alpha_w(z) &= - \lim_{n \rightarrow \infty} \frac{1}{nt_0} \log A_w(z; n) \quad \text{and} \\ \beta_w(z) &= - \lim_{n \rightarrow \infty} \frac{1}{nt_0} \log b_w(z; n) \end{aligned}$$

are well defined (as elements of $[-\infty, \infty]$) and that

$$(10) \quad A_w(z; n) \leq \frac{1}{z^2} e^{-\alpha_w(z)t_0(n+2)} \quad \text{and} \quad b_w(z; n) \geq e^{-\beta_w(z)t_0(n+1)}$$

for every $0 \leq z \leq 1$ and $n \geq 0$. Note that we trivially have that $A_w(z; n) \leq b_w(z; n)$, and hence that

$$(11) \quad \alpha_w(z) \geq \beta_w(z)$$

for every $0 \leq z \leq 1$.

LEMMA 2.3. *Let G be a connected, locally finite graph, let $\Gamma \subseteq \text{Aut}(G)$ be transitive and nonunimodular, and let $w : \Omega \rightarrow [0, \infty)$ be a good weight function. Then $\chi_w(z, \lambda) < \infty$ if and only if $\beta_w(z) > \max\{\lambda, 1 - \lambda\}$. In particular, $\beta_w(z_{c,\lambda}(w)) \leq \max\{\lambda, 1 - \lambda\}$.*

PROOF. By definition,

$$\chi_w(z, \lambda) = \sum_{v \in V} \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\omega : 0 \rightarrow v) \Delta^\lambda(0, v)$$

for every $\lambda \in \mathbb{R}$ and $z \geq 0$. Using the assumptions that Γ is transitive and w is Γ -invariant and reversible, we apply the the tilted mass-transport principle to obtain that

$$\begin{aligned} &\sum_{v \in V} \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\Delta(0, v) < 1, \omega : 0 \rightarrow v) \Delta^\lambda(0, v) \\ &= \sum_{v \in V} \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\Delta(0, v) > 1, \omega : 0 \rightarrow v) \Delta^{1-\lambda}(0, v), \end{aligned}$$

and hence that

$$\begin{aligned} \chi_w(z, \lambda) &\asymp \sum_{v \in V} \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\Delta(0, v) \geq 1, \omega : 0 \rightarrow v) [\Delta^\lambda(0, v) + \Delta^{1-\lambda}(0, v)] \\ &\asymp \sum_{v \in V} \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\Delta(0, v) \geq 1, \omega : 0 \rightarrow v) \Delta^{\max\{\lambda, 1-\lambda\}}(0, v). \end{aligned}$$

Indeed, $\chi_w(z, \lambda)$ is upper bounded by twice the right-hand side of the second line and lower bounded by half the right-hand side of the second line. We deduce that

$$\chi_w(z, \lambda) \leq \sum_{n \geq 0} e^{\max\{\lambda, 1-\lambda\}t_0 n} b_w(z; n),$$

and consequently that $\chi_w(z, \lambda) < \infty$ if $\beta_w(z) > \max\{\lambda, 1 - \lambda\}$. Conversely, if $\chi_w(z, \lambda) < \infty$ then we have that

$$\begin{aligned} & \limsup_{n \rightarrow \infty} e^{\max\{\lambda, 1-\lambda\}t_0 n} b_w(z; n) \\ & \leq \limsup_{n \rightarrow \infty} \sum_{v \in H_n^+} \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\omega : 0 \rightarrow v) \Delta^{\max\{\lambda, 1-\lambda\}}(0, v) = 0, \end{aligned}$$

and hence by (10) that $\beta_w(z) > \max\{\lambda, 1 - \lambda\}$ whenever $\chi_w(z, \lambda) < \infty$.

Finally, the bound $\beta_w(z_{c,\lambda}(w)) \leq \max\{\lambda, 1 - \lambda\}$ follows since $\chi_w(z_{c,\lambda}, \lambda) = \infty$ by submultiplicativity of $Z_w(\lambda, n)$. \square

Moreover, we have the following.

LEMMA 2.4. *Let G be a connected, locally finite graph, let $\Gamma \subseteq \text{Aut}(G)$ be transitive and nonunimodular, and let $w : \Omega \rightarrow [0, \infty)$ be a good weight function. Then $\alpha_w(z)$ is left continuous on $(0, \infty)$ and $\beta_w(z)$ is right continuous on $(0, z_t(w))$. Moreover, both $\alpha_w(z)$ and $\beta_w(z)$ are strictly decreasing when they are positive.*

PROOF. Both $\alpha_w(z)$ and $\beta_w(z)$ are clearly decreasing in z for $z > 0$. For each $n \geq 0$, $A_w(z; n)$ and $b_w(z; n)$ are both defined as power series in z with nonnegative coefficients. It follows that they are each left continuous in z for $z > 0$ and are continuous in z within their respective radii of convergence, which are always at least $z_t(w)$ by the trivial bound

$$A_w(z; n) \leq b_w(z; n) \leq \chi_w(z, \lambda),$$

which holds for every $z, n, \lambda \geq 0$. If $z > 0$ and we define

$$\alpha_w(z-) := \inf_{0 < \varepsilon < z} \alpha_w(z - \varepsilon) = \lim_{\varepsilon \downarrow 0} \alpha_w(z - \varepsilon),$$

then (10) implies that

$$A_w(z - \varepsilon; n) \leq \frac{1}{(z - \varepsilon)^2} e^{-\alpha_w(z-\varepsilon)t_0(n+2)} \leq \frac{1}{(z - \varepsilon)^2} e^{-\alpha_w(z-)t_0(n+2)}$$

for every $n \geq 0$. It follows by left continuity of $A_w(z; n)$ that the bound

$$A_w(z; n) \leq \frac{1}{z^2} e^{-\alpha_w(z-)t_0(n+2)}$$

also holds. This implies that $\alpha_w(z-) \leq \alpha_w(z)$ for every $z > 0$, which is equivalent to left continuity since $\alpha_w(z)$ is decreasing. The proof of the claim concerning right continuity of $\beta_w(z)$ on $(0, z_t(w))$ is similar.

The claim that $\alpha_w(z)$ and $\beta_w(z)$ are strictly decreasing when they are positive follows from the trivial inequalities

$$A_w(z'; n) \geq \left(\frac{z'}{z}\right)^n A_w(z; n) \quad \text{and} \quad b_w(z'; n) \geq \left(\frac{z'}{z}\right)^n b_w(z; n),$$

which hold for every $n \geq 0$ and $z' \geq z > 0$. \square

Lemma 2.4 has the following very useful consequence.

LEMMA 2.5. *Let G be a connected, locally finite graph, let $\Gamma \subseteq \text{Aut}(G)$ be transitive and nonunimodular, and let $w : \Omega \rightarrow [0, \infty)$ be a good weight function. Then $\alpha_w(z_{c,\lambda}(w)) \geq \max\{\lambda, 1 - \lambda\}$ for every $\lambda \in \mathbb{R}$.*

PROOF. We trivially have that $\alpha_w(z) \geq \beta_w(z)$ for every $z > 0$, and hence by Lemma 2.3 that $\alpha_w(z) > \max\{\lambda, 1 - \lambda\}$ for every $0 < z < z_{c,\lambda}(w)$. Thus, the claim follows by left continuity of $\alpha_w(z)$. \square

2.1. *Relating walks and bridges.* We now wish to relate the quantities $\alpha_w(z)$ and $\beta_w(z)$. The following proposition, which is the central idea behind this proof, is a tilted analogue of a well-known inequality relating generating functions for walks and bridges in \mathbb{Z}^d due to Madras and Slade [31], Corollary 3.1.8, and related to the work of Hammersley and Welsh [21]. Indeed, the idea of the proof here is to combine that proof with a judicious use of the tilted mass-transport principle.

PROPOSITION 2.6. *Let G be a connected, locally finite graph, let $\Gamma \subseteq \text{Aut}(G)$ be transitive and nonunimodular, and let w be a good weight function. Then the estimate*

$$\chi_w(z, 1/2) \leq \frac{1}{z} \exp\left[2 \sum_{t \in \mathbb{H}_+} a_w(z; t) e^{t/2}\right] \leq \frac{1}{z} \exp\left[2 \sum_{n \geq 0} A_w(z; n) e^{(n+1)t_0/2}\right]$$

holds for every $z \geq 0$.

PROOF. We say that ω is a *upper half-space walk* if $\Delta(\omega^-, \omega(i)) > 1$ for every $i > 0$, that is, if ω is strictly higher than its starting point at every positive time. Similarly, we call ω a *reverse descent* if $\Delta(\omega^-, \omega(i)) \geq 1$ for every $i > 0$, that is, if ω is at least as high as its starting point at every positive time. A path ω is a *descent* if its reversal is a reverse descent. Define

$$h_w(z; t) = \sum_{v \in V} \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\omega : 0 \rightarrow L_t \text{ is an upper half-space walk}),$$

$$r_w(z; t) = \sum_{v \in V} \sum_{\omega \in \Omega} w(\omega) z^{|\omega|} \mathbb{1}(\omega : 0 \rightarrow L_t \text{ is a reverse descent})$$

and

$$\mathcal{H}_w(z, \lambda) = \sum_{t \in \mathbb{H}_{\geq 0}} h_w(z; t) e^{\lambda t}.$$

Note that $h_w(z; 0) = 1$, as the only upper half-space walk ending in L_0 is the trivial path at 0.

Let η be a path of length 1 ending in 0 whose starting point $\eta^- = v$ has height $-t_0$. Then for every $t \geq 0$ and every reverse descent $\omega : 0 \rightarrow L_t$, the composition

$\eta \circ \omega$ is an upper half-space walk $\eta \circ \omega : v \rightarrow L_t$. Using the fact that w is zero-range, nondegenerate and Γ -invariant, this yields the inequality

$$(12) \quad h_w(z; t + t_0) \geq z r_w(z; t)$$

for every $t \in \mathbb{H}_{\geq 0}$.

Let $\omega \in \Omega$. Let ω_1 be the portion of ω up until the last time that it visits a point of minimal height, and let ω_2 be the remaining portion of ω , so that $\omega = \omega_1 \circ \omega_2$. This decomposition is defined in such a way that ω_1 is a descent and ω_2 is an upper half-space walk. Thus, using repulsivity, Γ -invariance, and reversibility of w , we have that

$$\begin{aligned} \chi_w(z, \lambda) &\leq \sum_{t \in \mathbb{H}_{\geq 0}} w(\omega_1) z^{|\omega_1|} \mathbb{1}(0 \xrightarrow{d.} L_{-t}) \sum_{s \in \mathbb{H}_{\geq 0}} w(\omega_2) z^{|\omega_2|} \mathbb{1}(\omega_1^+ \xrightarrow{\text{u.h.s.}} L_s(\omega_1^+)) e^{\lambda(s-t)} \\ &= \sum_{t \in \mathbb{H}_{\geq 0}} w(\omega_1) z^{|\omega_1|} \mathbb{1}(0 \xrightarrow{d.} L_{-t}) e^{-\lambda t} \sum_{s \in \mathbb{H}_{\geq 0}} h_w(z; s) e^{\lambda s} \\ &= \sum_{t \in \mathbb{H}_{\geq 0}} r_w(z; t) e^{(1-\lambda)t} \sum_{s \in \mathbb{H}_{\geq 0}} h_w(z; s) e^{\lambda s}, \end{aligned}$$

where the superscripts d. and u.h.s. denote descents and upper half-space walks respectively and where the tilted mass-transport principle is used in the final equality. Applying (12) we obtain that

$$\chi_w(z, \lambda) \leq \frac{1}{z} \mathcal{H}_w(z, \lambda) \mathcal{H}_w(z, 1 - \lambda)$$

for every $z > 0$ and $\lambda \in \mathbb{R}$. Thus, to conclude the proof of the present proposition, it suffices to prove that the inequality

$$(13) \quad \mathcal{H}_w(z, 1/2) \leq \exp \left[\sum_{t \in \mathbb{H}_{\geq 0}} a_w(z; t) e^{t/2} \right]$$

holds for every $z \geq 0$.

Let $t \in \mathbb{H}_+$ and let $\omega : 0 \rightarrow L_t$ be an upper half-space walk. We decompose $\omega = \omega_1 \circ \omega_2 \circ \dots \circ \omega_k$ for some $k \geq 1$ recursively as follows: We first define ω_1 to be the portion of ω up until the last time it attains its maximum height. Now suppose that $i \geq 2$. If $\omega_1 \circ \dots \circ \omega_{i-1} = \omega$, we stop. Otherwise, consider the piece of ω that remains after $\omega_1 \circ \dots \circ \omega_{i-1}$. If i is odd, let ω_i be the portion of this piece up to the last time it attains its maximum height. If i is even, let ω_i be the portion of this piece up to the last time it attains its minimum height.

Let $s(\omega_i)$ be the absolute value of the height difference between ω_i^- and ω_i^+ . Observe that for each $i \leq k$, ω_i is an up-bridge if i is odd and a down-bridge if i is even. Moreover, the sequence $s(\omega_i)$ is decreasing and satisfies

$\sum_{i=1}^k (-1)^{i+1} s(\omega_i) = t$. This leads to the bound

$$h_w(z; t) \leq \sum_{k \geq 1} \sum_{s \in S_{t,k}} \prod_{i=0}^{\lfloor (k-1)/2 \rfloor} a_w(z; s_{2i+1}) \prod_{i=1}^{\lfloor k/2 \rfloor} d_w(z; s_{2i}),$$

where we define $S_{t,k}$ to be the set of decreasing sequences s_1, \dots, s_k in \mathbb{H}_+ such that $\sum_{i=1}^k (-1)^{i+1} s_i = t$. Now, observe that

$$a_w(z; t) = e^{-t/2} \sqrt{a_w(z; t) d_w(z; t)} \quad \text{and} \quad d_w(z; t) = e^{t/2} \sqrt{a_w(z; t) d_w(z; t)},$$

by (6), and hence that

$$\begin{aligned} h_w(z; t) &\leq \sum_{k \geq 1} \sum_{s \in S_{t,k}} \prod_{i=1}^k e^{(-1)^i s_i/2} \sqrt{a_w(z; s_i) d_w(z; s_i)} \\ &= e^{-t/2} \sum_{k \geq 1} \sum_{s \in S_{t,k}} \prod_{i=1}^k \sqrt{a_w(z; s_i) d_w(z; s_i)} \end{aligned}$$

for every $t \in \mathbb{H}_+$. Now, let S_k be the set of all decreasing sequences $s = s_1, \dots, s_k$ of elements of \mathbb{H}_+ , and observe that for any nonnegative function $f : \mathbb{H}_+ \rightarrow [0, \infty]$ we have that

$$\prod_{t \in \mathbb{H}_+} (1 + f(t)) = 1 + \sum_{k \geq 1} \sum_{s \in S_k} \prod_{i=1}^k f(s_i) = 1 + \sum_{t \in \mathbb{H}_+} \sum_{k \geq 1} \sum_{s \in S_{t,k}} \prod_{i=1}^k f(s_i).$$

Applying this equality with $f(t) = \sqrt{a_w(z; t) d_w(z; t)}$, we obtain that

$$\begin{aligned} \mathcal{H}_w(z, 1/2) &= 1 + \sum_{t \in \mathbb{H}_+} h_w(z; t) e^{t/2} \\ &\leq 1 + \sum_{t \in \mathbb{H}_+} \sum_{k \geq 1} \sum_{s \in S_{t,k}} \prod_{i=1}^k \sqrt{a_w(z; s_i) d_w(z; s_i)} \\ &= \prod_{t \in \mathbb{H}_+} (1 + \sqrt{a_w(z; t) d_w(z; t)}) = \prod_{t \in \mathbb{H}_+} (1 + e^{t/2} a_w(z; t)). \end{aligned}$$

Using the elementary inequality $1 + x \leq e^x$ concludes the proof. \square

The following is an immediate consequence of Lemma 2.5, Proposition 2.6 and the estimate (10).

COROLLARY 2.7. *If $z \geq 0$ is such that $\alpha = \alpha_w(z) > 1/2$, then*

$$\chi_w(z, 1/2) \leq \frac{1}{z} \exp \left[\frac{2}{z^2 e^{\alpha t_0} [e^{(\alpha-1/2)t_0} - 1]} \right] < \infty.$$

In particular, $z_{c,\lambda}(w) < z_{c,1/2}(w) = z_t(w)$ for every $\lambda \neq 1/2$.

We now apply Corollary 2.7 to prove Theorems 1.5 and 1.8.

PROOFS OF THEOREMS 1.5 AND 1.8. For each $v \in V$, let $\gamma_v \in \Gamma$ be an automorphism with $\gamma_v 0 = v$, and for each $t \in \mathbb{R}$ and $v \in V$ let

$$S_t(v) = \gamma_v S_t = \{u \in V : t \leq \Delta(v, u) < t + t_0\}.$$

Write $z_{c,\lambda} = z_{c,\lambda}(w)$. It follows from Corollary 2.7 and Lemma 2.5 that

$$\sum_{\omega \in \Omega} w(\omega) z_{c,\lambda}^{|\omega|} \mathbb{1}[\omega : 0 \rightarrow S_0 \cup S_{t_0}] \leq \chi_w(z_{c,\lambda}, 1/2) < \infty.$$

Now suppose that $t > 0$ and that $\omega : 0 \rightarrow S_t$ is a path. Then we can decompose $\omega = \omega_1 \circ \omega_2 \circ \omega_3$, where $\omega_1 : 0 \rightarrow S_0$, the path $\omega_2 : \omega_1^+ \rightarrow S_{t-t_0} \subseteq S_{t-t_0}(\omega_1^+) \cup S_{t-2t_0}(\omega_1^+)$ is an up-bridge, and $\omega_3 : \omega_2^+ \rightarrow S_t \subseteq S_0(\omega_2^+) \cup S_{t_0}(\omega_2^+)$. Indeed, simply take ω_1 to be the portion of ω_1 up to the last visit to S_0 , take ω_2 to be the portion of ω between the last visit to S_0 and the first subsequent visit to S_{t-t_0} , and take ω_3 to be the remaining final piece (it is possible that some of these paths have length zero, but this is not a problem). By summing over possible choices of ω_1, ω_2 and ω_3 , and using both transitivity of Γ and Γ -invariance and repulsivity of w , we obtain that

$$\begin{aligned} & \sum_{\omega \in \Omega} w(\omega) z_{c,\lambda}^{|\omega|} \mathbb{1}[\omega : 0 \rightarrow S_t] \\ & \leq \left(\sum_{\omega \in \Omega} w(\omega) z_{c,\lambda}^{|\omega|} \mathbb{1}[\omega : 0 \rightarrow S_0 \cup S_{t_0}] \right)^2 [A_w(z; t - t_0) + A_w(z; t - 2t_0)]. \end{aligned}$$

When $\lambda \neq 1/2$ the prefactor on the right-hand side is finite and does not depend on t , and we deduce easily that $\beta_w(z_{c,\lambda}) \geq \alpha_w(z_{c,\lambda})$, and hence that $\beta_w(z_{c,\lambda}) = \alpha_w(z_{c,\lambda})$ by (11). It then follows from Lemmas 2.3 and 2.5 that

$$(14) \quad \alpha_w(z_{c,\lambda}) = \beta_w(z_{c,\lambda}) = \max\{\lambda, 1 - \lambda\}$$

for every $\lambda \neq 1/2$. Using left continuity of $\alpha_w(z)$ and right continuity of $\beta_w(z)$ from Lemma 2.4, this implies that $\alpha_w(z)$ is a continuous, strictly decreasing function $(0, z_t) \mapsto [1/2, \infty)$ whose inverse is given by $\lambda \mapsto z_{c,\lambda}$. This implies that the latter function is continuous and strictly increasing on $(-\infty, 1/2]$ as claimed. \square

3. Critical exponents, two-point function decay and ballisticity.

3.1. *Counting walks.* Let G be a connected, locally finite graph, let $\Gamma \subseteq \text{Aut}(G)$ be transitive and nonunimodular, and let $w : \Omega \rightarrow [0, \infty)$ be a good weight function. For each $z \geq 0$, the *bubble diagram* is defined to be the ℓ^2 -norm of the two-point function, that is,

$$B_w(z) = \sum_{x \in V} G_w(x)^2.$$

Convergence of the bubble diagram at z_c is well known to be a signifier of mean-field behaviour for the self-avoiding walk; see [31], Section 1.5.

The following lemma allows us to easily deduce the convergence of the bubble diagram at z_c from Theorems 1.5 and 1.8.

LEMMA 3.1. *Let G be a connected, locally finite graph, let Γ be a transitive nonunimodular subgroup of $\text{Aut}(G)$, and let $w : \Omega \rightarrow [0, \infty)$ be a good weight function. Then*

$$B_w(z) \leq \chi_w(z, \lambda)^2$$

for every $\lambda \in \mathbb{R}$. In particular, if $0 \leq z < z_t(w)$ then $B_w(z) < \infty$.

Note that it is always best to take $\lambda = 1/2$ when applying this bound.

PROOF OF LEMMA 3.1. We can express

$$B_w(z) = \sum_{x \in V} \sum_{\omega:0 \rightarrow x} w(\omega_1)z^{|\omega_1|} \sum_{\omega:x \rightarrow 0} w(\omega_1)z^{|\omega_1|}.$$

Since $\Delta^\lambda(0, 0) = 1$, we have the trivial bound

$$\begin{aligned} B_w(z) &\leq \sum_{x \in V} \sum_{\omega:0 \rightarrow x} w(\omega_1)z^{|\omega_1|} \sum_{y \in V} \sum_{\omega:x \rightarrow y} w(\omega_1)z^{|\omega_1|} \Delta^\lambda(0, y) \\ &= \sum_{x \in V} \sum_{\omega:0 \rightarrow x} w(\omega_1)z^{|\omega_1|} \Delta^\lambda(0, x) \sum_{y \in V} \sum_{\omega:x \rightarrow y} w(\omega_1)z^{|\omega_1|} \Delta^\lambda(x, y) \\ &= \chi_w(z, \lambda)^2, \end{aligned}$$

where the co-cycle identity was used in the second equality and Γ -invariance of w was used in the third. \square

The following differential inequality, which is classical for self-avoiding walk on \mathbb{Z}^d [31], Lemma 1.5.2, allows us to deduce Theorem 1.3 and item (1) of both Theorems 1.6 and 1.9 from Theorems 1.5 and 1.8.

LEMMA 3.2. *Let G be a connected, locally finite graph, let $\Gamma \subseteq \text{Aut}(G)$ be a transitive group of automorphisms and let $w : \Omega \rightarrow [0, \infty)$ be a good weight function. Then for every $\lambda \in \mathbb{R}$ and $z \in [0, z_{c,\lambda}(w))$, we have that*

$$\frac{\chi_w(z, \lambda)^2}{B_w(z)} \leq \frac{\partial}{\partial z} [z\chi_w(z, \lambda)] \leq \chi_w(z, \lambda)^2.$$

The proof is closely adapted from the proof given in [31], Lemma 1.5.2. We simply use the co-cycle identity and the tilted mass-transport principle to “take the modular function along for the ride”.

PROOF OF LEMMA 3.2. For every $0 \leq z < z_{c,\lambda}(w)$, we have that

$$\frac{\partial}{\partial z}[z\chi_w(z, \lambda)] = \sum_{v \in V} \sum_{\omega:0 \rightarrow v} (|\omega| + 1)w(\omega)z^{|\omega|} \Delta^\lambda(0, v).$$

Since $|\omega| + 1$ is the number of ways to split ω into two (possibly length zero) subpaths, we deduce that

$$\frac{\partial}{\partial z}[z\chi_w(z, \lambda)] = \sum_{u, v \in V} \sum_{\omega_1:0 \rightarrow u} \sum_{\omega_2:u \rightarrow v} w(\omega_1 \circ \omega_2)z^{|\omega_1|+|\omega_2|} \Delta^\lambda(0, v).$$

For the upper bound, we use repulsivity and the co-cycle identity to write

$$\begin{aligned} \frac{\partial}{\partial z}[z\chi_w(z, \lambda)] &\leq \sum_{u \in V} \sum_{\omega_1:0 \rightarrow u} w(\omega_1)z^{|\omega_1|} \Delta^\lambda(0, u) \sum_{u \in V} \sum_{\omega_2:u \rightarrow v} w(\omega_2)z^{|\omega_2|} \Delta^\lambda(u, v) \\ &= \chi_w(z, \lambda)^2, \end{aligned}$$

where the equality on the second line follows by transitivity of Γ and Γ -invariance of w .

We now turn to the lower bound. We begin by applying the tilted mass-transport principle to the sum over u to deduce that

$$\begin{aligned} \frac{\partial}{\partial z}[z\chi_w(z, \lambda)] &= \sum_{u, v \in V} \sum_{\omega_1:0 \rightarrow u} \sum_{\omega_2:0 \rightarrow v} w(\omega_1 \circ \omega_2)z^{|\omega_1|+|\omega_2|} \Delta^\lambda(u, v) \Delta(0, u) \\ &= \sum_{u, v \in V} \sum_{\omega_1:u \rightarrow 0} \sum_{\omega_2:0 \rightarrow v} w(\omega_1 \circ \omega_2)z^{|\omega_1|+|\omega_2|} \Delta^{1-\lambda}(0, u) \Delta^\lambda(0, v), \end{aligned}$$

where the co-cycle identity has been used in the second line. Since w is zero-range, we can bound

$$\begin{aligned} w(\omega_1 \circ \omega_2) &\geq w(\omega_1)w(\omega_2)\mathbb{1}(\omega_1, \omega_2 \text{ disjoint}) \\ &= w(\omega_1)w(\omega_2)[1 - \mathbb{1}(\omega_1, \omega_2 \text{ not disjoint})]. \end{aligned}$$

If $\omega_1 : u \rightarrow 0$ and $\omega_2 : 0 \rightarrow v$ are *not* disjoint, then there exists $x \in V$ and paths $\omega_{1,1} : u \rightarrow x$, $\omega_{1,2} : x \rightarrow 0$, $\omega_{2,1} : 0 \rightarrow x$ and $\omega_{2,2} : x \rightarrow v$ such that $\omega_1 = \omega_{1,1} \circ \omega_{1,2}$, $\omega_2 = \omega_{2,1} \circ \omega_{2,2}$, such that $\omega_{1,1}$ and $\omega_{2,2}$ are disjoint, and such that neither $\omega_{1,2}$ or $\omega_{2,1}$ is trivial. Indeed, simply take $\omega_{1,1}$ to be the portion of ω_1 up until the first time it intersects ω_2 , and let $\omega_{2,1}$ be the portion of ω_2 up until the last time it visits $\omega_{1,2}^+$. It follows that

$$\begin{aligned} &\sum_{u, v \in V} \sum_{\omega_1:u \rightarrow 0} \sum_{\omega_2:0 \rightarrow v} \mathbb{1}[\omega_1, \omega_2 \text{ not disjoint}]w(\omega_1)w(\omega_2)z^{|\omega_1|+|\omega_2|} \Delta^{1-\lambda}(0, u) \\ &\leq \sum_{x, u, v \in V} \sum_{\substack{\omega_{1,1}:u \rightarrow x \\ \omega_{1,2}:x \rightarrow 0 \\ |\omega_{1,2}| \geq 1}} \sum_{\substack{\omega_{2,1}:0 \rightarrow x \\ \omega_{2,2}:x \rightarrow v \\ |\omega_{2,1}| \geq 1}} \mathbb{1}[\omega_{1,1}, \omega_{2,1} \text{ disjoint}]w(\omega_{1,1} \circ \omega_{1,1})w(\omega_{2,2} \circ \omega_{2,2}) \\ &\quad \times z^{|\omega_{1,1}|+|\omega_{1,2}|+|\omega_{2,1}|+|\omega_{2,2}|} \Delta^{1-\lambda}(0, u) \Delta^\lambda(0, v). \end{aligned}$$

Using repulsivity of w , the zero-range property, and the co-cycle identity, we can bound

$$\begin{aligned} & \mathbb{1}[\omega_{1,1}, \omega_{2,2} \text{ disjoint}]w(\omega_{1,1} \circ \omega_{1,2})w(\omega_{2,1} \circ \omega_{2,2})\Delta^{1-\lambda}(0, u)\Delta^\lambda(0, v) \\ & \leq w(\omega_{1,1} \circ \omega_{2,2})w(\omega_{1,2})w(\omega_{2,1})\Delta(0, x)\Delta^{1-\lambda}(x, u)\Delta^\lambda(x, v). \end{aligned}$$

This leads to the bound

$$\begin{aligned} & \sum_{u, v \in V} \sum_{\omega_1: u \rightarrow 0} \sum_{\omega_2: 0 \rightarrow v} \mathbb{1}[\omega_1, \omega_2 \text{ not disjoint}]w(\omega_1)w(\omega_2)z^{|\omega_1|+|\omega_2|}\Delta^{1-\lambda}(0, u) \\ & \leq \sum_{x \in V} [G_w(x) - \mathbb{1}(x = 0)]^2 \Delta(0, x) \\ & \quad \times \sum_{u, v \in V} w(\omega_{1,1} \circ \omega_{2,2})z^{|\omega_{1,1}|+|\omega_{2,2}|}\Delta^{1-\lambda}(0, u)\Delta^\lambda(0, v) \\ & = \frac{\partial}{\partial z} [z\chi_w(z, \lambda)] \sum_{x \in V} [G_w(x)^2 - \mathbb{1}(x = 0)]\Delta(0, x) \\ & = (B_w(z) - 1) \frac{\partial}{\partial z} [z\chi_w(z, \lambda)], \end{aligned}$$

where the tilted mass-transport principle is used in the final equality. (The $-\mathbb{1}(x = 0)$ arises from the restriction that neither $\omega_{1,2}$ or $\omega_{2,1}$ is trivial.) On the other hand, similar manipulations to those used in the upper bound, above, yield that

$$\begin{aligned} & \sum_{u, v \in V} \sum_{\omega_1: 0 \rightarrow u} \sum_{\omega_2: 0 \rightarrow v} w(\omega_1)w(\omega_2)z^{|\omega_1|+|\omega_2|}\Delta^{1-\lambda}(0, u)\Delta^\lambda(0, v) \\ & = \chi_w(z; \lambda)\chi_w(z; 1 - \lambda) = \chi_w(z; \lambda)^2. \end{aligned}$$

Combining these inequalities, we deduce that

$$\frac{\partial}{\partial z} [z\chi_w(z, \lambda)] \geq \chi_w(z, \lambda)^2 - (B_w(z) - 1) \frac{\partial}{\partial z} [z\chi_w(z, \lambda)],$$

which rearranges to give the desired inequality. \square

Integrating this differential inequality yields the following estimates; see [31], Theorem 1.5.3.

COROLLARY 3.3. *Let G be a connected, locally finite graph, let $\Gamma \subseteq \text{Aut}(G)$ be a transitive group of automorphisms, and let $w : \Omega \rightarrow [0, \infty)$ be a good weight function. Then*

$$\frac{z_{c,\lambda}}{z_{c,\lambda} - z} \leq \chi_w(z; \lambda) \leq B_w(z_{c,\lambda}) \frac{z_{c,\lambda}}{z_{c,\lambda} - z} + B_w(z_{c,\lambda})$$

for every $\lambda \in \mathbb{R}$ and $0 \leq z < z_{c,\lambda} = z_{c,\lambda}(w)$.

We now have everything we need to deduce the upper bound on the susceptibility for Theorems 1.3, 1.6 and 1.9. To use the susceptibility bounds to deduce the bounds on $Z_w(\lambda; n)$, we use the following lemma. It replaces the Tauberian theory that is typically used in the literature.

LEMMA 3.4. *Let $(c_n)_{n \geq 0}$ be a nonnegative submultiplicative sequence with generating function $\Phi(x) = \sum_{n \geq 0} x^n c_n$. Then*

$$x^n c_n \leq \left[\frac{\Phi(y)}{n+1} \right]^2 \left(\frac{x}{y} \right)^{2n}$$

for every $m \geq 1$ and $x \geq y > 0$.

When we apply this lemma, we will take $y = nx/(n+1)$ so that $(x/y)^{2n} \approx e^2$ is of constant order.

PROOF OF LEMMA 3.4. We clearly have that

$$\sum_{k=0}^n \frac{c_{n-k}x^{n-k} + c_kx^k}{2} = \sum_{k=0}^n c_kx^k \leq \left(\frac{x}{y} \right)^n \Phi(y),$$

and so there must exist $0 \leq k \leq n$ such that

$$\frac{c_{n-k}x^{n-k} + c_kx^k}{2} \leq \left(\frac{x}{y} \right)^n \frac{\Phi(y)}{n+1}.$$

For this k , we have that

$$c_nx^n \leq c_{n-k}x^{n-k}c_kx^k \leq \left[\frac{c_{n-k}x^{n-k} + c_kx^k}{2} \right]^2 \leq \left[\frac{\Phi(y)}{n+1} \right]^2 \left(\frac{x}{y} \right)^{2n}$$

by submultiplicativity and the inequality of arithmetic and geometric means. \square

PROOFS OF THEOREM 1.3 AND PART (I) OF THEOREMS 1.6 AND 1.9. Theorem 1.8 implies that the tilted susceptibility $\chi_w(z_{c,\lambda}, 1/2)$ is finite for every $\lambda \neq 1/2$, and it follows from Lemma 3.1, and Corollary 3.3 that

$$(15) \quad \chi_w(z; \lambda) \leq \chi_w(z_{c,\lambda}; 1/2)^2 \frac{z_{c,\lambda}}{z_{c,\lambda} - z} + \chi_w(z_{c,\lambda}; 1/2)^2.$$

for every $0 \leq z < z_{c,\lambda} = z_{c,\lambda}(w)$. Thus, applying Lemma 3.4 with $c_n = Z_w(\lambda; n)$, $x = z_{c,\lambda}(w)$, and $y = nz_{c,\lambda}(w)/(n+1)$ yields that

$$(16) \quad Z_w(\lambda; n) \leq [e^2 \chi_w(z_{c,\lambda}, 1/2)^4 + o(1)] \mu_{c,\lambda}^n$$

as $n \rightarrow \infty$. \square

REMARK 3.5. In the untilted case $\lambda = 0$, Corollary 2.7 together with (15) and (16) yield that

$$(17) \quad \chi(z) \leq \mu_c^2 \exp\left[\frac{4\mu_c^2}{e^{3t_0/2} - e^{t_0}}\right] \frac{z_c}{z_c - z} + O(1), \quad z \nearrow z_c,$$

$$(18) \quad Z(n) \leq \mu_c^4 \exp\left[\frac{8\mu_c^2}{e^{3t_0/2} - e^{t_0}} + 2\right] \mu_c^n + o(\mu_c^n), \quad n \nearrow \infty.$$

In particular, these estimates hold with $t_0 \geq \log(k - 1)$ for the product of the k -regular tree T_k with an arbitrary transitive graph G . We have not attempted to optimize these constants.

3.2. *Ballisticity and two-point function decay.*

PROOFS OF ITEMS 2 AND 3 OF THEOREMS 1.6 AND 1.9. Observe that the trivial inequality

$$G_w(z; x) \leq \left(\frac{z}{z'}\right)^{d(0,x)} G_w^k(z'; x)$$

holds for every $z \geq 0$ and $x \in V$. On the other hand, for $z < z_t$ we have that

$$G_w(z; x) \leq \chi_w(z; 1/2) \Delta^{-1/2}(0, x)$$

for every $x \in V$, and it follows by symmetry that

$$G_w(z, x) \leq \chi_w(z; 1/2) [\min\{\Delta(0, x), \Delta(x, 0)\}]^{1/2} \leq \chi_w(z; 1/2)$$

for every $0 < z < z_t$ and $x \in V$. Thus, we have that

$$G_w(z, x) \leq \chi_w(z'; 1/2) \left(\frac{z}{z'}\right)^{d(0,x)}$$

for every $x \in V$ and $0 < z < z' < z_t$, which implies the claim of item 2.

We now prove item 3. We prove the claim in the case $\lambda > 1/2$, the case $\lambda < 1/2$ being similar. Let $\lambda > \lambda' > 1/2$. Then we have that

$$\begin{aligned} \mathbb{P}_{w,\lambda,n}(\log \Delta(0, X_n) \leq cn) &= Z_w(\lambda; n)^{-1} \sum_{x \in H_{cn}^-} \sum_{\omega \in \Omega} w(\omega) \Delta^\lambda(0, x) \mathbb{1}[\omega : 0 \rightarrow H_{cn}^-, |\omega| = n] \\ &\leq Z_w(\lambda, n)^{-1} Z_w(\lambda'; n) e^{(\lambda - \lambda')cn}. \end{aligned}$$

We deduce from item 1 of Theorem 1.9 that

$$\mathbb{P}_{w,\lambda,n}(\log \Delta(0, X_n) \leq cn) \leq C_{\lambda'} \left(\frac{z_{c,\lambda}}{z_{c,\lambda'}}\right)^n e^{(\lambda - \lambda')cn}.$$

The result follows by fixing $\lambda > \lambda' > 1/2$ and letting $c = c_{\lambda,\lambda'}$ be sufficiently small that $z_{c,\lambda}/z_{c,\lambda'} < e^{-c(\lambda - \lambda')}$. \square

4. Remarks and open problems. Several interesting questions remain open concerning the behaviour of the *critically tilted SAW*, that is, the case $\lambda = 1/2$.

4.1. *Exponents at $\lambda = 1/2$ are graph dependent.* In this subsection, we briefly outline an example that shows that the exponent governing the critically tilted susceptibility depends on the choice of G and Γ , and in particular that Theorem 1.6 cannot always be generalised to $\lambda = 1/2$. This follows from a related analysis for percolation on trees with respect to two different choices of nonunimodular automorphism group that we performed in [26]. (Note that on trees the tilted susceptibilities for SAW and percolation are equal for $0 \leq z = p \leq 1$.)

Let T be the k -regular tree. The most obvious choice of a nonunimodular transitive subgroup of T is the group Γ_ξ consisting of those automorphisms of T that fix some given end ξ of T . For the pair (T, Γ_ξ) , we can easily compute

$$\alpha(z) = -\log_{k-1}(z) \quad \text{and} \quad z_{c,\lambda} = (k-1)^{-\max\{\lambda, 1-\lambda\}}.$$

Moreover, for $z < z_{c,\lambda}$ we can compute the tilted susceptibility to be

$$\chi(z, \lambda) = \frac{1 - z^2}{(1 - (k-1)^{1-\lambda}z)(1 - (k-1)^\lambda z)}.$$

Thus, we see that for $\lambda \neq 1/2$, $\chi(z_{c,\lambda} - \varepsilon, \lambda)$ grows like ε^{-1} as $\varepsilon \rightarrow 0$, as stated in Theorem 1.6, while at $\lambda = 1/2$ the denominator has a double root and we have instead that

$$\chi_{z_{c,1/2-\varepsilon},1/2} = \frac{k-2}{k-1} \varepsilon^{-2}.$$

This shows that Theorem 1.6 cannot be extended in general to the case $\lambda = 1/2$.

We now describe a different transitive nonunimodular group of automorphisms on the four-regular tree. We define a $(1, 1, 2)$ -orientation of T to be a (partial) orientation of the edge set of T such that every vertex has one oriented edge emanating from it, two oriented edges pointing into it, and one unoriented edge incident to it. Fix one such orientation of T , and let Γ' be the group of automorphisms of T that preserve the orientation. In [26], we compute that

$$z_{c,\lambda} = \frac{2^\lambda + 2^{1-\lambda} + 1 - \sqrt{(2^\lambda + 2^{1-\lambda} + 1)^2 - 12}}{6}, \quad \lambda \in \mathbb{R},$$

and

$$\chi(z, \lambda) = \frac{1 - 3z^2}{1 - (2^\lambda + 2^{1-\lambda} + 1)z + 3z^2}, \quad \lambda \in \mathbb{R}, 0 \leq z < z_{c,\lambda}.$$

The denominator of this expression never has a double root, so that, in contrast to the previous example,

$$\chi(z_{c,\lambda} - \varepsilon, \lambda) \asymp \varepsilon^{-1}, \quad \lambda \in \mathbb{R}, \varepsilon \downarrow 0$$




FIG. 1. Comparison of $\alpha(z)$ and $z_{c,\lambda}$ for the 4-regular tree with respect to the automorphism group fixing an end (blue) and the automorphism group fixing a $(1, 1, 2)$ -orientation (red). The second figure is formed by reflecting the first around the line $\alpha(z) = 1/2$ and then rotating. The intersection of the two curves on the left occurs at $(z_c, 1)$. This intersection must occur since $z_{c,0} = z_{c,1} = z_c$ does not depend on the choice of automorphism group.

for every $\lambda \in \mathbb{R}$. Since z_t for this example is smaller than z_t for the previous example, we have by Lemma 3.1 that the bubble diagram converges at z_t , so that we could also have deduced this behaviour from Corollary 3.3. Furthermore, it follows from our analysis of percolation in [26] that $\alpha(z)$ has a jump discontinuity from $1/2$ to $-\infty$ at z_t . Indeed,

$$\alpha(z) = \begin{cases} \log_2 \left(\frac{3z^2 - z + 1 + \sqrt{9z^4 - 6z^3 - z^2 - 2z + 1}}{2z} \right) & z \leq z_t \\ -\infty & z > z_t. \end{cases}$$

A plot of this function is given in Figure 1. As a further point of contrast to the previous example, we note that in this example the function $\lambda \mapsto z_{c,\lambda}$ is real-analytic, whereas in the previous example it was not differentiable at $\lambda = 1/2$.

4.2. *Hammersley–Welsh-type bounds for critically tilted SAW.* Consider SAW on \mathbb{Z}^d . The Hammersley–Welsh inequality [21] states that

$$Z(n) \leq \exp[O(n^{1/2})] \mu_c^n.$$

See [31], Section 3.1, for background and [27] for a small improvement.

We now briefly outline how an analogous inequality may be obtained for critically tilted ($\lambda = 1/2$) self-avoiding walk in the nonunimodular context. It can be deduced from Corollary 2.7 that

$$\chi(z_t - \varepsilon, 1/2) \leq \exp[O(\varepsilon^{-1})],$$

and applying Lemma 3.4 with $c_m = Z(1/2; n)$, $x = z_t$, and $y = z_t(1 - m^{-1/2})$ yields that

$$Z(1/2; n) \leq \exp[O(n^{1/2})] \mu_{c,1/2}^n,$$

which is an exact analogue of the Hammersley–Welsh bound.

4.3. *Questions.*

QUESTION 4.1. Let T_k be a k -regular tree, let $d \geq 1$ and consider the group of automorphisms $\Gamma_\xi \times \text{Aut}(\mathbb{Z}^d) \subseteq \text{Aut}(T_k \times \mathbb{Z}^d)$ of $T_k \times \mathbb{Z}^d$, where Γ_ξ is the group of automorphisms that fix some specified end ξ of T .

1. Is $B(z_t) < \infty$?
2. What are the asymptotics of $a(z_t; n)$ and $b(z_t; n)$ as defined in Section 2?
3. What is the behaviour of $\chi(z_t - \varepsilon, 1/2)$ as $\varepsilon \rightarrow 0$? What about $Z(n, 1/2)$ as $n \rightarrow \infty$?
4. What is the typical displacement of a SAW sampled from $\mathbb{P}_{1/2, n}$?
5. For which of these questions does the answer depend on d ?

QUESTION 4.2. Let G be a connected locally finite graph and let $\Gamma \subseteq \text{Aut}(G)$ be transitive and nonunimodular. Does there exist $C = C(G, \Gamma) < \infty$ such that

$$Z(1/2; n) = O(n^C \mu_c^n)$$

for every $n \geq 1$? Is there a universal choice of this C ? Does $C = 1$ always suffice?

The question concerning $C = 1$ arises from the guess that the pair (T_k, Γ_ξ) has the largest subexponential correction to $Z(1/2; n)$ among all pairs (G, Γ) .

QUESTION 4.3. Let G be a connected locally finite graph and let $\Gamma \subseteq \text{Aut}(G)$ be transitive and nonunimodular. What asymptotics are possible for the typical displacement of a sample from $\mathbb{P}_{n, 1/2}$? Is it always of order at least $n^{1/2}$?

Acknowledgements. This work was carried out while the author was an intern at Microsoft Research, Redmond. We thank Omer Angel for improving Lemma 3.4 by a factor of 4. We also thank Tyler Helmuth for helpful discussions, and thank Gordon Slade and Hugo Duminil-Copin for comments on an earlier draft and thank the anonymous referee for several helpful suggestions.

REFERENCES

[1] BAUERSCHMIDT, R., BRYDGES, D. C. and SLADE, G. (2015). Logarithmic correction for the susceptibility of the 4-dimensional weakly self-avoiding walk: A renormalisation group analysis. *Comm. Math. Phys.* **337** 817–877. [MR3339164](#)

[2] BAUERSCHMIDT, R., DUMINIL-COPIN, H., GOODMAN, J. and SLADE, G. (2012). Lectures on self-avoiding walks. In *Probability and Statistical Physics in Two and More Dimensions*. *Clay Math. Proc.* **15** 395–467. Amer. Math. Soc., Providence, RI. [MR3025395](#)

[3] BENJAMINI, I. (2016). Self avoiding walk on the seven regular triangulation. Preprint. Available at [arXiv:1612.04169](#).

[4] BENJAMINI, I. and CURIEN, N. (2012). Ergodic theory on stationary random graphs. *Electron. J. Probab.* **17** no. 93. [MR2994841](#)

[5] DIESTEL, R. and LEADER, I. (2001). A conjecture concerning a limit of non-Cayley graphs. *J. Algebraic Combin.* **14** 17–25. [MR1856226](#)

- [6] DOMB, C. and JOYCE, G. (1972). Cluster expansion for a polymer chain. *J. Phys. C, Solid State Phys.* **5** 956.
- [7] DUMINIL-COPIN, H., GLAZMAN, A., HAMMOND, A. and MANOLESCU, I. (2016). On the probability that self-avoiding walk ends at a given point. *Ann. Probab.* **44** 955–983. [MR3474464](#)
- [8] DUMINIL-COPIN, H. and HAMMOND, A. (2013). Self-avoiding walk is sub-ballistic. *Comm. Math. Phys.* **324** 401–423. [MR3117515](#)
- [9] DUMINIL-COPIN, H. and SMIRNOV, S. (2012). The connective constant of the honeycomb lattice equals $\sqrt{2 + \sqrt{2}}$. *Ann. of Math. (2)* **175** 1653–1665. [MR2912714](#)
- [10] GILCH, L. A. and MÜLLER, S. (2017). Counting self-avoiding walks on free products of graphs. *Discrete Math.* **340** 325–332. [MR3584819](#)
- [11] GRIMMETT, G. and LI, Z. (2013). Self-avoiding walks and the Fisher transformation. *Electron. J. Combin.* **20** Paper 47. [MR3118955](#)
- [12] GRIMMETT, G. R., HOLROYD, A. E. and PERES, Y. (2014). Extendable self-avoiding walks. *Ann. Inst. Henri Poincaré D* **1** 61–75. [MR3166203](#)
- [13] GRIMMETT, G. R. and LI, Z. (2014). Strict inequalities for connective constants of transitive graphs. *SIAM J. Discrete Math.* **28** 1306–1333. [MR3252803](#)
- [14] GRIMMETT, G. R. and LI, Z. (2015). Bounds on connective constants of regular graphs. *Combinatorica* **35** 279–294. [MR3367126](#)
- [15] GRIMMETT, G. R. and LI, Z. (2016). Cubic graphs and the golden mean. Available at [arXiv:1610.00107](#).
- [16] GRIMMETT, G. R. and LI, Z. (2017). Connective constants and height functions for Cayley graphs. *Trans. Amer. Math. Soc.* **369** 5961–5980. [MR3646785](#)
- [17] GRIMMETT, G. R. and LI, Z. (2017). Self-avoiding walks and amenability. *Electron. J. Combin.* **24** Paper 4.38. [MR3743101](#)
- [18] GRIMMETT, G. R. and LI, Z. (2017). Self-avoiding walks and connective constants Available at [arXiv:1704.05884](#).
- [19] GRIMMETT, G. R. and LI, Z. (2018). Locality of connective constants. *Discrete Math.* **341** 3483–3497. [MR3862645](#)
- [20] HAMMERSLEY, J. M. and MORTON, K. W. (1954). Poor man’s Monte Carlo. *J. Roy. Statist. Soc. Ser. B* **16** 23–38; discussion 61–75. [MR0064475](#)
- [21] HAMMERSLEY, J. M. and WELSH, D. J. A. (1962). Further results on the rate of convergence to the connective constant of the hypercubical lattice. *Q. J. Math.* **13** 108–110. [MR0139535](#)
- [22] HARA, T. (2008). Decay of correlations in nearest-neighbor self-avoiding walk, percolation, lattice trees and animals. *Ann. Probab.* **36** 530–593. [MR2393990](#)
- [23] HARA, T. and SLADE, G. (1992). The lace expansion for self-avoiding walk in five or more dimensions. *Rev. Math. Phys.* **4** 235–327. [MR1174248](#)
- [24] HARA, T. and SLADE, G. (1992). Self-avoiding walk in five or more dimensions. I. The critical behaviour. *Comm. Math. Phys.* **147** 101–136. [MR1171762](#)
- [25] HUTCHCROFT, T. (2016). Critical percolation on any quasi-transitive graph of exponential growth has no infinite clusters. *C. R. Math. Acad. Sci. Paris* **354** 944–947. [MR3535351](#)
- [26] HUTCHCROFT, T. (2017). Non-uniqueness and mean-field criticality for percolation on nonunimodular transitive graphs. Available at [arXiv:1711.02590](#).
- [27] HUTCHCROFT, T. (2018). The Hammersley-Welsh bound for self-avoiding walk revisited. *Electron. Commun. Probab.* **23** Paper No. 5. [MR3771763](#)
- [28] KESTEN, H. (1964). On the number of self-avoiding walks. II. *J. Math. Phys.* **5** 1128–1137. [MR0166845](#)
- [29] LI, Z. (2016). Positive speed self-avoiding walks on graphs with more than one end. Preprint. Available at [arXiv:1612.02464](#).

- [30] LYONS, R. and PERES, Y. (2016). *Probability on Trees and Networks. Cambridge Series in Statistical and Probabilistic Mathematics* **42**. Cambridge Univ. Press, New York. [MR3616205](#)
- [31] MADRAS, N. and SLADE, G. (2013). *The Self-Avoiding Walk. Modern Birkhäuser Classics*. Birkhäuser/Springer, New York. Reprint of the 1993 original. [MR2986656](#)
- [32] MADRAS, N. and WU, C. C. (2005). Self-avoiding walks on hyperbolic graphs. *Combin. Probab. Comput.* **14** 523–548. [MR2160417](#)
- [33] NACHMIAS, A. and PERES, Y. (2012). Non-amenable Cayley graphs of high girth have $p_c < p_u$ and mean-field exponents. *Electron. Commun. Probab.* **17** Paper No. 57. [MR3005730](#)
- [34] NIENHUIS, B. (1982). Exact critical point and critical exponents of $O(n)$ models in two dimensions. *Phys. Rev. Lett.* **49** 1062–1065. [MR0675241](#)
- [35] PAK, I. and SMIRNOVA-NAGNIBEDA, T. (2000). On non-uniqueness of percolation on non-amenable Cayley graphs. *C. R. Acad. Sci., Sér. 1 Math.* **330** 495–500. [MR1756965](#)
- [36] SOARDI, P. M. and WOESS, W. (1990). Amenability, unimodularity, and the spectral radius of random walks on infinite graphs. *Math. Z.* **205** 471–486. [MR1082868](#)
- [37] TIMÁR, Á. (2006). Percolation on nonunimodular transitive graphs. *Ann. Probab.* **34** 2344–2364. [MR2294985](#)
- [38] YAMAMOTO, K. (2017). An upper bound for the critical probability on the cartesian product graph of a regular tree and a line. Preprint. Available at [arXiv:1705.06873](#).

STATSLAB, DPMMS
UNIVERSITY OF CAMBRIDGE
CAMBRIDGE, CB3 0WA
UNITED KINGDOM
E-MAIL: t.hutchcroft@maths.cam.ac.uk