



Nigro, Steven J; Wick, Ryan; Holt, Kathryn E; Hall, Ruth M (2018) Complete Genome Sequence of WM99c, an Antibiotic-Resistant *Acinetobacter baumannii* Global Clone 2 (GC2) Strain Representing an Australian GC2 Lineage. *MICROBIOLOGY RESOURCE ANNOUNCEMENTS*, 7 (22). ISSN 2576-098X DOI: <https://doi.org/10.1128/MRA.01199-18>


Downloaded from: <http://researchonline.lshtm.ac.uk/4651020/>

DOI: [10.1128/MRA.01199-18](https://doi.org/10.1128/MRA.01199-18)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by/2.5/>



Complete Genome Sequence of WM99c, an Antibiotic-Resistant *Acinetobacter baumannii* Global Clone 2 (GC2) Strain Representing an Australian GC2 Lineage

Steven J. Nigro,^a Ryan Wick,^{b,c} Kathryn E. Holt,^{b,c} Ruth M. Hall^a

^aSchool of Life and Environmental Sciences, University of Sydney, Sydney, Australia

^bCentre for Systems Genomics, University of Melbourne, Melbourne, Australia

^cDepartment of Biochemistry and Molecular Biology, Bio21 Molecular Science and Biotechnology Institute, University of Melbourne, Melbourne, Australia

ABSTRACT The extensively antibiotic-resistant *Acinetobacter baumannii* isolate WM99c recovered in Sydney, Australia, in 1999 is an early representative of a distinct lineage of global clone 2 (GC2) seen on the east coast of Australia. We present the complete 4.121-Mbp genome sequence (chromosome plus 2 plasmids), generated via long-read sequencing (PacBio).

The *Acinetobacter baumannii* isolate WM99c was collected in 1999 at Westmead Hospital in Sydney, Australia (1), and is one of the earliest global clone 2 (GC2) isolates recorded in Australia (2, 3). It is resistant to several current antibiotics, including imipenem, meropenem, ceftazidime, cefotaxime, fluoroquinolones, ciprofloxacin, and levofloxacin, as well as older antibiotics, including sulfonamides, tetracycline, minocycline, and the aminoglycosides gentamicin, kanamycin, and neomycin (4). It is also resistant to tigecycline (5). It was previously shown to be representative of a distinct GC2 lineage that has the *oxa23* carbapenem resistance gene in transposon Tn2006 and an ISAb17 in the AbGRI1-2 variant (formerly Tn6167 [6]) of the AbGRI1 resistance island, which is located in the *comM* gene (4). This variant has been seen only in isolates from hospitals on the east coast of Australia (3). A characteristic variant of the AbGRI2 resistance island, AbGRI2-1, is also present in the chromosome (4). Third-generation cephalosporin resistance is due to increased transcription of the *ampC* gene from an upstream ISAb1, and fluoroquinolone resistance arose via mutations in *gyrA* and *parC*.

Whole-cell DNA was prepared (7) from cells grown in LB, quality controlled for length using agarose gel electrophoresis, and quantified with the Qubit (Invitrogen) system. The DNA was subjected to library preparation and sequencing on 2 PacBio single-molecule real-time (SMRT) cells (chemistry version C2-P4) at DNA Link (South Korea). A total of 164,293 reads were obtained with an average length of 12,816 bp and average quality of 0.823. The PacBio reads (2.1 Gbp; SRA accession number [SRR8162697](https://www.ncbi.nlm.nih.gov/SRA/acc.cgi?acc=SRR8162697)) were retrieved from the raw data using PacBio's SMRT Analysis software and quality control (QC) filters of a PacBio quality score of 85 or greater, and a length of 13 kbp or greater was used to take the read set to a manageable size of 300 Mbp. These reads were combined with available Illumina HiSeq data (SRA accession number [ERR110084](https://www.ncbi.nlm.nih.gov/SRA/acc.cgi?acc=ERR110084)) with Unicycler version 0.4.4 (8) with default parameters, and the resulting assembly was polished with both read sets with unicycler-polish.

The assembled contiguous circular sequence of the chromosome was 4,016,817 bp and had a 39.12% GC content. Two plasmid contigs represented pWM99c-1 (11 kb), which is identical to pD72-1 (GenBank accession number [KM051986](https://www.ncbi.nlm.nih.gov/GenBank/acc.cgi?acc=KM051986)), and pWM99c-2 (110 kb), which is identical to pC20-3 (9) and >99.9% identical (6 single-base-pair differences) to pABTJ2 (GenBank accession number [CP004359](https://www.ncbi.nlm.nih.gov/GenBank/acc.cgi?acc=CP004359)). A 13.3-kb integrative

Received 30 September 2018 Accepted 10 November 2018 Published 6 December 2018

Citation Nigro SJ, Wick R, Holt KE, Hall RM. 2018. Complete genome sequence of WM99c, an antibiotic-resistant *Acinetobacter baumannii* global clone 2 (GC2) strain representing an Australian GC2 lineage. *Microbiol Resour Announc* 7:e01199-18. <https://doi.org/10.1128/MRA.01199-18>.

Editor Irene L. G. Newton, Indiana University, Bloomington

Copyright © 2018 Nigro et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ruth M. Hall, ruth.hall@sydney.edu.au.

element encoding a site-specific tyrosine recombinase was detected in the excised free circular form as well as in the chromosome (bp 992828 to 1006173). Protein coding, rRNA, and tRNA genes were annotated with Prokka version 1.12 (10), and the antibiotic resistance regions, transposons, insertion sequences, and plasmids were annotated manually. Polysaccharide biosynthesis loci for capsule (KL) and outer core of lipooligosaccharide (OCL) were annotated according to published nomenclature (11). Four potential integrated phage genomes of 52.6 kb, 49.5 kb, 37 bp, and 36.1 kb were identified with PHASTER (12).

WM99c belongs to sequence type 208 (ST208) (Oxford scheme; <https://pubmlst.org/abaumannii/>) and carries the KL2 capsule cluster and the OCL1 lipooligosaccharide outer core cluster (11). In addition to the IS26 copies in AbGRI2-1, the chromosome includes 20 copies of the insertion sequence ISAbA1, including 4 associated with expression of antibiotic resistance genes, namely, the 2 that are part of Tn2006 and the single copies upstream of the *ampC* and *sul2* resistance genes. Two copies each of ISAbA17 and ISAbA22 and 1 of ISAbA26 were also detected using ISfinder (<https://isfinder.biotoul.fr/>).

The genome sequence of WM99c will underpin studies of the origin and evolution of the unique GC2 lineage found in hospitals on the east coast of Australia.

Data availability. The complete genome sequence of the *Acinetobacter baumannii* isolate WM99c has been deposited in DDBJ/ENA/GenBank under the accession numbers CP031743 (chromosome), CP031745 (pWM99c-1), and CP031744 (pWM99c-2). The versions described in this paper are the first versions, CP031743.1 to CP031745.1. The PacBio reads have been deposited in the SRA under accession number SRR8162697. Illumina HiSeq data are available under SRA accession number ERR110084.

ACKNOWLEDGMENTS

This work was supported by the NHMRC of Australia (project grant number 1026189 to R.M.H. and fellowship number 1061409 to K.E.H.).

We thank Jon Iredell for supplying isolate WM99c.

REFERENCES

- Valenzuela JK, Thomas L, Partridge SR, van der Reijden T, Dijkshoorn L, Iredell J. 2007. Horizontal gene transfer in a polyclonal outbreak of carbapenem-resistant *Acinetobacter baumannii*. *J Clin Microbiol* 45: 453–460. <https://doi.org/10.1128/JCM.01971-06>.
- Hamidian M, Wynn M, Holt KE, Pickard D, Dougan G, Hall RM. 2014. Identification of a marker for two lineages within the GC1 clone of *Acinetobacter baumannii*. *J Antimicrob Chemother* 69:557–558. <https://doi.org/10.1093/jac/dkt379>.
- Nigro SJ, Hall RM. 2016. Loss and gain of aminoglycoside resistance in global clone 2 *Acinetobacter baumannii* in Australia via modification of genomic resistance islands and acquisition of plasmids. *J Antimicrob Chemother* 71:2432–2440. <https://doi.org/10.1093/jac/dkw176>.
- Nigro SJ, Farrugia DN, Paulsen IT, Hall RM. 2013. A novel family of genomic resistance islands, AbGRI2, contributing to aminoglycoside resistance in *Acinetobacter baumannii* isolates belonging to global clone 2. *J Antimicrob Chemother* 68:554–557. <https://doi.org/10.1093/jac/dks459>.
- Iredell J, Thomas L, Power D, Mendes E. 2007. Tigecycline resistance in Australian antibiotic-resistant Gram-negative bacteria. *J Antimicrob Chemother* 59:816–818. <https://doi.org/10.1093/jac/dkm002>.
- Nigro SJ, Hall RM. 2012. Tn6167, an antibiotic resistance island in an Australian carbapenem resistant *Acinetobacter baumannii* GC2, ST92 isolate. *J Antimicrob Chemother* 67:1342–1346. <https://doi.org/10.1093/jac/dks037>.
- Wilson K. 2001. Preparation of genomic DNA from bacteria. *Curr Protoc Mol Biol* 56:2.4.1–2.4.5. <https://doi.org/10.1002/0471142727.mb0204s56>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Nigro SJ, Hall RM. 2014. Amikacin resistance plasmids in extensively antibiotic-resistant GC2 *Acinetobacter baumannii* from two Australian hospitals. *J Antimicrob Chemother* 69:3435–3437. <https://doi.org/10.1093/jac/dku310>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- Kenyon JJ, Hall RM. 2013. Variation in the complex carbohydrate biosynthesis loci of *Acinetobacter baumannii* genomes. *PLoS One* 8:e62160. <https://doi.org/10.1371/journal.pone.0062160>.
- Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.